

Na Huang and [Piotr Fryzlewicz](#)

NOVELIST estimator of large correlation and covariance matrices and their inverses

**Article (Accepted version)
(Refereed)**

Original citation:

Huang, Na and Fryzlewicz, Piotr (2018) *NOVELIST estimator of large correlation and covariance matrices and their inverses*. [TEST](#). ISSN 1133-0686 (In Press)

DOI: [10.1007/s11749-018-0592-4](https://doi.org/10.1007/s11749-018-0592-4)

© 2018 Sociedad de Estadística e Investigación Operativa

This version available at: <http://eprints.lse.ac.uk/89055/>

Available in LSE Research Online: July 2018

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

NOVELIST estimator of large correlation and covariance matrices and their inverses

Na Huang · Piotr Fryzlewicz

Received: date / Accepted: date

Abstract We propose a “NOVEL Integration of the Sample and Thresholded covariance estimators” (NOVELIST) to estimate the large covariance (correlation) and precision matrix. NOVELIST performs shrinkage of the sample covariance (correlation) towards its thresholded version. The sample covariance (correlation) component is non-sparse and can be low-rank in high dimensions. The thresholded sample covariance (correlation) component is sparse, and its addition ensures the stable invertibility of NOVELIST. The benefits of the NOVELIST estimator include simplicity, ease of implementation, computational efficiency and the fact that its application avoids eigenanalysis. We obtain an explicit convergence rate in the operator norm over a large class of covariance (correlation) matrices when the dimension p and the sample size n satisfy $\log p/n \rightarrow 0$, and its improved version when $p/n \rightarrow 0$. In empirical comparisons with several popular estimators, the NOVELIST estimator performs well in estimating covariance and precision matrices over a wide range of models and sparsity classes. Real data applications are presented.

Keywords covariance regularisation · high-dimensional covariance · long memory · non-sparse modelling · singular sample covariance · high dimensionality.

1 Introduction

Estimating the covariance matrix and its inverse, also known as the concentration or precision matrix, has always been an important part of multivariate analysis, and arises prominently, for example, in financial risk management (Markowitz, 1952; Longerstaeey et al., 1996), linear discriminant analysis (Fisher, 1936; Guo et al., 2007), principal component analysis (Pearson, 1901; Croux & Haesbroeck, 2000) and network science (Jeong et al., 2001; Gardner et al., 2003). Naturally, this is also true of the correlation matrix, and the following discussion applies to it, too. The sample covariance matrix is a straightforward and often used estimator of the covariance matrix. However, when the dimension p of the data is larger than the sample size n , the sample covariance matrix is singular. Even if p is smaller than but of the same order of magnitude as n , the number of parameters to estimate is $p(p+1)/2$, which can significantly exceed n . In this case, the sample covariance matrix is not reliable, and alternative estimation methods are needed.

Piotr Fryzlewicz
Department of Statistics, London School of Economics
Tel.: +123-45-678910
Fax: +123-45-678910
E-mail: p.fryzlewicz@lse.ac.uk

Na Huang
Department of Statistics, London School of Economics

We would categorise the most commonly used alternative covariance estimators into two broad classes. Estimators in the first class rely on various structural assumptions on the underlying true covariance. One prominent example is ordered covariance matrices, often appearing in time series analysis, spatial statistics and spatio-temporal modelling; these assume that there is a metric on the variable indices. Bickel & Levina (2008a) use banding to achieve consistent estimation in this context. Furrer & Bengtsson (2007) and Cai et al. (2010) regularise estimated ordered covariance matrices by tapering. Cai et al. (2010) derive the optimal estimation rates for the covariance matrix under the operator and Frobenius norms, a result which implies sub-optimality of the convergence rate of the banding estimator of Bickel & Levina (2008a) in the operator norm. The estimator of Cai et al. (2010) only achieves the optimal rate if the bandwidth parameter is chosen optimally; however, the optimal bandwidth depends crucially on the underlying unknown covariance matrix and therefore this estimator's optimality is only oracular. The banding technique is also applied to the estimated Cholesky factorisation of the covariance matrix (Bickel & Levina, 2008a; Wu & Pourahmadi, 2003).

Another important example of a structural assumption on the true covariance or precision matrices is sparsity; it is often made e.g. in the statistical analysis of genetic regulatory networks (Gardner et al., 2003; Jeong et al., 2001). El Karoui (2008) and Bickel & Levina (2008b) regularise the estimated sparse covariance matrix by universal thresholding. Adaptive thresholding, in which the threshold is a random function of the data (Cai & Liu, 2011; Fryzlewicz, 2013), leads to more natural thresholding rules and hence, potentially, more precise estimation. The Lasso penalty is another popular way to regularise the covariance and precision matrices (Zou, 2006; Rothman et al., 2008; Friedman et al., 2008). Focusing on model selection rather than parameter estimation, Meinshausen & Bühlmann (2006) propose the neighbourhood selection method. One other commonly occurring structural assumption in covariance estimation is the factor model, often used e.g. in financial applications. Fan, Fan & Lv (2008) impose sparsity on the covariance matrix via a factor model. Fan et al. (2013) propose the POET estimator, which assumes that the covariance matrix is the sum of a part derived from a factor model, and a sparse part.

Estimators in the second broad class do not assume a specific structure of the covariance or precision matrices, but shrink the sample eigenvalues of the sample covariance matrix towards an assumed shrinkage target (Ledoit & Wolf, 2012). A considerable number of shrinkage estimators have been proposed along these lines. Ledoit & Wolf (2004) derive an optimal linear shrinkage formula, which imposes the same shrinkage intensity on all sample eigenvalues but leave the sample eigenvectors unchanged. Non-linear shrinkage is considered in Ledoit & Péché (2011) and Ledoit & Wolf (2012, 2015). Lam (2016) introduces a Nonparametric Eigenvalue-Regularized Covariance Matrix Estimator (NERCOME) through subsampling of the data, which is asymptotically equivalent to the nonlinear shrinkage method of Ledoit & Wolf (2012). Shrinkage can also be applied on the sample covariance matrix directly. Ledoit & Wolf (2003) propose a weighted average estimator of the covariance matrix with a single-index factor target. Schäfer & Strimmer (2005) review six different shrinkage targets. Naturally related to the shrinkage approach is Bayesian estimation of the covariance and precision matrices. Evans (1965), Chen (1979), and Dickey et al. (1985) use possibly the most natural covariance matrix prior, the inverted Wishart distribution. Other notable references include Leonard & John (2012) and Alvarez et al. (2014).

The POET method of Fan et al. (2013) proposes to estimate the covariance matrix as the sum of a non-sparse, low-rank matrix coming from the factor model part, and a certain sparse matrix, added on to ensure invertibility of the resulting covariance estimator. In this paper, we are motivated by the general idea of building a covariance estimator as the sum of a non-sparse and a sparse part. By following this route, the resulting estimator can be hoped to perform well in estimating both non-sparse and sparse covariance matrices if the amount of sparsity is chosen well. At the same time, the addition of the sparse part can guarantee stable invertibility of the estimated covariance, a pre-requisite for the successful estimation of the precision matrix. On the other hand, we wish to move away from the heavy modelling assumptions used by the POET estimator: indeed, our empirical results presented later suggest that POET can underperform if the factor model assumption does not hold.

Motivated by this observation, this paper proposes a simple, practically assumption-free estimator of the covariance and correlation matrices, termed NOVELIST (NOVEL Integration of the Sample and Thresholded covariance/correlation estimators). NOVELIST arises as the linear combination of two parts: the sample covariance (correlation) estimator, which is always non-sparse and has low rank if $p > n$, and its thresholded version, which is sparse. The inclusion of the sparse thresholded part means that NOVELIST can always be made stably invertible. NOVELIST can be viewed as a shrinkage estimator where the sample covariance (correlation) matrix is shrunk towards a flexible, non-parametric, sparse target. By selecting the appropriate amount of contribution of either of the two components, NOVELIST can adapt to a wide range of underlying covariance structures, including sparse but also non-sparse ones. In the paper, we show consistency of the NOVELIST estimator in the operator norm uniformly under a class of covariance matrices introduced by Bickel & Levina (2008b), as long as $\log p/n \rightarrow 0$, and offer an improved version of this result if $p/n \rightarrow 0$. The benefits of the NOVELIST estimator include simplicity, ease of implementation, computational efficiency and the fact that its application avoids eigenanalysis, which is unfamiliar to some practitioners. In our simulation studies, NOVELIST performs well in estimating both covariance and precision matrices for a wide range of underlying covariance structures, benefitting from the flexibility in the selection of its shrinkage intensity and thresholding level.

The rest of the paper is organised as follows. In Section 2 we introduce the NOVELIST estimator and its properties. Section 3 discusses the case where the two components of the NOVELIST estimator are combined in a non-convex way. Section 4 describes the procedure for selecting its parameters. Section 5 shows empirical improvements of NOVELIST. Section 6 exhibits practical performance of NOVELIST in comparison with the state of the art. Section 7 presents real-data performance in portfolio optimisation problems and concludes the paper, and proofs appear in the appendix. The R package ‘‘novelist’’ is available on CRAN.

2 Method, motivation and properties

2.1 Notation and Method

We observe n i.i.d. p -dimensional observations $\mathbf{X}_1, \dots, \mathbf{X}_n$, distributed according to a distribution F , with $E(\mathbf{X}) = 0$, $\Sigma = \{\sigma_{ij}\} = E(\mathbf{X}\mathbf{X}^T)$, and $R = \{\rho_{ij}\} = D^{-1}\Sigma D^{-1}$, where $D = (\text{diag}(\Sigma))^{1/2}$. In the case of heteroscedastic data, we apply NOVELIST to the sample correlation matrix and only then obtain the corresponding covariance estimator. The NOVELIST estimator of the correlation matrix is defined as

$$\hat{R}^N(\hat{R}, \lambda, \delta) = \underbrace{(1 - \delta) \hat{R}}_{\text{non-sparse part}} + \underbrace{\delta T(\hat{R}, \lambda)}_{\text{sparse part}}, \quad (1)$$

and the corresponding covariance estimator is defined as $\hat{\Sigma}^N = \hat{D} \hat{R}^N \hat{D}$, where $\hat{\Sigma} = \{\hat{\sigma}_{ij}\}$ and $\hat{R} = \{\hat{\rho}_{ij}\}$ are the sample covariance and correlation matrices respectively, $\hat{D} = (\text{diag}(\hat{\Sigma}))^{1/2}$, δ is the weight or shrinkage intensity, which is usually within the range $[0, 1]$ but can also lie outside it, λ is the thresholding value, which is a scalar parameter in $[0, 1]$, and $T(\cdot, \cdot)$ is a function that applies any generalised thresholding operator (Rothman et al., 2009) to each off-diagonal entry of its first argument, with the threshold value equal to its second argument. The generalised thresholding operator refers to any function satisfying the following conditions for all $z \in \mathbb{R}$, (i) $|T(z, \lambda)| \leq |z|$; (ii) $T(z, \lambda) = 0$ for $|z| \leq \lambda$; (iii) $|T(z, \lambda) - z| \leq \lambda$. Typical examples of T include soft thresholding T_s with $T(z, \lambda) = (z - \text{Sign}(z)\lambda) \mathbb{1}(|z| > \lambda)$, hard thresholding T_h with $T(z, \lambda) = z \mathbb{1}(|z| > \lambda)$, and SCAD (Fan and Li, 2001). Note that $\hat{\Sigma}^N$ can also be written directly as a NOVELIST estimator with a $p \times p$ adaptive threshold matrix A , $\hat{\Sigma}^N = (1 - \delta) \hat{\Sigma} + \delta T(\hat{\Sigma}, A)$, where $A = \{\lambda \hat{\sigma}_{ii} \hat{\sigma}_{jj}\}$.




Fig. 1 Left: Illustration of NOVELIST operators for any off-diagonal entry of the correlation matrix $\hat{\rho}_{ij}$ with soft thresholding target T_s ($\lambda = 0.5$, $\delta = 0.1, 0.5$ and 0.9). Right: ranked eigenvalues of NOVELIST plotted versus ranked eigenvalues of the sample correlation matrix.

NOVELIST is a shrinkage estimator, in which the shrinkage target is assumed to be sparse. The degree of shrinkage is controlled by the δ parameter, and the amount of sparsity in the target by the λ parameter. Numerical results shown in Figure 1 suggest that the eigenvalues of the NOVELIST estimator arise as a certain non-linear transformation of the eigenvalues of the sample correlation (covariance) matrix, although the application of NOVELIST avoids explicit eigenanalysis.

2.2 Motivation: link to ridge regression

In this section, we show how the NOVELIST estimator can arise in a penalised solution to the linear regression problem, which is linked to ridge regression. For linear regression $Y = \tilde{X}\beta + \varepsilon$, the traditional OLS solution $(\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T Y$ cannot be used if $p > n$ because of the non-invertibility of $\tilde{X}^T \tilde{X}$. The OLS solution rewrites as $[(1 - \delta)\tilde{X}^T \tilde{X} + \delta\tilde{X}^T \tilde{X}]^{-1} \tilde{X}^T Y$, where $\delta \in [0, 1]$. Using this as a starting point, we consider a regularised solution

$$[(1 - \delta)\tilde{X}^T \tilde{X} + \delta f(\tilde{X}^T \tilde{X})]^{-1} \tilde{X}^T Y \doteq A^{-1} \tilde{X}^T Y \quad (2)$$

where $f(\tilde{X}^T \tilde{X})$ is any elementwise modification of the matrix $\tilde{X}^T \tilde{X}$ designed (a) to make A invertible and (b) to ensure adequate estimation of β . The expression in (2) is the minimiser of a generalised ridge regression criterion

$$(1 - \delta) \|Y - \tilde{X}\beta\|_2^2 + \delta \beta^T f(\tilde{X}^T \tilde{X})\beta, \quad (3)$$

where δ acts as a tuning parameter. If $f(\tilde{X}^T \tilde{X}) = I$, (3) is reduced to ridge regression and A is the shrinkage estimator with the identity matrix target. If $f(\tilde{X}^T \tilde{X}) = T(\tilde{X}^T \tilde{X}, \lambda \hat{\sigma}_{ii} \hat{\sigma}_{jj})$, A is the NOVELIST estimator of the covariance matrix.

From formula (3), NOVELIST penalises the regression coefficients in a pairwise manner which can be interpreted as follows: for a given threshold λ , we place a penalty on the products $\beta_i \beta_j$ of those coefficients of β for which the sample correlation between \tilde{X}_i and \tilde{X}_j , the i th and j th column of \tilde{X} (respectively), exceeds λ . In other words, if the sample correlation is high, we penalise the product of the corresponding β 's, hoping that the resulting estimated β_i and β_j are not simultaneously large.

2.3 Asymptotic properties of NOVELIST

2.3.1 Consistency of the NOVELIST estimators. In this section, we establish consistency of NOVELIST in the operator norm and derive the rates of convergence under different scenarios. Bickel & Levina (2008b) introduce a uniformity class of covariance matrices invariant under permutations as

$$\mathcal{U}(q, c_0(p), M, \epsilon_0) = \left\{ \Sigma : \sigma_{ii} \leq M, \sum_{j=1}^p |\sigma_{ij}|^q \leq c_0(p), \text{ for all } i \text{ and } \lambda_{\min}(\Sigma) \geq \epsilon_0 > 0 \right\}, \quad (4)$$

where $0 \leq q < 1$, c_0 is a function of p , the parameters M and ϵ_0 are constants, and $\lambda_{\min}(\cdot)$ is the smallest eigenvalue operator. Analogously, we define a uniformity class of correlation matrices as

$$\mathcal{V}(q, s_0(p), \epsilon_0) = \left\{ R : \sum_{j=1}^p |\rho_{ij}|^q \leq s_0(p), \text{ for all } i \text{ and } \lambda_{\min}(R) \geq \epsilon_0 > 0 \right\}, \quad (5)$$

where $0 \leq q < 1$ and ϵ_0 is a constant. The parameters q and $s_0(p)$ (equiv. $c_0(p)$) together control the permitted degree of ‘‘sparsity’’ of the members of the given class. In the remainder of the paper, where it does not cause confusion, we mostly work with $s_0(p)$ rather than $c_0(p)$, noting that these two parameterisations are equivalent.

Next, we establish consistency of the NOVELIST estimator in the operator norm, $\|A\|_2^2 = \lambda_{\max}(AA^T)$, where $\lambda_{\max}(\cdot)$ is the largest eigenvalue operator.

Proposition 1 *Let F satisfy $\int_0^\infty \exp(\gamma t) dG_j(t) < \infty$ for $0 < |\gamma| < \gamma_0$, where $\gamma_0 > 0$ and G_j is the cdf of X_{1j}^2 . Let $R = \{\rho_{ij}\}$ and $\Sigma = \{\sigma_{ij}\}$ be the true correlation and covariance matrices with $1 \leq i, j \leq p$, and $\sigma_{ii} \leq M$, where $M > 0$. Then, uniformly on $\mathcal{V}(q, s_0(p), \epsilon_0)$, for sufficiently large M' , if $\lambda = M' \sqrt{\log p/n}$ and $\log p/n = o(1)$,*

$$\|\hat{R}^N - R\| = \underbrace{O_p((1-\delta)p\sqrt{\log p/n})}_{(A)} + \underbrace{O_p(\delta s_0(p)(\log p/n)^{(1-q)/2})}_{(B)} = \|(\hat{R}^N)^{-1} - R^{-1}\| \quad (6)$$

$$\|\hat{\Sigma}^N - \Sigma\| = O_p((1-\delta)p\sqrt{\log p/n}) + O_p(\delta s_0(p)(\log p/n)^{(1-q)/2}) = \|(\hat{\Sigma}^N)^{-1} - \Sigma^{-1}\|. \quad (7)$$

Proposition 2 *Let the length- p column vector \mathbf{X}_i satisfy the sub-Gaussian condition $P(|v^T \mathbf{X}_i| > t) \leq \exp(-t^2 \rho/2)$ for a certain $\rho > 0$, for all $t > 0$ and $\|v\|_2 = 1$. Let $R = \{\rho_{ij}\}$ and $\Sigma = \{\sigma_{ij}\}$ be the true correlation and covariance matrices with $1 \leq i, j \leq p$, and $\sigma_{ii} \leq M$, where $M > 0$. Then, uniformly on $\mathcal{V}(q, s_0(p), \epsilon_0)$, for sufficiently large M' , if $\lambda = M' \sqrt{\log p/n}$ and $p = o(n)$,*

$$\|\hat{R}^N - R\| = \underbrace{O_p((1-\delta)\sqrt{(p+\log n)/n})}_{(A)} + \underbrace{O_p(\delta s_0(p)(\log p/n)^{(1-q)/2})}_{(B)} = \|(\hat{R}^N)^{-1} - R^{-1}\| \quad (8)$$

$$\|\hat{\Sigma}^N - \Sigma\| = O_p((1-\delta)\sqrt{(p+\log n)/n}) + O_p(\delta s_0(p)(\log p/n)^{(1-q)/2}) = \|(\hat{\Sigma}^N)^{-1} - \Sigma^{-1}\|. \quad (9)$$

The proofs are given in the Appendix. The NOVELIST estimators of the correlation and covariance matrices and their inverses yield the same convergence rate.

We now discuss the optimal asymptotic δ under the settings of Propositions 1 and 2. Proposition 1 can be thought of as a ‘‘large p ’’ setting, while Proposition 2 applies to moderately large and small p .

2.3.2 Optimal δ and rate of convergence in Proposition 1. Proposition 1 corresponds to ‘‘large p ’’ scenarios, in which p can be thought of as being $O(n)$ or larger (indeed, the case $p = o(n)$ is covered by Proposition 2). For such a large p , the pre-condition for the consistency of the NOVELIST estimator is that $\delta \rightarrow 1$, i.e. that the estimator asymptotically degenerates to the thresholding estimator. To see this, take $p = n^{1+\Delta}$ with $\Delta \geq 0$. If $\delta \not\rightarrow 1$, the error in part (A) of formula (6) would be of order $n^{1/2+\Delta} \sqrt{\log n^{1+\Delta}}$ and therefore would not converge to zero.

Focusing on \hat{R}^N without loss of generality, the optimal rate of convergence is obtained by equating parts (A) and (B) in formula (6). The resulting optimal shrinkage intensity $\tilde{\delta}$ is

$$\tilde{\delta} = \frac{p(\log p/n)^{q/2}}{s_0(p) + p(\log p/n)^{q/2}} = \frac{(\log p/n)^{q/2}}{s_0(p)/p + (\log p/n)^{q/2}}. \quad (10)$$

In typical scenarios, bearing in mind that p is at least of order n or larger, and that $q < 1$, the term $s_0(p)/p$ will tend to zero much faster than the term $(\log p/n)^{q/2}$, which will result in $\tilde{\delta} \rightarrow 1$ and in the rate of convergence of NOVELIST being $O_p(s_0(p)(\log p/n)^{(1-q)/2})$. Examples of such scenarios are given directly below.

Scenario 1 $q = 0$, $s_0(p) = o((n/\log p)^{1/2})$.

When $q = 0$, the uniformity class of correlation matrices controls the maximum number of non-zero entries in each row. The typical example is the moving-average (MA) autocorrelation structure in time series.

Scenario 2 $q \neq 0$, $s_0(p) \leq C$ as $p \rightarrow \infty$.

A typical example of this scenario is the auto-regressive (AR) autocorrelation structure.

We now show a scenario in which NOVELIST is inconsistent, under the setting of Proposition 1. Consider the long-memory autocorrelation matrix, $\rho_{ij} \sim |i - j|^{-\alpha}$, $0 < \alpha \leq 1$, for which $s_0(p) = \max_{1 \leq i \leq p} \sum_{j=1}^p \max(1, |i - j|)^{-\alpha q} = O(p^{1-\alpha q})$. Take $q \neq 0$. Note a sufficient condition for $\tilde{\delta}$ to tend to 1 is that $(\log p)^{(1/2)} n^{-1/2} p^\alpha \rightarrow \infty$. This more easily happens for larger α 's, i.e. for ‘‘less long’’-memory processes. However, considering the implied rate of convergence, we have $s_0(p)(\log p/n)^{(1-q)/2} = p^{1-\alpha q}(\log p/n)^{(1-q)/2}$, which is divergent even if $\alpha = 1$.

2.3.3 Optimal δ and rate of convergence in Proposition 2. Similarly, in the setting of Proposition 2, the resulting optimal shrinkage intensity $\tilde{\delta}$ is

$$\tilde{\delta} = \frac{((p + \log n)/n)^{1/2}}{((p + \log n)/n)^{1/2} + s_0(p)(\log p/n)^{(1-q)/2}}. \quad (11)$$

We now highlight a few special-case scenarios.

Scenario 3 p fixed (and hence $q = 0$).

Note that in the case of p being fixed or bounded in n , one can take $q = 0$ (to obtain as fast a rate for the thresholding part as possible) as the implied $s_0(p)$ will also be bounded in n . In this case, we have $\tilde{\delta} \rightarrow 1$ (and hence NOVELIST degenerates to the thresholding estimator with its corresponding speed of convergence), but the speed at which $\tilde{\delta}$ approaches 1 is extremely slow ($O(\log^{-1/2} n)$).

Scenario 4 $p \rightarrow \infty$ with n , and $q = 0$.

In this case, the quantity $\{(p + \log n)/\log p\}^{1/2}$ acts as a transition phase: if $s_0(p)$ is of a larger order, then we have $\tilde{\delta} \rightarrow 0$; if it is of a smaller order, then $\tilde{\delta} \rightarrow 1$; if it is of this order and if $\tilde{\delta}$ has a limit, then its limit lies in $(0, 1)$. Therefore NOVELIST will be closer to the sample covariance (correlation) if the truth is dense (i.e. if $s_0(p)$ is large), and closer to the thresholding estimator if $s_0(p)$ is small.

Scenario 5 $p \rightarrow \infty$ with n , and $q \neq 0$.

Here, the transition phase quantity is $\frac{(p+\log n)^{1/2}}{(\log p)^{\frac{1-q}{2}} n^{q/2}}$ and conclusions analogous to those of the preceding Scenario can be formed.

In the context of Scenario 5, we now revisit the long-memory example from before. The most “difficult” case still included in the setting of Proposition 2 is when p is “almost” the size of n ; therefore, we assume $p = n^{1-\Delta}$, with Δ being a small positive constant. Neglecting the logarithmic factors, the transition-phase quantity $\frac{(p+\log n)^{1/2}}{(\log p)^{\frac{1-q}{2}} n^{q/2}}$ reduces to $n^{\frac{1-\Delta-q}{2}}$. We have $s_0(p) = O(n^{(1-\Delta)(1-\alpha q)})$, and therefore $s_0(p)$ is of a larger order than $n^{\frac{1-\Delta-q}{2}}$ if $\alpha < \frac{1-\Delta+q}{2q(1-\Delta)}$; in this case, $\tilde{\delta} \rightarrow 0$, and the NOVELIST estimator degenerates to the sample covariance (correlation) estimator, which is consistent in this setting at the rate of $n^{-\Delta/2}$ (neglecting the log-factors). The other case, $\alpha \geq \frac{1-\Delta+q}{2q(1-\Delta)}$, is impossible as we must have $\alpha \leq 1$. Therefore, the NOVELIST estimator is consistent for the long-memory model under the setting of Proposition 2, i.e. when $p = o(n)$ (and degenerates to the sample covariance estimator). This is in contrast to the setting of Proposition 1, where, as argued before, the consistency of NOVELIST in the long-memory model cannot be shown.

3 δ outside $[0, 1]$

Some authors (Ledoit & Wolf, 2003; Schäfer & Strimmer, 2005; Savic & Karlsson, 2009), more or less explicitly, discuss the issue of the shrinkage intensity (for other shrinkage estimators) falling within versus outside the interval $[0, 1]$. Ledoit & Wolf (2003) “expect” it to lie between zero and one, Schäfer & Strimmer (2005) truncate it at zero or one, and Savic & Karlsson (2009) view negative shrinkage as a “useful signal for possible model misspecification”. We are interested in the performance of the NOVELIST estimator with $\delta \notin [0, 1]$, and have reasons to believe that $\delta \notin [0, 1]$ may be a good choice in certain scenarios.

We use the diagrams below to briefly illustrate this point. When the target T is appropriate, the “oracle” NOVELIST estimator (by which we mean one where δ is computed with the knowledge of the true R by minimising the spectral norm distance to R) will typically be in the convex hull of \hat{R} and T , i.e. $\delta \in [0, 1]$ as shown in the left graph. However, the target may also be misspecified. For example, if the true correlation matrix is highly non-sparse, the sparse target may be inappropriate, to the extent that R will be further away from T than from \hat{R} , as shown in the middle graph. In that case, the optimal δ should be negative to prevent NOVELIST being close to the target. By contrast, when the sample correlation is far from the (sparse) truth, perhaps because of high dimensionality, the optimal delta may be larger than one.




Diagram 1: Geometric illustration of shrinkage estimators. R is the truth, T is the target, \hat{R} is the sample correlation, \hat{R}_{opt}^N is the “oracle” NOVELIST estimator defined as the linear combination of T and \hat{R} with minimum spectral norm distance to R . LEFT: $\delta \in (0, 1)$ if target T is appropriate; MIDDLE: $\delta < 0$ if target T is misspecified; RIGHT: $\delta > 1$ if \hat{R} is far from R .

4 Empirical choices of (λ, δ) and algorithm

The choices of the shrinkage intensity (for shrinkage estimators) and the thresholding level (for thresholding estimators) are intensively studied in the literature. Bickel & Levina (2008b) propose a cross-validation method for choosing the threshold value for their thresholding estimator. However, NOVELIST requires simultaneous selection of the two parameters λ and δ , which makes straight cross-validation computationally intensive. Ledoit & Wolf (2003), and Schäfer & Strimmer (2005) give an analytic solution to the problem of choosing the optimal shrinkage level, under the Frobenius norm, for any shrinkage estimator. Since NOVELIST can be viewed as a shrinkage estimator, we borrow strength from this result and proceed by selecting the optimal shrinkage intensity $\delta^*(\lambda)$ in the sense of Ledoit & Wolf (2003) for each λ , and then perform cross-validation to select the best pair $(\lambda', \delta^*(\lambda'))$. This process significantly accelerates computation.

Cai & Liu (2011) and Fryzlewicz (2013) use adaptive thresholding for covariance matrices, in order to make thresholding insensitive to changes in the variance of the individual variables. This, effectively, corresponds to thresholding sample correlations rather than covariances. In the same vein, we apply NOVELIST to sample correlation matrices. We use soft thresholding as it often exhibits better and more stable empirical performance than hard thresholding, which is partly due to its being a continuous operation. Let $\hat{\Sigma}$ and \hat{R} be the sample covariance and correlation matrices computed on the whole dataset, and let $T = \{T_{ij}\}$ be the soft-thresholding estimator of the correlation matrix. The algorithm proceeds as follows.

For estimating the covariance matrix,

LW (Ledoit-Wolf) step: Using all available data, for each $\lambda \in (0, 1)$ chosen from a uniform grid of size m , find the optimal empirical δ as

$$\delta^*(\lambda) = \frac{\sum_{1 \leq i \neq j \leq n} \text{Var}(\hat{R}_{ij}) - \text{Cov}(\hat{R}_{ij}, T_{ij})}{\sum_{1 \leq i \neq j \leq n} (\hat{R}_{ij} - T_{ij})^2} = \frac{\sum_{1 \leq i \neq j \leq n} \text{Var}(\hat{R}_{ij}) \mathcal{I}(\hat{R}_{ij} < \lambda)}{\sum_{1 \leq i \neq j \leq n} (\hat{R}_{ij} - T_{ij})^2}, \quad (12)$$

to obtain the pair $(\lambda, \delta^*(\lambda))$.

The first equality comes from Ledoit & Wolf (2003), and the second follows because of the fact that our shrinkage target T is the soft-thresholding estimator with threshold λ (applied to the off-diagonal entries only).

CV (Cross-Validation) step: For each $z = 1, \dots, Z$, split the data randomly into two equal-size parts A (training data) and B (test data), letting $\hat{\Sigma}_A^{(z)}$ and $\hat{\Sigma}_B^{(z)}$ be the sample covariance matrices of these two datasets, and $\hat{R}_A^{(z)}$ and $\hat{R}_B^{(z)}$ – the sample correlation matrices.

1. For each λ , obtain the NOVELIST estimator of the correlation matrix $\hat{R}_A^{N^{(z)}}(\lambda) = \hat{R}^N(\hat{R}_A^{(z)}, \lambda, \delta^*(\lambda))$, and of the covariance matrix $\hat{\Sigma}_A^{N^{(z)}}(\lambda) = \hat{D}_A \hat{R}_A^{N^{(z)}}(\lambda) \hat{D}_A$, where $\hat{D}_A = (\text{diag}(\hat{\Sigma}_A^{(z)}))^{1/2}$.

2. Compute the spectral norm error $Err(\lambda)^{(z)} = \|\hat{\Sigma}_A^{N^{(z)}}(\lambda) - \hat{\Sigma}_B^{(z)}\|_2^2$.

3. Repeat steps 1 and 2 for each z and obtain the averaged error $Err(\lambda) = \frac{1}{Z} \sum_{z=1}^Z Err(\lambda)^{(z)}$. Find $\lambda' = \min_{\lambda} Err(\lambda)$, then obtain the optimal pair $(\lambda', \delta') = (\lambda', \delta^*(\lambda'))$.

4. Compute the cross-validated NOVELIST estimators of the correlation and covariance matrices as

$$\hat{R}_{cv}^N = \hat{R}^N(\hat{R}, \lambda', \delta'), \quad (13)$$

$$\hat{\Sigma}_{cv}^N = \hat{D} \hat{R}_{cv}^N \hat{D}, \quad (14)$$

where $\hat{D} = (\text{diag}(\hat{\Sigma}))^{1/2}$.

For estimating the inverses of the correlation and the covariance matrices, the difference lies in step 2, where the error measure is adjusted as follows. If $n > 2p$ (i.e. in the case when $\hat{\Sigma}_B^{(z)}$ is invertible), we use the measure $Err(\lambda)^{(z)} = \|(\hat{\Sigma}_A^{N^{(z)}}(\lambda))^{-1} - (\hat{\Sigma}_B^{(z)})^{-1}\|_2^2$; otherwise, use $Err(\lambda)^{(z)} = \|\|$

$(\hat{\Sigma}_A^{N(z)}(\lambda))^{-1} \hat{\Sigma}_B^{(z)} - \mathcal{I} \|_2^2$, where \mathcal{I} is the identity matrix. In step 4, we compute the cross-validated NOVELIST estimators of the inverted correlation and covariance matrices as

$$(\hat{R}_{cv}^N)^{-1} = (\hat{R}^N(\hat{R}, \lambda', \delta'))^{-1}, \quad (15)$$


$$(\hat{\Sigma}_{cv}^N)^{-1} = (\hat{D} \hat{R}_{cv}^N \hat{D})^{-1}. \quad (16)$$

We note that a closely related procedure for choosing δ has also been described in Lam & Feng (2017).

5 Empirical improvements of NOVELIST

5.1 Fixed parameters

As shown in the simulation study of Section 6.2, the performance of cross validation is generally adequate, except in estimating large precision matrices with highly non-sparse covariance structures, such as in factor models and long-memory autocovariance structures. To remedy this problem, we suggest that fixed, rather than cross-validated parameters be used, if the eigenanalysis of the sample correlation matrix indicates that there are prominent principal components, when $p > 2n$ or close. We suggest the following rules of thumb: first, we look for the evidence of “elbows” in the scree plot of eigenvalues, by examining if $\sum_{k=1}^p \mathbb{I}\{\gamma(k) + \gamma(k+2) - 2\gamma(k+1) > 0.1p\} > 0$, where $\gamma(k)$ is the k th principal component. If so, then we look for the evidence of long-memory decay, by examining if the off-diagonals of the sample correlation matrix follow a high-kurtosis distribution. If the sample kurtosis ≤ 3.5 , this suggests that the factor structure may be present, and we use the fixed parameters $(\lambda'', \delta'') = (0.90, 0.50)$; if the sample kurtosis > 3.5 , this may point to long memory, and we use the fixed parameters $(\lambda'', \delta'') = (0.50, 0.25)$. The above decision procedure, including all the specific parameter values, has been obtained through extensive numerical experiments not shown in this paper. It is sketched in the following flowchart.



Flowchart 1: Decision procedure for using cross-validated or fixed parameters in estimating precision matrices.

5.2 Principal-component-adjusted NOVELIST

NOVELIST can further benefit from any prior knowledge about the underlying covariance matrix, such as the factor model structure. If the underlying correlation matrix follows a factor model, we can decompose the sample correlation matrix as

$$\hat{R} = \sum_{k=1}^K \hat{\gamma}_{(k)} \hat{\xi}_{(k)} \hat{\xi}_{(k)}^T + \hat{R}_{rem}, \quad (17)$$

where $\hat{\gamma}_{(k)}$ and $\hat{\xi}_{(k)}$ are the k th eigenvalue and eigenvector of sample correlation matrix, K is the number up to which the principal components are considered to be “large”, and \hat{R}_{rem} is the sample correlation

matrix after removing the first K principal components. Instead of applying NOVELIST on \hat{R} directly, we keep the first K components unchanged and only apply NOVELIST to \hat{R}_{rem} . Principal-component-adjusted NOVELIST estimators are obtained by

$$\hat{R}_{rem}^N = \sum_{k=1}^K \hat{\gamma}_{(k)} \hat{\xi}_{(k)} \hat{\xi}_{(k)}^T + \hat{R}^N(\hat{R}_{rem}, \lambda, \delta), \quad (18)$$

$$\hat{\Sigma}_{rem}^N = \hat{D} \hat{R}_{rem}^N \hat{D}. \quad (19)$$

In the remainder of the paper, we always use the not-necessarily-optimal value $K = 1$. We suggest that PC-adjusted NOVELIST should only be used with prior knowledge or if empirical testing indicates that there are prominent principal components.

6 Simulation study

In this section, we investigate the performance of the NOVELIST estimator of covariance and precision matrices based on optimal and data-driven choices of (λ, δ) for seven different models and in comparison with five popular competitors. According to the algorithm in Section 4, the NOVELIST estimator of the correlation is obtained first; the corresponding estimator of the covariance follows by formula (14) and the inverse of the covariance estimator is obtained by formula (16). In all simulations, the sample size $n = 100$, and the dimension $p \in \{10, 100, 200, 500\}$. We perform $N = 50$ repetitions.

6.1 Simulation models

We use the following models for Σ .

- (A) *Identity*. $\sigma_{ij} = \mathbb{1}\{i = j\}$, for $1 \leq i, j \leq p$.
- (B) *MA(1) autocovariance structure*.

$$\sigma_{ij} = \begin{cases} 1, & \text{if } i = j; \\ \rho, & \text{if } |i - j| = 1; \\ 0, & \text{otherwise} \end{cases} \quad (20)$$

for $1 \leq i, j \leq p$. We set $\rho = 0.5$.

- (C) *AR(1) autocovariance structure*.

$$\sigma_{ij} = \rho^{|i-j|}, \quad \text{for } 1 \leq i, j \leq p, \quad (21)$$

with $\rho = 0.9$.

- (D) *Non-sparse covariance structure*. We generate a positive-definite matrix as

$$\Sigma = Q\Lambda Q^T, \quad (22)$$

where Q has iid standard normal entries and Λ is a diagonal matrix with its diagonal entries drawn independently from the χ_5^2 distribution. The resulting Σ is non-sparse and lacks an obvious pattern.

- (E) *Factor model covariance structure*. Let Σ be the covariance matrix of $\mathbf{X} = \{X_1, X_2, \dots, X_p\}^T$, which follows a 3-factor model

$$\mathbf{X}_{p \times n} = \mathbf{B}_{p \times 3} \mathbf{Y}_{3 \times n} + \mathbf{E}_{p \times n}, \quad (23)$$

where

$\mathbf{Y} = \{Y_1, Y_2, Y_3\}^T$ is a 3-dimensional factor, generated independently from the standard normal distribution, i.e. $\mathbf{Y} \sim \mathcal{N}(\mathbf{0}, \mathcal{I}_3)$,

$\mathbf{B} = \{\beta_{ij}\}$ is the coefficient matrix, $\beta_{ij} \stackrel{i.i.d.}{\sim} U(0, 1)$, $1 \leq i \leq p$, $1 \leq j \leq 3$,

$\mathbf{E} = \{\epsilon_1, \epsilon_2, \dots, \epsilon_p\}^T$ is p -dimensional random noise, generated independently from the standard normal distribution, $\epsilon \sim \mathcal{N}(\mathbf{0}, \mathbf{1})$.

Based on this model, we have $\sigma_{ij} = \begin{cases} \sum_{k=1}^3 \beta_{ik}^2 + 1 & \text{if } i = j; \\ \sum_{k=1}^3 \beta_{ik} \beta_{jk} & \text{if } i \neq j. \end{cases}$

(F) *Long-memory autocovariance structure.* We use the autocovariance matrix of the Fractional Gaussian Noise (FGN) process, with

$$\sigma_{ij} = \frac{1}{2} [|i-j| + 1 |^{2H} - 2 |i-j|^{2H} + |i-j| - 1 |^{2H}] \quad 1 \leq i, j \leq p. \quad (24)$$

The model is taken from Bickel & Levina (2008a), Section 6.1, and is non-sparse. We take $H = 0.9$ in order to investigate the case with strong long memory.

(G) *Seasonal covariance structure.*

$$\sigma_{ij} = \rho^{|i-j|} \mathbb{1}\{|i-j| = l\mathbb{Z}_{\geq 0}\}, \quad \text{for } 1 \leq i, j \leq p, \quad (25)$$

where $\mathbb{Z}_{\geq 0}$ is the set of non-negative integers. We take $l = 3$ and $\rho = 0.9$.

The models can be broadly divided into 3 groups. (A)-(C) and (G) are sparse, (D) is non-sparse, and (E) and (F) are highly non-sparse. In models (B), (C) (F) and (G), the covariance matrix equals the correlation matrix. In order to depart from the case of equal variances, we also work with modified versions of these models, denoted by (B*), (C*) (F*) and (G*), in which the correlation matrix $\{\rho_{ij}\}$ is generated as in (B), (C) (F) and (G), respectively, and which have unequal variances independently generated as $\sigma_{ii} \sim \chi_5^2$. As a result, in the ‘starred’ models, we have $\sigma_{ij} = \rho_{ij} \sqrt{\sigma_{ii} \sigma_{jj}}$, $i, j \in (1, p)$.

The performance of the competing estimators is presented in two parts. In the first part, we compare the estimators with optimal parameters identified with the knowledge of the true covariance matrix. These include (a) the soft thresholding estimator T_s , which applies the soft thresholding operator to the off-diagonal entries of \hat{R} only, as described in Section 2.1, (b) the banding estimator B (Section 2.1 in Bickel & Levina (2008a)), (c) the optimal NOVELIST estimator $\hat{\Sigma}_{opt}^N$ and (d) the optimal PC-adjusted NOVELIST estimator $\hat{\Sigma}_{opt.rem}^N$. In the second part, we compare the data-driven estimators including (e) the linear shrinkage estimator S (Target D in Table 2 from Schäfer & Strimmer (2005)), which estimates the correlation matrix by “shrinkage of the sample correlation towards the identity matrix” and estimates the variances by “shrinkage of the sample variances towards their median”, (f) the POET estimator P (Fan et al., 2013), (g) the cross-validated NOVELIST estimator $\hat{\Sigma}_{cv}^N$, (h) the PC-adjusted NOVELIST $\hat{\Sigma}_{rem}^N$, and (i) the nonlinear shrinkage estimator NS (Ledoit & Wolf, 2015). The sample covariance matrix $\hat{\Sigma}$ is also listed for reference. We use the R package *corpcor* to compute S , and the R package *POET* to compute P . In the latter, we use $k = 7$ as suggested by the authors, and use soft thresholding in NOVELIST and POET as it tends to offer better empirical performance. We use $Z = 50$ for $\hat{\Sigma}_{cv}^N$, and extend the interval for δ to $[-0.5, 1.5]$. $\hat{\Sigma}_{cv}^N$ with fixed parameters are only considered for estimating precision matrix under model (E), (F) and (F*) when $p = 100, 200, 500$. We use $K = 1$ for $\hat{\Sigma}_{opt.rem}^N$ and $\hat{\Sigma}_{rem}^N$. NS is performed by using the commercial package SNOPT for Matlab (Ledoit & Wolf, 2015).

6.2 Simulation results

Performance of $\hat{\Sigma}^N$ as a function of (λ, δ) . Examining the results presented in Figures 2 and 3 and Table 1, it is apparent that the performance of NOVELIST depends on the combinations of λ and δ used. Generally speaking, the average operator norm errors increase as sparsity decreases and dimension p increases. The positions of empirically optimal λ^* and δ^* are summarised below.




Fig. 2 Image plots of operator norm errors of NOVELIST estimators of Σ with different λ and δ under Models (A)-(C) and (G), $n = 100$, $p = 10$ (Left), 100 (Middle), 200 (Right), simulation times=50. The darker the area, the smaller the error.

1. The higher the degree of sparsity, the closer δ^* is to 1. The δ^* parameter tends to be close to 1 or slightly larger than 1 for the sparse group, around 0.5 for the non-sparse group, and about 0 or negative for the highly non-sparse group.
2. δ^* moves closer to 1 as p increases. This is especially true for the sparse group.
3. Unsurprisingly, the choice of λ is less important when δ is closer to 0.
4. Occasionally, $\delta^* \notin [0, 1]$. In particular, for the AR(1) and seasonal models, $\delta^* \in (1, 1.5]$, while in the highly non-sparse group, δ^* can take negative values, which is a reflection of the fact that $\hat{\Sigma}_{opt}^N$ attempts to reduce the effect of the strongly misspecified sparse target.

Performance of cross-validated choices of (λ, δ) . Table 1 shows that the cross-validated choices of the parameter (λ', δ') for $\hat{\Sigma}_{cv}^N$ are close to the optimal (λ^*, δ^*) for most models when $p = 10$, but there are bigger discrepancies between (λ', δ') and (λ^*, δ^*) as p increases, especially for the highly non-sparse group. Again, Figure 4, which only includes representative models from each sparsity category, shows that the choices of (λ', δ') are consistent with (λ^*, δ^*) in most of the cases. For models (A) and (C), cross




Fig. 3 Image plots of operator norm errors of NOVELIST estimators of Σ with different λ and δ under Models (D)-(F), $n = 100$, $p = 10$ (Left), 100 (Middle), 200 (Right), simulation times=50. The darker the area, the smaller the error.

validation works very well: the vast majority of (λ', δ') lead to the error lying in the 1st decile of the possible error range, whereas for models (D) and (G) with $p = 10$, in the 1st or 2nd decile.

However, as shown in Tables 3 and 5, the performance of cross validation in estimating Σ^{-1} with highly non-sparse covariance structures, such as in factor models and long-memory autocovariance structures, is less good (a remedy to this was described in Section 5.1).

Comparison with competing estimators. For the estimators with the optimal parameters, NOVELIST performs the best for $p = 10$ for both Σ and Σ^{-1} , and beats the competitors across the non-sparse and highly non-sparse model classes when $p = 100, 200$ and 500 . The banding estimator beats NOVELIST in covariance matrix estimation in the homoscedastic sparse models by a small margin in the higher-dimensional cases. For the identity matrix, banding, thresholding and the optimal NOVELIST attain the same results. Optimal PC-adjusted NOVELIST achieves better relative results for estimating Σ^{-1} than for Σ .

In the competitions based on the data-driven estimators, when $p = 10$, the cross-validation NOVELIST is the best for most of the models with heteroscedastic variances, and only slightly worse than linear or nonlinear shrinkage estimator for the other models. When $p = 100, 200$ or 500 , the cross-validation NOVELIST is the best for most of the models in the sparse and the non-sparse groups (more so for heteroscedastic models) for both Σ and Σ^{-1} , but is beaten by POET for the factor model and the FGN model by a small margin, and is slightly worse than nonlinear shrinkage for homoscedastic sparse models. However, POET underperforms for the sparse and non-sparse models for Σ , and nonlinear shrinkage does worse than NOVELIST for heteroscedastic sparse models. The cases where the cross-validation NOVELIST performs the worst are rare. NOVELIST with fixed parameters as in Flowchart 1 for highly

Table 1 Choices of (λ^*, δ^*) and (λ', δ') for $\hat{\Sigma}^N$ (50 replications).

	$\hat{\Sigma}_{opt}^N$		$\hat{\Sigma}_{cv}^N$		$\hat{\Sigma}_{opt}^N$		$\hat{\Sigma}_{cv}^N$	
	λ^*	δ^*	λ'	δ'	λ^*	δ^*	λ'	δ'
	p=10, n=100				p=100, n=100			
(A) Identity	(0.50, 1.00)	1.00	0.60	1.00	(0.50, 1.00)	1.00	0.60	1.00
(B) MA(1)	0.15	1.00	0.25	0.80	0.20	1.00	0.20	0.95
(B*) MA(1)*	0.15	0.95	0.30	0.65	0.15	1.00	0.30	0.90
(C) AR(1)	0.50	0.00	0.40	0.15	0.15	0.50	0.10	0.70
(C*) AR(1)*	0.50	0.05	0.40	0.00	0.30	0.60	0.30	0.85
(D) Non-sparse	0.40	0.50	0.55	0.40	0.45	0.60	0.35	0.80
(E) Factor	0.40	0.00	0.65	0.10	0.20	-0.15	0.50	0.05
(F) FGN	0.50	-0.05	0.50	0.00	0.30	-0.10	0.55	0.05
(F*) FGN*	0.50	-0.05	0.50	0.00	0.40	-0.05	0.65	0.05
(G) Seasonal	0.15	0.75	0.15	0.70	0.10	1.30	0.05	1.50
(G*) Seasonal*	0.25	0.75	0.20	0.65	0.10	1.30	0.05	1.50
	p=200, n=100				p=500, n=100			
(A) Identity	0.55	1.00	0.60	1.00	0.55	1.00	0.60	1.00
(B) MA(1)	0.25	1.00	0.20	1.00	0.30	1.00	0.25	1.00
(B*) MA(1)*	0.25	1.00	0.25	0.95	0.25	1.00	0.20	1.00
(C) AR(1)	0.05	1.00	0.05	1.00	0.10	1.10	0.05	0.80
(C*) AR(1)*	0.05	1.10	0.05	1.30	0.10	0.95	0.10	1.10
(D) Non-sparse	0.30	0.65	0.55	0.40	0.40	0.75	0.40	0.90
(E) Factor	0.10	-0.10	0.60	0.05	0.20	-0.10	0.50	0.05
(F) FGN	0.30	0.05	0.65	0.10	0.35	0.10	0.40	0.10
(F*) FGN*	0.25	0.05	0.50	0.05	0.15	-0.10	0.35	0.10
(G) Seasonal	0.10	1.10	0.05	1.50	0.10	1.30	0.10	1.20
(G*) Seasonal*	0.10	1.10	0.05	1.50	0.10	1.30	0.10	1.20

non-sparse cases improves the results for Σ^{-1} . PC-adjusted NOVELIST can further improve the results for estimating Σ^{-1} but not for Σ . We would argue that NOVELIST is the overall best performer, followed by nonlinear shrinkage, linear shrinkage and POET.

7 Portfolio selection

In this section, we apply the NOVELIST algorithm and the competing methods to share portfolios composed of the constituents of the FTSE 100 index. Similar competitions were previously conducted to compare the performance of different covariance matrix estimators (Ledoit & Wolf, 2003; Lam, 2016). We compare the performance for risk minimisation purposes. The data were provided by Bloomberg.

Daily returns. Our first dataset consists of $p = 85$ stocks of FTSE 100 (we removed all those constituents that contained missing values) and 2606 daily returns $\{r_t\}$ for the period January 1st 2005 to December 31st 2015. We use data from the first $n = 120$ days to estimate the initial covariance matrices of the returns based on 6 different competing covariance matrix estimators, and create 6 portfolios with weights given by the well known weight formula

$$\hat{w}_t = \frac{\{\hat{\Sigma}_t^{(120)}\}^{-1} \mathbf{1}_p}{\mathbf{1}_p^T \{\hat{\Sigma}_t^{(120)}\}^{-1} \mathbf{1}_p}, \quad (26)$$

where $\hat{\Sigma}_t^{(120)}$ is an estimator of the $p \times p$ covariance matrix of the past 120-trading-day returns on trading day t (i.e. computed over days $t - 119$ to t) and $\mathbf{1}_p$ is the column vector of p ones. We hold these




Fig. 4 50 replicated cross validation choices of (δ', λ') (green circles) against the background of contour lines of operator norm distances to Σ under model (A), (C), (D) and (F) [equivalent to Figures 2 and 3], $n = 100$, $p = 10$ (Left), 100 (Middle), 200 (Right). The area inside the first contour line contains all combinations of (λ, δ) for which $\|\hat{\Sigma}^N(\lambda, \delta) - \Sigma\|$ is in the 1st decile of $[\min_{(\lambda, \delta)} \|\hat{\Sigma}^N(\lambda, \delta) - \Sigma\|, \max_{(\lambda, \delta)} \|\hat{\Sigma}^N(\lambda, \delta) - \Sigma\|]$.

portfolios for the next 22 trading days and compute their out-of-sample standard deviations as (Ledoit & Wolf, 2003)

$$STD = \left\{ \hat{w}_t \frac{1}{22} \sum_{i=1}^{22} r_{t+i} r_{t+i}^T \hat{w}_t^T \right\}^{1/2}, \quad (27)$$

which is a measure of risk. On the 23rd day, we liquidate the portfolios and start the process all over again based on the past 120 trading days. The dataset is composed of 113 instances of such 22-trading-day blocks and the average STD of each portfolio is computed.

5-minute returns. The second dataset consists of $p = 100$ constituents of FTSE 100 and 13770 five-minute returns $\{y_t\}$ for the period March 2nd 2015 to September 4th 2015 (135 trading days). The procedure is similar to the one above and only the differences are explained here. We use the first 2 days

Table 2 Average operator norm error to Σ for competing estimators with optimal parameters (50 replications). The best results and those up to 5% worse than the best are boxed. The worst results are in bold.

	$\hat{\Sigma}$	T_s	B	$\hat{\Sigma}_{opt}^N$	$\hat{\Sigma}_{opt.rem}^N$	$\hat{\Sigma}$	T_s	B	$\hat{\Sigma}_{opt}^N$	$\hat{\Sigma}_{opt.rem}^N$
	p=10, n=100					p=100, n=100				
(A) Identity	0.578	0.246	0.246	0.246	—	2.946	0.436	0.436	0.436	—
(B) MA(1)	0.623	0.447	0.361	0.435	—	3.055	0.670	0.554	0.668	—
(B*) MA(1)*	1.400	1.008	0.871	0.988	—	6.458	1.890	1.370	1.800	—
(C) AR(1)	1.148	0.762	1.072	0.475	—	6.112	4.977	3.999	4.703	—
(C*) AR(1)*	2.010	1.707	2.004	1.020	—	16.338	8.353	8.786	7.992	—
(D) Non-sparse	3.483	2.954	3.127	2.812	—	25.844	11.302	11.539	10.717	—
(E) Factor	1.811	1.462	1.742	1.120	1.221	14.350	13.675	13.993	9.881	9.921
(F) FGN	1.110	0.751	0.970	0.527	0.711	7.824	6.777	7.478	5.135	7.033
(F*) FGN*	2.239	1.617	2.108	1.129	1.683	15.666	13.383	15.147	10.878	13.782
(G) Seasonal	0.850	0.564	0.797	0.527	—	4.290	2.493	2.205	2.460	—
(G*) Seasonal*	1.664	1.228	1.594	1.158	—	6.694	3.028	2.362	2.959	—
	p=200, n=100					p=500, n=100				
(A) Identity	4.661	0.440	0.440	0.440	—	9.321	0.467	0.467	0.467	—
(B) MA(1)	4.886	0.717	0.626	0.716	—	9.828	0.761	0.729	0.761	—
(B*) MA(1)*	10.727	1.884	1.545	1.881	—	21.233	2.041	1.775	2.041	—
(C) AR(1)	10.291	6.922	4.898	6.768	—	17.877	9.311	5.584	9.261	—
(C*) AR(1)*	20.277	14.691	14.943	14.426	—	39.241	18.780	11.738	18.728	—
(D) Non-sparse	26.729	10.990	11.240	10.322	—	50.915	13.917	13.284	12.913	—
(E) Factor	31.183	28.053	29.819	20.463	20.432	82.451	65.234	73.807	48.104	48.928
(F) FGN	14.732	12.729	13.877	9.906	15.881	35.041	30.201	31.272	23.939	30.782
(F*) FGN*	32.370	26.692	29.862	20.357	28.983	68.154	66.833	66.320	49.853	55.998
(G) Seasonal	6.913	2.961	2.418	2.930	—	13.157	3.582	2.499	3.460	—
(G*) Seasonal*	14.709	6.427	5.171	6.350	—	27.627	7.873	5.660	7.538	—

Note: The results of $\hat{\Sigma}_{opt.rem}^N$ are only presented for the highly non-sparse group, i.e. Models (E), (F) and (F*).

($n = 204$) to estimate the initial covariance matrices of the returns and create portfolios with weights given by

$$\hat{y}_t = \frac{\{\hat{\Sigma}_t^{(204)}\}^{-1} \mathbf{1}_p}{\mathbf{1}_p^T \{\hat{\Sigma}_t^{(204)}\}^{-1} \mathbf{1}_p}, \quad (28)$$

where $\hat{\Sigma}_t^{(204)}$ is an estimator of the $p \times p$ covariance matrix of the 5-minute returns over the past 204 data points (2 days) at trading time t . We hold them for the next day and the out-of-sample standard deviations are calculated by

$$STD = \{\hat{w}_t \frac{1}{102} \sum_{i=1}^{102} r_{t+i} r_{t+i}^T \hat{w}_t^T\}^{1/2}. \quad (29)$$

We rebalance the portfolios every day and compute the sum of out-of-sample STD's over the 133 trading days.

Following the advice from Section 5.1, we apply fixed parameters for both NOVELIST and PC-adjusted NOVELIST. Table 6 shows the results. NOVELIST has the lowest risk for both daily and 5-minute portfolios, followed by PC-adjusted NOVELIST and nonlinear shrinkage in the low-frequency

Table 3 Average operator norm error to Σ for competing estimators with data-driven parameters (50 replications). The best results and those up to 5% worse than the best are boxed. The worst results are in bold.

	p=10, n=100						p=100, n=100					
	$\hat{\Sigma}$	S	P	$\hat{\Sigma}_{cu}^N$	$\hat{\Sigma}_{rem}^N$	NS	$\hat{\Sigma}$	S	P	$\hat{\Sigma}_{cu}^N$	$\hat{\Sigma}_{rem}^N$	NS
(A) Identity	0.578	0.084	0.823	0.263	—	0.116	2.946	0.088	3.657	0.446	—	0.087
(B) MA(1)	0.623	0.444	0.732	0.493	—	0.481	3.055	0.670	3.730	0.704	—	0.694
(B*) MA(1)*	1.400	1.165	1.546	1.159	—	1.191	6.458	1.985	8.015	1.877	—	2.449
(C) AR(1)	1.148	1.013	1.135	1.153	—	1.017	6.112	5.423	6.257	5.390	—	5.892
(C*) AR(1)*	2.010	2.190	2.291	2.114	—	2.190	16.338	8.878	19.468	8.446	—	12.095
(D) Non-sparse	3.483	3.120	3.860	3.046	—	2.934	25.844	12.453	29.355	11.739	—	11.730
(E) Factor	1.811	1.793	1.866	1.741	1.763	1.537	14.350	17.681	14.304	16.497	16.438	15.285
(F) FGN	1.110	0.849	1.020	1.021	1.024	0.980	7.824	6.628	7.798	7.799	7.732	7.554
(F*) FGN*	2.239	2.218	2.221	2.222	2.227	1.960	15.666	14.795	15.611	15.225	15.254	16.561
(G) Seasonal	0.850	0.666	0.852	0.687	—	0.659	4.290	3.200	4.826	2.534	—	3.098
(G*) Seasonal*	1.664	1.647	1.652	1.452	—	1.480	6.694	4.268	7.171	3.016	—	6.979
	p=200, n=100						p=500, n=100					
(A) Identity	4.661	0.058	5.414	0.443	—	0.067	9.321	0.064	10.076	0.468	—	0.047
(B) MA(1)	4.886	0.658	5.615	0.744	—	0.694	9.828	0.645	10.566	0.819	—	0.683
(B*) MA(1)*	10.727	2.094	12.458	1.956	—	2.729	21.233	2.060	23.034	2.116	—	3.004
(C) AR(1)	10.291	8.123	11.446	8.217	—	7.759	17.877	12.785	18.496	12.484	—	12.036
(C*) AR(1)*	20.277	18.172	23.721	16.251	—	18.751	39.241	26.571	40.903	18.903	—	24.581
(D) Non-sparse	26.729	11.920	30.108	11.220	—	10.993	50.915	13.758	54.462	13.636	—	12.996
(E) Factor	31.183	34.237	31.064	33.224	33.194	31.020	82.451	83.101	81.489	81.697	81.382	80.852
(F) FGN	14.732	12.961	14.376	14.640	14.593	14.125	35.041	26.672	34.344	31.296	30.992	36.299
(F*) FGN*	32.370	31.165	30.263	31.470	31.042	32.188	68.154	84.958	69.133	75.546	75.377	74.432
(G) Seasonal	6.913	4.126	7.403	2.972	—	4.016	13.157	4.994	13.722	3.471	—	4.949
(G*) Seasonal*	14.709	9.225	15.855	6.494	—	9.064	27.627	11.030	28.949	7.561	—	11.132

Note: For $\hat{\Sigma}_{rem}^N$, (λ'', δ'') is fixed to be (0.10, 0.30) in (E), and (0.30, 0.50) in (F) and (F*).

case, and by POET and nonlinear shrinkage in the high-frequency case. In summary, NOVELIST offers the best option in terms of risk minimisation.

8 Discussion

As many other covariance (correlation) matrix estimators which incorporate thresholding, the NOVELIST estimator is not guaranteed to be positive-definite in finite samples. To remedy this, our advice is similar to other authors' (e.g. Cai et al. (2010), Fan et al. (2013), Bickel & Levina (2008b)): we propose to diagonalise the NOVELIST estimator and replace any eigenvalues that fall under a certain small positive threshold by the value of that threshold. How to choose the threshold is, of course, an important matter, and we do not believe there is a generally accepted solution in the literature, partly because the value of the "best" such threshold will necessarily be problem-dependent. Denoting the such-corrected estimator by $\hat{\Sigma}^N(\zeta)$ (in the covariance case) and $\hat{R}^N(\zeta)$ (in the correlation case), where ζ is the eigenvalue-threshold, one possibility would be to choose the lowest possible ζ for which the matrix $\hat{\Sigma}^N(\hat{\Sigma}^N(\zeta))^{-1}$ (and analogously for the correlation case) resembles the identity matrix, in a certain user-specified sense.

We also note that either part of the NOVELIST estimator can be replaced by a banding-type estimator, for example as defined by Cai et al. (2010). In this way, we would depart from the particular construction of the NOVELIST estimator towards the more general idea of using convex combinations of two (or more) covariance estimators, which is conceptually and practically appealing but lies outside the scope of the current work.

Table 4 Average operator norm error to Σ^{-1} for competing estimators with optimal parameters (50 replications). The best results and those up to 5% worse than the best are boxed. The worst results are in bold.

	$\hat{\Sigma}$	T_s	B	$\hat{\Sigma}_{opt}^N$	$\hat{\Sigma}_{opt.rem}^N$	$\hat{\Sigma}$	T_s	B	$\hat{\Sigma}_{opt}^N$	$\hat{\Sigma}_{opt.rem}^N$	
p=10, n=100						p=100, n=100					
(A) Identity	0.917	0.281	0.281	0.281	—	—	0.469	0.469	0.469	—	
(B) MA(1)	1.177	0.681	0.656	0.605	—	—	1.244	1.300	1.166	—	
(B*) MA(1)*	0.626	0.489	0.732	0.442	—	—	0.846	0.779	0.745	—	
(C) AR(1)	9.078	7.751	9.078	5.502	—	—	14.313	18.064	10.792	—	
(C*) AR(1)*	4.491	2.736	4.491	2.339	—	—	8.915	7.298	6.001	—	
(D) Non-sparse	0.378	0.256	0.297	0.210	—	—	2.670	2.775	1.793	—	
(E) Factor	0.846	0.403	0.610	0.370	0.400	—	0.712	0.715	0.653	0.518	
(F) FGN	2.995	1.727	2.980	1.560	1.535	—	3.585	4.650	3.112	2.734	
(F*) FGN*	1.571	1.193	1.212	1.001	1.018	—	2.029	2.038	1.948	1.761	
(G) Seasonal	2.688	1.538	2.685	1.302	—	—	3.806	5.444	3.260	—	
(G*) Seasonal*	1.340	1.091	1.726	0.827	—	—	2.526	4.345	1.971	—	
p=200, n=100						p=500, n=100					
(A) Identity	—	0.527	0.527	0.527	—	—	0.599	0.599	0.599	—	
(B) MA(1)	—	1.358	1.530	1.258	—	—	1.405	1.562	1.377	—	
(B*) MA(1)*	—	1.100	0.795	0.850	—	—	1.040	1.145	0.962	—	
(C) AR(1)	—	15.023	18.122	11.469	—	—	15.622	18.136	11.064	—	
(C*) AR(1)*	—	14.509	20.358	7.362	—	—	18.392	23.740	7.155	—	
(D) Non-sparse	—	2.460	2.016	1.459	—	—	5.986	5.896	4.289	—	
(E) Factor	—	0.711	0.711	0.677	0.537	—	0.744	0.744	0.730	0.557	
(F) FGN	—	3.972	4.658	3.317	3.024	—	4.267	4.737	3.527	3.306	
(F*) FGN*	—	2.974	4.096	2.083	1.849	—	4.426	5.674	2.250	2.083	
(G) Seasonal	—	4.029	5.469	3.538	—	—	4.188	5.477	3.673	—	
(G*) Seasonal*	—	3.328	4.885	2.259	—	—	3.726	5.479	2.358	—	

Note: The results of $\hat{\Sigma}_{opt.rem}^N$ are only presented for the highly non-sparse group, i.e. Models (E), (F) and (F*). The worst results for model (A) with $p = 100, 200$ and 500 are not labelled, as T_s , B and $\hat{\Sigma}_{opt}^N$ obtain exactly the same results.

To summarise, the flexible control of the degree of shrinkage and thresholding offered by NOVELIST means that it is able to offer competitive performance across most models, and in situations in which it is not the best, it tends not to be much worse than the best performer. We recommend NOVELIST as a simple, good all-round covariance, correlation and precision matrix estimator ready for practical use across a variety of models and data dimensionalities.

9 Appendix

9.1 Additional lemmas and proofs

Firstly, we briefly introduce two lemmas that will be used in the proof of Proposition 1.

Lemma 1 *If F satisfies $\int_0^\infty \exp(\gamma t) dG_j(t) < \infty$, for $0 < |\gamma| < \gamma_0$, for some $\gamma_0 > 0$, where G_j is the cdf of X_{1j}^2 , $R = \{\rho_{ij}\}$ and $\Sigma = \{\sigma_{ij}\}$ are the true correlation and covariance matrices, $1 \leq i, j \leq p$, and $\sigma_{ii} \leq M$, where M is a constant, then, for sufficiently large M' , if $\lambda = M' \sqrt{\log p/n}$ and $\log p/n = o(1)$, we have $\max_{1 \leq i, j \leq p} |\hat{\rho}_{ij} - \rho_{ij}| = O_p(\sqrt{\log p/n})$, for $1 \leq i, j \leq p$.*

Table 5 Average operator norm error to Σ^{-1} for competing estimators with data-driven parameters (50 replications). The best results and those up to 5% worse than the best are boxed. The worst results are in bold.

	$\hat{\Sigma}^N$, $\hat{\Sigma}_{rem}^N$, NS						$\hat{\Sigma}^N$, $\hat{\Sigma}_{rem}^N$, NS					
	$\hat{\Sigma}$	S	P	$\hat{\Sigma}_{cv}^N$	$\hat{\Sigma}_{rem}^N$	NS	$\hat{\Sigma}$	S	P	$\hat{\Sigma}_{cv}^N$	$\hat{\Sigma}_{rem}^N$	NS
(A) Identity	0.917	0.090	4.472	0.469	—	0.146	—	0.045	0.882	0.472	—	0.109
(B) MA(1)	1.123	0.799	6.474	0.824	—	0.780	—	1.273	1.403	1.439	—	1.405
(B*) MA(1)*	0.626	0.526	4.892	0.448	—	0.440	—	1.358	0.993	0.935	—	1.748
(C) AR(1)	9.078	7.309	40.142	8.574	—	5.396	—	13.410	15.704	12.605	—	12.272
(C*) AR(1)*	4.941	5.390	27.593	4.841	—	3.264	—	12.508	13.649	10.167	—	13.446
(D) Non-sparse	0.378	0.500	1.705	0.328	—	0.340	—	2.937	2.916	2.910	—	2.979
(E) Factor	0.846	1.142	1.806	0.864	—	0.296	—	2.603	0.893	1.608	—	0.343
(F) FGN	2.995	1.864	16.530	2.097	—	1.701	—	4.565	3.060	4.212	—	3.122
(F*) FGN*	1.571	1.174	10.284	2.017	—	1.101	—	4.474	2.965	3.431	—	4.432
(G) Seasonal	2.688	1.897	13.175	2.103	2.115	1.687	—	4.229	4.721	3.839	—	3.947
(G*) Seasonal*	1.340	1.284	8.436	1.143	—	1.219	—	3.510	3.799	2.743	—	4.538
	$p=100, n=100$						$p=100, n=100$					
	$p=200, n=100$						$p=500, n=100$					
(A) Identity	—	0.046	0.930	0.529	—	0.136	—	0.078	0.923	0.601	—	0.139
(B) MA(1)	—	1.449	1.371	1.401	—	1.463	—	1.473	1.445	1.540	—	1.487
(B*) MA(1)*	—	1.293	1.256	1.169	—	1.906	—	1.914	1.140	1.221	—	2.463
(C) AR(1)	—	15.066	17.128	14.125	—	13.907	—	16.526	17.700	16.025	—	15.924
(C*) AR(1)*	—	17.480	18.286	13.201	—	19.037	—	22.833	23.053	19.169	—	23.740
(D) Non-sparse	—	2.602	2.842	2.563	—	3.206	—	5.998	6.171	5.994	—	5.660
(E) Factor	—	3.701	0.892	1.450	—	0.348	—	5.672	0.962	4.106	—	0.347
(F) FGN	—	9.397	3.552	5.670	—	3.434	—	8.621	3.933	6.652	—	3.752
(F*) FGN*	—	6.649	2.765	4.024	—	5.519	—	6.241	3.083	5.442	—	6.519
(G) Seasonal	—	4.676	5.019	4.176	—	4.526	—	5.045	5.256	4.548	—	5.001
(G*) Seasonal*	—	4.540	4.643	3.514	—	6.068	—	5.632	5.254	4.489	—	6.988

Note: For models (E), (F) and (F*), results by both cross validation and fixed parameters (in brackets) are presented for NOVELIST when $n < 2p$. For $\hat{\Sigma}_{cv}^N$, fixed parameters (λ'' , δ'') are (0.90, 0.50) for Model (E), and (0.50, 0.25) for Models (F) and (F*). For $\hat{\Sigma}_{rem}^N$, (λ'' , δ'') is fixed to be (0.50, 0.90) for (E), and (0.25, 0.65) for (F) and (F*).

Proof of Lemma 1: By the sub-multiplicative norm property $\|AB\| \leq \|A\| \|B\|$ (Golub & Van Loan, 1989), we write

$$\begin{aligned}
& \max_{1 \leq i, j \leq p} |\hat{\rho}_{ij} - \rho_{ij}| \\
&= \max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij}/(\hat{\sigma}_{ii}\hat{\sigma}_{jj})^{1/2} - \sigma_{ij}/(\sigma_{ii}\sigma_{jj})^{1/2}| \\
&\leq \max_{1 \leq i \leq p} |\hat{\sigma}_{ii}^{-1/2} - \sigma_{ii}^{-1/2}| \max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij} - \sigma_{ij}| \max_{1 \leq j \leq p} |\hat{\sigma}_{jj}^{-1/2} - \sigma_{jj}^{-1/2}| \\
&+ \max_{1 \leq i \leq p} |\hat{\sigma}_{ii}^{-1/2} - \sigma_{ii}^{-1/2}| \max_{1 \leq i, j \leq p} (|\hat{\sigma}_{ij}| |\sigma_{jj}^{-1/2}| + |\hat{\sigma}_{ii}^{-1/2}| |\sigma_{ij}|) \\
&+ \max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij} - \sigma_{ij}| \max_{1 \leq i \leq p} |\hat{\sigma}_{ii}^{-1/2}| \max_{1 \leq i \leq p} |\sigma_{ii}^{-1/2}| \\
&= O_p(\sqrt{\log p/n}) \tag{30}
\end{aligned}$$

Table 6 Standard deviation of minimum variance portfolios in percentage (daily and 5-minute returns).

	STD (Daily returns)	STD (5-min returns)
Sample	1.256	10.675
Linear shrinkage	0.851	7.809
Nonlinear shrinkage	0.733	7.670
POET	0.760	7.253
NOVELIST	0.709	6.987
PC-adjusted NOVELIST	0.715	8.577

The last equality holds as we have $\max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij} - \sigma_{ij}| = O_p(\sqrt{\log p/n}) = \max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij}^{-1} - \sigma_{ij}^{-1}|$ (Bickel & Levina, 2008b), and $\max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij}| = O_p(\sqrt{\log p/n}) = \max_{1 \leq i, j \leq p} |\hat{\sigma}_{ij}^{-1}|$, and $\sigma_{ii} \leq M$, $1 \leq i, j \leq p$. ■

Lemma 2 *If F satisfies $\int_0^\infty \exp(\gamma t) dG_j(t) < \infty$, for $0 < |\gamma| < \gamma_0$, for some $\gamma_0 > 0$, where G_j is the cdf of X_{1j}^2 , $R = \{\rho_{ij}\}$ is the true correlation matrix, $1 \leq i, j \leq p$, then, uniformly on $\mathcal{V}(q, s_0(p), \varepsilon_0)$, for sufficiently large M' , if $\lambda = M' \sqrt{\log p/n}$ and $\log p/n = o(1)$,*

$$\|T(\hat{R}, \lambda) - R\| = O_p(s_0(p)(\log p/n)^{(1-q)/2}). \quad (31)$$

where T is any kind of generalised thresholding estimator.

Lemma 2 is a correlation version of Theorem 1 in Rothman et al. (2009) and follows in a straightforward way by replacing $\hat{\Sigma}, \Sigma, \mathcal{U}(q, c_0(p), M, \varepsilon_0)$ and $c_0(p)$ by $\hat{R}, R, \mathcal{V}(q, s_0(p), \varepsilon_0)$ and $s_0(p)$ in the proof of the theorem.

Proof of Proposition 1:

We first show the result for \hat{R}^N . By the triangle inequality,

$$\begin{aligned} \|\hat{R}^N - R\| &= \|(1 - \delta)\hat{R} + \delta T(\hat{R}, \lambda) - R\| \\ &\leq (1 - \delta)\|\hat{R} - R\| + \delta\|T(\hat{R}, \lambda) - R\| \\ &= I + II. \end{aligned} \quad (32)$$

Using Lemma 2, we have

$$II = O_p\{\delta s_0(p)(\log p/n)^{(1-q)/2}\}. \quad (33)$$

For symmetric matrices M , Corollary 2.3.2 in Golub & Van Loan (1989) states that

$$\|M\| \leq (\|M\|_{(1,1)}\|M\|_{(\infty,\infty)})^{1/2} = \|M\|_{(1,1)} = \max_{1 \leq i \leq p} \sum_{j=1}^p |m_{ij}|. \quad (34)$$

Then by Lemma 1,

$$\|\hat{R} - R\| \leq \max_{1 \leq i \leq p} \sum_{j=1}^p |\hat{R}_{ij} - R_{ij}| \leq p \max_{1 \leq i, j \leq p} |\hat{\rho}_{ij} - \rho_{ij}| = O_p(p\sqrt{\log p/n}). \quad (35)$$

Thus, we have

$$I = (1 - \delta)\|\hat{R} - R\| \leq O_p((1 - \delta)p\sqrt{\log p/n}). \quad (36)$$

Combining formulae (33) and (36) yields the first equality. The second equality follows because

$$\|(\hat{R}^N)^{-1} - R^{-1}\| \asymp \|\hat{R}^N - R\| \quad (37)$$

uniformly on $\mathcal{V}(q, s_0(p), \varepsilon_0)$.

For the $\hat{\Sigma}^N$ estimator, recalling that $T = T(\hat{R}, \lambda)$ and $D = (\text{diag}(\Sigma))^{1/2}$, we have

$$\begin{aligned} \|\hat{\Sigma}^N - \Sigma\| &= \|\hat{D}\hat{R}^N\hat{D} - DRD\| \\ &= \|\hat{D}((1 - \delta)\hat{R} + \delta T)\hat{D} - DRD\| \\ &\leq (1 - \delta)\|\hat{\Sigma} - \Sigma\| + \delta\|\hat{D}T\hat{D} - DRD\| \\ &= III + IV. \end{aligned} \quad (38)$$

Similarly as in 36, we obtain $III = O_p((1 - \delta)p\sqrt{\log p/n})$. For IV , we write

$$\begin{aligned} & \|\hat{D}T\hat{D} - DRD\| \\ & \leq \|\hat{D} - D\| \|T - R\| \|\hat{D} - D\| + \|\hat{D} - D\| (\|T\| \|D\| + \|\hat{D}\| \|R\|) \\ & \quad + \|T - R\| \|\hat{D}\| \|D\| \\ & = O_p((1 + s_0(p)(\log p/n)^{-q/2})\sqrt{\log p/n}). \end{aligned} \quad (39)$$

The last equality holds as we have $\|T - R\| = O_p(s_0(p)(\log p/n)^{(1-q)/2})$, $\|\hat{D} - D\| = O_p(\sqrt{\log p/n})$, $\|\hat{D}\| = O_p(1) = \|T\|$, and $\|D\| = O(1)$ as $\sigma_{ii} < M$. Because $(\log p/n)^{q/2}(s_0(p))^{-1}$ is bounded from above by the assumption that $\log p/n = o(1)$ and $\|(\hat{\Sigma}^N)^{-1} - \Sigma^{-1}\| \asymp \|\hat{\Sigma}^N - \Sigma\|$ uniformly on $\mathcal{V}(q, s_0(p), \varepsilon_0)$, the result follows. ■

Proof of Proposition 2:

We only need to show the rate for the sample covariance (correlation) part as the arguments for the thresholding part are identical to those in Proposition 1. We first collect the relevant arguments from the proof of Lemma 3 in Cai et al. (2010). Let $\|\cdot\|$ denote the spectral norm of a matrix. From the proof of Lemma 3 in Cai et al. (2010), there exist vectors $v_1, v_2, \dots, v_{5^m} \in S^{m-1}$, where S^{m-1} is the unit sphere in the Euclidean distance in \mathbb{R}^m , such that

$$\|A\| \leq 4 \sup_{j \leq 5^m} |v_j^T A v_j|$$

for all $m \times m$ symmetric matrices A .

Consider now the sample covariance matrix $\hat{\Sigma} = \frac{1}{n} \sum_{i=1}^n \mathbf{X}_i \mathbf{X}_i^T$ (recall that $E(\mathbf{X}) = 0$), satisfying a sub-Gaussian condition in the sense that the length- p column vector \mathbf{X}_i satisfies

$$P(|v^T \mathbf{X}_i| > t) \leq \exp(-t^2 \rho/2)$$

for a certain $\rho > 0$, for all $t > 0$ and $\|v\|_2 = 1$.

Then, by the same arguments as in the proof of Lemma 3 in Cai et al. (2010), there exists $\rho_1 > 0$ such that

$$P\left\{\left|\frac{1}{n} \sum_{i=1}^n v^T (\mathbf{X}_i \mathbf{X}_i^T - \Sigma) v\right| > x\right\} \leq \exp(-nx^2 \rho_1/2),$$

where Σ is the population covariance matrix, for all $0 < x < \rho_1$ and $\|v\| = 1$.

We then bound

$$\begin{aligned} P(\|\hat{\Sigma} - \Sigma\| > x) & \leq P(4 \sup_{j \leq 5^p} |v_j^T (\hat{\Sigma} - \Sigma) v_j| > x) \\ & \leq 5^p \sup_{v_j} P(|v_j^T (\hat{\Sigma} - \Sigma) v_j| > y) \\ & \leq 5^p \exp(-ny^2 \rho_1/2) \\ & = \exp(p \log 5 - ny^2 \rho_1/2). \end{aligned}$$

with $y = x/4$.

As ρ_1 is unknown, the only ‘‘safe’’ y ’s to consider are such that $y \rightarrow 0$ as $n \rightarrow \infty$, uniformly over all permitted p . We now want

$$\exp(p \log 5 - ny^2 \rho_1/2) \leq \frac{C}{n} = \exp(\log C - \log n),$$

which leads to

$$y \geq \sqrt{\frac{2(p \log 5 + \log n - \log C)}{n\rho_1}}.$$

This can only converge to zero if $p = o(n)$. Under this assumption, we therefore indeed have

$$\|\hat{\Sigma} - \Sigma\| = O_P\left(\sqrt{\frac{p + \log n}{n}}\right),$$

which completes the proof. ■

References

- ALVAREZ, I., NIEMI, J. & SIMPSON, M. (2014). Bayesian inference for a covariance matrix. Preprint.
- BICKEL, P. & LEVINA, E. (2008a). Regularized estimation of large covariance matrices. *Annals of Statistics* **36**, 199–227.
- BICKEL, P. & LEVINA, E. (2008b). Covariance regularization by thresholding. *Annals of Statistics* **36**, 2577–2604.
- CAI, T. & LIU, W. (2011). Adaptive thresholding for sparse covariance matrix estimation. *Journal of the American Statistical Association* **106**, 672–684.
- CAI, T. T., ZHANG, C. & ZHOU, H. H. (2010). Optimal rates of convergence for covariance matrix estimation. *Annals of Statistics* **38**, 2118–2144.
- CHEN, C. (1979). Bayesian inference for a normal dispersion matrix and its application to stochastic multiple regression analysis. *Journal of the Royal Statistical Society Series B* **41**, 235–248.
- CROUX, C. & HAESBROECK, G. (2000). Principal component analysis based on robust estimators of the covariance or correlation matrix: influence functions and efficiencies. *Biometrika* **87**, 603–618.
- DICKEY, J. M., LINDLEY, D. V. & PRESS, S. J. (1985). Bayesian estimation of the dispersion matrix of a multivariate normal distribution. *Communications in Statistics-Theory and Methods* **14**, 1019–1034.
- EL KAROUI, N. (2008). Operator norm consistent estimation of large-dimensional sparse covariance matrices. *Annals of Statistics* **36**, 2717–2756.
- EVANS, I. G. (1965). Bayesian estimation of parameters of a multivariate normal distribution. *Journal of the Royal Statistical Society Series B* **27**, 279–283.
- FAN, J., FAN, Y. & LV, J. (2008). High dimensional covariance matrix estimation using a factor model. *Journal of Econometrics* **147**, 186–197.
- FAN, J., LIAO, Y. & MINCHEVA, M. (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society Series B* **75**, 603–680.
- FISHER, R. A. (1936). The use of multiple measurements in taxonomic problems. *Annals of Eugenics* **7**, 179–188.
- FRIEDMAN, J., HASTIE, T. & TIBSHIRANI, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics* **9**, 432–441.
- FRYZLEWICZ, P. (2013). High-dimensional volatility matrix estimation via wavelets and thresholding. *Biometrika* **100**, 921–938.
- FURRER, R. & BENGTTSSON, T. (2007). Estimation of high-dimensional prior and posterior covariance matrices in kalman filter variants. *Journal of Multivariate Analysis* **98**, 227–255.
- GARDNER, T. S., DI BERNARDO, D., LORENZ, D. & COLLINS, J. J. (2003). Inferring genetic networks and identifying compound mode of action via expression profiling. *Science* **301**, 102–105.
- GOLUB, G. H. & VAN LOAN, C. F. (1989). *Matrix Computations, 2nd ed.* Baltimore, MD: Johns Hopkins University Press.

- GUO, Y. Q., HASTIE, T. & TIBSHIRANI, R. (2007). Regularized linear discriminant analysis and its application in microarrays. *Biostatistics* **8**, 86–100.
- JEONG, H., MASON, S. P., BARABÁSI, A.-L. & N., O. Z. (2001). Lethality and centrality in protein networks. *Nature* **411**, 41–42.
- LAM, C. (2016). Nonparametric eigenvalue-regularized precision or covariance matrix estimation. *Annals of Statistics* **44**, 928–953.
- LAM, C. & FENG, P. (2017). Integrating regularized covariance matrix estimators. Preprint.
- LEDOIT, O. & PÉCHÉ, S. (2011). Eigenvectors of some large sample covariance matrix ensembles. *Probability Theory and Related Fields* **151**, 233–264.
- LEDOIT, O. & WOLF, M. (2003). Improved estimation of the covariance matrix of stock returns with an application to portfolio selection. *Journal of Empirical Finance* **10**, 603–621.
- LEDOIT, O. & WOLF, M. (2004). A well-conditioned estimator for large-dimensional covariance matrices. *Journal Multivariate Analysis* **88**, 365–411.
- LEDOIT, O. & WOLF, M. (2012). Nonlinear shrinkage and estimation of large-dimensional covariance matrices. *Annals of Statistics* **4**, 1024–1060.
- LEDOIT, O. & WOLF, M. (2015). Spectrum estimation: A unified framework for covariance matrix estimation and PCA in large dimensions. *Journal of Multivariate Analysis* **139**, 360–384.
- LEONARD, T. & JOHN, S. J. H. (2012). Bayesian inference for a covariance matrix. *Annals of Statistics* **20**, 1669–1696.
- LONGERSTAEY, J., ZANGARI, A. & HOWARD, S. (1996). Risk MetricsTM-technical document. Technical Document. J. P. Morgan, New York.
- MARKOWITZ, H. (1952). Portfolio selection. *The Journal of Finance* **7**, 77–91.
- MEINSHAUSEN, N. & BÜHLMANN, P. (2006). High-dimensional graphs and variable selection with the lasso. *Annals of Statistics* **34**, 1436–1462.
- PEARSON, K. (1901). On lines and planes of closest fit to systems of points in space. *Philosophical Magazine* **2**, 559–572.
- ROTHMAN, A. J., BICKEL, P., LEVINA, E. & ZHU, J. (2008). Sparse permutation invariant covariance estimation. *Electronic Journal of Statistics* **2**, 494–515.
- ROTHMAN, A. J., LEVINA, E. & ZHU, J. (2009). Generalized thresholding of large covariance matrices. *Journal of the American Statistical Association* **104**, 177–186.
- SAVIC, R. M. & KARLSSON, M. O. (2009). Importance of shrinkage in empirical bayes estimates for diagnostics: problems and solutions. *American Association of Pharmaceutical Scientists* **11**, 558–569.
- SCHÄFER, J. & STRIMMER, K. (2005). A shrinkage approach to large-scale covariance matrix estimation and implications for functional genomic. *Statistical Applications in Genetics and Molecular Biology* **4**, 1544–6115.
- WU, W. B. & POURAHMADI, M. (2003). Nonparametric estimation in the gaussian graphical model. *Biometrika* **90**, 831–844.
- ZOU, H. (2006). The adaptive lasso and its oracle properties. *Journal of the American Statistical Association* **101**, 1418–1429.

Acknowledgements Piotr Fryzlewicz's work has been supported by the Engineering and Physical Sciences Research Council grant no. EP/L014246/1.