

Effects of *Chlorella vulgaris* on blood and immunological parameters of Caspian Sea salmon (*Salmo trutta caspius*) fry exposed to Viral Nervous Necrosis (VNN) virus

Saberi A.¹; Zorriehzahra M.J.^{2*}; Emadi H.¹; Kakoolaki S.²,
Fatemi S.M.R.¹

Received: August 2015

Accepted: November 2015

Abstract:

In the present study, the effects of *Chlorella vulgaris* on blood and immunological parameters of Caspian salmon (*Salmo trutta caspius*) before and after exposure to Viral Nervous Necrosis (VNN) virus were examined. In this regard, four treatments in triplicate were chosen. Groups included one control and 3 treatments (T₁, T₂ and T₃). Fish in control group, T₁, T₂ and T₃ were fed diets supplemented with 0, 1×10⁸, 2×10⁷ and 3×10⁶ chlorella/450 g of food respectively, for sixty days. In addition, a virus supernatant was prepared from infected wild golden grey mullet (*Liza auratus*) and used for virus challenge of *S. trutta caspius*. Virus was injected intraperitoneally and blood samples were collected before and 14 days after the challenge. Immunological (IgM, C₃, C₄, total protein, respiratory burst, albumin and lysozyme) and changes in blood parameters (RBC, WBC, Htc, Hb, MCH, MCHC and MCV) were also measured. Results showed that *C. vulgaris* could act as a natural immunostimulant. Also, the alteration trend in hematological and immunological parameters showed that experimental fish could be considered to be resistant to VNN virus after exposure and fish treated with *C. vulgaris* were more resistant in comparison to those in the control group. The dose used in T₁ (1×10⁸ chlorella/450 g food) was the most effective approach with significant differences.

Keywords: *Chlorella vulgaris*, Blood parameters, Immunological parameters, *Salmo trutta caspius*, Viral Nervous Necrosis virus

1-Dept. of Marine Biology, Science and Research Branch, Islamic Azad University, Hesarak, Tehran, I.R. Iran

2-Aquatic Animal health and Diseases Dept., Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Tehran, I.R. Iran

* Corresponding author's Email: zorrieh@yahoo.com

Introduction

Viral Nervous Necrosis (VNN) (Yoshikoshi and Inoue, 1990), also known as viral encephalopathy and retinopathy (VER; OIE, 2003), is caused by piscine nodaviruses. It was first described in 1990 in hatchery-reared Japanese parrotfish (*Oplegnathus fasciatus*) in Japan (Yoshikoshi and Inoue, 1990) and after that in more than 50 species all around the world (Johnsen *et al.*, 2002; Munday *et al.*, 2002; Nakai *et al.*, 2009; Crane and Hyatt, 2011) including wild golden grey mullet (*Liza auratus*) in the Caspian Sea in Iran (Zorriehzahra *et al.*, 2005 & 2016). Affected juveniles and older fishes show a variety of erratic swimming behaviors such as spiral, whirling, belly-up at rest with inflation of swim bladder (Zorriehzahra, 2004). Histopathologically, the disease is characterized by severely extended necrosis and vacuolation of the central nervous system (brain, spinal cord) and retina, but sometimes fish in early larval stages lack tissue vacuolation (Munday *et al.*, 2002). VNN is highly resistant to various environmental conditions and can survive for a long time in sea water (Arimoto *et al.*, 1996). The disease is easily reproduced in healthy fish by co-habitation with infected fish, immersion in a virus suspension, or injection (Neguyen *et al.*, 1996; Grotmol *et al.*, 1999; Peducasse *et al.*, 1999). Therefore occurrence of VNN in the Caspian Sea raised concerns about the infection of other fish in this region including Caspian Sea salmon (*Salmo trutta caspius*).

Caspian salmon is considered as an endangered species and has been listed as a protected species in the Iranian environmental regulations since August 1999 and is also included in the Red Data Books of Kazakhstan and Turkmenistan (Vera *et al.*, 2010). Its natural populations have declined drastically in recent decades as a result of over-fishing, poaching, river pollution, destruction of natural spawning areas, and drought (Abdoli, 2000; Niksirat and Abdoli, 2009). It is anadromous and migrates up rivers to spawn. Captive breeding and conservation programs have been initiated to produce, restore, and protect populations (Sarvi *et al.*, 2006; Jalali and Amiri, 2009). More than 900,000 fingerlings are produced annually through mixed milt fertilization of wild breeders and also from long-time hatchery breeders due to the decline of wild populations (Annual report of IFO, 2010). It takes 2 years before the fingerlings reach a weight of 15–20 g, corresponding to a length of 10–15 cm, which is considered suitable for releasing (Vera *et al.*, 2010). It is important to know if Caspian Sea Salmon is at risk or if there is any approach to improve fish immune system against VNN. Furthermore, natural immunestimulants such as microalgae are always an appropriate option to test. Nowadays, these microscopic organisms are consumed as food supplements such as *Chlorella vulgaris* (Fradique *et al.*, 2010). *C. vulgaris* is a spherical microscopic cell with a 2–10 μm diameter (Yamamoto *et al.*, 2004). This microalga has a rapid

growth rate and is ideal for production because it is remarkably resistant against harsh conditions and invaders (Safi *et al.*, 2014). It's dry weight contains (42–58% of protein), (5–40% of lipid), and (12–55% of carbohydrate), pigments such as chlorophyll and carotenoid, vitamins and minerals which are used for different purposes like biofuels, human nutrition, wastewater treatment and animal feed (Safi *et al.*, 2014). Consequently, the effect of *C. vulgaris*, as a natural immunestimulator was examined in Caspian Sea salmon exposed to VNN virus.

Materials and methods

Fish

A total of 800 Caspian Sea salmon fingerlings (mean individual initial weight of 8–10 g) were transferred from center of propagation and culture of Salmonids (Kelardasht, Iran) to the Caspian Sea Ecology Research Center (Sari, Iran). Fish were adapted to new conditions for a week and they were fed on a basal diet during acclimation.

Fish health examination

Prior to the experimental challenge, 40 fish specimen that were selected randomly from the mentioned population were screened for nodavirus by RT-PCR. All samples were negative and no VNN infection was observed (Gomez *et al.*, 2004). The fish were acclimatised for one week and normal feeding behaviour was observed before the start of the challenge.

Microalgae

A pure culture of *C. vulgaris* was provided by the Caspian Sea Ecology Research Center. *C. vulgaris* cells was counted using a Neubauer slide and light microscopy (Liu *et al.*, 2007) and was added to fish food according to determined concentrations for treatments.

Treatments design

720 fish were divided into 12 groups, in the form of four treatments (one control group and three experimental groups) with three replicates, each containing 60 fish. Treatments were named control, T₁, T₂ and T₃ and were fed on a diet supplemented with different concentrations of *C. vulgaris*.

Rearing condition

The mean water pH, dissolved oxygen (DO) and temperature were 7.5, 8 mg/L and 15°C respectively during the course of the experiment.

Food and feeding

Food was prepared once every ten days. To prepare the food, at first *C. vulgaris* cells were counted using a Neubauer slide and were then added to ground food at a certain concentration for different treatments. The mixture of food and microalgae were dried in an oven. The control group was fed on the basal diet without *C. vulgaris*, while fish in T₁, T₂ and T₃ were fed with the diets supplemented with 1×10^8 , 2×10^7 and 3×10^6 chlorella/450 g food respectively. The feeding trial lasted for 60 days. Fish were feed two times a day (8:00 a.m, and 14:00 p.m) at a daily

feeding rate of 8% of body weight in the first 2 weeks and then 6% and 4% of body weight in the second and final three weeks, respectively.

Supernatant preparation

A supernatant of the virus was prepared using brains and eyes of infected wild golden grey mullet. According to protocol provided by Kokawa *et al.* (2008), brain tissues were mixed with HBSS (Hanks balanced salt solution), then centrifuged at 1610 g for 20 min at 15°C and filtered through a 0.45 nm filter. The prepared supernatant was stored at -80°C for use to challenge with *S. trutta caspius*.

Virus challenge

Sixty days after the start of the feeding experiments, 10 fish were selected randomly from each treatment. In this study, the mentioned treatment groups kept at 18°C were challenged with an intraperitoneal injection (IP) and 1 group was kept as a control. The RGNNV used for the challenge was propagated in a cell line (SSN-1) derived from the striped snakehead *Channa striatus*. The Caspian salmon were challenged with 100 µL RGNNV infected SSN-1 cell culture supernatant (1.0×10^7 TCID₅₀ mL⁻¹). The control groups were mock challenged with an injection of supernatant from uninfected SSN-1 cells (Húsgaro, *et al.*, 2001). Each experimental group consisted of 60 fish. Virulence of mentioned supernatant was examined in a trial challenge in guppy fish (*Poecilia reticulata*), (Nazari, 2014).

Blood sampling and measurements

Blood samples were collected before and 14 days after the challenge. Fish were anaesthetized with clove extract (Farahi *et al.*, 2011). Blood was collected from the caudal vein using 1mL sterile disposable plastic syringes. Blood was transferred into 1.5 mL heparinated tubes (Trittau, Germany) for hematological study and also in non-heparinated tubes for plasma biochemistry analysis. Heparinated blood samples were placed in a refrigerator at 4°C. Non-heparinated samples were immediately centrifuged at 4°C at 1500 ×g for 5 minutes. Plasma was collected with a micropipette and stored at -80°C until analysis. Blood parameters were measured which included Red Blood Cells (RBC), White Blood Cells (WBC) (Torfi Moazenzadeh *et al.*, 2015), Haematocrit (Hct; %) (Barros *et al.*, 2002), Haemoglobin (Hb; gL⁻¹) (Drabkin, 1945), MCH (pg/cell), Mean Corpuscular Volume (MCV) (fL) and Mean corpuscular hemoglobin concentration (MCHC) (g/dL) (Dacie and Lewis, 2001). Immune parameters were analysed using an autoanalyser (Mindray BS-200, China), with commercial clinical investigation kits (Pars Azmoon Kit, Tehran, Iran). Measured immunological parameters included total protein (TP), albumin (ALB), IgM, C₃, C₄, respiratory burst and lysozyme (Torfi Moazenzadeh *et al.*, 2015).

Statistical analysis

The mean and standard error of three replicates and control were reported for

all parameters at the level of $\alpha=0.05$. Paired sample t-test was used to determine the effect of each concentration of *C. vulgaris* on mean values of hematological and serum parameters in comparison to the control groups before and after virus exposure. The Pearson ratio was used to analyze the correlation of whole parameters together. Pearson's correlation coefficient was used to determine an association between the hematological and immunological parameters, pairwise. All statistical analyses were performed using the SPSS ver. 18.0.

Results

The results of measured hematological and immunological parameters and paired comparisons of their alteration trends before and after exposure to VNN virus are presented in Tables 1 and 2 and also in Figs. 1-4. No significant differences were observed in amount of Hct, Hb, RBC, WBC, MCHC, respiratory burst, C_3 , albumin (ALB), total protein (TP) and lysozyme in control groups and all experimental groups before the virus challenge ($p>0.05$) (Table 1). The highest amount of MCH was observed in T_1 but it showed no significant differences with control groups. Amount of IgM in T_1 was significantly more than other treatments. In addition, the amount of C_4 in T_1 , T_2 and T_3 were significantly more than in the control group ($p<0.05$), yet no significant differences were observed between those treatments ($p>0.05$).

Results of measured parameters after VNN virus exposure showed that MCV

in T_1 and T_2 were significantly more than T_3 and control group ($p<0.05$). The amount of MCHC and IgM in T_1 , T_2 and T_3 were significantly less than control group ($p<0.05$), yet no significant differences were observed between those treatments ($p>0.05$). Also, C_3 in control group was significantly more than experimental groups ($p<0.05$). In addition, significant differences were observed in C_3 of all experimental groups, so that C_3 increased follow by increasing the amount of *C. vulgaris* used in treatments.

Table 1: The effect of different concentrations of *Chlorella vulgaris* on some hematological parameters of Caspian salmon before and after VNN virus exposure ($n=36$) and paired test.

		Control	T ₃	T ₂	T ₁
Hct (%)	Before	36.33±0.88	38.5±1.72	35.82±2.44	35.83±1.99
	After	33.33±1.174	30.17±0.872	31.5±0.992	33.5±1.47
	Sig.	0.045	0.001	0.203	0.462
	R	0.429	0.761	0.43	-0.414
Hb (gL ⁻¹)	Before	6.93±0.25	6.83±0.30	6.56±0.32	6.68±0.30
	After	6.83±0.152	6.03±0.264	5.83±1.66	6.41±0.452
	Sig.	0.679	0.010	0.149	0.704
	R	0.479	0.761	-0.460	-0.527
RBC (×10 ⁶ μL)	Before	0.858±0.04	0.96±0.05	0.90±0.075	0.83±0.63
	After	1.06±0.041	0.99±0.067	0.91±0.024	0.99±0.068
	Sig.	0.022	0.624	0.921	0.195
	R	0.706	0.528	-0.044	-0.278
WBC (μL)	Before	7066±877	8433±1819	7633±743	7483±727
	After	8200±302	8116±675	9366±1213	10666±1630
	Sig.	0.333	0.877	0.321	0.163
	R	-0.486	-0.013	-0.249	-0.2.6
MCV* (fL)	Before	427±16.56	400±7.39	399±11.99	435±16.66
	After	312±5.59 ^b	306±11.47 ^b	345±3.40 ^a	341±11.14 ^a
	Sig.	0.000	0.001	0.005	0.005
	R	0.604	-0.100	0.334	0.334
MCH* (pg cell ⁻¹)	Before	81.41±2.58 ^a	71.18±1.12 ^b	73.88±3.59 ^b	81.03±2.63 ^a
	After	64.13±1.18	60.85±1.82	63.96±0.58	64.81±1.11
	Sig.	0.548	0.879	0.046	0.001
	R	0.311	-0.081	-0.096	0.460
MCHC* (g L ⁻¹)	Before	19.05±0.29	17.78±0.29	18.46±0.44	18.65±0.32
	After	20.55±0.36 ^b	19.91±0.40 ^a	18.55±0.16 ^a	19.05±0.59 ^a
	Sig.	0.009	0.019	0.888	0.464
	R	0.425	-0.611	-0.670	0.523

* In each row, shows significant difference at least between 2 groups, before or after VNN virus exposure or both of them ($p < .05$). In each row, different superscripts show significant difference ($\alpha=0.05$). Sig.s less than 0.05 ($p < 0.05$), in paired compare between a parameter before and after VNN virus exposure shows significant difference. "r" shows Pearson correlation of a parameter before and after VNN virus exposure.

Table 2: The effect of different concentrations of *Chlorella vulgaris* on some immunological parameters of Caspian salmon before and after VNN virus exposure ($n=36$) and paired test.

		Control	T ₃	T ₂	T ₁
Respiratory Burst	Before	1482±168	1279±91	1491±119	1314±65
	After	3064±102	2744±324	2129±211	2830±206
	Sig.	0.000	0.009	0.002	0.000
	r	0.282	-0.172	-0.095	0.430
IgM* (mg L ⁻¹)	Before	107.26±10.2 ^{cd}	129.36±9.45 ^{bcd}	142.23±11.45 ^{bc}	170.46±5.90 ^a
	After	131.16±9.58 ^b	83.20±6.34 ^a	82.18±5.23 ^a	83.96±4.69 ^a
	Sig.	0.224	0.024	0.009	0.000
	r	-0.668	-0.670	-0.434	0.714

Continued Table 2:

	Control	T ₃	T ₂	T ₁	Control
C3* (mg L ⁻¹)	Before	29.95±3.32	29.60±1.45	35.35±3.42	28.81±2.57
	After	50.40±3.95 ^{bd}	20.93±1.34 ^{bc}	27.55±1.38 ^{bc}	40.81±4.22 ^a
	Sig.	0.002	0.008	0.068	0.010
	R	0.538	-0.059	0.244	0.710
C4* (mg L ⁻¹)	Before	19.51±1.77 ^{bc}	24.50±1.06 ^{abc}	29.25±4.84 ^{abc}	34.88±5.29 ^{abc}
	After	17.91±1.80	17.23±1.85	16.46±1.19	16.28±1.65
	Sig.	0.558	0.015	0.028	0.024
	R	-0.013	0.129	0.639	-0.179
ALB (g L ⁻¹)	Before	2.13±0.21	1.98±0.07	1.90±0.06	2.23±0.15
	After	2.84±0.27	1.95±0.22	2.08±0.10	2.68±0.21
	Sig.	0.135	0.842	0.150	0.108
	R	0.704	0.945	0.241	0.285
TP (g L ⁻¹)	before	3.63±0.19	3.95±0.45	3.66±0.12	3.86±0.26
	After	3.88±0.37	3.03±0.10	3.41±0.21	4.33±0.60
	Sig.	0.432	0.101	0.269	0.389
	R	0.620	0.114	0.389	0.594
Lysozyme (IU/ml)	Before	1.23±0.15	1.61±0.21	1.31±0.37	1.24±0.17
	After	1.70±0.15	1.59±0.49	2.18±0.58	3.18±0.82
	Sig.	0.033	0.979	0.143	0.074
	R	0.490	-0.358	0.534	-0.096

*In each row, shows significant difference at least between 2 groups, before or after VNN virus exposure or both of them ($p < 0.05$). In each row, different superscripts show significant difference ($\alpha = 0.05$). Sig.s less than 0.05 ($p < 0.05$), in paired compare between a parameter before and after VNN virus exposure shows significant difference. "r" shows Pearson correlation of a parameter before and after VNN virus exposure.




Figure 1: Effect of different concentrations of *Chlorella vulgaris* on blood parameters before exposure to VNN virus.




Figure 2: Effect of different concentrations of *Chlorella vulgaris* on blood parameters after exposure to VNN virus.




Figure 3: Effect of different concentrations of *Chlorella vulgaris* on immunological parameters before exposure to VNN virus.




Figure 4: Effect of different concentrations of *Chlorella vulgaris* on immunological parameters after exposure to VNN virus.

Discussion

Research in fish immunostimulants is developing and many of these agents are currently in use in the aquaculture industry. Immunostimulants such as synthetic chemicals or natural ingredient like microalgae, increase resistance to infectious diseases, not by enhancing specific immune responses, but by enhancing non-specific immune defence mechanisms. Use of these immunostimulants is an effective means of increasing the immunocompetency and disease resistance of fish (Sakai, 1999; Klesius *et al.*, 2001; Subasinghe, 2009). Results of the present study showed that *C. vulgaris* could have acted as a natural immunestimulator in Caspian salmon. Different studies have been shown that microalgae such as *Spirulina platensis* (Ibrahim *et al.*, 2013), *Nannochloropsis oculata* (Yanuhar *et al.*, 2011), *C. minutissima*

(Katharios *et al.*, 2005) and also *C. vulgaris* (Xu *et al.*, 2014) are effective agents in improving fish immune system. Crude polysaccharide extracts from the microalgae *Chlorella stigmatophora* (Chlorophyceae) and *Phaeodactylum tricornutum* (Bacillariophyceae) have shown strong anti-inflammatory and immunomodulatory activities both *in vivo* and *in vitro* (Guzman *et al.*, 2003).

No mortality or disease symptoms were observed in fish 14 days after they were challenged with VNN virus which indicates that Caspian salmon (*S. trutta caspius*) could be considered as a resistant species to VNN virus or fish need longer time to show probable disease symptoms. Results of cell culture, IFAT and RT-PCR of challenged fish were also negative (Data will be revealed in the next article

soon) that confirmed *S. trutta caspius* is resistant to VNN virus.

However, the most effective concentration of *C. vulgaris* to improve fish immune system was observed in T₁ (1×10^8), so that IgM and C₄ in *C. vulgaris* treated fish were significantly more than that in the control group and IgM in T₁ was significantly more than that in other groups.

C. vulgaris had a definite influence on the fish immune system in that it increased in all experimental groups even in T₃ with the lowest *C. vulgaris* concentration. Paired comparison of IgM levels before and after the challenge in treatments showed that the factor decreased after challenge in fish treated with *C. vulgaris* but increased in the control group. It seems that *C. vulgaris* treated fish were prepared against VNN virus in terms of IgM and the challenge caused IgM consumption, yet the humoral system was not active in the control group. IgM was produced just after the challenge and it would be consumed in a longer time (longer than the duration of the experiment).

In teleost fish, evaluating the complement system as a humoral component is an essential part of the innate immune system. A specific immunoglobulin or IgM is triggered by the binding of an antibody to the cell surface but can also be activated by acute phase proteins such as ligand-bound CRP or directly by viruses, bacteria and virus-infected cells (Balfry and Higgs, 2001; Aoki *et al.*, 2008).

C₄ is a Non-specific humoral molecule and is a part of complement system that has an effective role in the

fish immune system (Kum and Sekkin, 2011).

Results showed that *C. vulgaris* has no effect on blood parameters before VNN virus exposure; however it seems that higher the amounts of MCV in T₁ and T₂ after virus exposure is related to consuming more *C. vulgaris* before the challenge. In addition, MCHC increased in all treatments after exposure in that significant differences were observed between the control group and other experimental treatments but significant alteration of MCHC before and after the challenge were only observed in the control group and T₃ that were treated with no and a low concentration of *C. vulgaris*, respectively. It seems that consuming higher concentrations of *C. vulgaris* helps in reducing changes in MCV and MCHC after virus exposure.

C. vulgaris has no significant effect on Hct and Hb of fish before the challenge. Similarly, spirulina causes no significant differences in Hct and Hb levels in olive flounder (*Paralichthys olivaceus*) (Kim *et al.*, 2013). Paired test results showed that higher concentrations of microalgae (concentration used in T₁ and T₂) cause lesser impact of virus on Hct and no significant differences were observed in T₁ and T₂ Hct levels not only before but also, after virus exposure. In addition, VNN virus affected Hct levels in fish in T₃ and in the control group which received less concentration or no *C. vulgaris*. Significant differences in Hb levels were only observed in T₃ suggesting no effect of *C. vulgaris* on Hb during the experiment.

In addition, fish start showing resistance to the virus after virus injection in that significant increase was observed in the respiratory burst trend before and after the challenge in all treatments that is in agreement with the results of studies by Kim *et al.*, (2013), Watanuki *et al.* (2006) and Abdel-Tawwab and Ahmad (2009). They found spirulina causes significantly higher respiratory burst activity in olive flounder (*P. olivaceus*), common carp (*Cyprinus carpio*), and Nile tilapia (*Oreochromis niloticus*), respectively.

Also, changes in other immune parameters such as C₄, C₃, and IgM showed that fish had begun to show resistance against VNN virus. The alteration trend in C₃ and following that in MCV and IgM showed that fish in T₁ were more ready than other treatments to deal with the virus. If fish blood and immunological parameters had been studied for a longer time, it would have been possible to see significant increase in WBC, TP and ALB that would suggest that *C. vulgaris* concentration in T₁ had the best effects against VNN virus.

It seems that C₃ and lysozyme alteration trends were not caused by the virus but may have been accidental.

Lysozyme activity was not changed by *C. vulgaris* supplementation. Paired tests showed that lysozyme in T₁ and in the control group increased significantly and that it was accidental and not due to *C. vulgaris* or the virus effect. On the contrary, Kim *et al* (2013) suggested spirulina's influence on lysozyme activity of olive flounder (*P. olivaceus*) and Promya and

Chitmanat (2011) showed a significant increase in serum lysozyme activity in African sharptooth catfish (*Clarias gariepinus*) fed 3% or 5% dietary spirulina. Differences observed in the present study and other mentioned studies might depend on fish species, nutrition, environmental condition, age (Adams *et al.*, 2004; Adams and Thompson, 2008) and many other factors that were dissimilar in special experimental condition.

Immune systems affected by drugs such as immunostimulants, should act through the enhancement of the innate immune response (Austin and Brunt, 2009; Nayak, 2010) that was also observed in many measured factors in the present study.

The antiviral effect of microalgae is due to the existence of sulfated polysaccharides, which can inhibit viral infection and/or replicate (Fabregas *et al.*, 1999). Some functions of immune parameters like C₄, IgM, lysozym, etc include promoted binding of microbes to phagocytes, promoted inflammation at the complement activation, that cause osmotic lysis or apoptotic death. Also, change in the surface charge of microbes to facilitate phagocytosis, and haemolytic and antivirucidal effects (Kum and Sekkin, 2011) get improved due to the use of natural immunestimolants such as *C.vulgaris*.

In conclusion, by experimentation done in this study we can suggest that Caspian salmon should be considered as resistant to VNN virus. Also we can not deny the fact that since the exposure time was only 14 days in the present study, it might be possible that the virus

might be able to cause symptoms in this fish in longer periods of time. However, most of the studies about the possible outbreak of VNN in challenged fishes were done for longer than 14 days. For example after 20 days in Sevenband grouper (*Epinephelus septemfasciatus*) (Tanaka *et al.*, 1998) and after 30 days in guppy, zebra, oscar, and gold fish (Zorriehzadra *et al.*, 2013). However other than time, there were other differences in these experiments such as fish species, mode of challenge, virus titration and water temperature that should be considered. However Lopez-Jimena *et al* (2012) focused on time and sampled European seabass (*Dicentrarchus labrax*) on days 3, 10, 17, 24, 31 and also 2 months after the challenge. Although VNN was previously reported in European seabass (Breton *et al.*, 1997), they reported antigenic proteins detected by the use of immunohistochemistry and also vacuolation in brains and retina tissues in the fish 3 days after the challenge.

Furthermore, it is important to emphasize that the trial infection was performed at a very low temperature (15°C), and according to recommendation of other researchers challenge trials at higher temperature should be recommendable before making final conclusions (A. Toffan, personal communication, May 18, 2015). On the other hand, the experimental species Caspian salmon (*S. trutta caspius*) is a coldwater species and the optimum temperature for them would be 10-16°C (trial temperature was 15°C). When water temperature

reaches above 18 °C, some physiological disorders may happen leading to undesirable mortality that would occur naturally. In fact, high temperatures more than 20-22°C would be hazardous for them.

Finally, according to the obtained results in this study, and the lack of findings from similar examinations in Caspian salmon, based on the effects of time and rearing temperature, which was set at 15°C, it may be concluded that the Viral Nervous Necrosis Virus (VNNV) cannot cause morbidity and mortality in Caspian Salmon under these conditions. However, we cannot definitely claim that Caspian salmon is a resistant species to VNN at higher degrees of temperature or longer time of exposure. Therefore, it is suggested to consider the effect of longer time on morbidity and mortality of VNN in challenged *S. trutta caspius* in future studies.

Acknowledgments

We would like to express our gratitude to the head and staff of Caspian Sea Ecology Research Center in Mazandaran Province for their technical and laboratory assistance.

References

- Abdel-Tawwab, M. and Ahmad, M.H., 2009.** Live spirulina (*Arthrospira platensis*) as a growth and immunity promoter for Nile tilapia, *Oreochromis niloticus* (L.), challenged with pathogenic *Aeromonas hydrophila*. *Aquaculture Research*, 40, 1037-1046. <http://dx.doi>.

- Abdoli, A., 2000.** Inland water fishes of Iran, 1st ed. Iranian Museum of Nature and Wild Life, Tehran.
- Adams, A., 2004.** Immunodiagnostics in aquaculture. *Bulletin of the European Association of Fish Pathologists*, 24, 33-37.
- Adams, A. and Thompson, K.D., 2008.** Recent applications of biotechnology to novel diagnostics for aquatic animals. *Revue Scientifique et Technique-Office International des Epizooties*, 27(1), 197-209.
- Annual Report of IFRO, 2010.** Iran Fisheries Organization, 45P.
- Aoki, T., Takano, T., Santos, M.D. and Kondo, H., 2008.** Molecular innate immunity in teleost fish: Review and future perspectives. In: fisheries for global welfare and environmental, memorial book of the 5th world fisheries congress, K. Tsukamoto, T. Kawamura, T. Takeuchi, T.D. Beard, Jr. and M.J. Kaiser (Eds.), 263-276, ISBN 978-4-88704-144-8, Terrapub, Setagaya-ku, Japan.
- Arimoto, M., Sato, J., Maruyama, K., Mimura, G. and Furusawa, I., 1996.** Effect of chemical and physical treatments on the inactivation of striped jack nervous necrosis virus (SJNNV). *Aquaculture*, 143, 15-22.
- Austin, B. and Brunt, J.W., 2009.** The use of probiotics in aquaculture, chapter 7. In: Aquaculture microbiology and biotechnology: Volume 1, D. Montet and R.C. Ray (Eds), 185-207, Science Publishers, ISBN 978-1-57808-574-3, Enfield, New Hampshire, USA.
- Balfry, S.K. and Higgs, D.A., 2001.** Influence of dietary lipid composition on the immune system and disease resistance of fish, chapter 11, In: Nutrition and fish health, L. Chhorn and C.D. Webster (Eds.), 213-234, The Haworth Press, Inc., ISBN 1-56022-887-3, Binghamton, New York, USA.
- Barros, M.M. Lim, C. and Klesius, P.H., 2002.** Effect of iron supplementation to cotton seed meal diets on growth performance of channel catfish, *Ictalurus punctatus*. *Journal of Applied Aquaculture*, 10(65), 86-92.
- Breton, A.L., Grisez, L., Sweetman, J. and Ollevier, F., 1997.** Viral nervous necrosis (VNN) associated with mass mortalities in cage-reared sea bass, *Dicentrarchus labrax* (L.). *Journal of Fish Diseases*, 20(2), 145-151.
- Crane, M. and Hyatt, A., 2011.** Viruses of fish: An overview of significant pathogens. *Viruses*, 3, 2025-2046.
- Dacie, J.V. and Lewis, S.M., 2001.** Practical hematology, 9th edn. Churchill Livingstone, London, 633P.
- Drabkin, D.R., 1945.** Crystallographic and optical properties of human hemoglobin: a proposal for the standardization of hemoglobin. *American Journal of Medicine Science*, 209, 268-270.
- Fabregas, J., Garcia, D., Fernandez-Alonso, M., Rocha, A.I., Gomez-Puertas, P., Escribano, J.M.,**

- Otero, A. and Coll, J.M., 1999.** In vitro inhibition of the replication of Haemorrhagic septicaemia virus and African swine fever virus by extracts from marine microalgae. *Antiviral Research*, 44(1), 67-73.
- Farahi, A., Kasiri, M., Talebi, A. and Sudagar, M., 2011.** Effects of clove extract as an anesthetic on sperm motility traits and some hematological parameters in prussian Carp *Carassius gibelio*. *Advances in Environmental Biology*, 5(6), 1406-1410.
- Fradique, M., Batista, A.P., Nunes, M.C., Gouveia, L., Bandarra, N.M. and Raymundo, A., 2010.** Incorporation of *Chlorella vulgaris* and *Spirulina maxima* biomass in pasta products. Part1: preparation and evaluation. *Journal of the Science of Food and Agriculture*, 90, 1656–1664.
- Gomez, D.K., Sato, J., Mushiake, K., Isshiki, T., Okinaka, Y. and Nakai, T., 2004.** PCR-based detection of betanodaviruses from cultured and wild marine fish with no clinical signs. *Journal of Fish Diseases*, 27, 603–608
- Grotmol S., Bergh, O. and Totland, G.K., 1999.** Transmission of viral encephalopathy and retinopathy (VER) to yolk-sac larvae of the Atlantic halibut *Hippoglossus hippoglossus*: occurrence of nodavirus in various organs and a possible route of infection. *Diseases of Aquatic Organisms*, 36, 95-106.
- Guzman, S., Gato, A., Lamela, M., Freire-Garabal, M. and Calleja, J.M., 2003.** Anti-inflammatory and immunomodulatory activities of polysaccharide from *Chlorella stigmatophora* and *Phaeodactylum tricornutum*. *Phytotherapy Research*, 17, 665-670.
- Husgaro, S., Grotmol, S., Hjeltnes, B.K., Rødseth, O.M. and Biering, E., 2001.** Immune response to a recombinant capsid protein of striped jack nervous necrosis virus (SJNNV) in turbot *Scophthalmus maximus* and Atlantic halibut *Hippoglossus hippoglossus*, and evaluation of a vaccine against SJNNV. *Diseases of Aquatic Organisms*, 45, pp. 33–44.
- Ibrahim, M.D., Mohamed, M.F. and Ibrahim, M.A., 2013.** The role of *Spirulina platensis* (*Arthrospira platensis*) in growth and immunity of Nile tilapia (*Oreochromis niloticus*) and its resistance to bacterial infection. *Journal of Agricultural Science*, 5(6), 109-117.
- Jalali, M.A. and B.M. Amiri, 2009.** Threatened fishes of the world: *Salmo trutta caspius* (Kessler, 1877) (Salmoniforms: Salmonidae). *Environmental Biology of Fishes*, 86, 375–376.
- Johansen, R., Sommerset, I., Tørud, B., Korsnes, K., Hjortaa, M.J., Nilsen, F., Nerland, A.H. and Dannevig, B.H., 2002.** Characterization of nodavirus and viral encephalopathy and retinopathy in farmed turbot, *Scophthalmus maximus* (L.). *Fish Diseases*, 27, 591–601.
- Katharios P., Papadakis, I.E., Prapas, A., Dermon, C.R., Ampatzis, K. and Divanach, P., 2005.** Mortality control of VNN disease in grouper,

- (*Epinephelus marginatus*) after prolonged bath in dense *Chlorella minutissima* culture. European aquaculture society, biotechnologies for quality, barcelona, Spain, 20-23 October 2004. pp. 455-456.
- Kim, S., Rahimnejad, S., Kim, K.W., Lee, B.J. and Lee, K.J., 2013.** Effects of dietary supplementation of Spirulina and Quercetin on growth, innate immune responses, disease resistance against *Edwardsiella tarda*, and dietary antioxidant capacity in the juvenile olive flounder *Paralichthys olivaceus*. *Fisheries Aquatic Science*, 16(1), 1-8.
- Klesius, P.H., Shoemaker, C.A., Evans, J.J. and Lim, C., 2001.** Vaccines: Prevention of diseases in aquatic animals, Chapter 17, In: *Nutrition and Fish Health*, L. Chhorn and C.D. Webster (Eds.), 317-335, The Haworth Press, Inc., ISBN 1-56022-887-3, Binghamton, New York, USA.
- Kokawa, Y., Takami, I., Nishizawa, T. and Yoshimizu, M., 2008.** A mixed infection in sevenband grouper *Epinephelus septemfasciatus* affected with viral nervous necrosis (VNN). *Aquaculture*, 284, pp. 41-45.
- Kum, C. and Sekkin, S., 2011.** The immune system drugs in fish: Immune function, immunoassay, drugs, recent advances in fish farms, Dr. Faruk Aral (Ed.), ISBN: 978-953-307-759-8, InTech, Available from: <http://www.intechopen.com/books/recent-advances-in-fish-farms/the-immune-system-drugs-in-fish-immune-function-immunoassay-drugs>.
- Liu, W., Au, D.W.T., Anderson, D.M., Lam, P.K.S. and Wu, R.S.S., 2007.** Effects of nutrients, salinity, pH and light: dark cycle on the production of reactive oxygen species in the alga *Chattonella marina*. *Journal of Experimental Marine Biology and Ecology*, 1-11. doi:10.1016/j.jembe.2007.03.007.
- Lopez-Jimena, B., Gasia-Rosado, E., Thompson, K.D., Adams, A., Infante, C., Borrego, J.J. and Alonso, M.C., 2012.** Distribution of red-spotted grouper nervous necrosis virus (RGNNV) antigens in nervous and non-nervous organs of European sea bass (*Dicentrarchus labrax*) during the course of an experimental challenge. *Journal of Veterinary Science*, 13 (4), 355-362.
- Munday, B.L., Kwang, J. and Moody, N., 2002.** Betanodavirus infections of teleost fish: A review. *Journal of Fish Diseases*, 25, 127-142.
- Nakai, T., Mori, K., Sugaya, T., Nishioka, T., Mushiake, K. and Yamashita, H., 2009.** Current knowledge on Viral Nervous Necrosis (VNN) and its causative betanodaviruses. *The Israeli Journal of Aquaculture – Bamidgah*, 61(3), 198-207.
- Nayak, S.K., 2010.** Probiotics and immunity: A fish perspective. *Fish and Shellfish Immunology*, 29, 2-14.
- Nazari, A., Hassan, M. D., Bovo, G., Zorriehzahra, M. J., Azmi, T.I. and Arshad, S.S., 2014.** Pathogenicity of viral nervous necrosis virus for Guppy fish,

- Poecilia reticulata*. *Iranian Journal of Fisheries Sciences*, 13(1), 168-177.
- Neguyen, H.D., Nakai, T. and Muroga, K., 1996.** Progression of striped jack nervous necrosis virus (SJNNV) infection in naturally and experimentally infected striped jack Larvae. *Diseases of Aquaculture Organisms*, 24, 99-105.
- Niksirat, H. and Abdoli, A., 2009.** On the status of the critically endangered Caspian brown trout, *Salmo trutta caspius*, during recent decades in the southern Caspian Sea basin. *Zoology in the Middle East*, 46, 55-60.
- Office International Epizooties (OIE), 2003.** Chapter 2.1.7. Viral encephalopathy and retinopathy. pp. 135-141. In: Manual of diagnostic tests for aquatic animals, 4th ed. Office International des Epizooties, Paris. 358P.
- Peducasse, S., Castric, J., Thiery, R., Jeffroy, J., Le Ven, A. and Baudin Laurencin, F., 1999.** Comparative study of viral encephalopathy and retinopathy in juvenile sea bass *Dicentrarchus labrax* infected in different ways. *Diseases of Aquatic Organisms*. 36, 11-20.
- Promya, J. and Chitmanat, C., 2011.** The effects of *Spirulina platensis* and *Cladophora* algae on the growth performance, meat quality and immunity stimulating capacity of the African sharptooth catfish (*Clarias gariepinus*). *International Journal of Agriculture & Biology*, 13, 77-82.
- Safi, C., Zebib, B., Merah, O., Pontalier, P. and Vaca-Garsia, C., 2014.** Morphology, composition, production, processing and applications of *Chlorella vulgaris*: A review. *Renewable and Sustainable Energy Reviews*, 35, 265-278.
- Sakai, M., 1999.** Current research status of fish immunostimulants. *Aquaculture*. 172, pp. 63-92.
- Sarvi, K.H., Niksirat, B.M. Amiri, Mirtorabi, S.M., Rafiee G.R. and Bakhtiyari, M., 2006.** Cryopreservation of semen from the endangered Caspian brown trout (*Salmo trutta caspius*). *Aquaculture*, 256, 564-569.
- Subasinghe, R., 2009.** Disease control in aquaculture and the responsible use of veterinary drugs and vaccines: The issues; prospects and challenges. In: Options Méditerranéennes, Series A, No. 86: The Use of Veterinary Drugs and Vaccines in Mediterranean Aquaculture, C. Rodgers and B. Basurco (Eds.), 5-11, CIHEAM/FAO, ISBN 2-85352-422-1, Zaragoza, Spain.
- Tanaka, Sh., Aoki, H. and Nakai, T., 1998.** Pathogenicity of the Nodavirus detected from diseased Sevenband grouper, *Epinephelus septemfasciatus*. *Fish Pathology*, 33 (1), 31-36.
- Torfi Moazenzadeh, M., Yaghoubi, M., Yavari, V., Agh, N., Marammazi, J.G. and Popovic, N.T., 2015.** Reference intervals for haematological and plasma biochemical parameters in sobaity sea bream juveniles (*Sparidentex hasta*, Valenciennes 1830). *Comparative Clinical Pathology*. DOI: 10.1007/s00580-015-2107-y

- Vera, M., Sourinejad, I., Bouza, C., Vilas, R., Pino-Querido, A., Kalbassi, M.R. and Martinez, P., 2010.** Phylogeography, genetic structure, and conservation of the endangered Caspian brown trout, *Salmo trutta caspius* (Kessler, 1877), from Iran. *Journal of Hydrobiologia*.
- Watanuki, H., Ota, K., Tassakka, A.C.M.A.R., Kato, T. and Sakai, M., 2006.** Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157-163.
- Xu, W., Gao, Qi, Zh., Qiu, Zh., Peng, M.J. and Shao, R., 2014.** Effect of dietary chlorella on the growth performance and physiological parameters of gibel carp, *Carassius auratus gibelio*. *Turkish Journal of Fisheries and Aquatic Sciences*, 14, 53-57.
- Yamamoto, M., Fujishita, M., Hirata, A. and Kawano, S., 2004.** Regeneration and maturation of daughter cell walls in the auto spore-forming green alga *Chlorella vulgaris* (Chlorophyta, Trebouxiophyceae). *Journal of Plant Research*, 117, 257-64.
- Yanhuar, U., Nurdiani, R. and Hertika, A.M.S., 2011.** Potency of *Nannochloropsis oculata* as antibacterial, antioxidant and antiviral on humpback grouper infected by *Vibrio alginolyticus* and Viral Nervous Necrotic. *Journal of Food science and Engineering*, 1, 323-330.
- Yoshikoshi K. and Inoue, K., 1990.** Viral nervous necrosis in hatchery-reared larvae and juveniles of Japanese parrotfish, *Oplegnathus fasciatus* (Temminck & Schlegel). *Journal of Fish Diseases*, 13, 69-77.
- Zorriehzahra, M.E.J., 2004.** Professional report of first occurrence of VNN in golden mullet (*Liza auratus*) in Mazandaran Province. Iran Fisheries Research Organization (IFRO). 45P. in Persian Language.
- Zorriehzahra, M.E.J., Nakai, T., Sharifpour, I., Gomes, D.K., Chi, S.C., Soltani, M., Mohd, D., Hj, H., Sharif Rohani, M. and Saidi, A.A., 2005.** Mortality of wild golden mullet (*Liza auratus*) in Iranian waters of the Caspian Sea associated with viral nervous necrosis like agent. *Iranian Journal of Fisheries Science*, 45, 43-58.
- Zorriehzahra, M.E.J., Ghasemi, M., Mehrabi, M.R., Kakoolaki, Sh., Radkhah, K., Nazari, A., Rohani, M.S. and Haghghi Karsidani, S., 2013.** The investigation on histopathological changes of four ornamental fish species due to expose of causative virus of viral nervous necrosis (VNN) disease. 16th international conference on diseases of fish and shellfish, 2-6 September 2013, At Tampere, Finland.
- Zorriehzahra, M.E.J., Ghasemi, M., Ghiasi, M., Karsidani, S. H., Bovo, G., Nazari, A. and Dhama, K., 2016.** Isolation and confirmation of viral nervous necrosis (VNN) disease in golden grey mullet (*Liza aurata*) and leaping mullet (*Liza saliens*) in the Iranian waters of the Caspian Sea. *Veterinary microbiology*, 190, 27-37.