

Technical University of Denmark



Tissue Translocation of Polystyrene Micro- and Nanoparticles in *Daphnia magna*?

Schuer, C.; Rist, Sinja; Hartmann, Nanna B.; Wagner, Martin

Published in:
SETAC Europe 28th Annual Meeting - Abstract book

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Schuer, C., Rist, S., Hartmann, N. B., & Wagner, M. (2018). Tissue Translocation of Polystyrene Micro- and Nanoparticles in *Daphnia magna*? In SETAC Europe 28th Annual Meeting - Abstract book (pp. 279-279). Brussels, Belgium: Society of Environmental Toxicology and Chemistry.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

TU183 Tissue Translocation of Polystyrene Micro- and Nanoparticles in *Daphnia magna*?

C. Schuer, Goethe University Frankfurt / Dpt. Aquatic Ecotoxicology; **S. Rist**, DTU (Technical University of Denmark) / Department of Environmental Engineering; **N.B. Hartmann**, Technical University of Denmark (DTU) / DTU Environment; **M. Wagner**, Norwegian University of Science and Technology / Department of Biology. The last decade has seen a surge in research investigating various aspects of micro- and nanoplastics originating from plastic pollution in aquatic ecosystems. Aspects include occurrence, uptake, and potential effects in biota. Working with particles in a laboratory setting bears its own kinds of challenges, some of which had already been faced by researchers in the realm of nanotoxicology. Our knowledge about biota-particle-interactions is still limited and often based on early studies that – due to the infancy of the field – may have deficiencies in the experimental design and quality controls. One such example relates to the potential of plastic particles to cross the gut epithelium and translocate to other tissues. This phenomenon has been reported in the literature for the freshwater cladoceran *Daphnia magna* and – if true – is of toxicological relevance. To substantiate the limited available data, the aim of our study was to replicate these findings. We exposed neonate daphnids in a number of scenarios regarding particle concentration and exposure duration at two independent geographical locations using animals from two separate cultures. We expanded on the previous experiment by improving imaging through the addition of a fructose-based-clearing followed by investigation through confocal laserscan microscopy. We additionally applied the lipophilic dye Nile red to localize lipid droplets. This step facilitated the identification of lipid droplets inside the tissue and could therefore associate fluorescence detected before staining to a respective tissue. Our findings potentially challenge previous publications that reported the translocation of both micro- and nanoparticles. This discrepancy may be based on false-negative results on our side or false-positive results in the earlier reports, both potentially caused by inadequate exposure settings during the investigative parts of the studies. We were unable to replicate these findings implying a tissue translocation of nano- and microplastics under conditions closely resembling those reported in the literature. This highlights that the replication of nano- and microplastics studies is important, especially if these have a high impact on the body of knowledge. Our study also demonstrates that attempts of replication are inhibited by a lack of transparency in reporting methodology and results. We were able to adapt a fructose-based clearing protocol to the use with high amounts of *Daphnia* samples.