

Technical University of Denmark



Integrated Assessment of Anaerobic Reductive Dechlorination of Chlorinated Ethenes – by Stable Isotope Analysis and Microbial Techniques

Ottosen, Cecilie Bang; Broholm, Mette Martina; Tsitonaki, Katerina; Zimmermann, Jeremy; Hunkeler, Daniel; Tuxen, Nina

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Ottosen, C. B., Broholm, M. M., Tsitonaki, K., Zimmermann, J., Hunkeler, D., & Tuxen, N. (2018). Integrated Assessment of Anaerobic Reductive Dechlorination of Chlorinated Ethenes – by Stable Isotope Analysis and Microbial Techniques. Abstract from 11th International Conference on Remediation of Chlorinated and Recalcitrant Compounds, Palm springs, United States.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Integrated Assessment of Anaerobic Reductive Dechlorination of Chlorinated Ethenes – by Stable Isotope Analysis and Microbial Techniques

Cecilie Bang Ottosen (ceot@env.dtu.dk),

Mette Martina Broholm (Technical University of Denmark, Kongens Lyngby, Denmark),

Katerina Tsitonaki (Orbicon, Taastrup, Denmark),

Jeremy Zimmermann, Daniel Hunkeler (University of Neuchâtel, Neuchâtel, Switzerland),

Nina Tuxen (Capital Region of Denmark, Hillerød, Denmark)

Background/Objectives: Natural degradation is a cost effective alternative to traditional remediation methods for chlorinated solvents. However, the degradation products formed through natural degradation processes are also hazardous, and complete dechlorination is necessary to reduce the risk. Documentation of the occurrence, extent and rate of the complete dechlorination process is therefore vital in producing reliable risk assessments. At the current knowledge level, no stand-alone method can provide this necessary information. The scope of this project was therefore to document and quantify the natural reductive dechlorination of chlorinated ethenes at a contaminated site by an integrated approach based on a variety of analytical and conceptual tools. Field investigations were carried out at a site in Denmark, where degradation occurs in the source zone and plume in a geologically complex and strongly reduced subsurface.

Approach/Activities: An extensive integrated approach including traditional methods as hydrogeological, geochemical and chemical measurements as well as the new promising techniques, microarrays e.g. QuantArray-Chlor (specific degraders and functional genes) and compound specific isotope analysis (CSIA) was applied at field scale, where 21 boreholes (several with two screens) were sampled.

Results/Lessons Learned: Measurements of specific degradation products can provide documentation for the occurrence of the natural degradation. However, as the contaminant distribution is affected by transport processes (cis-DCE and VC migrate further than PCE and TCE), it cannot alone be used to determine where the biodegradation occurs, for this advanced techniques can be integrated. The biodegradation potential, determined by the abundance of specific degraders and functional genes, was highest near the source zone and less abundant down-gradient. Carbon isotope results further revealed the highest enrichment in the $\delta^{13}\text{C}$ value for both cis-DCE and VC near the source zone, indicating currently higher degradation near the source zone than in the past and less potential for further degradation down-gradient. At sites, where there is a temporal and spatial variation in the conditions and degradation, a combination of microbial analysis and CSIA thus provide a more robust characterization of the biodegradation (areas, extent and rate). Application of isotope mass balances and a lumped variable enrichment factor further facilitate the quantification of biodegradation, even at sites where the initial isotopic signature is uncertain. The integrated approach provided essential information for the evaluation of the occurrence, extent and rate of the natural biodegradation at the complex site. This is promising for the future evaluation of natural biodegradation in site risk assessment, and for this specific project, the results have influenced the Regions prioritizing of the site.