

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Root-colonizing bacteria enhance the levels of (E)- β -caryophyllene produced by maize roots in response to rootworm feeding.

Authors: Chiriboga M X, Guo H, Campos-Herrera R, Röder G, Imperiali N, Keel C, Maurhofer M, Turlings TCJ

Journal: Oecologia

Year: 2018 Jun

Issue: 187

Volume: 2

Pages: 459-468

DOI: 10.1007/s00442-017-4055-5

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Oecologia

Root-colonizing bacteria enhance the levels of (E)- β -caryophyllene produced by maize roots in response to rootworm feeding --Manuscript Draft--

Manuscript Number:					
Full Title:	Root-colonizing bacteria enhance the levels of (E)- β -caryophyllene produced by maize roots in response to rootworm feeding				
Article Type:	Special Topic				
Corresponding Author:	Ted Turlings, PhD Universite de Neuchatel Neuchâtel , Neuchâtel SWITZERLAND				
Order of Authors:	Xavier M Chiriboga Huijuan Guo, PhD Raquel Campos-Herrera, PhD Gregory Röder, PhD Nicola Imperiali, PhD Christoph Keel, PhD Monika Maurhofer, PhD Ted Turlings, PhD				
Suggested Reviewers:	Li Chen, PhD Associate Professor, State Key Lab of Integrated Management of Pest Insects and Rodents, Institute of Biology chenli@ioz.ac.cn chemically mediated insect-plant relationships Nicole van Dam, PhD Professor, German Centre for Integrative Biodiversity Research nicole.vandam@idiv.de chemically mediated insect-plant relationships Jared G Ali, PhD Assistant Professor of Entomology, Pennsylvania State University jga8@psu.edu chemically mediated insect-plant relationships				
Opposed Reviewers:					
Funding Information:	<table border="1"> <tr> <td>Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_161904 (NRP 68))</td> <td>Prof. Ted Turlings</td> </tr> <tr> <td>Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_143141)</td> <td>Dr. Monika Maurhofer</td> </tr> </table>	Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_161904 (NRP 68))	Prof. Ted Turlings	Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_143141)	Dr. Monika Maurhofer
Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_161904 (NRP 68))	Prof. Ted Turlings				
Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (406840_143141)	Dr. Monika Maurhofer				
Abstract:	When larvae of rootworms feed on maize roots they induce the emission of the sesquiterpene (E)- β -caryophyllene (E β C). E β C is attractive to entomopathogenic nematodes, which parasitize and rapidly kill the larvae, thereby protecting the roots from further damage. Certain root-colonizing bacteria of the genus <i>Pseudomonas</i> also benefit plants by promoting growth, suppressing pathogens or inducing systemic resistance (ISR), and some strains also have insecticidal activity. It remains unknown how these bacteria influence the emissions of root volatiles. In this study, we evaluated how colonization by the growth-promoting and insecticidal bacteria <i>Pseudomonas protegens</i> CHA0 and <i>Pseudomonas chlororaphis</i> PCL1391 affects the production of E β C upon feeding by larvae of the banded cucumber beetle, <i>Diabrotica balteata</i> Le Conte (Coleoptera: Chrysomelidae). Using a combination of chemical analysis and				

gene expression measurements, we found that E β C emission and the expression of the E β C synthase gene (TPS23) was enhanced in Pseudomonas-colonized roots after 72 hours of *D. balteata* feeding. Undamaged roots colonized by Pseudomonas spp. showed no measurable increase in E β C production, but a slight increase in TPS23 expression. Pseudomonas colonization did not affect root biomass, but larvae that fed on roots colonized by *P. protegens* CHA0 tended to gain more weight than larvae that fed on roots colonized by *P. chlororaphis* PCL1391. Larvae mortality on Pseudomonas spp. colonized roots was slightly, but not significantly higher. The observed enhanced production of E β C upon Pseudomonas spp. colonization may enhance the protective role of entomopathogenic nematodes and other soil beneficial organisms.

[Click here to view linked References](#)

1 **Root-colonizing bacteria enhance the levels of (*E*)- β -caryophyllene**
2 **produced by maize roots in response to rootworm feeding**

3

4 Xavier Chiriboga M.¹, Huijuan Guo^{1,2}, Raquel Campos-Herrera^{1,3}, Gregory Röder¹,
5 Nicola Imperiali⁴, Christoph Keel⁴, Monika Maurhofer⁵, Ted C.J. Turlings¹

6 ¹ *Fundamental and Applied Research in Chemical Ecology Laboratory, University of*
7 *Neuchâtel, Emile-Argand 11, Neuchâtel CH 2000 (Switzerland)*

8 ² *State Key Laboratory of Integrated Management of Insect Pests and Rodents, Institute*
9 *of Zoology, Chinese Academy of Sciences, Beijing 100101 (China)*

10 ³ *Current address: Centro para os Recursos Biológicos e Alimentos Mediterrânicos*
11 *(MeditBio), Campus Gambelas, Edif. 8, FCT, Universidade do Algarve, 8005-139, Faro*
12 *(Portugal)*

13 ⁴ *Department of Fundamental Microbiology, University of Lausanne (Switzerland)*

14 ⁵ *Plant Pathology, Institute of Integrative Biology, Swiss Federal Institute of*
15 *Technology, Zurich (Switzerland)*

16 Authors:

17 Xavier Chiriboga M. email address: xavier.chiriboga@unine.ch

18 Huijuan Guo email address: guohj@ioz.ac.cn

19 Raquel Campos-Herrera email address: rcherrera@ualg.pt

20 Gregory Röder email address: gregory.roeder@unine.ch

21 Nicola Imperiali email address: nicola.imperiali@unil.ch

22 Christoph Keel email address: christoph.keel@unil.ch

23 Monika Maurhofer email address: monika.maurhofer@usys.eth.ch

24 Corresponding author:

25 Ted C. J. Turlings email address: ted.turlings@unine.ch

26 Fundamental and Applied Research in Chemical Ecology (FARCE lab)
27 Institute of Biology, University of Neuchâtel, Emile-Argand 11, Neuchâtel CH-2000,
28 Switzerland
29 Phone: 0041327183000; Fax: 0041327183001

30

31

32 **Author contribution statement**

33 XCM, HG, TCJT and RC-H conceived the experiments, XCM and RC-H analyzed the
34 data and wrote the paper, NI and GR provide technical assistance for microbiology
35 techniques and GC-MS analyses and, respectively. CK, MM and TCJT edited the text
36 and approve the paper for publication.

37

38 **Abstract**

39 When larvae of rootworms feed on maize roots they induce the emission of the
40 sesquiterpene (*E*)- β -caryophyllene (*E* β C). *E* β C is attractive to entomopathogenic
41 nematodes, which parasitize and rapidly kill the larvae, thereby protecting the roots
42 from further damage. Certain root-colonizing bacteria of the genus *Pseudomonas* also
43 benefit plants by promoting growth, suppressing pathogens or inducing systemic
44 resistance (ISR), and some strains also have insecticidal activity. It remains unknown
45 how these bacteria influence the emissions of root volatiles. In this study, we evaluated
46 how colonization by the growth-promoting and insecticidal bacteria *Pseudomonas*
47 *protegens* CHA0 and *Pseudomonas chlororaphis* PCL1391 affects the production of
48 *E* β C upon feeding by larvae of the banded cucumber beetle, *Diabrotica balteata* Le
49 Conte (Coleoptera: Chrysomelidae). Using a combination of chemical analysis and gene
50 expression measurements, we found that *E* β C emission and the expression of the *E* β C
51 synthase gene (TPS23) was enhanced in *Pseudomonas*-colonized roots after 72 hours of
52 *D. balteata* feeding. Undamaged roots colonized by *Pseudomonas* spp. showed no
53 measurable increase in *E* β C production, but a slight increase in TPS23 expression.
54 *Pseudomonas* colonization did not affect root biomass, but larvae that fed on roots
55 colonized by *P. protegens* CHA0 tended to gain more weight than larvae that fed on
56 roots colonized by *P. chlororaphis* PCL1391. Larvae mortality on *Pseudomonas* spp.
57 colonized roots was slightly, but not significantly higher. The observed enhanced
58 production of *E* β C upon *Pseudomonas* spp. colonization may enhance the protective
59 role of entomopathogenic nematodes and other soil beneficial organisms.

60

61 **Key words:** Root-colonizing bacteria, *Diabrotica balteata*, (*E*)- β -caryophyllene,
62 terpene synthase, maize

63

64 **Introduction**

65 During the past decade it has been found that insect-damaged roots emit volatile
66 compounds that may serve as attractants for the natural enemies of the damaging insects
67 (Rasmann et al. 2005; Ali et al. 2010; Tonelli et al. 2016). The first such attractant was
68 identified for maize roots, which respond to feeding by larvae of the beetle *Diabrotica*
69 *virgifera virgifera* Le Conte (Coleoptera: Chrysomelidae) with the release of the
70 sesquiterpene (*E*)- β -caryophyllene (*E* β C). This herbivore-induced volatile (HIPV)
71 attracts entomopathogenic nematodes (EPNs) and, thereby, helps to protect maize roots
72 against herbivore damage (Rasmann et al. 2005; Degenhardt et al. 2009). Although
73 similar root-produced EPN attractants have been identified for several other plants (Boff
74 et al. 2001; Ali et al. 2011), it is still poorly understood how other soil organisms affect
75 the production or may respond to these signals.

76 Besides root herbivores, numerous other organisms that live in the rhizosphere
77 may form associations with a plant. Their effects may be beneficial (e.g. mycorrhizal
78 fungi, N-fixing bacteria) or detrimental (e.g. pathogenic fungi or bacteria) to plant
79 performance (Brussaard 1998; Rasmann and Turlings, 2016). There is increasing
80 interest in some strains of root-associated bacteria of the genus *Pseudomonas* that have
81 plant-beneficial properties. They can promote plant growth, suppress pathogens and/or
82 induce systemic plant defenses (Kupferschmied et al. 2013; Lugtenberg and Kamilova
83 2009; Van Oosten et al. 2008). Recent studies have also revealed that specific
84 *Pseudomonas* strains possess insecticidal activity against several insect herbivore
85 species (Ruffner et al. 2013). It has become increasingly evident that natural isolates of
86 *Pseudomonas fluorescens* and *Pseudomonas chlororaphis* (γ -Proteobacteria:
87 Pseudomonaceae) have a high potential to be applied as plant protection products
88 against various insect pests (Kupferschmied et al. 2013). Since many strains of the *P.*

89 *fluorescens* group are adapted to live on plant roots, show environmental persistence
90 and are competitive and strong root colonizers, they may be ideal not only to enhance
91 plant growth, but also to control insects pests (Lugtenberg & Kamilova 2009;
92 Kupferschmied et al. 2013). The current study is part of an interdisciplinary effort to
93 explore potential synergies in applying combinations of plant beneficial soil organisms
94 (<http://www.nrp68.ch/en>).

95 Studies measuring the effects of root-associated bacteria on volatiles organic
96 compounds have been largely limited to aboveground volatiles (Ballhorn et al. 2013;
97 Pineda et al., 2013; Pangesti et al., 2015a) and the reported effects are greatly
98 contrasting. Pineda et al. (2013) and Pangesti et al. (2015a) both used the bacterium *P.*
99 *fluorescens* WCS417r to colonize *Arabidopsis thaliana* roots, but they employed
100 aboveground herbivores of different feeding guilds to induce the leaves. It was found
101 that *Myzus persicae* (Homoptera: Aphididae), a phloem feeder, induced increased levels
102 of volatiles in colonized plants (Pineda et al., 2013), whereas colonized-plants that were
103 damaged by leaf chewing caterpillars of *Mamestra brassicae* (Lepidoptera: Noctuidae)
104 had reduced levels of HIPVs (Pangesti et al., 2015a). These differences can be
105 explained by the different hormonal pathways that are activated by different plant
106 antagonists. Chewing insects and necrotrophic pathogens typically induced the jasmonic
107 acid pathway, whereas phloem-feeding insects and biotrophic pathogens usually
108 upregulate the salicylic acid pathway (Zarate et al. 2006; Thaler et al., 2012; Jacobs et
109 al. 2011; Pieterse et al. 2012). Thus, crosstalk between the two pathways may result in
110 their mutual suppression (Zhang et al., 2009; Thaler et al., 2012). This is also a possible
111 explanation for the results found by Ballhorn et al. (2013), who compared volatile
112 emissions by rhizobia-colonized lime bean plants after experimental induction with
113 jasmonic acid. Colonized plants produced higher amounts of shikimic acid-derived

114 compounds than non-colonized plants, whereas the emission of compounds produced
115 via the octadecanoid, mevalonate and non-mevalonate pathways was reduced.

116 We are aware of only one study that looked at the effects of root-colonizing
117 bacteria on root-produced HIPVs. Santos et al. (2014) found that maize root
118 colonization by *Azospirillum brasilense* (α -Proteobacteria: Rhodospirillaceae) produced
119 higher amounts of $E\beta C$ compared to non-colonized maize roots, in this case without
120 insect damage. They further found that larvae of the generalist root feeder *Diabrotica*
121 *speciosa* (Coleoptera: Chrysomelidae) oriented preferentially towards non-inoculated
122 maize roots *versus* inoculated roots and gained less weight when feeding on inoculated
123 roots. Interestingly, larvae of the maize specialist *D. virgifera virgifera*, which were
124 initially studied in the context of inducible $E\beta C$ (Rasmann et al., 2005), are attracted to
125 $E\beta C$ and perform better on already infested root systems (Robert et al., 2012a).

126 It remains unknown how root-associated bacteria affect the induction of
127 belowground volatiles in response to root herbivory. This prompted the current study in
128 which we studied these effects in maize roots damaged by larvae of another generalist
129 *Diabrotica* beetle, the banded cucumber beetle *Diabrotica balteata* Le Conte
130 (Coleoptera: Chrysomelidae). *D. balteata* larvae induce lesser amounts of $E\beta C$ in maize
131 roots than *D. virgifera* larvae, but this still results in some attraction of EPN (Rasmann
132 and Turlings 2008). *D. balteata* is an important agricultural pest in Central and North
133 America (Capinera 2011), attacking a broad spectrum of crops, including cucumber,
134 squash, beet, bean, soybean, pea, sweet potato, okra, maize, lettuce, onion, and various
135 cabbages (Saba, 1970; Chittenden, 1992; Capinera, 2011). It may damage all parts of a
136 plant, but the most serious injury caused by *D. balteata* is to the roots (Capinera 2011).
137 Enhancing $E\beta C$ emissions in maize roots damaged by *D. balteata* might render EPN
138 more effective in finding and killing the larvae of this important generalist root pest.

139 This pest is therefore a good model to test the possible effects of plant-beneficial root
140 colonizing bacteria on *EβC* emissions.

141 In the present study, we used a chemical as well as a molecular approach to
142 evaluate the effects of maize root colonization by the bacteria *P. chlororaphis* PCL1391
143 and *P. protegens* CHA0 on the emission of (*E*)- β -caryophyllene. Roots were inoculated
144 (or not) by one of the bacteria and infested or not by *D. balteata* larvae. We then
145 collected and analyzed the volatiles emissions from the roots and we measured the
146 expression of the maize *EβC* synthase gene (TPS23) (Köllner et al. 2008).

147 The species *P. protegens* CHA0 is a root-associated bacterium that not only
148 produces antifungal metabolites, but also an insecticidal protein. This protein is very
149 similar to the potent insect toxin Mcf1 of the entomopathogen *Photorhabdus*
150 *luminescens* (γ -Proteobacteria: Enterobacteriaceae) (Péchy-Tarr et al., 2008). *P.*
151 *protegens* CHA0 causes insect toxicity in experimental infections of aboveground
152 feeding insect larvae (Péchy-Tarr et al., 2008) and also in feeding assays with artificial
153 diets or leaves treated with the bacteria (Ruffner et al., 2013). It is unknown how these
154 root-associated bacteria affect root feeding insect larvae. We therefore also studied the
155 effect of the bacteria on the performance and mortality of *D. balteata* larvae.

156 Hence, we studied if colonization by *P. protegens* CHA0 or *P. chlororaphis*
157 PCL1391: i) induces a change in the production of *EβC* after *D. balteata* attack in maize
158 roots, ii) changes the expression of the maize *EβC* synthase gene TPS23, iii) affects root
159 growth in maize plants, and iv) affects the performance and mortality of *D. balteata*
160 larvae. We discuss our results in terms of the physiological changes that may occur in
161 plants upon *Pseudomonas* colonization and how these changes may influence HIPVs.
162 We further address the possibility of applying the bacteria in combination with EPNs

163 for the effective control of diabroticine beetle larvae in maize and other cropping
164 systems.

165

166 **Materials and methods**

167

168 **Soil, plants and insect larvae**

169

170 A substrate containing potting soil (Terreau semis Capito, Landi-Switzerland,
171 pH = 5.8-6.8) and white sand (Migros, Switzerland) in proportion 1:1 was used to grow
172 the plants. The substrate was autoclaved twice at 120 °C for 120 min. Plastic pots (11
173 cm, height x 4 cm, diameter) were autoclaved once at 120 °C for 120 min before each
174 sowing.

175 Maize seeds (var. Delprim and var. F268) were surface sterilized by washing
176 them with ethanol 70% for 2 min and sodium hypochlorite 3% for 2 minutes and rinsing
177 them with sterile water. Plants were watered with 20 mL of sterile distilled water every
178 2-3 days. Plants were grown either in a greenhouse (30±5 °C, 8:16 h dark:light
179 photoperiod) in summer or in a phytotron (30±2 °C, 8:16 h dark:light photoperiod, 300
180 $\mu\text{mol m}^{-2} \text{s}^{-1}$, CLF Plant Climatics, Germany) in winter.

181 Second instar larvae of *D. balteata* were reared from eggs provided by Syngenta
182 (Stein, Switzerland) and they were fed with maize germinate. Larvae were used to infest
183 11 days old maize plants (after a period of 6 days of roots colonization by bacteria), by
184 burying them in small holes in the soil. Each plant was infested with six *D. balteata*
185 larvae.

186

187

188 **Bacteria cultures and inoculation**

189

190 The bacteria *P. protegens* CHA0 and *P. chlororaphis* PCL1391 (Department of
191 Fundamental Microbiology, University of Lausanne) were cultured in LB agar (Miller,
192 Sigma-Aldrich) supplemented with 100 µg/mL of rifampicin ($\geq 97\%$ powder, Sigma-
193 Aldrich) for 48 hours in 9 cm diam. Petri dishes at 30 °C. Bacteria were scratched from
194 the plates under sterile conditions and transferred to 100 mL of sterile rifampicin
195 supplemented-LB broth. Both species were cultivated independently in an orbital
196 agitator (IKA-KS 4000) at 30 °C and 190 rpm for 16 hours. Bacterial cultures were then
197 centrifuged at 6846 x g for 10 minutes to separate bacterial cells from the liquid culture
198 media. Resulting bacterial cell pellets were diluted again in sterile distilled water.
199 Standard bacteria concentrations (1×10^6 CFU ml⁻¹) were obtained, calibrating the
200 inoculum with a spectrophotometer at an optical density of 0.2A at 600 nm.

201 After 4-5 days of sowing, at the shoot emergence stage, plants were selected for
202 the application of different treatments: a) inoculated with *P. protegens* CHA0, and
203 infested with *D. balteata* (CHA0+Db), b) inoculated with *P. chlororaphis* PCL1391,
204 and infested with *D. balteata* (PCL+Db), c) not inoculated with bacteria, infested with
205 *D. balteata* (Db), d) control healthy plants (Healthy), e) only inoculated with *P.*
206 *protegens* CHA0 (CHA0), and f) only inoculated with *P. chlororaphis* PCL1391 (PCL).
207 Plants treated with root-colonizing bacteria were inoculated with 20 mL of *P. protegens*
208 CHA0 or *P. chlororaphis* PCL1391 inoculum prepared as described above. Plants
209 infested only with *D. balteata* and control-healthy were watered with 20 mL of sterile
210 water. Preliminary experiments were performed before, measuring production of EβC
211 after 72 hours of insect feeding, with six replicates per treatment (n = 6). Nine replicates
212 (n = 9) per treatment were done in a final time-course experiment. Plants of different

213 treatments were kept separated in different plastic trays to avoid cross-contamination
214 and kept either in a greenhouse or a phytotron for 6 days during the root colonization
215 period.

216 Colonization of maize roots with *P. protegens* CHA0 or *P. chlororaphis*
217 PCL1391 was verified for a subset of plants of the same batch used for the volatiles and
218 gene expression analysis. For this, roots of inoculated plants were harvested and the soil
219 was gently removed and roots were weighed. Then the roots were suspended in flasks
220 with 40 mL of sterile water and the flasks were shaken vigorously for 10 minutes to
221 wash off the bacteria from the roots. Serial dilutions of the washed roots were prepared
222 and plated on rifampicin-LB agar Petri dishes. Plates were incubated at 30 °C and after
223 24 h the numbers of colony-forming units (CFU) were counted and CFU per gram of
224 root calculated.

225

226 **Volatile extraction and analyses**

227

228 In preliminary experiments, we analyzed volatiles produced by the whole root
229 system after 72 hours of *D. balteata* infestation, whereas in the final time-course
230 experiment, we standardized the amount of ground root sample per vial for volatile
231 analysis. We quantified the amount of *EβC* produced by roots of maize plants var.
232 Delprim after 6 and 72 hours of insect infestation.

233 Roots were harvested and washed gently with tap water 6 and 72 hours after
234 insect infestation and immediately frozen in liquid nitrogen for grinding. Roots were
235 ground in a frozen mortar with liquid nitrogen. Root volatiles were extracted following
236 the standard procedure by Rasmann (2005): 500 mg of ground root material were
237 weighed and transferred to 10-mL glass vials sealed with a Teflon-coated septum and

238 stored at -80 °C for analysis. A 100 µm polydimethylsiloxane SPME fiber (Supelco,
239 Sigma-Aldrich Chemie SA, Buchs, Switzerland) was inserted through the septum and
240 exposed in the headspace for 60 min at 40 °C. The compounds adsorbed onto the fiber
241 were analyzed with an Agilent 7890a Series GC system coupled to mass-selective
242 detector (Agilent 5975c, transfer line 280 °C, source 230 °C, quadrupole 150 °C,
243 ionization potential 70 eV) (Palo Alto CA, USA). The fiber was inserted into the
244 injector port (250 °C), desorbed and the volatile compounds were separated on a non-
245 polar column (HP1-MS; 30 m, 0.25 mm internal diameter, 0.25 mm film thickness; J &
246 W Scientific, Agilent Technologies SA, Basel, Switzerland). Helium at a constant flow
247 mode of 0.9 mL min⁻¹ (127.9 kPa) was used as a carrier gas. After fiber insertion, the
248 column temperature was maintained at 50 °C for 3 min, then increased to 180 °C at 5 °C
249 min⁻¹, before a final ramp at 8 °C min⁻¹ to reach 250 °C (hold 3 min). Chromatograms
250 processing were carried out with ChemStation software (Agilent Technologies SA,
251 Basel, Switzerland). Relative abundance of the root volatiles was calculated by
252 integrating peaks and values were corrected for sample weight to calculate relative
253 abundance of the volatile per gram of root.

254

255 **cDNA synthesis and gene expression analysis**

256

257 Approximately 60 mg of ground root material was used for the analysis of Zm-
258 TPS23 gene expression. RNA from roots was extracted using the Isolate II RNA Plant
259 Kit (Bioline, Germany), and RNA concentration was determined using a Nanodrop
260 (Control Program ND-1000 v.3.3.0., ThermoScientific, Wilmington, DE). cDNA was
261 synthesized using Sunscript RT RNase H+ (Bioline, Germany). Real-time qPCR was
262 performed in 100-well gene discs reaction plates (Biolabo, Scientific Instruments,

263 Switzerland) in the Corbett Research real-time qPCR using Zm-TPS23 specific primers
264 (F: GTGGGCCTCTACCTATCCA, R: CTGTGGTGGTGCCGTATTT) and Zm-actin
265 specific primers (F: CAGTGGTCGAACAACGGGTA, R:
266 GGTAAGGTCACGACCAGCAA) as a reference gene (Köllner et al. 2008). The qPCR
267 mix was adjusted to a final volume of 10 µL, using RNA-free water, specific primers
268 (either for TPS23 or for actin detection) both forward and reverse (0.05 µM) and SYBR
269 Green (Bioline, Germany) and 1 µL of DNA template. Negative control contained free
270 RNAase water instead of DNA template, to verify there is not contamination in the
271 reactions. A qPCR analysis was carried out using the following thermal cycling
272 conditions: a hold at 95 °C for 10 min and 40 cycles, at 95 °C for 10 s and at 60 °C for
273 45 s. Relative expressions of the genes TPS23 and actin for different treatments were
274 obtained using the correction method $2^{-\Delta\Delta C_t}$ (Livak and Schmittgen 2001).

275

276 **Assessment of larvae weight gain and mortality**

277

278 For this evaluation, we used the same set of plants that we used for volatile
279 extraction in the time-course experiment. We weighed *D. balteata* larvae (Mettler
280 Toledo MX5 microbalance) before placing them on the plants and we recorded weight
281 gain of the larvae after 6 hours, 48 hours and 72 hours of feeding. We also recorded the
282 number of dead larvae per treated plant.

283

284 **Statistical analysis**

285

286 Relative abundance of volatiles per gram of root values ($E\beta C$) were normalized
287 prior statistical analysis by log transformation. We employed a Linear mixed-effects

288 model, each time-point was analyzed separately. Relative expression of terpene
289 synthase gene data was analyzed with a Generalized linear model with a quasi-Poisson
290 distribution. Tukey method was used to compare Least square means in both cases and
291 T-test was used to compare differences between time-points. Root growth data was
292 analyzed with One-way ANOVA. Larvae weight gain data were analyzed with Two-
293 Way ANOVA. Mortality data were arcsin transformed and analyzed with Two-way
294 ANOVA, differences between means were obtained with the Tukey method in all cases.
295 All data were analyzed using R 3.3.2. (2016). Data is presented as mean \pm SEM of
296 untransformed values.

297

298 **Results**

299 **Maize root colonization by *Pseudomonas* spp. and production of (E)- β -** 300 **caryophyllene after *Diabrotica balteata* damage**

301

302 The root colonization by *Pseudomonas* spp. was similar for all bacterial
303 treatments (ANOVA, $F_{3,4} = 1.4$, $P > 0.1$) (Table 1). Our preliminary experiments, in
304 which we analyzed the roots from two maize genotypes (var. Delprim and inbred line
305 F268), showed a trend of higher production of $E\beta C$ in response to *D. balteata* feeding
306 on *Pseudomonas*-colonized roots as compared to non-colonized roots (72 h post-attack)
307 (Supplementary Fig.1). However, variability within the treatments was high and no
308 significant differences were detected.

309 The subsequent experiments showed that the production of $E\beta C$ in maize roots
310 was affected by treatment after 6 hours ($F_{5,40} = 9.12$, $P < 0.001$) and 72 hours ($F_{5,7} =$
311 10.9 , $P < 0.01$) of insect feeding (Fig. 1; Supplementary Table 1). After 6 hours, non-
312 inoculated roots attacked by the insects produced significantly larger amounts of $E\beta C$

313 than control healthy roots ($P < 0.01$). There was a marginal difference in $E\beta C$ quantities
314 between insect-damaged roots colonized by any of the bacteria species and control
315 healthy roots. However, there was no difference between insect-damaged roots
316 colonized by any of the bacteria species and non-colonized roots attacked by the insect
317 ($P > 0.1$) (Fig.1).

318 Seventy two hours after *D. balteata* attack, roots colonized by *P. protegens*
319 CHA0 produced significantly larger amounts of $E\beta C$ ($P < 0.05$) than non-colonized
320 roots attacked by the insects whereas roots colonized by *P. chlororaphis* PCL produced
321 similar ($P > 0.1$) amounts of $E\beta C$ than non-colonized roots attacked by *D. balteata*.
322 Control healthy roots produced the same amounts of $E\beta C$ ($P > 0.1$) as undamaged roots
323 colonized by either bacterium (Fig. 1). We found a significant higher production of $E\beta C$
324 ($P < 0.05$) after 72 hours than after 6 hours of insect damaged in roots colonized by *P.*
325 *protegens* CHA0. For the other treatments, there were no differences between the two
326 time points, neither for insect-damaged plants colonized by *P. chlororaphis* PCL1391
327 ($P > 0.1$).

328

329 **Expression of the terpene synthase-TPS23 after *Diabrotica balteata* damage in** 330 **maize roots colonized by *Pseudomonas protegens* CHA0 and *Pseudomonas*** 331 ***chlororaphis* PCL1391**

332 The treatments also affected the expression of TPS23 (after 6 hours: $F_{5,37} = 3.27$,
333 $P < 0.05$; after 72 hours: $F_{5,28} = 18.32$, $P < 0.001$). After 6 hours of insect feeding, the
334 expression of the gene was significantly higher in roots colonized by *P. chlororaphis*
335 PCL1391 and attacked by *D. balteata* ($P < 0.05$), and in non-colonized roots attacked
336 by the insect ($P < 0.05$), as compared to healthy control roots (Fig. 2; Supplementary
337 Table 2).

338 After 72 hours of *D. balteata* attack, gene expression in insect-damaged roots
339 colonized by *P. protegens* CHA0 ($P < 0.01$) and *P. chlororaphis* PCL1391 ($P < 0.05$)
340 was significantly higher than in insect-damaged non-colonized roots. The expression in
341 the latter roots was not different from the expression in control healthy roots ($P = 0.1$),
342 nor from the expression in undamaged roots colonized by either one of the bacteria
343 species ($P > 0.1$) (Fig. 2; Supplementary Table 2). As found for the release of $E\beta C$ (Fig.
344 1), TPS23 expression was significantly higher ($P < 0.01$) after 72 hours of insect attack
345 than after 6 hours in insect-damaged roots colonized by *P. protegens* CHA0 and *P.*
346 *chlororaphis* PCL. In all of the other treatments, the expression was not statistically
347 different ($P > 0.01$) between the two time-points.

348

349 **Root colonization does not change roots biomass**

350 We did not find an effect of any of the treatments on root fresh weight ($P =$
351 0.09), measured after the 72 hours of *D. balteata* feeding (Fig. 3A). However, there was
352 a trend that biomass of insect-damaged roots was higher for plants colonized by *P.*
353 *chlororaphis* PCL as compared to the insect-damaged roots grown in presence of *P.*
354 *protegens* CHA0 or in absence of bacterial inoculants.

355

356 **Effects of bacterial colonization on the weight gain and mortality of *Diabrotica*** 357 ***balteata* larvae**

358 Overall, there was no effect of the treatments on larval weight gain ($F_{2,72} = 1.72,$
359 $P = 0.18$), but there was a trend of better weight gain when larvae were feeding on *P.*
360 *protegens* CHA0 colonized roots than when feeding on *P. chlororaphis* PCL-colonized
361 roots (Fig. 3b), and this correlates with differences in root biomass (Fig. 3a and
362 Supplementary Fig. 2). We measured an overall increase in weight over time ($F_{2,72} =$

363 8.59, $P < 0.001$) (Fig. 3b), but no significant interaction between time and treatment
364 ($F_{4,72} = 0.72$, $P = 0.57$). Within each treatment, weight over time varied only significant
365 for larvae that had fed on roots colonized by *P. protegens* CHA0.

366 In a preliminary experiment with maize plants var. F268, we found a similar
367 pattern of weight gain for *D. balteata* feeding on roots colonized by *P. protegens*
368 CHA0, *P. chlororaphis* PCL1391 and non-colonized roots (Supplementary Fig. 3). In
369 this experiment, we detected a significant effect of time ($F_{4,123} = 10.85$, $P < 0.01$), but
370 no obvious effect of the treatment ($F_{2,123} = 1.11$, $P > 0.1$), nor an interaction between
371 time and treatment ($F_{5,123} = 0.26$, $P > 0.1$).

372 For the main experiment, we also found an effect of time on the mortality of *D.*
373 *balteata* larvae ($F_{2,72} = 21.76$, $P < 0.001$), but no effect of the treatment ($F_{2,72} = 2.03$, $P >$
374 0.1), nor an interaction between time and treatment ($F_{4,72} = 0.98$, $P > 0.1$) (Fig. 3C).

375

376 **Discussion**

377

378 We found quantitative but no qualitative differences in the volatile profiles for
379 the different treatments. Maize roots colonized by *P. protegens* CHA0 and *P.*
380 *chlororaphis* PCL1391 bacteria without insect infestation produced only minor
381 quantities of the root volatile $E\beta C$ (Fig.1 and Supplementary Fig.1.), but colonization by
382 *P. protegens* CHA0 significantly enhanced the production of the sesquiterpene in maize
383 after 72 hours of *D. balteata* feeding. To our knowledge, ours is the first study that
384 evaluates how root-associated bacteria affect the emissions of a belowground HIPV
385 upon root herbivory. Yet, Santos et al. (2014), using the same maize variety (Delprim),
386 showed that the plant-beneficial bacterium *Azospirillum brasilense* affects $E\beta C$

387 emissions in plants without insect damage. They found that colonized roots released
388 more *EβC* and repelled larvae of *Diabrotica speciosa*.

389 Other studies on how root-associated bacteria affect volatile emissions have
390 focused on volatiles released from aboveground plant parts, and show contrasting
391 results. Root colonization by pseudomonads can decrease (Pangesti et al. 2015a) or
392 increase (Pineda et al., 2013) aboveground HIPVs. *Arabidopsis thaliana* plants
393 colonized by *Pseudomonas fluorescens* WCS417r and subsequently attacked by
394 *Mamestra brassicae* caterpillars, produced lower amounts of methyl salicylate, linal and
395 the terpene (*E*)- α -bergamotene in comparison with non-colonized plants infested with
396 caterpillars (Pangesti et al. 2015a). In contrast, Pineda et al. (2013) showed with the
397 same plant-bacteria system, but using the aphid *Myzus persicae* as herbivore, that the
398 aphid-induced production of eight leaf volatiles (2-nonenal, isovaleric acid, dimethyl
399 sulfoxide, 2-cyclopente-1-one, (*R*)-verbenone, (*E*)-2-heptanal, 1-pentanol and 5,5
400 dimethyl-2(5H)-furanone) was enhanced in soil bacteria-colonized plants compared
401 with non-colonized plants. Some other volatiles were produced in high quantities in
402 plants colonized by *P. fluorescens* even without insect damage in the same study.
403 Hence, effects of root colonizing bacteria on inducible volatiles appear to vary strongly,
404 depending on the plants species, root-associated bacteria and on the insect herbivores.

405 Our findings on *EβC* emissions correlate nicely with the results for the
406 expression of the terpene synthase gene Zm-TPS23. In roots colonized by *P. protegens*
407 CHA0 and *P. chlororaphis* PCL1391, the expression was enhanced after 72 hours of *D.*
408 *balteata* infestation in comparison with non-colonized roots attacked by the insect (Fig.
409 2). Interestingly, we also found a higher expression of the gene TPS23 in undamaged
410 roots colonized by *P. chlororaphis* PCL1391 than in control healthy roots at the second
411 time-point (after 72 hours). This is again different from Pangesti et al. (2015a), who

412 reported a negative effect of *P. fluorescens* colonization on the expression of the terpene
413 synthases TPS03 and TPS04 in *Arabidopsis* upon insect leaf herbivory. These
414 contrasting results confirm, as mentioned above, that the effects of root-associated
415 bacteria on volatile emissions may vary depending on the system under study.

416 Inducible plant defenses, including volatile emissions, are mediated by wound-
417 induced jasmonic acid (JA), which is derived from the lipoxygenase (*LOX*) pathway
418 (Turner et al. 2002; Schmelz et al. 2003; Maffei et al. 2011; Dudareva et al, 2013).
419 Previous studies found that *Pseudomonas* colonization of *A. thaliana* plants promotes
420 the expression of the gene *LOX2* (Pineda et al. 2012) and JA-responsive genes (Oosten
421 et al. 2008), and results in stronger JA-signaling (Pangesti et al., 2015b) after insect
422 attack. We also know that the gene *Zm-TPS23* is locally and systemically induced in
423 maize roots in response to feeding by *D. virgifera*. This appears to be triggered by local
424 induction of jasmonic acid (JA) and its isoleucine conjugate (JA-Ile) after 30 minutes,
425 resulting in an exponentially increasing production of *EβC* over 48 hours of feeding
426 (Erb 2009; Hiltpold et al. 2011). Taking all together, we can hypothesize that
427 belowground enhanced production of *EβC* in maize roots colonized by *P. protegens*
428 *CHA0* and *P. chlororaphis* PCL1391 might be mediated by increased JA-signaling.

429 Pangesti et al. (2015b) point out that differences in soil composition may explain
430 some of the variable outcomes of plant-mediated effects of root-associated microbes on
431 volatile signals and insect performance. It remains to be investigated if the effects of *P.*
432 *protegens* *CHA0* and *P. chlororaphis* PCL-1391 on the enhanced production of the root
433 sesquiterpene *EβC* are consistent in different types of soils. We previously showed the
434 importance of studying the dynamics of *EβC* production and diffusion under different
435 soil conditions (Chiriboga M. et al. 2017).

436 It has also been proposed that the effect of root-associated microbes on insect
437 herbivores is different for specialist and generalist herbivores and for insects with
438 different modes of feeding. Pineda et al. (2010) expect a negative effect on generalist
439 chewing insects and mesophyll feeders, and positive or neutral on specialist chewing
440 insects and phloem feeders. The effects on herbivore performance are directly related to
441 the activation of defensive responses in the plant, including the production of HIPVs. It
442 is pertinent to investigate what additional volatiles are produced upon root-colonization
443 by bacteria, also by the bacteria themselves (D'Alessandro et al., 2014), and how these
444 affect the interactions with other soil organisms.

445 We did not measure a clear effect of any treatment on root biomass (Fig. 3A),
446 but there was a trend of lower biomass for insect-damaged roots that were colonized by
447 *P. protegens* CHA0 compared to insect-damaged roots colonized by *P. chlororaphis*
448 PCL (Fig. 3A). The poorer performance of the larvae on PLC-colonized plants may
449 have contributed to this trend (Fig. 3B and Supplementary Fig 2.). Indeed, *D. balteata*
450 larvae feeding on maize roots colonized by *P. protegens* CHA0 tended to gain more
451 weight than larvae feeding in roots colonized by *P. chlororaphis* PCL1391 after 72
452 hours of feeding. Possibly, the increased emissions of $E\beta C$ in roots colonized by *P.*
453 *protegens* CHA0 stimulated feeding and/or benefitted *D. balteata* weight gain. This has
454 been shown for larvae of the maize specialist *D. virgifera*, which are attracted to $E\beta C$
455 (Robert et al. 2012a) and perform better on already infested roots (Robert et al. 2012b).
456 In contrast, larvae of the generalist *D. speciosa* larvae gained less weight on and are less
457 attracted to roots that produce increased amounts of $E\beta C$ (Santos et al. 2014).

458 It is further possible that the differences in weight gain on roots with different
459 treatments were due to differences in nutritional quality and/or biomass of the roots.
460 Mutualistic microorganisms are known to influence plant tolerance to herbivory

461 (Strauss and Agrawal 1999). *Diabrotica* feeding also triggers tolerance responses,
462 including regrowth of roots and resource reallocation in maize (Erb, 2009) and it would
463 be worthwhile to determine if PCL1391-colonization has an effect on these responses.

464 There were no significant differences in mortality among different treatments
465 (Fig. 3C), but there was a trend for higher mortality in larvae feeding 72 h on *P.*
466 *chlororaphis* PCL-treated plants. If we had let the larvae feed longer this might have
467 resulted in clearer effects, as pathogenicity of *Pseudomonas* bacteria can be rather a
468 long process that involves several steps: bacteria ingestion, release of the toxin, toxin
469 binding, breaking of the gut wall and insect death (Kupferschmied et al. 2013, Keel
470 2016). The observed enhanced signaling ability and possible higher larval mortality on
471 *Pseudomonas*-colonized roots imply that the application of the bacteria in combination
472 with EPNs might be a highly effective strategy for the control of root herbivores in
473 maize production. This compatibility was confirmed in a field study, in which two
474 species of *Pseudomonas* in combination with the EPN *Heterorhabditis bacteriophora*
475 were found to be best in enhancing wheat plant performance (Imperiali et al., under
476 review). How the application of such combinations plays out against *Diabrotica* pest
477 under realistic field condition remains to be determined.

478

479 **Conclusions**

480

481 Colonization of maize roots by *P. protegens* CHA0 was found to enhance the
482 emission of $E\beta C$ after 72 h of feeding by *D. balteata* larvae. Consistent with this
483 enhanced emission of the EPN attractant, we found a higher expression of the terpene
484 synthase gene Zm-TPS23 after 72 h of insect infestation in colonized roots. The gene
485 expression data revealed a positive effect of both *Pseudomonas* strains. Undamaged

486 roots colonized by *P. protegens* CHA0 and *P. chlororaphis* PCL1391 also had a slightly
487 enhanced expression of the terpene synthase gene. The mechanisms that are involved in
488 this enhanced production of $E\beta C$ are still unclear. The same is true for the observed
489 differences in larval growth and mortality on roots of the different treatments. Yet, it is
490 evident from this study that the application of beneficial *Pseudomonad* bacteria and
491 EPN is compatible and may be a highly complementary strategy for the control of soil
492 pests and to enhance crop performance.

493

494 **Acknowledgements**

495 We thank Jean-Marc Freyermuth for statistical advice, Geoffrey Jaffuel and Allan
496 Kergunteuil for fruitful discussions, and Angela Köhler and Pamela Bruno for technical
497 assistance. This study was supported by the NRP68 program “Sustainable use of soil as
498 a resource” (projects n° 406840_161904 and 406840_143141) from the Swiss National
499 Science Foundation awarded to TCJT, CK, and MM. XCM was funded by an
500 Excellence Scholarship of the Swiss Confederation, HG was supported with a Post-Doc
501 Grant from the Chinese Academy of Sciences and RC-H was supported with an
502 Investigator Program Award (IF/00552/2014) from the government of Portugal.

503

504 **REFERENCES**

505 Ali JG, Alborn HT, Stelinski LL (2010) Subterranean herbivore-induced volatiles
506 released by citrus roots upon feeding by *Diaprepes abbreviatus* recruit
507 entomopathogenic nematodes. *J Chem Ecol* 36:361–368. doi: 10.1007/s10886-
508 010-9773-7

509 Ali JG, Alborn HT, Stelinski LL (2011) Constitutive and induced subterranean plant
510 volatiles attract both entomopathogenic and plant parasitic nematodes. *J Ecol*

511 99:26–35. doi: 10.1111/j.1365-2745.2010.01758.x

512 Ballhorn DJ, Kautz S, Schädler M (2013) Induced plant defense via volatile production
513 is dependent on rhizobial symbiosis. *Oecologia* 172:833–846. doi:
514 10.1007/s00442-012-2539-x

515 Boff MIC, Van Tol RWHM, Smits PH (2002) Behavioural response of *Heterorhabditis*
516 *megidis* (strain NLH-E87.3) towards plant roots and insect larvae. *BioControl* 47,
517 67-83

518 Brooks DM, Bender CL, Kunkel BN (2005) The *Pseudomonas syringae* phytotoxin
519 coronatine promotes virulence by overcoming salicylic acid-dependent defences in
520 *Arabidopsis thaliana*. *Mol Plant Pathol* 6:629–639. doi: 10.1111/j.1364-
521 3703.2005.00311.x

522 Brussaard L (1998) Soil fauna, guilds, functional groups and ecosystem processes. *Appl*
523 *Soil Ecol* 9:123–135. doi: 10.1016/S0929-1393(98)00066-3

524 Capinera JL (2011) Banded Cucumber Beetle , *Diabrotica balteata* LeConte (*Insecta* :
525 *Coleoptera* : *Chrysomelidae*) 1. *Inst Food Agric* 93:1–3.

526 Chiriboga M. X, Campos-Herrera R, Jaffuel G, et al (2017) Diffusion of the maize root
527 signal (E)- β -caryophyllene in soils of different textures and the effects on the
528 migration of the entomopathogenic nematode *Heterorhabditis megidis*.
529 *Rhizosphere* 3:53–59. doi: 10.1016/j.rhisph.2016.12.006

530 Degenhardt J, Hiltbold I, Kollner TG, et al (2009) Restoring a maize root signal that
531 attracts insect-killing nematodes to control a major pest. *Proc Natl Acad Sci*
532 106:17606–17606. doi: 10.1073/pnas.0909073106

533 Hiltbold I, Erb M, Robert C a M, Turlings TCJ (2011) Systemic root signalling in a
534 belowground, volatile-mediated tritrophic interaction. *Plant Cell Environ* 34:1267–
535 75. doi: 10.1111/j.1365-3040.2011.02327.x

536 Jacobs S, Zechmann B, Molitor A, et al (2011) Broad-spectrum suppression of innate
537 immunity is required for colonization of *Arabidopsis* roots by the fungus
538 *Piriformospora indica*. *Plant Physiol* 156:726–740. doi: 10.1104/pp.111.176446

539 Keel C (2016) A look into the toolbox of multi-talents: insect pathogenicity
540 determinants of plant-beneficial pseudomonads. *Environ Microbiol* 18:3207–3209.
541 doi: 10.1111/1462-2920.13462

542 Köllner TG, Held M, Lenk C, et al (2008) A maize (E)- β -Caryophyllene synthase
543 implicated in indirect defense responses against herbivores is not expressed in most
544 American maize varieties. *Plant Cell Online* 20:482–494. doi:
545 10.1105/tpc.107.051672

546 Kupferschmied P, Maurhofer M, Keel C (2013) Promise for plant pest control: root-
547 associated pseudomonads with insecticidal activities. *Front Plant Sci* 4:287. doi:
548 10.3389/fpls.2013.00287

549 Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-
550 time quantitative PCR and. *Methods* 25:402–408. doi: 10.1006/meth.2001.1262

551 Lugtenberg B, Kamilova F (2009) Plant-growth-promoting rhizobacteria. *Annu Rev*
552 *Microbiol* 63:541–556. doi: 10.1146/annurev.micro.62.081307.162918

553 Pangesti N, Pineda A, Dicke M, van Loon JJA (2015a) Variation in plant-mediated
554 interactions between rhizobacteria and caterpillars: potential role of soil
555 composition. *Plant Biol* 17:474–483. doi: 10.1111/plb.12265

556 Pangesti N, Weldegergis BT, Langendorf B, et al (2015b) Rhizobacterial colonization
557 of roots modulates plant volatile emission and enhances the attraction of a
558 parasitoid wasp to host-infested plants. *Oecologia* 178:1169–1180. doi:
559 10.1007/s00442-015-3277-7

560 Pieterse CMJ, Van der Does D, Zamioudis C, et al (2012) Hormonal modulation of

561 plant immunity. *Annu Rev Cell Dev Biol* 28:489–521. doi: 10.1146/annurev-
562 cellbio-092910-154055

563 Pineda A, Soler R, Weldegergis BT, et al (2013) Non-pathogenic rhizobacteria interfere
564 with the attraction of parasitoids to aphid-induced plant volatiles via jasmonic acid
565 signalling. *Plant, Cell Environ* 36:393–404. doi: 10.1111/j.1365-
566 3040.2012.02581.x

567 Pineda A, Zheng SJ, van Loon JJA, Dicke M (2012) Rhizobacteria modify plant-aphid
568 interactions: a case of induced systemic susceptibility. *Plant Biol* 14:83–90. doi:
569 10.1111/j.1438-8677.2011.00549.x

570 Rasmann S, Köllner TG, Degenhardt J, et al (2005) Recruitment of entomopathogenic
571 nematodes by insect-damaged maize roots. *Nature* 434:732–7. doi:
572 10.1038/nature03451

573 Rasmann S, Turlings TCJ (2008) First insights into specificity of belowground
574 tritrophic interactions. *Oikos* 117:362–369. doi: 10.1111/j.2007.0030-
575 1299.16204.x

576 Ruffner B, Péchy-Tarr M, Ryffel F, et al (2013) Oral insecticidal activity of plant-
577 associated pseudomonads. *Environ Microbiol* 15:751–763. doi: 10.1111/j.1462-
578 2920.2012.02884.x

579 Saba F (1970) Host plant spectrum and temperature limitations of *Diabrotica balteata*.
580 *Can Entomol* 102:684–691. doi: 10.4039/Ent102684-6

581 Santos F, Peñaflor MFG V, Paré PW, et al (2014) A novel interaction between plant-
582 beneficial rhizobacteria and roots: Colonization induces corn resistance against the
583 root herbivore *diabrotica speciosa*. *PLoS One*. doi: 10.1371/journal.pone.0113280

584 Schmelz E a, Alborn HT, Banchio E, Tumlinson JH (2003) Quantitative relationships
585 between induced jasmonic acid levels and volatile emission in *Zea mays* during

586 *Spodoptera exigua* herbivory. *Planta* 216:665–673. doi: 10.1007/s00425-002-0898-y

587 Strauss S, Agrawal A (1999) The ecology and evolution of plant tolerance to herbivory.

588 *Trends Ecol Evol* 14:179–185. doi: 10.1016/S0169-5347(98)01576-6

589 Tonelli M, Peñaflor MFGV, Leite LG, et al (2016) Attraction of entomopathogenic

590 nematodes to sugarcane root volatiles under herbivory by a sap-sucking insect.

591 *Chemoecology* 26:59–66. doi: 10.1007/s00049-016-0207-z

592 Van Oosten VR, Bodenhausen N, Reymond P, et al (2008) Differential effectiveness of

593 microbially induced resistance against herbivorous insects in *Arabidopsis*. *Mol*

594 *Plant-Microbe Interact* 21:919–930. doi: 10.1094/MPMI-21-7-0919

595 Zarate SI, Kempema LA, Walling LL (2006) Silverleaf whitefly induces salicylic acid

596 defenses and suppresses effectual jasmonic acid defenses. *Plant Physiol* 143:866–

597 75. doi: 10.1104/pp.106.090035

598 Zhang P-J, Zheng S-J, van Loon JJA, et al (2015) Whiteflies interfere with indirect

599 plant defense against spider mites in Lima bean. *Proc Natl Acad Sci USA*

600 106:21202–21207. doi: 10.1073/pnas.0907890106

601

602

603 **Figure Legends**

604 **Fig. 1** Relative abundance of $E\beta C$ (mean \pm SE) released by maize roots *var.*
605 Delprim after different treatments: inoculated with *P. protegens* CHA0 and
606 infested with *D. balteata* (CHA0+Db), inoculated with *P. chlororaphis* PCL1391
607 and infested with *D. balteata* (PCL+Db), infested with *D. balteata* (Db), control
608 healthy plants, inoculated with *P. protegens* CHA0 (CHA0), and inoculated with
609 *P. chlororaphis* PCL1391 (PCL), (N=9). Lower case letters indicate significant
610 differences between treatments after 6 hours of feeding. Capital letters indicate
611 significant differences between treatments after 72 hours of feeding. Stars
612 indicate significant differences between times. N.S. indicate not significant
613 differences between times.

614

615 **Fig. 2** Relative expression (calculated in relation to actin relative expression) of
616 the terpene synthase gene *Zm-TPS23* (mean \pm SE) in maize roots *var.* Delprim
617 after treatments: inoculated with *P. protegens* CHA0 and infested with *D.*
618 *balteata* (CHA0+Db), inoculated with *P. chlororaphis* PCL1391 and infested with
619 *D. balteata* (PCL+Db), infested with *D. balteata* (Db), control healthy plants,
620 inoculated with *P. protegens* CHA0 (CHA0), and inoculated with *P. chlororaphis*
621 PCL1391 (PCL), (N=9). Lower case letters indicate significant differences
622 between treatments after 6 hours of feeding. Capital letters indicate significant
623 differences between treatments after 72 hours of feeding. Stars indicate
624 significant differences between times. N.S. indicate not significant differences
625 between times.

626

627 **Fig. 3a** Root fresh weight (mean \pm SE) of 14-days-old maize plants *var.*
628 Delprim: inoculated with *P. protegens* CHA0 and infested with *D. balteata*
629 (CHA0+Db), inoculated with *P. chlororaphis* PCL1391 and infested with *D.*
630 *balteata* (PCL+Db), infested with *D. balteata* (Db), control healthy plants,
631 inoculated with *P. protegens* CHA0 (CHA0), and inoculated with *P. chlororaphis*
632 PCL1391 (PCL), (N=12) **b** Weight gain (percentage, mean \pm SE) of *D. balteata*
633 larvae after 6 hours, 48 hours and 72 hours of feeding on maize roots *var.*
634 Delprim with different treatments, (N=9) **c** Percentage of mortality of *D. balteata*
635 larvae after 6 hours, 48 hours and 72 hours of feeding on roots with different
636 treatments, (N=9). Different letters show significant differences between
637 treatments. N.S. not significant differences.

Figure 1




Figure 2




Figure 3






Table 1. Quantification of root colonization by *P. protegens* CHA0 and *P. chloraphis* PCL1391 in different treatments

Treatment	C.F.U. / g of root (\pmSEM)
<i>P. protegens</i> CHA0 + <i>D.balteata</i>	$5.7 \times 10^7 \pm 0.20$ a
<i>P. chloraphis</i> PCL + <i>D.balteata</i>	$1.3 \times 10^8 \pm 0.07$ a
<i>P. protegens</i> CHA0	$2.4 \times 10^8 \pm 1.70$ a
<i>P. chloraphis</i> PCL	$3.5 \times 10^7 \pm 0.65$ a
Control healthy	0