

Reg Environ Change (2011) 11:203–206
DOI 10.1007/s10113-010-0185-6

RAPID COMMUNICATION

Landscape multifunctionality: a powerful concept to identify effects of environmental change

Janine Bolliger · Michèle Bättig · Justus Gallati ·
Andreas Kläy · Michael Stauffacher · Felix Kienast

Received: 22 April 2010 / Accepted: 9 November 2010 / Published online: 1 December 2010
© Springer-Verlag 2010

Abstract The interdisciplinary concept of landscape multifunctionality provides a suitable platform to combine or disentangle effects of multiple environmental stressors acting on the landscape. The concept allows mapping of trade-offs, synergies, and priority conflicts between individual landscape functions, thus providing easily accessible, hands-on means to communicate findings of environmental research to decision makers and society. This rapid communication provides an overview of current developments and potential future research avenues in landscape multifunctionality.

Keywords Environmental change assessments · Landscape multifunctionality · Regional environmental change

J. Bolliger, M. Bättig, J. Gallati, A. Kläy, M. Stauffacher are members of the SAGUF workgroup “Climate change and local action”.

J. Bolliger (✉) · F. Kienast
Swiss Federal Research Institute WSL, Zürcherstrasse 111,
8903 Birmensdorf, Switzerland
e-mail: janine.bolliger@wsl.ch

M. Bättig
Econcept AG, Gerechtigkeitsgasse 20, 8002 Zürich, Switzerland

J. Gallati · M. Stauffacher
Institute for Environmental Decisions (IED), Natural and Social
Science Interface (NSSI), ETH Zurich Universitätsstrasse 22,
8092 Zürich, Switzerland

A. Kläy
Centre for Development and Environment,
Institute of Geography, University of Bern,
Hallerstrasse 10, 3012 Bern, Switzerland

Landscapes and ecosystems provide benefits (goods and services) to society which include both physical and cultural resources (e.g., environmental quality, food, esthetic qualities) which are of great societal value (de Groot 2006; MA 2005). These services are the annual flow of benefits provided by the “natural capital” (e.g., de Groot 2006) or “stock” (Kienast et al. 2009) of a landscape or ecosystem. These “stocks” are the capacity of a landscape to provide services. They are represented in the landscape either by landscape structures (e.g., vegetation, topographical constraints, land cover) or by ecosystem processes (e.g., net primary productivity). It has been suggested to use the term landscape functions (de Groot 2006; Haines-Young and Potschin 2009) to describe the composite nature of the landscape’s capital stocks. The literature (Bolliger and Kienast 2010; Costanza et al. 1997; de Groot et al. 2002; Hein et al. 2006; Kienast et al. 2009; MA 2005) suggests that a wide range of landscape functions (stocks) and associated services (flows) can be identified belonging to four major groups: (1) production functions, (2) regulation functions, (3) habitat functions for maintaining ecological structures/processes, and (4) information functions. Overall, the topic of landscape functions currently seems to gain increasing attention across a broad range of communities dealing with environmental issues (e.g., http://www.eea.europa.eu/publications/briefing_2008_2).

Typically, landscapes are characterized by a variety of overlapping functions. Multifunctionality assessments of landscapes are therefore needed to allow identification of the potential benefits that a landscape can provide to society (Helming and Wiggering 2003; Brandt and Vejre 2004). In a given region, some landscape functions may be beneficial, some adverse, some may be synergetic, and some may be sources of conflicts. Some functions may be spatially and temporally segregated; others may

become effective at the same location at the same time. Hence, landscape functions may be particularly useful for public policy making which often relies on efforts to negotiate trade-offs between potentially conflicting demands of various stakeholders (Gimona and van der Horst 2007).

Research needs

Analyzing trade-offs between landscape functions and related services

It has been recognized that multifunctionality assessments of landscapes may be a powerful tool to analyze trade-offs between landscape services (Brandt and Vejre 2004; Helming and Wiggering 2003; Kienast et al. 2009; Potschin and Haines-Young 2006; Zebisch et al. 2004). For example, trends in land-use change lead to an increase in forest coverage (Albert et al. 2008; Tasser and Tappeiner 2002) with largely negative consequences for open-land species (Bolliger et al. 2007). This is particularly true for marginal and mountainous European regions where economic development has resulted in abandonment of low-intensity agriculture (Bätzig 1996; Maurer et al. 2006; Meeus et al. 1991). As a consequence, land-use changes are generally considered to have a negative impact on grassland species (Bolliger et al. 2007; Dullinger et al. 2003; Laiolo et al. 2004). Contrarily to grassland species (representing habitat functions), the effects on C stocks in terrestrial ecosystems (representing regulation functions) are considered positive as forests are the most important C sink (Janssens et al. 2003). Thus, reforestation may cause severe concerns for conservation (representing habitat functions) while, at the same time, provide an important mitigation strategy for reducing the increase in atmospheric CO₂ concentrations (Bolliger et al. 2008).

Mapping multifunctionality

Visualizing landscape functions as spatially explicit entities adds an important component to research conducted in the context of landscape multifunctionality. With maps, policy makers and planners are able to fully assess the set of landscape capacities at specific locations in an intuitively and comprehensive way. An increasing number of authors try to map landscape functions and trade-offs across large areas (Egoh et al. 2008; Eigenbrod et al. 2010; Kienast et al. 2009; Naidoo et al. 2008; Nelson et al. 2009; Swetnam et al. 2010). This attempt is challenging since landscape functions are rarely a direct outcome of one precisely measured variable in space and time. They rather relate to a blend of different drivers with a high spatial and

temporal data heterogeneity (e.g., resolutions, time windows). Thus, data interpolation is an important prerequisite of landscape function mapping. Often, spatial data are used in combination with decision rules (look-up tables) for mapping landscape functions (e.g., Gimona and van der Horst 2007; Kienast et al. 2009). Decision rules rely on available expert knowledge and usually express assumptions that cannot be related spatially explicitly to sites. To overcome this drawback, promising approaches have recently been presented which explicitly consider the spatial heterogeneity and complexity of landscape characteristics (e.g., Willemen et al. 2008). Given the large spatial GIS databases that are currently available, future methodological progress will hopefully allow to use a broad range of the currently available spatial GIS datasets for multifunctionality assessments.

Fighting mono-causality

Current assessments often provide mono-causal information on individual functions or individual stressors, and it remains largely unknown how the different processes interact. For example, many studies deal with effects of climate change (Beniston 2007; Beniston and Garcia-Herrera 2008; Bonnard et al. 2008; IPCC 2007; Lopez-Moreno et al. 2008; Metzger et al. 2006) and land-use change separately (Bolliger et al. 2007, 2008; Laiolo et al. 2004; Lütolf et al. 2009; Menzel 2000; Steck et al. 2007; Verburg et al. 2006, 2009; Verburg and Overmars 2009). Promising approaches for integrated effects of both climate and land-use change are presented by e.g., Metzger et al. (2008) or by Bradley (2010) who separate impacts caused by land use from impacts caused by climate change.

Emphasizing the temporal dimension of multifunctionality

It has been shown that climate change leads to changes in habitat functions, e.g., the distribution range of communities/species (Thomas et al. 2004; Thuiller et al. 2005). Thus, it may well be that currently protected species (e.g., in dry grasslands) may be driven out of their current reserves in the future (Araujo et al. 2004; Kienast et al. 1998). Additionally, invasive species may affect current species compositions (Bohren et al. 2008; Scharfy et al. 2009), may become problematic for public health (Ackermann-Liebrich et al. 2009), or species and populations may face extinction (Thomas et al. 2004; Thuiller et al. 2005). Promising examples of dynamic multifunctionality assessments are given by e.g., Hasselmann et al. (2010), Hersperger and Bürgi (2009), Houet et al. (2010), or Verburg et al. (2010).

Emphasizing the anthropogenic dimension

Production, regulation, and habitat functions appear to dominate the literature dealing with environmental issues. However, the anthropogenic dimension requires increasing consideration in scientific environmental change studies, since land management and policies may overrule landscape-scale effects of environmental change to a considerable extent (Willemen et al. 2008). For example, if former fields and pastures are increasingly covered by shrubs and eventually by forest, scenic beauty, an important landscape resource, may be at risk (Buijs et al. 2006). Previous studies show that a low to medium degree of spontaneous reforestation (patches or bosques of forest cover dispersed throughout a matrix of open space) is often found particularly attractive (Hunziker 1995; Hunziker and Kienast 1999), but see Gehring (2006). It has also been shown that conflicts may arise when the visual appearance of landscapes opposes public preferences (De la Fuente de Val et al. 2006). Thus, increasing attention should be given to the anthropogenic dimension of landscape-change assessments by addressing and developing the yet limited availability of information functions such as esthetic, recreation, cultural and artistic, or historic information.

Landscape multifunctionality is a challenging field to explore in current and future landscapes subject to a changing environment. Landscape multifunctionality can only be addressed if larger, interdisciplinary research consortia are willing to collaborate, merge current knowledge, and conduct new or supplementary research. No doubt, however, that the current manifold environmental issues are pressing and that scientific results are not easily communicated to land managers, politicians, and other decision makers. Visualization of landscape multifunctionality through maps of landscape function (changes) may provide helpful baseline information for discussions with stakeholders if tackling (future) requirements to support a dynamic environment is a priority for environmental research.

References

- Ackermann-Liebrich U, Schindler C, Frei P et al (2009) Sensitisation to ambrosia in Switzerland: a public health threat in waiting. *Swiss Med Wkly* 139:70–75
- Albert CH, Thuiller W, Lavorel S et al (2008) Land-use change and subalpine tree dynamics: colonization of *Larix decidua* in French subalpine grasslands. *J Appl Ecol* 45:659–669
- Araujo MB, Cabeza M, Thuiller W et al (2004) Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Glob Chang Biol* 10:1618–1626
- Bätzig W (1996) Landwirtschaft im Alpenraum unverzichtbar, aber zukunftslos? Eine alpenweite Bilanz der aktuellen Probleme und der möglichen Lösungen. In: Bätzig W (ed) *Landwirtschaft im Alpenraum—unverzichtbar, aber zukunftslos?* Europäische Akademie Bozen, Fachbereich Alpine Umwelt, Blackwell, pp 9–11
- Beniston M (2007) Linking extreme climate events and economic impacts: examples from the Swiss Alps. *Energy Policy* 35:5384–5392
- Beniston M, Garcia-Herrera R (2008) Climate change and health. In: Thomson MC et al (eds) *Seasonal forecasts, climatic change and human health—health and climate*. Springer, Berlin, pp 131–133
- Bohren C, Delabays N, Rometsch S (2008) Invasive plants: agricultural aspects. *Agrarforschung* 15:314–319
- Bolliger J, Kienast F (2010) Landscape functions in a changing environment. *Landscape Online*
- Bolliger J, Kienast F, Soliva R et al (2007) Spatial sensitivity of species habitat patterns to scenarios of land-use change (Switzerland). *Landscape Ecol* 22:773–789
- Bolliger J, Hagedorn F, Leifeld J et al (2008) Potential carbon-pool changes under various scenarios of land-use change in a mountainous region (Switzerland). *Ecosystems* 11:895–907
- Bonnard C, Tacher L, Beniston M (2008) Prediction of landslide movements caused by climate change: modelling the behaviour of a mean elevation large slide in the Alps and assessing its uncertainties. *Crc Press-Taylor and Francis Group*, Boca Raton
- Bradley BA (2010) Assessing ecosystem threats from global and regional change: hierarchical modeling of risk to sagebrush ecosystems from climate change, land use and invasive species in Nevada, USA. *Ecography* 33:198–208
- Brandt J, Vejre H (2004) *Multifunctional landscapes: theory, values and history*. WIT Press, Southampton
- Buijs AE, Pedroli B, Luginbühl Y (2006) From hiking through farmland to farming in a leisure landscape: changing social perception of the European landscape. *Landsc Ecol* 21:375–389
- Costanza R, D'Arge R, de Groot R et al (1997) The value of the world's ecosystem services and natural capital. *Nature* 387:253–260
- de Groot R (2006) Function-analysis and valuation as a tool to assess land use conflicts in planning for sustainable, multi-functional landscapes. *Landsc Urban Plan* 75:175–186
- de Groot RS, Wilson M, Boumans R (2002) A typology for the description, classification and valuation of ecosystem functions, goods and services. *Ecol Econ* 41:393–408
- De la Fuente de Val G, Atauri JA, de Lucio JV (2006) Relationship between landscape visual attributes and spatial pattern indices: a test study in Mediterranean-climate landscapes. *Landsc Urban Plan* 77:393–407
- Dullinger S, Dirnböck T, Grabherr G (2003) Patterns of shrub invasion into high mountain grasslands of the Northern calcareous Alps, Austria. *Arct Antarct Alp Res* 35:434–441
- Egoh B, Reyers B, Rouget M et al (2008) Mapping ecosystem services for planning and management. *Agric Ecosyst Environ* 127:135–140
- Eigenbrod F, Armsworth PR, Anderson BJ et al (2010) The impact of data quality on the spatial congruence of ecosystem services. *J Appl Ecol* 47:377–385
- Gehring K (2006) Landscape needs and notions. Preferences, expectations, leisure motivation, and the concept of landscape from a cross-cultural perspective. *Swiss Federal Research Institute WSL, Zurich*
- Gimona A, van der Horst D (2007) Mapping hotspots of multiple landscape functions: a case study on farmland afforestation in Scotland. *Landsc Ecol* 22:1255–1264
- Haines-Young R, Potschin M (2009) The links between biodiversity, ecosystem services and human well-being. In: Raffaelli D, Frid C (eds) *Ecosystem ecology: a new synthesis*. BES Ecological Reviews Series, CUP
- Hasselmann F, Csaplovics E, Falconer I et al (2010) Technological driving forces of LUCC: conceptualization, quantification, and

- the example of urban power distribution networks. *Land Use Policy* 27:628–637
- Hein L, van Koppen K, de Groot RS et al (2006) Spatial scales, stakeholders and the valuation of ecosystem services. *Ecol Econ* 57:209–228
- Helming K, Wiggering H (2003) Sustainable development of multifunctional landscapes. Springer, Berlin
- Hersperger AM, Burgi M (2009) Going beyond landscape change description: quantifying the importance of driving forces of landscape change in a Central Europe case study. *Land Use Policy* 26:640–648
- Houet T, Verburg PH, Loveland TR (2010) Monitoring and modelling landscape dynamics. *Landscape Ecol* 25:163–167
- Hunziker M (1995) The spontaneous reafforestation in abandoned agricultural lands: perception and aesthetic assessment by locals and tourists. *Landscape Urban Plan* 31:399–410
- Hunziker M, Kienast F (1999) Potential impacts of changing agricultural activities on scenic beauty—a prototypical technique for automated rapid assessment. *Landscape Ecol* 14:161–176
- IPCC, 2007: Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007, Parry ML et al, Eds., Cambridge University Press, Cambridge
- Janssens IA, Freibauer A, Ciais P et al (2003) Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic CO₂ emissions. *Science* 300:1538–1542
- Kienast F, Wildi O, Brzeziecki B (1998) Potential impacts of climate change on species richness in mountain forests—an ecological risk assessment. *Biol Conserv* 83:291–305
- Kienast F, Bolliger J, Potschin M et al (2009) Assessing landscape functions with broad-scale environmental data: insights gained from a prototype development for Europe. *Environ Manage* 44:1099–1120
- Laiolo P, Dondero F, Ciliento E et al (2004) Consequences of pastoral abandonment for the structure and diversity of the alpine avifauna. *J Appl Ecol* 41:294–304
- Lopez-Moreno JJ, Goyette S, Beniston M (2008) Climate change prediction over complex areas: spatial variability of uncertainties and predictions over the Pyrenees from a set of regional climate models. *Int J Climatol* 28:1535–1550
- Lütolf M, Bolliger J, Kienast F et al (2009) Scenario-based assessment of future land-use change on butterfly species distributions. *Biodivers Conserv* 18:1329–1347
- MA (2005) Millennium ecosystem assessment, Business and industry. Synthesis report. Island Press, Washington
- Maurer S, Weyand A, Fischer M et al (2006) Old cultural traditions, in addition to land use and topography, are shaping plant diversity of grasslands in the Alps. *Biol Conserv* 130:438–445
- Meeus J, Van Der Ploeg JD, Wijermans M (1991) Changing agricultural landscapes in Europe: continuity, deterioration or rupture? IFLA conference, Rotterdam
- Menzel A (2000) Trends in phenological phases in Europe between 1951 and 1996. *Int J Biometeorol* 44:76–81
- Metzger MJ, Rounsevell MDA, Acosta-Michlik L et al (2006) The vulnerability of ecosystem services to land use change. *Agric Ecosyst Environ* 114:69–85
- Metzger MJ, Schroter D, Leemans R et al (2008) A spatially explicit and quantitative vulnerability assessment of ecosystem service change in Europe. *Reg Environ Change* 8:91–107
- Naidoo R, Balmford A, Costanza R et al (2008) Global mapping of ecosystem services and conservation priorities. *PNAS* 105:9495–9500
- Nelson E, Mendoza G, Regetz J et al (2009) Modeling multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs at landscape scales. *Front Ecol Environ* 7:4–11
- Potschin M, Haines-Young R (2006) 'Rio +10', sustainability science and landscape ecology. *Landscape Urban Plan* 75:162–174
- Scharfy D, Eggenschwiler H, Venterink HO et al (2009) The invasive alien plant species *Solidago gigantea* alters ecosystem properties across habitats with differing fertility. *J Vegetat Sci* 20:1072–1085
- Steck C, Bürgi M, Bolliger J et al (2007) Conservation of grasshopper diversity in a changing environment. *Biol Conserv* 138:360–370
- Swetnam RD, Fisher B, Mbilinyi BP et al (2010) Mapping socio-economic scenarios of land cover change: A GIS method to enable ecosystem service modelling. *J Environ Manage* 1–12
- Tasser E, Tappeiner U (2002) Impact of land use changes on mountain vegetation. *Appl Vegetat Sci* 5:173–184
- Thomas CD, Cameron A, Green RE et al (2004) Extinction risk from climate change. *Nature* 427:145–148
- Thuiller W, Lavorel S, Araujo MB et al (2005) Climate change threats to plant diversity in Europe. *Proc Natl Acad Sci USA* 102:8245–8250
- Verburg PH, Overmars KP (2009) Combining top-down and bottom-up dynamics in land use modeling: exploring the future of abandoned farmlands in Europe with the Dyna-CLUE model. *Landscape Ecol* 24:1167–1181
- Verburg P, Schulp CJE, Witte N et al (2006) Downscaling of land use change scenarios to assess the dynamics of European landscapes. *Agric Ecosyst Environ* 114:39–56
- Verburg P, van Berkel D, Willemsen L et al (2009) Exploration of the influence of land use and land cover change on landscape functions. In: Breuste J et al (eds) European landscapes in transformation—challenges for landscape ecology and management. University of Salzburg, Salzburg, pp 236–237
- Verburg PH, van Berkel DB, van Doorn AM et al (2010) Trajectories of land use change in Europe: a model-based exploration of rural futures. *Landscape Ecol* 25:217–232
- Willemsen L, Verburg PH, Hein L et al (2008) Spatial characterization of landscape functions. *Landscape Urban Plan* 88:34–43
- Zebisch M, Wechsung F, Kenneweg H (2004) Landscape response functions for biodiversity—assessing the impact of land-use changes at the county level. *Landscape Urban Plan* 67:157–172