



## SHORT COMMUNICATION

# Megalin in normal tissues and carcinoma cells carries oligo/poly $\alpha$ 2,8 deaminoneuraminic acid as a unique posttranslational modification

Martin Ziak, Mirjam Meier and Jürgen Roth\*

*Division of Cell and Molecular Pathology, Department of Pathology, University of Zürich, CH-8091 Zürich, Switzerland*

**In rat kidney, megalin, a member of the low density lipoprotein receptor gene family, is the sole glycoprotein which carries oligo/poly  $\alpha$ 2,8 deaminoneuraminic acid (KDN) as a posttranslational modification. We have investigated immunoprecipitated megalin from rat brain, lung and placenta, mouse yolk sac carcinoma and megalin synthesizing carcinoma cell lines, for presence of this unique glycan structure. Our immunoblot analysis revealed the presence of oligo/poly  $\alpha$ 2,8 KDN on megalin in all the studied normal tissues and carcinoma cells. Furthermore, it is demonstrated to be part of oligosaccharides *O*-glycosidically linked to megalin.**

**Keywords:** oligo/poly  $\alpha$ 2,8 deaminoneuraminic acid, megalin, lung, brain, placenta, F9 cells, L2 cells

### Introduction

In previous studies, we demonstrated the presence of oligo/poly  $\alpha$ 2,8 linked deaminoneuraminic acid (oligo/poly  $\alpha$ 2,8 KDN) in various embryonic, postnatal developing and adult mammalian tissues by immunohistochemistry and immunoblot analysis [1,2] with the use of the monoclonal antibody mAb.kdn8kdn [3]. The existence of KDN in mammalian tissues was furthermore confirmed by gas liquid chromatography analysis [1], and a sensitive fluorescent probe for HPLC [4] detection of KDN in tissues [5]. Western blot analysis of extracts of various rat tissues revealed the presence of a single reactive band in each tissue studied. The oligo/poly  $\alpha$ 2,8 KDN was found on a single 150 kDa glycoprotein except for a single >350 kDa glycoprotein in kidney [1,2].

Recently, we isolated and purified the oligo/poly  $\alpha$ 2,8 KDN bearing glycoprotein from rat kidney [6] and identified it as being megalin, a member of the low-density lipoprotein receptor gene family [7]. The presence of oligo/poly  $\alpha$ 2,8 KDN on kidney megalin was confirmed by combined immunoprecipitation / immunoblot analysis and on RAP-

affinity purified megalin. Further supporting evidence was obtained by immunoelectron microscopy revealing an identical subcellular distribution of oligo/poly  $\alpha$ 2,8 KDN and megalin in rat kidney proximal tubules [6].

Megalin represents a major membrane protein of kidney proximal tubules but is additionally found in rat lung, choroid plexus and microvasculature of brain, placenta, yolk sac epithelia, ciliary epithelium of the eye, parathyroid as well as inner ear [8]. Furthermore, by immunoblotting megalin was demonstrated in mouse F9 teratocarcinoma cells [9] and L2 rat yolk sac carcinoma cells [10].

Here, we report that megalin from various normal rat tissues, carcinoma tissue and carcinoma cell lines carries oligo/poly  $\alpha$ 2,8 KDN which is part of *O*-glycosidically linked oligosaccharides.

### Materials and methods

#### Materials

For the detection of oligo/poly  $\alpha$ 2,8 KDN, purified IgM of the mouse monoclonal antibody mAb.kdn8kdn was used [3]. The hybridoma cell line 2G-5 [3] was kindly supplied by Dr. Ken Kitajima (University of Nagoya, Nagoya Japan). Polyclonal antibodies against purified rat kidney megalin were raised in rabbits as described previously [11] and IgG

\*To whom correspondence should be addressed: Tel: +41 1 255 50 90. Fax: +41 1 255 44 07. E-mail: [juergen.roth@pty.usz.ch](mailto:juergen.roth@pty.usz.ch)

fractions prepared using Protein A Sepharose CL-4B (Pharmacia, Uppsala, Sweden). Alkaline phosphatase-conjugated donkey anti-mouse IgM (affinity-purified) was purchased from Jackson ImmunoResearch Laboratories, Inc. (West Grove, PA, USA) and frozen rat tissues from Pel Freez Biologicals (Rogers, AR, USA). Mouse yolk sac tumor tissue and the L2 cell line were kindly supplied by Dr. Ulla Wewer (University Institute of Pathological Anatomy, Copenhagen, Denmark). Mouse F9 teratocarcinoma cells were obtained from the American Type Culture Collection (Rockville, MD). Protease inhibitor cocktail tablets, recombinant N-Glycosidase F and digoxigenin-conjugated Concanavalin A were from Boehringer (Mannheim, Germany). Sephacryl S-400, XK 16/100 column were purchased from Pharmacia (Uppsala, Sweden). All other reagents were of analytical grade and purchased from Fluka (Buchs, Switzerland).

### Cell cultures

L2 cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with high glucose and 10% fetal calf serum at 37°C in a humid atmosphere containing 5% CO<sub>2</sub>. Mouse F9 teratocarcinoma cells were cultured in gelatin coated (0.1%) flasks in DMEM with high glucose and 15% fetal calf serum. To induce differentiation, F9 cells were incubated for 72 hours with 1 μM retinoic acid and 1 mM dibutyryl cyclic AMP.

### Megaline immunoprecipitation

Protein A Sepharose CL-4B was incubated with PBS containing 1% BSA and 0.05% of Triton X-100 and Tween 20 for 1 h at 4°C, followed by incubation with anti-megaline antibodies for 18 h at 4°C. Rat kidney (0.7 g), lung (4.2 g), brain (25 g), placenta (4.2g), mouse yolk sac carcinoma tissue (4.2 g) and pellets of L2 and F9 cells (200 μg each) were homogenized in PBS containing protease inhibitors and Triton X-100 (final concentration 1%). After 60 min on ice, the homogenate was centrifuged at 100,000 × g for 60 min. The soluble extract was incubated with the anti-megaline / Protein A Sepharose CL-4B overnight at 4°C. Following two washes with PBS containing 0.1% Triton X-100, the Protein A Sepharose CL-4B was pelleted, placed in SDS-PAGE sample buffer and heated. The supernatant was analyzed by SDS-PAGE and Western blotting using anti-oligo/poly α<sub>2,8</sub> KDN or anti-megaline antibodies as described previously [6].

### Partial purification of megaline from mouse yolk sac carcinoma tissue

A mouse yolk sac carcinoma tissue homogenate was applied onto a XK 16/100 gel filtration column packed with Sephacryl S-400. Fractions, immunoreactive for oligo/poly

α<sub>2,8</sub> KDN were pooled and used for Western blot and lectin blot analysis.


### β-elimination, N-Glycosidase F treatment and lectin blotting

For β-elimination, PVDF strips containing partially purified megaline from yolk sac carcinoma tissue were incubated with 0.1 N sodium hydroxide for 24 h at 37°C. As control, strips were incubated with PBS under the same conditions as described above. For N-Glycosidase F treatment, nitrocellulose strips with megaline were blocked with 1% BSA in 50 mM sodium acetate buffer (pH 5.5) followed by incubation with 5 U of N-Glycosidase F for 16 h at 37°C. For lectin blotting, strips were incubated with digoxigenin-conjugated Concanavalin A for 1 h followed by alkaline phosphatase-conjugated polyclonal sheep anti-digoxigenin Fab' fragments (150 U/ml). Color reaction was performed using nitroblue tetrazolium/BCIP-phosphate as substrates.


### Results and discussion

In our previous studies [6] the >350 kDa oligo/poly α<sub>2,8</sub> KDN bearing glycoprotein from rat kidney was identified as being megaline. Megaline is the most abundant membrane protein in proximal tubules of rat kidney but it is also expressed in other specialized epithelia [8] albeit at much lower levels. Immunoprecipitated megaline from rat lung, brain and placenta, from mouse yolk sac carcinoma tissue as well as L2 cells and from mouse F9 teratocarcinoma cells, induced to differentiate (data not shown) exhibited immunoreactivity for oligo/poly α<sub>2,8</sub> KDN (Fig. 1). Thus, megaline present in these normal rat tissues and in carcinoma cells carries oligo/poly α<sub>2,8</sub> KDN as a unique posttranslational modification. These findings on megaline are similar to those for a 150 kDa glycoprotein in lung which also carries oligo/poly α<sub>2,8</sub> KDN both in normal human and rat embryonic and postnatal lung and in human lung carcinomas [2]. Furthermore, KDN has been detected in both normal human ovary and ovarian carcinoma cells but no data have been reported regarding its protein carrier [12].

β-elimination on stripes containing partially purified megaline from mouse yolk sac tissue resulted in the absence in immunoreactivity for oligo/poly α<sub>2,8</sub> KDN. In contrast N-glycosidically linked oligosaccharides on megaline are not affected by this treatment as demonstrated by the reactivity with Concanavalin A (Fig. 2). Furthermore, N-Glycosidase F treatment increased the immunoreactivity for oligo/poly α<sub>2,8</sub> KDN (Figure 2) indicating that oligo/poly α<sub>2,8</sub> KDN is part of O-glycosidically linked oligosaccharides on megaline from mouse yolk sac carcinoma tissue. These data are in agreement with previously described results obtained from purified rat kidney megaline [6].



**Figure 1.** Immunoprecipitated megalin from rat kidney (lane 1), lung (lane 2) brain (lane 3) and placenta (lane 4), mouse yolk sac carcinoma (lane 5) and L2 cells (lane 6) exhibited immunoreactivity for oligo/poly  $\alpha$ 2,8 KDN. Molecular mass standards were as follows: myosin (200 kDa),  $\beta$ -galactosidase (116 kDa), phosphorylase b (97 kDa), bovine serum albumin (69 kDa).



**Figure 2.**  $\beta$ -elimination and N-Glycosidase F treatment on partially purified megalin from yolk sac carcinoma tissue.  $\beta$ -elimination resulted in the absence of immunoreactivity for oligo/poly  $\alpha$ 2,8 KDN (lane 1). Lectin blot using Concanavalin A after  $\beta$ -elimination (lane 2). Immunoblot for oligo/poly  $\alpha$ 2,8 KDN (lane 3), Concanavalin A blot (lane 4) and immunoblot for megalin (lane 5) after incubation of the stripes with PBS at 37°C for 24 h. Increased immunoreactivity for oligo/poly  $\alpha$ 2,8 KDN after N-Glycosidase F treatment (lane 6) and absence of labeling for Concanavalin A (lane 7). Immunoblot for megalin after N-Glycosidase F treatment (lane 8).

Megalyn has been found to bind a number of different ligands (for a review see [13]) *in vitro*. What may be a possible function of oligo/poly  $\alpha$ 2,8 KDN on megalyn? Due to its localization in clathrin-coated pits megalyn was suggested to act as an endocytic receptor in kidney. Dependent on the organ localization of megalyn and the composition of the surrounding fluids it can be anticipated that the nature of the ligands will vary from one organ to another. For example, in proximal tubules it was suggested that megalyn is involved in the reabsorption of filtered proteins [14] and  $\text{Ca}^{2+}$ -ions [15]. In type II pneumocytes, megalyn seems to be responsible for the clearance of protease / protease inhibitor complexes [16,17] from the alveolar space [18]. Charge interactions between negatively charged complement-type repeats and basic regions of the ligands seems to be important for receptor / ligand interaction. Furthermore, megalyn is not only a  $\text{Ca}^{2+}$  binding protein but  $\text{Ca}^{2+}$ -ions are essential for ligand binding. *In vitro* oligo/poly  $\alpha$ 2,8 KDN binds  $\text{Ca}^{2+}$ -ions highly preferentially [19]. Therefore, we speculate that the polyanionic nature of oligo/poly  $\alpha$ 2,8 KDN together with the negatively charged complement-type repeats may be important for ligand binding to megalyn and for the receptor activity in the various tissues. Studies are in progress to clarify this proposal.

## References

- Ziak M, Qu B, Zuo X, Zuber C, Kanamori A, Kitajima K, Inoue S, Inoue Y, Roth J (1996) *Proc Natl Acad Sci USA* **93**: 2759–63.
- Qu B, Ziak M, Zuber C, Roth J (1996) *Proc Natl Acad Sci USA* **93**: 8995–8.
- Kanamori A, Inoue S, Xulei Z, Zuber C, Roth J, Kitajima K, Ye J, Troy FA, Inoue Y (1994) *Histochemistry* **101**: 333–40.
- Hara S, Takemori Y, Yamaguchi M, Nakamura M, Ohkura Y (1987) *Anal Biochem* **164**: 138–45.
- Inoue S, Kitajima K, Inoue Y (1996) *J Biol Chem* **271**: 24341–4.
- Ziak M, Kerjaschki D, Farquhar M, G., Roth J (1999) *J Am Soc Nephrol*. **10**: 203–09.
- Saito A, Pietromonaco S, Loo AK, Farquhar MG (1994) *Proc Natl Acad Sci USA* **91**: 9725–9.
- Zheng G, Bachinsky DR, Stamenkovic I, Strickland DK, Brown D, Andres G, McCluskey RT (1994) *J Histochem Cytochem* **42**: 531–42.
- Czekay RP, Orlando RA, Woodward L, Adamson ED, Farquhar MG (1995) *J Cell Sci* **108**: 1433–41.
- Lundstrom M, Orlando RA, Saedi MS, Woodward L, Kurihara H, Farquhar MG (1993) *Am J Pathol* **143**: 1423–35.
- Guhl B, Ziak M, Roth J (1998) *Histochem Cell Biol* **110**: 603–11.
- Inoue S, Lin SL, Chang T, Wu SH, Yao CW, Chu TY, Troy FA, Inoue Y (1998) *J Biol Chem* **273**: 27199–204.
- Farquhar MG, Saito A, Kerjaschki D, Orlando RA (1995) *J Amer Soc Nephrol* **6**: 35–47.

- 14 Moestrup SK, Christensen EI, Nielsen S, Jorgensen KE, Bjorn SE, Roigaard H, Gliemann J (1994) *Ann NY Acad Sci* **737**: 124–37.
- 15 Christensen EI, Gliemann J, Moestrup SK (1992) *J Histochem Cytochem* **40**: 1481–90.
- 16 Stefansson S, Kounnas MZ, Henkin J, Mallampalli RK, Chappell DA, Strickland DK, Argraves WS (1995) *J Cell Sci* **108**: 2361–8.
- 17 Poller W, Willnow TE, Hilpert J, Herz J (1995) *J Biol Chem* **270**: 2841–5.
- 18 Willnow TE, Hilpert J, Armstrong SA, Rohlmann, A, Hammer, RE, Burns DK, Herz J, (1996) *Proc Natl Acad Sci USA* **93**: 8460–4.
- 19 Shimoda Y, Kitajima K, Inoue S, Inoue Y (1994) *Biochemistry* **33**: 1202–8.

Received 4 November 1998, revised 19 February 1999, accepted 3 March 1999.