

J.F. von Raumer · G.M. Stampfli · G. Borel · F. Bussy

Organization of pre-Variscan basement areas at the north-Gondwanan margin

Received: 12 May 2000 / Accepted: 24 January 2001 / Published online: 12 May 2001
© Springer-Verlag 2001

Abstract Pre-Variscan basement elements of Central Europe appear in polymetamorphic domains juxtaposed through Variscan and/or Alpine tectonic events. Consequently, nomenclatures and zonations applied to Variscan and Alpine structures, respectively, cannot be valid for pre-Variscan structures. Comparing pre-Variscan relics hidden in the Variscan basement areas of Central Europe, the Alps included, large parallels between the evolution of basement areas of future Avalonia and its former peri-Gondwanan eastern prolongations (e.g. Cadomia, Intra-Alpine Terrane) become evident. Their plate-tectonic evolution from the Late Proterozoic to the Late Ordovician is interpreted as a continuous Gondwana-directed evolution. Cadomian basement, late Cadomian granitoids, late Proterozoic detrital sediments and active margin settings characterize the pre-Cambrian evolution of most of the Gondwana-derived microcontinental pieces. Also the Rheic ocean, separating Avalonia from Gondwana, should have had, at its early stages, a lateral continuation in the former eastern prolongation of peri-Gondwanan microcontinents (e.g. Cadomia, Intra-Alpine Terrane). Subduction of oceanic ridge (Proto-Tethys) triggered the break-off of Avalonia, whereas in the eastern prolongation, the presence of the ridge may have triggered the amalgamation of volcanic arcs and continental ribbons with Gondwana

(Ordovician orogenic event). Renewed Gondwana-directed subduction led to the opening of Palaeo-Tethys.

Keywords Rheic ocean · Peri-Gondwana evolution · Pre-Variscan basement · Palaeo-Tethys · Palaeoreconstruction · Cambro-Ordovician

Introduction

Pre-Variscan elements of Central Europe are mostly exposed as pre-Mesozoic basement areas that underwent the Variscan thermotectonic events, and occur as polymetamorphic domains which, after their respective drift and amalgamation, have been stacked and “shovelled” during Variscan (e.g. Bard 1997; Oncken 1997) and Alpine events one in front of the other to produce the present-day juxtaposition. During their large-scale accretion and strike-slip transport, many of these basement areas left behind their lithospheric roots. Consequently, in the European Variscan basement areas (Fig. 1; e.g. Iberia, Armorica, Moesia, French Massif Central, Saxothuringian and Moldanubian domains, External massifs, Penninic domain, parts of the southern Alps and the Austroalpine basement), nappe and strike-slip structures of different ages repeat elements which, initially, had a distinct pre-Variscan location, and a common evolution, comprising different steps since the Precambrian. The current nomenclatures and zonations applied to Variscan or Alpine regional structures cannot characterize the original location of pre-Variscan crustal elements and, necessarily, a new nomenclature has to be based on plate tectonic criteria considering the geotectonic evolution of these elements during the Neoproterozoic and Early Palaeozoic. As we are dealing mostly with lithostratigraphies of metamorphic areas, we try to maintain a standpoint independent of models of Variscan evolution of these areas.

J.F. von Raumer (✉)
Institut de Minéralogie et Pétrographie, Université Pérolles,
1700 Fribourg, Switzerland
E-mail: Juergen.vonRaumer@unifr.ch
Phone: +41-26-3008927
Fax: +41-26-3009765

G.M. Stampfli · G. Borel
Institut de Géologie, Université de Lausanne,
UNIL BFSH2 1015 Lausanne, Switzerland

F. Bussy
Institut de Minéralogie, Université de Lausanne,
UNIL BFSH2 1015 Lausanne, Switzerland




Fig. 1 Distribution of Pre-Mesozoic (Variscan and/or older) units in Europe (Zwart and Dornsiepen 1978) with indication of distinct microcontinents (see Fig. 7). AA Austro-Alpine; Ab Alboran; Am Armorica; Ap Apulia; AQ Aquitaine; AR Armoriga; Ch Channel terrane; Ct Cantabria; DH Dinarides-Hellenides; He Helvetic; iA intra-Alpine; Ib Iberia; MD Moldanubian; OM Ossa-Morena; Or Ortegale; Pe Penninic; sP south Portuguese; Sx Saxothuringian


Examples of the Early Palaeozoic evolution preserved in the Variscan mountain chain of Europe are numerous (see Dallmeyer et al. 1995; Keppie 1994; von Raumer and Neubauer 1993a, 1993b; Franke et al. 2000). Sedimentary and fossil records, as well as age determinations, show the overall presence of Neoproterozoic to Early Palaeozoic basements supposed to have evolved along the margin of Gondwana, and sharing specific structural elements and events. For the time period after the Ordovician, different plate-tectonic evolutions are proposed, favouring either one (Robardet et al. 1990) or two major mid-European

oceanic separations (Matte 1986), or the separation of a basement assemblage (“Hun-superterrane”; Stampfli 1996, 2000). Such divergent reconstructions need to be tested, and it is the aim of this paper to discuss, from a “Gondwanan” standpoint, and independently of the Variscan structures, parallels of pre-Variscan evolution among the polymetamorphic basement areas mentioned previously.

Pre-Ordovician peri-Gondwana evolution

Among the different pre-Variscan relics preserved in the Variscan basement areas – the pre-Mesozoic Alpine basement areas included – we can distinguish distinct geodynamic settings from the Precambrian to the Ordovician, comprising Precambrian basement blocks (Cadomian/PanAfrican basement) and their sedimentary cover, formation of Late Precambrian to Cambro-Ordovician oceanic crust, volcanic arcs, and active margin remnants, and zones of amalgamation (accretion, collision) of Early Palaeozoic age.

Fig. 2 Plate tectonic re-interpretation of pre-Variscan units in the Alps. Compared with Frisch and Neubauer (1989, terminology in *italic* with quotation marks), all Early Palaeozoic terranes of the Alps are re-interpreted in terms of their peri-Gondwanian origin. Pieces comparable to those of the “*Noric composite Terrane*” are interpreted to be related to the opening of Palaeotethys. The arc volcanic “*Celtic Terrane*” and its counterparts in the Western Alps represent units with Early Palaeozoic metamorphism. Terranes related to the opening of Early Palaeozoic oceanic domains (“*Speik Terrane*”) represent either contemporaneous or successor stages of Rheic ocean and/or its equivalents. Arc volcanics (“*Habach Terrane*”) represent Early Palaeozoic active margin settings. Other signatures in the domain of the Eastern Alps, not mentioned in the explanations, refer to Variscan elements and structures (see Frisch and Neubauer, 1989). *Inset* General situation of the Alpine pre-Mesozoic basement terranes (*black*) within their present-day Variscan framework. MC French Central Massif; BM Bohemian Massif



Rast 1983). It is not the scope of this paper to recall the complex evolution of these oldest elements which either appear as low-grade or as polymetamorphic domains (pan-African/Cadomian orogeny), and for which the relationship between Cadomian basement areas and Neoproterozoic to Early Palaeozoic sedimentary cover have been the subject of extensive research. The general picture of small continental blocks bordering the Gondwana margin (Nance and Murphy 1994) explains best the many relationships which existed between all these pieces. Large-scale magmatic activity is documented by syn- and postorogenic granitoids or acidic volcanites in the wider framework of Gondwanan evolution (see below). Also in the Alpine domain, a distinction can be made between a Proterozoic and Archean inheritance, Late Precambrian mafic and ultramafic rocks, and relics of Precambrian–Cambrian sediments (Schaltegger and Gebauer 1999). Widely distributed across the different Alpine realms, these sediments contain detrital zircons suggesting derivation from Cadomian–pan-African and older basement units. Comparable results were obtained from detrital micas (Dallmeyer and Neubauer 1994; Dallmeyer et al. 1996a, 1996b; Handler et al. 1997; Panwitz et al. 2000). Pre-Cambrian basement is discussed for a few regions in the Western and Central Alps (Gebauer 1993), including the Gotthard area of the External domain (Gebauer et al. 1988), the Berisal Complex (Stille and Tatsumoto 1985; Zingg 1989) and Siviez-Mischabel unit (Thélin et al. 1993) of the Penninic domain, probably the “ancient basement” in the internal zones of the Western Alps (Desmons et al. 1999), the Silvretta area of the Austroalpine basement (Schaltegger et al. 1997), and in the Eastern Alps (Neubauer 1991; von Quadt 1992; Thöni 1999). Sm–Nd data (e.g. Liebetrau and Nägler 1994; Böhm 1996; Poller et al. 1997) from Cambro-Ordovician or Variscan granitoids closely match values observed in the Bohemian Massif (Hegner and Kröner 2000), in NW Iberia (Fernández Suárez et al. 1998) and from West Avalonia (Nance and Murphy 1996). Findings of Cadomian ages in the Pontides (Üstaömer 1999) would allow extension of this domain eastward maybe up to the Caucasus.

Early Palaeozoic sedimentary troughs–oceanic settings

Subsidence analysis shows the large-scale opening of early Palaeozoic sedimentary basins for Laurentia and western Gondwana (Williams 1997), and contemporaneous events are registered in the Armorican Massif (Paris and Robardet 1990; Robardet et al. 1994), and in the Ossa Morena domain (Liñan and Quesada 1990). In an Early Cambrian reconstruction (McKerrow et al. 1992), Archaeocyathan assemblages indicate a climate-sensitive palaeogeographic location within 30° of the equator. Regular fracture patterns determine the direction of intra-continental sedimentary

troughs (Courjault-Radé et al. 1992), and delimit large areas with deposits of continental-margin environment in the Gondwana platform during the Cambrian. In fact, the most important lithology, locally highly metamorphosed association of metagreywackes, metapelites, a few quartzites, carbonates, and metavolcanic interlayers, is well known in the Central Iberian domain, and in the Montagne Noire area, but also from several Alpine domains (e.g. External Massifs, Southern Alps, von Raumer 1998; ancient basement areas of the Penninic domain, Desmons et al. 1999), or the Pyrenees (Laumonier 1998), and has astonishing parallels with fossil-bearing, Neoproterozoic to Cambrian lithostratigraphies from the Saxothuringian domain (e.g. Buschmann et al. 1995), with several localities around the Mediterranean Sea (Sdzuy et al. 1999), or well-preserved examples observed in the Anti-Atlas mountain chain of Morocco; however, in addition to the aforementioned parallels also discrepancies exist. Comparable lithologies may remain undiscovered in the polymetamorphic areas of the Variscan mountain chain, as the high plasticity of marbles during metamorphic overprint may reduce considerably original limestone occurrences, which may disappear completely during metamorphic evolution. On the other hand, a specific palaeogeographic configuration may influence carbonate distribution. They may have been deposited on the Gondwana platform, they may have been resedimented as reduced detrital carbonate layers near a rift shoulder (e.g. Freyer and Suhr 1987), and may have been replaced by contemporaneous clastic sequences in the domain of Cambrian rift zones. Extensive Ordovician areas of similar sedimentary facies at the Gondwana shelf (Noblet and Lefort 1990; Robardet 1996) may explain comparable facies evolution in the Saxothuringian areas (Linne-mann and Buschmann 1995), in the Armorican massif (Paris et al. 1999), and regions included in the polymetamorphic domains (e.g. Mingram 1996, 1998). Fossil findings confirm the presence of lower Palaeozoic metasediments even in Variscan high grade metamorphic rocks (e.g. Schwarzwald, Hanel et al. 1999) and in low-grade Alpine metamorphic series from the Southern Alps (Ordovician, Gansser and Pantic 1988) and Eastern Alps (Arenig, Grauwackenzone; Reitz and Höll 1991).

It is tempting to correlate the Cambrian period of rifting (see above) and the distribution of relics of oceanic domains. From the Late Precambrian to the Ordovician different generations of oceanic domains were involved in the plate-tectonic evolution. Late Precambrian oceanic crust and Cambrian volcanic arcs (Haydoutov 1989; von Quadt 1992; Schaltegger et al. 1997; Schaltegger and Gebauer 1999; Chen et al. 2000; Crowley et al. 2000; Fernández Suárez et al. 2000) confirm the relationship of several domains with the Neoproterozoic orogenic belt and the existence of different tectonic regimes along the Gondwana margin. Traces of what is assumed to represent the former

time equivalents of the Rheic ocean are found in the Cambrian mafic-ultramafic complexes such as the Chamrousse ophiolite of the Alps (Ménot 1987) and the Vesser zone of the Saxothuringian domain (H. Kemnitz et al., submitted), and can be followed through different metamorphic complexes in the Bohemian massif (Finger et al. 1998; Mingram et al. 1998; Stipska et al. 1998; Winchester et al. 1998; Kröner et al. 2000). Relics may occur in the French Central Massif (Pin 1991), and exist in the Iberian basement areas (Fernández-Suárez et al. 1998; Ordoñez Casado 1998; Abati et al. 1999; Andonaegui et al. 1999), and in the northern part of Sardinia (Carmignani et al. 1994). However, it cannot be excluded that relics of Cambro-Ordovician oceanic crust may represent one or more parallel “Rheic” rifts separated by ribbons of “Cadomian” or “Avalonian-type” basement areas (microcontinents) and, additionally, metabasites partly may represent either the Ordovician–Silurian successors of the Rheic ocean or time equivalents of the Proto-Tethys.

Comparing the peri-Gondwanan evolution

From the Neoproterozoic to the Ordovician detachment of Avalonia from Gondwana, striking parallels of plate-tectonic evolution become visible (Fig. 3) between Avalonia, Armorica and the peri-Gondwanan microcontinents located farther to the east, confirming in some way the lateral alignment of the microcontinents at the Gondwana margin. The rifting period accompanying the opening of Iapetus certainly left structures on either side of the future Iapetus ocean. Such intraplate rift systems may have guided the emplacement of Precambrian/Cambrian volcanites and intrusions, and also initial formation of oceanic crust in basement areas located at the eastern prolongation of future Avalonia. A general situation of Neoproterozoic active margin at the Gondwana border (see above) is documented (e.g. Iberia: Fernández-Suárez et al. 1999; Bohemian Massif: Zulauf et al. 1999, Kröner et al. 2000; Saxothuringian domain: Linnemann et al. 2000; in the Alps: Neubauer 1991; von Quadt 1992; Schaltegger et al. 1997), and the picture of widespread late Proterozoic sediments and volcanic rocks, and the formation of Cambrian sedimentary troughs and initial rift zones, become apparent for the Gondwana margin. The continuation of rifting, initially in a back-arc situation, triggered upwelling of asthenosphere, with contemporaneous Cambrian gabbros, granitoids and metamorphic conditions of lower crust (Abati et al. 1999) which announce the future opening of Rheic ocean. Comparable observations come from the Gotthard (Biino 1994) and Silvretta areas (Schaltegger et al. 1997). Pieces of oceanic crust, such as the Chamrousse area (Western Alps) or the Vesser zone (Saxothuringian zone), could represent the eastern continuation of the Rheic ocean (or pieces

of oceanic crust contemporaneous to the Rheic ocean), and corresponding highly transformed relics of the same could be hidden among the many pieces of “amphibolites” found in the polymetamorphic areas of the Variscan mountain chain, indicating the former suture of an eastern continuation of an early Rheic ocean. Also the eclogitized remnants of Early Palaeozoic oceanic crust (Pulo do Lobo, Cabo Ortegal; Ordoñez Casado 1998) have to be incorporated in the general pattern of the Rheic ocean. Interestingly, Kleinschrodt and Gayk (1999) underlined the striking parallels of structural evolution and exhumation ages of the eclogite complexes of the Münchberg Massif and from Cabo Ortegal, former protoliths of the Rheic ocean.

Also Ordovician events known from the Avalonian plate have their contemporaneous events in the adjacent microcontinental plates. During the Ordovician break-up of Avalonia (485 Ma, Van Staal et al. 1998; East Avalonia, 480–465 Ma, Prigmore et al. 1997), the birth of the Rheic ocean, contemporaneous rift-related magmatism is observed in NW Spain (Fig. 4; Santos Zalduegui et al. 1995; Ordoñez Casado 1998; Fernandez-Suarez et al. 1999; Montero et al. 2000; Valverde-Vaquero and Dunning 2000), in the French Central Massif (Pin and Marini 1993) and in the subterranean of the Saxothuringian domain (Linnemann et al. 1998a). Additional criteria for appreciating the general plate-tectonic situation are furnished by the widespread granitoid rocks of Early Palaeozoic age, by the evidence of an Ordovician orogenic event, and by the discussion of the possible continuation of Avalonian pieces preserved as relics in the former eastern continuation of Avalonia.

Parallels of granitoid evolution

During the plate-tectonic evolution of Avalonia (e.g. rifting related to the opening of Iapetus), accompanied by several magmatic cycles (e.g. early Cadomian, 580–570 Ma; Neoproterozoic, 550–540 Ma; Cambro-Ordovician, 500–480 Ma), and its break-off from Gondwana (formation of Rheic ocean, Cambro-Ordovician, 500–480 Ma), contemporaneous tectonic events and corresponding pulses of granitoids or their surface equivalents can be observed (Fig. 3) in most of the adjacent peri-Gondwanan continental blocks. Most Ordovician granitoids and acidic volcanic rocks from the Alps and areas outside the Alps (Fig. 4, 5), Avalonian outcrop areas (Tremblay et al. 1994; Keppie et al. 1997), the Himalayan domain (Girard and Bussy 1999), the Argentine Precordillera (Huff et al. 1998), and even from many Late Cadomian granitoids, independent of locality and age, show strong anomalies of Ba, Nb, Sr and Ti in mantle-normalized multi-element variation diagrams (von Raumer et al. 1999), and Late Proterozoic to Early Palaeozoic detrital sediments (e.g. Bollin 1994; Mingram 1996; Ugidos et al. 1997)

Fig. 4 Ages of Cambro-Ordovician granitoids and metamorphism

Alps	Helvetic Realm	
438±5 Ma U/Pb SHRIMP	meta-dacite, Arg.	Lombardo et al. 1997
489±22 Ma U/Pb Zrn	granite, Bell.	Barfy et al. 1997
496±6 Ma U/Pb Zrn	plagiogranite, Bell.	Menot et al. 1988:
453±3 Ma U/Pb Zrn	granitoids, Mt. Blanc	Bussy & von Raumer 1994
436±17 Ma Rb/Sr wr	granitoids, Gotthard	Arnold 1970
>440 Ma U/Pb Zrn	" "	Bossart et al. 1986
439±5 Ma U/Pb Zrn	" "	Sergeev & Steiger 1993
445±2 Ma U/Pb Zrn	migmatites, Aar massif	Schaltegger 1993
456±2 Ma U/Pb Zrn	granitic gneiss, Si.	Koppel et al. 1981
±450 U/Pb Zrn	eclogite, Gotthard	Gebauer et al. 1988:
461 Ma Sm/Nd grt wr	eclogite, Gotthard	
468 Ma U/Pb Zrn	island arc gabbro, Gotthard	Oberli et al. 1994:
467-475 Ma U/Pb Zrn	+ eclogite facies	
479±3 Ma U/Pb Zrn	island arc gabbro	Abrecht et al. 1995:
475-450 Ma U/Pb Zrn	metabasic protoliths	Paquette et al. 1989:
424 Ma U/Pb Zrn	eclogite, Argentera	Paquette et al. 1989:
	Penninic Realm	
507±9 U/Pb Zrn	granophyre	Guillot et al. 1991
507±9 U/Pb Zrn	Vanoise nord, granite	Guillot et al. 1998
482±5 SHRIMP	Pecllet granite	Guillot et al. 1998
451±5 SHRIMP	Modane granite	Guillot et al. 1998
500±10 SHRIMP	metarhyolite, Ambin	Guillot et al. 1998
500±4 U/Pb Zrn	alkaline granite	Bussy et al. 1996
497±4 U/Pb Zrn	Vanoise, granite	Bertrand & Leterrier 1997
±450 U/Pb Zrn	granitic gneiss, Berisal	Koppel et al. 1981
458±11 U-Pb SHRIMP	anatexis, Tauern	Eichhorn et al. 1999
	Austroalpine Realm	
420-450 U/Pb Zrn	Fluela-granite, Silv	Liebetau 1996
451±2 Rb/Sr WR	Fluela-granite, Silv	Flisch 1987
490±9 U/Pb EvZrn	Otztal, migmatite	Klotzli-Chovanetz et al. 1997
445±24 Rb/Sr WR	granitoid gneiss, TS.	Hammerschmidt 1981
494±73 Sm-Nd WR	Otztal meta-granite	Hoinkes et al. 1997
487±5 Sm-NdTit WR	Otztal meta-granite	Bernhard et al. 1996
485 Pb-Pb EvZrn	Otztal meta-granite	Bernhard et al. 1996
468 Ma U/Pb Zrn	Porphyroid Greywacke zone	Sollner et al. 1991
476 ±14 Ma U/Pb SHRIMP	Metagabbro Greywacke zone	Loth et al. 1999
	Southalpine Realm	
466±5 Rb/Sr WR	Ceneri gneiss	Boriani et al. 1981
457±7 U/Pb Zrn		Zurbruggen et al. 1997
457-450 U/Pb, Pb/Pb	Staur. formation	Romer et al. 1996
Ordovician, stratigraphy	Comelico-Porphyroid	Romer & Franz 1998
		Hubich & Loeschke 1993
	Saxothuringikum	
479±2 Ma ev Zrn	Barentiegel Porphyroid	Linnemann et al. 2000
487±6 Ma ev Zrn	Blambach Rhyolite	Linnemann et al. 2000
484±5 Ma ev Zrn	rhyolitic tuff Wurzelberg	Linnemann et al. 2000
485±6 Ma ev Zrn	Tourmaline granite	Linnemann et al. 2000
486±4 Ma ev Zrn	Metarhyolite	Linnemann et al. 2000
≈490 Ma ev Zrn	rhyolite pebbles	Linnemann et al. 2000
	Bohemian Massif + Schwarzwald	
±480	ZEV, pegmatoids, metamorphism	O'Brien et al. 1997
487±.8 Ma U/Pb SHRIMP	Metaeklogite	O'Brien and Kroner 1999
482±6 SHRIMP Zrn	Gfohl gneiss	Friedl et al. 1998
≈480 Ma Rb/Sr recalc	thermic event	Chen et al. 2000
501-504 Ma U/Pb EvZrn	granitoids	Kroner et al. 2000
469.3 ±3.8 Ma SHRIMP Zrn	inherited magm. Grains	Kroner et al. 2000
481.7±1.1 Ma Pb/PbEv Zrn	Frankenstein gneiss	Kroner and Willner 1998
485±12 Ma Pb/Pb EvZrn	Metarhyolith Ehrenfriedersdorf	Mingram & Rotzler 1999
480±12Ma Pb/Pb EvZrn	Augengneiss Schwarzenberg	Mingram & Rotzler 1999
	Polish Sudetes	
480-505 U/Pb Zrn	granitoids	Oliver et al. 1993
473-488 U/Pb EvZrn	Gory Sowie	Kroner & Hegner 1998
440 U/Pb EvZrn	anatectic granitoid	
	NWiberia +Ossa Morena	
500±2 U/Pb Zrn	orthogneiss	Abati et al. 1999
484, 493 U/Pb Zrn	migmatization	Gonzalez Cuadra et al. 1999
465-480 U/Pb Zrn	Olio de Sapo granitoid	Fernandez-Suarez et al. 1999
479.5 Ma U/Pb Zrn	meta-granodiorite	Santos Zalduegui et al. 1995
480±2 Ma U/Pb Zrn	Cardoso gneiss	Valverde-Vaquero & Dunning 2000
502±8MaPb/Pb Zrn	Castillo alkaline granite	Montero et al. 2000
480±7Ma Pb/Pb Zrn	Valuengo microgranite	Montero et al. 2000
	French Massif Central	
478±6 Ma U/Pb Zrn	orthogneiss, "leptynite"	Pin and Lancelot 1982
	Hungary	
462±6 Pb/Pb Ev Zrn	granitic gneiss, Sakarya	Ozmen and Reischmann 1999
	Turkey	
Ord. age, fossils	granite pebbles, Tisia	Kozur 1984 a,b




Fig. 5 Distribution of Ordovician granitoids in the Alpine domain – a tentative Silurian projection. Granitoids represent different stages of evolution of polymetamorphic domains at an active margin, and in relation with an Ordovician amalgamation, whereas late Ordovician volcanites represent the nearby situation of the opening Palaeotethys. Helvetic domain: *Aar* Aar massif; *Ad* Adula; *AR* Aiguilles Rouges massif; *Arg* different pieces of the Argentera massif, *Bell* Belledonne–Grandes Rousses; *Go* Gotthard massif; *L* Lebedun; *MB* Mont-Blanc; *ML* Monte Leone; *Si* Simano; *Tau* Tauern window units; *Ve* Verampio basement; *Ver* Carboniferous in the Verrucano unit. Penninic units: *Ac* Aceglgio unit; *Lig* different pieces in the Ligurian Alps (new Ordovician age data, oral comm. M. Molina, Genova); *SB* St. Bernhard nappe; *Su* Suretta; *Ta* Tambo; *Va* Vanoise unit; *ZH* Zone Houillère. Austroalpine units: *Am* Ambin; *Ca* Campo; *Gl* Gleinalm; *DM* different pieces in the Dora Maira unit; *EB* Err-Bernina units; *Gle* Gleinalm; *GP* Gran Paradiso; *Gr* Grobgneis unit; *GrP* Graz Palaeozoic; *Iv* Ivrea-Zone; *Ko* Koriden; *MR* Monte Rosa; *Mu* Murau nappe; *Oe* Oetztal; *Sa* Saualm; *Sil* Silvretta nappe; *St* Stangalm; *Sch* Schladminger Tauern; *Ve* Veitsch units; *We* Wechsel unit. Southalpine units: *Bo* Bozen; *DB* Dt. Blanche; *Iv* Ivrea zone; *Nö* Nötsch; *KW* Karawanken; *Ka* Karnische Alpen; *Se* Sesia-zone

Guillot et al. 1991, Bussy et al. 1996; Ossa Morena: Montero et al. 2000; Pyrenees: Navidad and Carreras 1995) may have formed in a thickened crust (see also NW Iberia; Santos Zalduegui et al. 1995). Equally, in the Cambro-Ordovician granitoid associations (e.g. Oetztal, Schindlmayr 1999; Silvretta, Liebetrau 1996; External massifs, Wirsing 1997), the relatively early calc-alkaline metaluminous trends can be related to a Cambro-Ordovician active margin setting (Schaltegger et al. 1997; Schermaier et al. 1998), whereas the corresponding relatively younger granitoids (± 450 Ma) have more or less calc-alkaline peraluminous compositions, carrying the characteristics of a late- to post-orogenic evolution. When plotted on a Rb/Y+Nb diagram (Pearce et al. 1984), older granitoids of Ordovician age in the Alps plot in the fields of VAG or intra-plate situation.

Post-Cambrian evolution of microcontinents

show similar normalized patterns indicating chemical homogeneity over wide areas. Such large-scale parallels, which may have resulted from comparable source areas of granitoids or sediments, show that normalized multi-element variation diagrams do not discriminate the plate-tectonic situation of the many granitoids.

Comparing trace elements, Linnemann et al. (2000) show that Cadomian detrital sediments of the Saxothuringian zone indicate the situation of a continental island arc, whereas Ordovician sediments carry the fingerprints of a passive margin. In geochemical plots (Maniar and Piccoli 1989), the Cambro-Ordovician granitoids follow, in some ways, distinct steps of a plate-tectonic evolution (Fig. 6). Local occurrences of Cambrian plagiogranites (Alps: Ménot et al. 1988, Müller et al. 1996; NW Iberia, Espasante) document evolution of oceanic crust, and Cambrian granitoids of more alkaline and metaluminous compositions (Alps:

The presence of Cambro-Ordovician active margin setting, Ordovician rifting and the evolution of Cambro-Ordovician granitoids raises the question of the existence of an orogenic event characterizing part of the Gondwana margin. The presence of so many Ordovician granites and their pre-Variscan organization (Fig. 5) and geochemical evolution leads, necessarily, to the question of their plate-tectonic situation (Schmidt 1977; Ebner et al. 1987; Biino 1995; Oberli et al. 1994; Abrecht et al. 1995; Handy et al. 1999; O'Brien et al. 1997). Ordovician ages of granitoids (Fig. 4, with references), of eclogites (Nussbaum et al. 1998, not precise age; Gebauer et al. 1988; O'Brien and Kröner 1999) as well as anatexis and metamorphism (Schaltegger 1992, 1993; Klötzli-Chovanetz et al. 1997; Ordovician subduction, Biino 1994; O'Brien et al. 1997; Zurbruggen et al. 1997; Weger et al. 1999; staurolite formation, Romer and Franz 1998)




Fig. 6A, B Distribution of Early Palaeozoic granitoids at the peri-Gondwanan margin (nomenclature and reconstructions: see Fig. 7). A Early Ordovician reconstruction (Fig. 7A) with location of granitoids emplaced before the opening of the Rheic ocean. B A Silurian projection (Fig. 7B) with granitoids and metamorphism related to the Ordovician orogenic event and to the opening of Palaeotethys

argue for the orogenic evolution from an active margin setting to a collision during a short time period (e.g. Gotthard massif, 5–20 Ma, Oberli et al. 1994; 467- to 475-Ma intrusion of metagabbros/440-Ma metamorphism of Streifengneis granitoids). Carmignani et al. (1994) discuss an evolution from Cambrian rift to an Ordovician Andean type margin for Sardinia. The coincidental evolution of certain types of granitoids through time mentioned previously confirms a plate-tectonic evolution including collisional and post-collisional events. We interpret the tectonic assemblage of Cadomian relics, Neoproterozoic arc setting, and/or Late Proterozoic to Early Palaeozoic sediments and oceanic crust, and Ordovician orogenic structures as indicating an Ordovician suture zone, where these elements were amalgamated to Gondwana during the Ordovician (Figs. 7, 8).

Ordovician palaeo-oceanographic reconstructions combine palaeocurrent data (Christiansen and Stouge 1999), palaeomagnetic criteria (MacNiocaill et al. 1997; Torsvik and Smethurst 1996) and faunal distributions (Harper et al. 1996; Paris 1993; Paris et al. 1999; Mélou et al. 1999) defining the Iapetus and Rheic oceans domains. The wide domain of Iapetus is characterized, since the Early Ordovician, by faunal provinces of brachiopods, where the peri-Gondwanan Avalonian border has cold-water Celtic faunas (Harper et al. 1996; Cocks 2000), and volcanic arc activity (MacNiocaill et al. 1997). Characteristics of Chitinozoan faunas give an idea about the early evolution of the Rheic ocean (Paris 1993; Paris et al. 1999) and, during the same time period, detrital sediments record maximum of subsidence and rift basin formation, accompanied by rift volcanism, in lateral continuity (e.g. Saxothuringian domain; Linnemann 1999).


Sedimentary basins, constituting the lower Palaeozoic cover series in the Eastern Alps, contain acidic volcanics among lower Palaeozoic sediments (Loeschke and Heinisch 1993) with a well-documented sedimentary evolution (Schönlaub 1997a, 1997b; Schönlaub and Heinisch 1993; Schönlaub and Histon 2000), representing a rifting environment since the Late Ordovician (Neubauer and von Raumer 1993; Neubauer and Sassi 1993; Sassi et al. 1994). Since the Late Ordovician, many areas are covered by the detrital sediments of Ordovician glaciation (e.g. Hamoumi 1999; Linnemann et al. 1999). It is only after the onset of the Avalonian drift that Armorica and the more easterly situated Intra-Alpine blocks followed a more independent plate-tectonic evolution.

Terrane detachment from Gondwana

Detachment from Gondwana started during the Mid-Ordovician (ATA; Tait et al. 1998), Caradoc/Ashgill (Saxothuringian domain, 445 Ma; Fig. 3; Linnemann et al. 1998b), and the latest Ordovician (Alpine domain, Ashgill/Llandovery, 435 Ma; Fig. 3; Stampfli 1996) when the Palaeo-Tethys rift opened. Sedimentological and faunistic criteria, and palaeomagnetic data, gave rise for models of detachment of Gondwana-derived continental blocks during the Ordovician and Silurian, favouring either the separation of a basement assemblage (Stampfli 1996) or a major mid-European oceanic separation (Robardet et al. 1990).

Stampfli (1996) introduced the Hun-Terrane model, a ribbon-like assemblage of continental blocks, not necessarily above sea level, or volcanic islands, including Armorica, the Intra-Alpine basement and microcontinents situated in the eastern prolongation. The observation of subsidence patterns in former peri-Gondwanan domains induced Stampfli (2000), and Stampfli and Mosar (1996), to reconstruct the contours of Palaeo-Tethys during the Early Silurian for the Intra-Alpine domains and the corresponding eastern prolongations. Prior to their Silurian separation from Gondwana, these Variscan terranes were located

Fig. 7 **A** Location of pre-Variscan basement units at the Gondwanan margin during the Early Ordovician (490 Ma), modified from Stampfli (2000), showing the early stages of the Rheic ocean spreading. **B** A Silurian projection (420 Ma), modified from Stampfli (2000), showing the early stages of opening of PalaeoTethys. Baltica: *Is* Istanbul; *Mo* Moesia; *Zg* Zonguldak. Avalonia: *MM* Meseta-Meguma. Cadomia: *AA* Austro-Alpine; *Cm* Cadomia s.str.; *He* Helvetic; *Lg* Ligerian; *MD* Moldanubian; *OM* Ossa-Morena; *Pe* Penninic; *SM* Serbo-Macedonian; *Sx* Saxothuringian. Serindia: *KT* Karakum-Turan; *Tn* north Tarim. Gondwana: *Ab* Alboran; *Ad* Adria; *Al* Alborz; *Am* Armorica; *Ap* Apulia; *Aq* Aquitaine; *Ct* Cantabria; *DH* Dinarides-Hellenides; *iA* intra-Alpine; *Ib* Iberia; *Ki* Kirshehir; *LT* Lut-Tabas; *Mn* Menderes; *Pr* Pamir; *SS* Sanadaj-Sirjan; *Ta* Taurus; *Ts* south-Tarim




along the margin of Gondwana and represent the future northern margin of Palaeo-Tethys. Consequently, a tentative Silurian palaeo-tectonic configuration has been derived by back-modelling the present day distribution of Variscan elements in Europe (Fig. 7), including also the basement areas of Armorica assumed to contain relics of a former rifting period since the Ordovician.

The corresponding palaeo-reconstructions are based on palaeomagnetic pole data and have been developed with the GMAP software package (Torsvik and Smethurst 1994; Torsvik and Smethurst 1999) using an orthographic (orthogonal) projection. The Early Ordovician (490 Ma) projection is centred on present-day latitude 60°N , longitude 80°E , the Late Silurian (420 Ma) projection is centered on present-day latitude 10°N , longitude 25°E . The palaeomagnetic pole data used for Baltica are from Torsvik et al. (1994)

for the Hun terrane elements and Siberia from Van der Voo (1993), and references therein. The position of Gondwana is constrained by the Australia data from Klootwijk (1996) and by Pangea tight-fit reconstruction. The Baltica poles are the absolute reference poles for all the reconstructions.

As a result of Ordovician to Early Devonian rifting events, the components of the Hun terrane were stepwise detached from Gondwana accompanying the opening of Palaeo-Tethys. Such reconstructions are not too different from palaeomagnetic reconstructions for the Armorican Terrane Assemblage (ATA; Tait et al. 1998), where distinct groups of microcontinents started to separate from Gondwana. The principle of a Hun-Terrane, which is grouping of certain microcontinental pieces, may thus be modified acknowledging a still more detailed individual separation from Gondwana. If we admit an opening of Palaeo-Tethys in a

Fig. 8 Cross sections model to illustrate the possible plate tectonic evolution between Baltica and Gondwana during the Early Palaeozoic. The Hun-superterrane resulted from the Ordovician collisional assemblage of elements possibly detached from Baltica-Siberia and areas comprising Cadomian basement (former eastern prolongation of Avalonia), Early Palaeozoic oceanic crust (Rheic), volcanic arcs and accretionary wedges, with the Gondwana margin. The opening of the Palaeotethys separated the composite Hun superterrane from Gondwana. This opening took place in a context of back-arc spreading of Palaeotethys and slab roll-back of the Mauritanian-Rheic ocean and was accompanied by the formation of volcanics



back-arc setting, the geometry of the detached upper plate is bound to be modified by differential slab roll-back behaviour of the lower plate as is observed presently along the Mariana trench, for example. It is clear that the opening of Palaeo-Tethys is diachronous (Stampfli 2000); however, subsidence curves alone cannot determine the onset of sea-floor spreading when very little Palaeotethyan sea-floor remnants are known. We propose a detachment of Gondwanan elements in three major steps: firstly, the Avalonia blocks; then, the mainly European Hun Terranes; and finally the Far-East Hun terranes. We assume that the latter could not be detached before the Silurian accretion to Gondwana of some exotic (Baltica-Siberia) elements grouped here as Serindia. There, the situation is quite different, amalgamation of terranes happened only in Silurian or Early Devonian times and larger continental fragments are involved such as the Pamir-south Tarim-Tsaidam block collision with the north Tarim-north China block (Yin and Nie 1996) or the north Qinling north China blocks collision (Meng and Zhang 1999). These drifting exotic blocks (Serindia terrane) followed the slab roll-back of the Proto-Tethys accompanied by the Asiatic ocean spreading in a back-arc position. An older western portion of the Asiatic ocean was involved in intra-oceanic subduction accompanied by the opening of the Khanti-Mansi back-arc ocean and development of the Kipchak arc (Sengör and Natal'in 1996). A southward connection between the Asiatic ocean mid-oceanic ridge and a remnant Proto-Tethys mid-oceanic ridge is suggested here. This Silurian collisional event affected also the

South China block and part of the Indochina block (or the north Qinling north China blocks collision; Meng and Zhang 1999; Findlay 1997) and possibly has been recognized westward up to eastern Turkey (Göncüoğlu and Kozur 1999).

Scenario of peri-Gondwanan evolution: a conclusion

When summing up the observations mentioned previously, we arrive, independently of Variscan tectonic complications, at the conclusion that most of pre-Variscan relics of the Variscan mountain chain had their origin at the Gondwana margin. Two evolution trends are noted:

1. Stampfli (2000) favoured the derivation of Cadomian basement blocks from Baltica triggered by a Baltica-directed subduction and island arc formation during the Early Ordovician which, during the Mid Ordovician, collided with the Gondwana margin (orogenic event). Inversion of subduction towards Gondwana produced the break-off of the Hun-superterrane and formation of Palaeo-Tethys as a back-arc ocean.
2. In our present model an opposite evolution is proposed for the Cadomian terrane. Assuming that already since the latest Precambrian subduction was generally directed towards Gondwana (Cadomian orogeny, Avalonian southern prong), we assume that most of the Cadomian basement blocks (exception Belka et al. 1998, 1999; Unrug et al. 1999) presently hidden in the Variscan mountain

chain should have derived from the Gondwana margin ("European margin"; Courjault-Radé et al. 1992). In this general picture, the Serindia terrane is still considered as derived from Baltica–Siberia.

Since the latest Cambrian (500 Ma) at several locations (see above) magmatic and metamorphic events are dated, presumably the consequence of general crustal extension and contemporaneous influence of rising asthenosphere. Such activity is accompanied, at several distant places, by formation of oceanic crust interpreted to represent the opening of the Rheic ocean. The alignment of the very different localities at the Gondwana margin suggests a contemporaneous opening over a large distance, from Ossa Morena, and Iberia to the Münchberg Massif. The question arises as to whether such oceanic trenches represent one uninterrupted long suture line, or if there existed parallel lines of opening (e.g. Rheic, Galicia-Massif Central; Matte 1986), compared with the parallel lines of Early Cambrian grabens intersecting the Gondwanan block. The large differences of oceanic crust at the different localities may indicate that oceanic crust did not result from cylindrical subduction and simultaneous slab roll-back, but from very oblique convergence, where either pull-aparts may have facilitated the appearance of true oceanic crust, accompanied laterally by only incomplete oceanic series, or simply volcanites and/or detrital sediments. Such strike-slip models have been proposed by Murphy and Nance (1989), Linnemann et al. (1998b) and Zulauf et al. (1999).

The striking parallels of evolution from the Late Neoproterozoic to the earliest Ordovician are best explained when using a ribbon-like lining-up model of pre-Variscan blocks along the Gondwana margin. Most of the continental pieces contain the relics of a Neoproterozoic active-margin setting and continental rift since the Late Cadomian, and early stages of Rheic ocean. Intrusion of granitoids and volcanites follow the regional tectonic pattern. We thus claim a common early evolution of Avalonia and its eastern continuation (e.g. Cadomia *sensu lato*, Intra-Alpine Terrane) at the Gondwana margin, and major differences of evolution between Avalonia and the remaining eastern microcontinents only after the successful break-off of Avalonia during the Early Ordovician (Figs. 3, 7, 8).

The independent drift story of the microcontinents situated in the eastern prolongation of Avalonia depended on the time of subduction of a remnant Proto-Tethys/Rheic mid-oceanic ridge; the latter was linked with the Asiatic ocean mid-oceanic and transform zone in between. The subsequent subduction/collision of the ridge/transform zone triggered the amalgamation of arc and continental ribbon with Gondwana leading to the Ordovician orogenic evolution. This collisional event produced a complex terrane configuration, the "Noric composite Terrane" (Eastern Alps; Frisch and Neubauer 1989) or "Hun superterrane" (Stampfli 1996). It comprises older

(former Gondwanan) basement blocks, accretionary wedges, volcanic arcs, and pieces of an aborted eastern Rheic ocean, and may have, locally, the aspect of an archipelago, but it also can be composed by large continental blocks. It is evident that there has been tectonic interaction between the different elements, thus complicating the orogenic assemblage, and our proposed distribution of terranes (Fig. 7) is only tentative. As traces of the Ordovician orogenic event are seen mainly in the Intra-Alpine domain, with some parallels in the Bohemian massif (Cambrian and Ordovician zircon ages of eclogites; O'Brien and Kröner 1999), the question arises as to whether the Armorican Terrane Assemblage (Tait et al. 1998) and more western situated microcontinents were really independent from the more eastern ones, and that the assemblage of Bohemian massif and the Intra-Alpine terranes had a distinct drift history, interrupted by this short event, or if the Devonian lower crustal events observed in the more western parts have completely erased the memory of such events. The evidence of Early Ordovician volcanites and/or granites, sometimes of alkaline tendencies, in the more western blocks just indicates the rifting related to the break-off of Avalonia.

The constituents of the future Hun-composite terrane are interpreted to represent an eastern prolongation of Avalonia (Figs. 3, 7), and the suture zone hidden in the superterrane are interpreted to represent the eastern prolongation of an early stage of the Rheic ocean (see also Finger et al. 1998), before the Ordovician amalgamation (Figs. 7, 8). This amalgamation would have been the result of a diachronous consumption of the mid-ocean ridge of Proto-Tethys which triggered the intrusion of late Cadomian granitoids and consequently the early separation (480 Ma) of Avalonia from its neighbouring blocks following the onset of generalized slab roll-back of the remaining ocean. The opening of the eastern branch of the Rheic was stopped by the contemporaneous spreading of the Asiatic ocean (Zonenshain et al. 1985) and partial collision with a major transformation zone at the southern end of the exotic Karakum–Tarim–north China terrane. It is only since the break-off of Avalonia that the plate-tectonic evolution of the more eastern-situated microcontinents (e.g. Cadomia, Intra Alpine Terrane) became different, and the following events could have been the consequence of accretion and shifting of subduction zones to external domains, as discussed for the contemporaneous evolution at the Pacific Gondwana margin (Rapela et al. 1992; Tessensohn 1999).

The time period after the Ordovician amalgamation is characterized by the successive stages of subduction, and the contemporaneous formation of accretionary wedges and arc-volcanics at the Silurian active margin of Gondwana. This is supported by Silurian eclogite ages from the Tauern area (von Quadt et al. 1997) and comparable ages from the Argentera External

Massif (Paquette et al. 1989), and the Massif Central domain (e.g. Ducrot et al. 1983; high-pressure event: Pin and Lancelot 1982). Relics of Silurian volcanic arcs have been recognized in the domain of the North German Crystalline Rise, interpreted as an active margin setting in the South of the Rhenohercynian domain (Altenberger and Besch 1993; Reischmann and Anthes 1996a, 1996b; Zeh et al. 1997), and in our model the active margin is located on the northern border of the Hun-superterrane (Figs. 7, 8). The diachronous Palaeo-Tethys opening in Silurian-Early Devonian times is viewed as a back-arc opening along the Gondwana margin (Stampfli et al. 1991; Stampfli 2000). Faunistic assemblages (Schönlaub 1997a, 1997b; Schönlaub 1993; Schönlaub and Histon 2000) and palaeomagnetic data (Bachtadse et al. 1995; Schätz et al. 1997; Tait et al. 1998; Torsvik et al. 1994; Torsvik and Eide 1998) support the migration of the Hun superterrane towards Laurussia. It is interesting that Noblet and Lefort (1990) admit in their lower Ordovician stratigraphy a “future opening of the Tethys I ocean”, and that in the Silurian palaeogeographic reconstruction (Paris and Robardet (1990) admit an opening oceanic domain as a branch of the Rheic ocean for the more eastern-situated domains of the large Gondwana shelf.

Difficulties arise when comparing drift models of continental pieces since the Silurian. Matte (1986) and Robardet (1996) propose the migration of Gondwana towards Laurussia, whereas Tait et al. (1998), based on palaeomagnetic data, discuss the migration of individual microcontinents or groups of continents, coinciding with predictions based on faunistic criteria (Schönlaub 1993), and such models could be correlated with a modified model of the Hun-superterrane and its migration (forthcoming discussion given by G.M. Stampfli et al., submitted).

Considering the parallels of pre-Variscan evolution of the different microcontinents preserved in the Variscan domain (Fig. 3), it is tempting to propose a nomenclature for the classical tectono-stratigraphic realms, taking into account the former geological evolution at the Gondwana margin. The assemblage of Cadomian basement, Neoproterozoic to Cambrian active margin setting, and Ordovician accretionary stages represent the leading edge of the Hun-superterrane and includes a Middle Ordovician suture of aborted Rheic ocean. Large areas of the Moldanubian, of the French Central Massif and from NW Iberia can be grouped under this assemblage. The units, characterized by the opening of the Paleo-Tethys which separated the Hun super-terrane from Gondwana in Late Silurian times, belong to the classical Saxothuringian domain and the southern part of Armorica s.l. As a consequence of the Variscan evolution since the Silurian (G.M. Stampfli et al., submitted), the classical Rhenohercynian zone together with the Lizard domain, and the South Portuguese and Moravo-Silesian zones, opened as a Devonian oceanic domain

within the Laurussia margin, due to Gondwana-directed slab pull. The contemporaneous leading edge of the Hun-superterrane is represented by the Silurian-Devonian active margin (part of the mid-German Crystalline Rise).

Although the proposed models do not contradict palaeoreconstructions in terms of palaeomagnetic information and faunal distributions mentioned previously, an interdisciplinary discussion is still needed to better understand plate-tectonic evolution during Early Paleozoic times. Important information on key areas are still lacking, and we view this paper as stimulation for further discussion and research, which has started already through the stimulating contribution from the reviewers. Little is known about the larger distribution of the Ordovician orogenic event, and precise age determinations of magmatic rocks or refined provenance studies of sedimentary rocks, even of high-grade metamorphic rocks, are greatly needed.

Acknowledgments We sincerely thank W. Franke (Giessen), D. Harper (Kopenhagen), E. Hegner (München), A. Kröner (Mainz), U. Linnemann (Dresden), M. Molina (Genova), D. Nance (Athens), F. Paris and M. Robardet (Rennes), P. Ryan (Galway), and Heinz Kozur for providing unpublished data or manuscripts or stimulating discussions. P. Matte (Montpellier), J. Mosar (Oslo), B. Murphy (Antigonish) and H.P. Schönlaub (Vienna) are thanked for their engaging reviews and the many suggestions which helped considerably to clear up differences of interpretations.

References

- Abati J, Dunning GR, Arenas R, Diaz Garcia F, Gonzalez Cuadra P, Martinez Catalan JR, Adonaegui P (1999) Early Ordovician orogenic event in Galicia (NW Spain): evidence from U–Pb ages in the uppermost unit of the Ordenes Complex. *Earth Planet Sci Lett* 165:213–228
- Abrecht J, Biino G, Schaltegger U (1995) Building the European continent: late Proterozoic–Early Palaeozoic accretion in the Central Alps of Switzerland. *Terra Nova* 5 Abstr (Suppl I):105
- Altenberger U, Besch T (1993) The Böllstein Odenwald: evidence for pre- to Early Variscan plate convergence in the Central European Variscides. *Geol Rundsch* 82:475–488
- Andonaegui P, Gonzalez del Tanago J, Arenas R, Abati J, Martinez Catalan JR, Peinado M, Diaz Garcia F (1999) Tectonic setting of the Monte Castelo gabbro (Ordenes Complex, NW Iberian Massif): evidence for an arc-related terrane in the hangingwall to the Variscan suture. *EUG10 Annual meeting* 11:400
- Arnold A (1970) On the history of the Gotthard Massif (Central Alps, Switzerland). *Eclog Geol Helv* 63:29–30
- Bachtadse V, Torsvik TH, Tait JA, Soffel HC (1995) Paleomagnetic constraints on the paleogeographic evolution of Europe during the Paleozoic. In: Dallmeyer RD, Franke W, Weber K (eds) *Pre-Permian geology of central and eastern Europe*. Springer, Berlin Heidelberg New York, pp 567–578
- Bard JP (1997) Démembrement anté-mésozoïque de la chaîne varisque d'Europe occidentale et d'Afrique du Nord: rôle essentiel des grands décrochements transpressifs dextres accompagnant la rotation-traduction horaire de l'Afrique durant le Stéphanien. *C R Acad Sci Paris* 324:693–704
- Barfély JC, Gidom M, Ménot RP, Debon F, Pecher A, Guillot S, Fourneaux JC, Gamond JP (1997) Notice de la feuille Domène (773)

- Belka Z, Ahrendt H, Franke W, Schäfer J, Wemmer K (1998) Early Palaeozoic accretion of Gondwana-derived terranes at the margin of Baltica: evidence from K/Ar ages of detrital muscovites and faunistic data. *Acta Univ Carolinae. Geologica* 42:211–212
- Belka Z, Ahrendt H, Franke W, Schäfer J, Wemmer K (1999) Accretion history of the Trans-European suture zone viewed from new provenance and biogeography data. *Terra Nostra* 99:60–61
- Bernhard F, Klötzli US, Thöni G, Hoinkes G (1996) Age, origin and geodynamic significance of a polymetamorphic felsic intrusion in the Ötztal crystalline basement. *Mineral Petrol* 58:171–196
- Bertrand JM, Leterrier J (1997) Granitoïdes d'âge Paléozoïque inférieur dans le socle de Vanoise méridionale: géochronologie U–Pb du métagranite de l'Arpont (Alpes de Savoie, France). *CR Acad Sci Paris* 325:839–844
- Biino G (1994) The pre-Late Ordovician metamorphic evolution of the Gotthard-Tavetsch massifs (Central Alps): from Lawsonite to kyanite eclogite to granulite retrogression. *Schweiz Mineral Petr Mitt* 74:87–104
- Biino G (1995) Pre-Variscan evolution of the eclogitized mafic rocks from the Helvetic basement of the Central Alps. *Eur J Mineral* 7:57–70
- Böhm CO (1996) Provenance and pre-Mesozoic evolution of the Lucomagno unit (Central Swiss Alps): a geochemical, isotopic and geochronological approach, Dissertation, ETH Zürich
- Bollin R (1994) Metamorphe Entwicklung und Ursprung der Paragneise (Biotitplagioklasgneise, Plagioklasnotengneise) der Silvretta. Dissertation, Fribourg 180 pp
- Boriani A, Giobbi Origoni E, Del Moro A (1981) Composition, level of intrusion and age of the "Serie dei Laghi" orthogneisses (northern Italy–Ticino, Switzerland). *Rend Soc Ital Min Petrol* 38:191–205
- Bossart PJ, Meier M, Oberli F, Steiger RH (1986) Morphology versus U–Pb systematics in zircon: a high resolution isotopic study of zircon population from a Variscan dike in the Central Alps. *Earth Planet Sci Lett* 78:339–354
- Brown M (1995) The late-Precambrian geodynamic evolution of the Armorican segment of the Cadomian belt (France): distortion of an active continental margin during south-west directed convergence and subduction of a bathymetric high. *Géol France* 3:3–22
- Buschmann B, Linnemann U, Schneider J, Stüss T (1995) Die cadomische Entwicklung im Untergrund der Torgau-Doberluger Synklinale. *Z Geol Wiss* 23:729–749
- Bussy F, Raumer J von (1994) U–Pb geochronology of Palaeozoic magmatic events in the Mont-Blanc Crystalline Massif, Western Alps. *Schweiz Mineral Petrol Mitt* 74:514–515
- Bussy F, Derron MH, Jaquod J, Sartori M, Thélin P (1996) The 500 Ma-old Thyon metagranite: a new A-type granite occurrence in the Western Penninic Alps (Wallis, Switzerland). *Eur J Mineral* 8:565–575
- Carmignani L, Carosi R, Pisa A di, Gattiglio M, Musumeci G, Oggiano G, Pertusati PC (1994) The hercynian chain in Sardinia (Italy). *Geodinam Acta* 7:31–47
- Chen F, Hegner E, Todt W (2000) Zircon ages and Nd isotopic and chemical compositions of orthogneisses from the Black Forest, Germany: evidence for a Cambrian magmatic arc. *Int J Earth Sci* 88:791–802
- Christiansen JL, Stouge S (1999) Oceanic circulation as an element in palaeogeographical reconstruction: the Arenig (early Ordovician) as an example. *Terra Nova* 11:73–78
- Cocks LRM (2000) The Early Palaeozoic geography of Europe. *J Geol Soc Lond* 157:1–10
- Cogné J, Wright AE (1980) L'orogène cadomien. *Int Geol Congr* 26 Sess C6:29–54
- Courjault-Radé P, Debrenne F, Gandin A (1992) Palaeogeographic and geodynamic evolution of the Gondwana continental margins during the Cambrian. *Terra Nova* 4:657–667
- Crowley QG, Floyd PA, Stodra V, Winchester JA, Holland JG (2000) The Marianske-Lazne Complex: remnants of a subducted early Palaeozoic seaway. In: Variscan–Appalachian dynamics: the building of the Upper Palaeozoic basement. *Basement Tectonics* 15:86–89
- Dallmeyer RD, Neubauer F (1994) Cadomian 40Ar/39 Ar apparent age spectra of detrital muscovites from the Eastern Alps. *J Geol Soc Lond* 151:591–598
- Dallmeyer RD, Franke W, Weber K (eds) (1995) Pre-Permian geology of central and eastern Europe. Springer, Berlin Heidelberg New York
- Dallmeyer RD, Neubauer F, Fritz H, Mocanu V (1996a) Variscan vs Alpine tectonothermal evolution of the Southern Carpathian orogen: constraints from 40Ar/39Ar ages. *Tectonophysics* 290:111–135
- Dallmeyer RD, Neubauer F, Fritz H, Müller W, Pana D, Putis M (1996b) Tectonothermal evolution of the internal Alps and Carpathians: evidence from 40Ar/39 Ar mineral and whole-rock data. *Eclog Geol Helv* 89:203–227
- Dalziel WD (1992) On the organization of American plates in the Neoproterozoic and the breakout of Laurentia. *Geol Soc Am Today* 2:237–241
- Desmons J, Compagnoni R, Cortesogno L, Frey M, Gaggero L (1999) Pre-Alpine metamorphism of the Internal zones of the Western Alps. *Schweiz Mineral Petr Mitt* 79:23–39
- Ducrot J, Lancelot JR, Marchand J (1983) Datation U–Pb sur zircons de l'éclogite de la Borie (Haut Allier, France) et conséquences sur l'évolution anté-hercynienne de l'Europe occidentale. *Earth Planet Sci Lett* 62:385–394
- Ebner F, Neubauer F, Statterger K (1987) The Caledonian event in the Eastern Alps: a review. In: Flügel HW, Greclia P, Sassi FP (eds) Pre-Variscan and Variscan events in the Alpine-Mediterranean mountain belts. Alfa, Bratislava, 169–182 pp
- Eichhorn R, Loth G, Höll R, Finger F, Schermaier A, Kennedy A (1999) U–Pb SHRIMP data of pre-alpine rocks of the Tauern Window (Austria): active margin and back-arc found, but where is the adjacent continental crust? *Terra Nostra* 99:79
- Fernández-Suárez J, Gutiérrez-Alonso J, Jenner GA, Jackson SE (1998) Geochronology and geochemistry of the Pola de Allande granitoids (northern Spain): their bearing on the Cadomian-Avalonian evolution of northwest Iberia. *Can J Earth Sci* 35:1–15
- Fernández-Suárez J, Gutiérrez-Alonso J, Jenner GA, Tubrett MN (1999) Crustal sources in Lower Palaeozoic rocks from NW Iberia: insights from laser ablation U–Pb ages of detrital zircons. *J Geol Soc Lond* 156:1065–1068
- Fernández-Suárez J, Gutiérrez-Alonso J, Jenner GA, Tubrett MN (2000) New ideas on the Proterozoic–Early Palaeozoic evolution of NW Iberia: insights from U–Pb detrital zircon ages. *Precamb Res* 102:185–206
- Findlay RH (1997) The Song Ma Anticlinorium, northern Vietnam: the structure of an allochthonous terrane containing an early Palaeozoic island arc sequence. *J Asian Earth Sci* 15:453–464
- Finger F, Quadt A von, Pin C, Steyrer HP (1998) The ophiolite chain along the western Moravo-Silesian plate margin: a trace of the Rheic suture? *Acta Univ. Carolinae Geol* 42:244–245
- Flisch M (1987) Geologische, petrographische und isotopengeologische Untersuchungen an Gesteinen des Silvretta-Kristallins. Dissertation, Univ Bern
- Franke W, Altherr R, Haak V, Oncken O, Tanner D (eds) (2000) Orogenic processes: quantification and modelling in the Variscan belt of Central Europe. *Geol Soc Lond Spec Publ* 179:1–456
- Freyer G, Suhr P (1987) Über Archaeocyathinen-Funde und den lithologischen Aufbau des Unterkambriums im Gebiet von Torgau. *Z Geol Wiss* 15:605–680

- Friedl G, McNaughton N, Fletcher IR, Finger F (1998) New SHRIMP-zircon ages for orthogneisses from the southeastern part of the Bohemian Massif (Lower Austria). *Acta Univ Carolinae Geol* 42:251–252
- Frisch W, Neubauer F (1989) Pre-Alpine terranes and tectonic zoning in the eastern Alps. *Geol Soc Am Spec Paper* 230:91–100
- Gansser A, Pantic N (1988) Prealpine events along the eastern Insubric Line (Tonale Line, northern Italy). *Eclog Geol Helv* 81:567–577
- Gebauer D (1993) The pre-Alpine evolution of the continental crust of the Central Alps: an overview. In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 93–117
- Gebauer D, Quadt A, Compston W, Williams IS, Grünenfelder M (1988) Archaean zircons in a retrograded, Caledonian eclogite of the Gotthard Massif (Central Alps, Switzerland). *Schweiz Mineral Petrogr Mitt* 68:485–490
- Girard M, Bussy F (1999) Late Pan-African magmatism in the Himalaya: new geochronological and geochemical data from the Ordovician Tso Moriri metagranites (Ladakh, NW India). *Schweiz Mineral Petrogr Mitt* 79:399–418
- Göncüoğlu Y, Kozur HW (1999) Palaeozoic stratigraphy and event succession in eastern Toros, Turkey. In: Talent J, Khan F, Mawson R (eds) *International Geological Correlation Program IGCP 421: North Gondwana Mid-Palaeozoic bioevent/biogeography patterns in relation to crustal dynamics*, 11–13, North Ryde (Macquarie University Printery)
- Gonzalez Cuadra P, Martinez Catalán JR, Arenas R, Diaz Garcia F, Abati J, Dunning GR (1999) Polyorogenic evolution of the uppermost units of the Ordenes Complex (NW Spain). *J Conf Abstr* 11:34
- Grunow A, Hanson R, Wilson T (1996) Were aspects of Pan-African deformation linked to Iapetus opening? *Geology* 24:1063–1066
- Guillot F, Liégeois JP, Fabre J (1991) Des granophyres du Cambrien terminal dans le Mont Pourri (Vanoise, zone Briançonnaise: première datation U–Pb sur zircon d'un socle des zones internes des Alpes françaises. *CR Acad Sci Paris* 313:239–244
- Guillot F, Bertrand JM, Pidgeon RT, Leterrier J, Gattiglio M (1998) Géochronologie U–Pb SHRIMP et IDTIMS de zircons des zones internes des Alpes occidentales. 17e RST98 Brest Abstract Volume, p 124
- Gurnis M, Torsvik TH (1994) Rapid drift of large continents during the late Precambrian and Palaeozoic: palaeomagnetic constraints and dynamic models. *Geology* 22:1023–1026
- Hammerschmidt K (1981) Isotopengeologische Untersuchungen am Augengneis vom Typ Campo Tures bei Rain in Taufers, Südtirol. *Mem Ist Geol Min Univ Padova* 34:273–300
- Hamoumi N (1999) Upper Ordovician glaciation spreading and its sedimentary record in Moroccan North Gondwana margin. *Acta Univ Carolinae Geol* 43:111–114
- Handler R, Dallmeyer RD, Neubauer F (1997) $^{40}\text{Ar}/^{39}\text{Ar}$ ages of detrital white mica from Upper Austroalpine units in the Eastern Alps, Austria: evidence for Cadomian and contrasting Variscan sources. *Geol Rundsch* 86:69–80
- Handy M, Franz L, Heller F, Janott B, Zurbriggen R (1999) Multistage accretion and exhumation of the continental crust (Ivrea crustal section, Italy and Switzerland). *Tectonics* 18:1154–1177
- Hanel M, Montenari M, Kalt A (1999) Determining sedimentation ages of high-grade metamorphic gneisses by their palynological record: a case study in the northern Schwarzwald (Variscan Belt, Germany). *Int J Earth Sci* 88:49–59
- Harper DAT, MacNiocaill CM, Williams SH (1996) The palaeogeography of early Ordovician Iapetus terranes: an integration of faunal and palaeomagnetic constraints. *Palaeogeogr Palaeoclimatol Palaeoecol* 121:297–312
- Haydoutov I (1989) Precambrian ophiolites, Cambrian island arc, and Variscan suture in the South Carpathian-Balkan region. *Geology* 17:905–908
- Hegner E, Kröner A (2000) Review of Nd isotopic data and xenocrystic and detrital zircon ages from the pre-Variscan basement in the eastern Bohemian Massif: speculation on palinspastic reconstructions. *Geol Soc Lond Spec Publ* 179:113–130
- Hoffman PF (1991) Did the breakout of Laurentia turn Gondwanaland inside-out? *Science* 252:1409–1412
- Hoinkes G, Thöni M, Lichem C, Bernhard F, Kaindl R, Schweigl J, Tropper P, Cosca M (1997) Metagranitoids and associated metasediments as indicators for the pre-Alpine magmatic and metamorphic evolution of the western Austroalpine Ötztal Basement (Kaunertal, Tirol). *Schweiz Mineral Petr Mitt* 77:299–314
- Hubich D, Loeschke J (1993) Der Comelico-Porphyröid der westlichen Karnischen Alpen (Österreich/Italien): Stratigraphie, Petrographie, Geochemie und Zirkonmorphologie. *Jahrb Geol BA* 136:351–374
- Huff WD, Bergström SM, Kolata DR, Cingolani CR, Astini RA (1998) Ordovician K-bentonites in the Argentine Precordillera: relations to Gondwana margin evolution. *Geol Soc Lond Spec Publ* 142:107–126
- Keppie JD (ed) (1994) *Pre-Mesozoic geology in France*. Springer, Berlin Heidelberg New York
- Keppie JD, Dostal J, Murphy JB, Cousins BL (1997) Palaeozoic within-plate volcanic rocks in Nova Scotia (Canada) reinterpreted: isotopic constraints on magmatic source and palaeocontinental reconstruction. *Geol Mag* 134:425–447
- Kleinschrodt R, Gayk T (1999) Analogies and synchronism in the exhumation of HP units in the Variscan orogen: a comparison of the Münchberg Complex and Cabo Ortegal Complex. *Terra Nostra* 99:119
- Klootwijk C (1996) Phanerozoic configurations of Greater Australia: evolution of the North West Shelf. Canberra Australian Geological Survey Organisation
- Klötzli-Chovanetz E, Klötzli U, Koller F (1997) Lower Ordovician migmatization in the Ötztal crystalline basement (Eastern Alps, Austria): linking U–Pb and Pb–Pb dating with zircon morphology. *Schweiz Mineral Petr Mitt* 77:315–324
- Köppel V, Günthert A, Grünenfelder M (1981) Patterns U–Pb zircon and monazite ages in polymetamorphic units of the Swiss central alps. *Schweiz Mineral Petr Mitt* 61:97–119
- Kozur H (1984a) Some new stratigraphical and paleogeographical data in the Paleozoic and Mesozoic of the Pannonian Median Massif and adjacent areas. *Acta Geodaetica Geophys Montan Hungar* 19:93–106
- Kozur H (1984b) New biostratigraphic data from the Bükk-, Uppony- and Mecsek Mountains and their tectonic implications. *Acta Geol Hungar* 27:307–319
- Kröner A, Hegner E (1998) Geochemistry, single zircon ages and Sm–Nd systematics of granitoid rocks from the Gory-Sowie (Owl Mts), Polish West Sudetes: evidence for Early Palaeozoic are-related plutonism. *J Geol Soc Lond* 155:711–724
- Kröner A, Willner AP (1998) Time of formation and peak of Variscan HP–HT metamorphism of quartz-feldspar rocks in the central Erzgebirge, Saxony, Germany. *Contrib Mineral Petrol* 132:1–20
- Kröner A, Stipská P, Schulmann K, Jaekel P (2000) Chronological constraints on the pre-Variscan evolution of the north-eastern margin of the Bohemian Massif, Czech Republic. In: Franke W, Altherr R, Haak V, Oncken O, Tanner D (eds) *Orogenic processes: quantification and modelling in the Variscan belt of Central Europe*. *Geol Soc Lond Spec Publ* 179:175–198
- Laumonier B (1998) Les Pyrénées centrales et orientales au début du Paléozoïque (Cambrien s.l.): évolution paléogéographique et géodynamique. *Geodin Acta* 11:1–11
- Liebetrau V (1996) *Petrographie, Geochemie und Datierung der "Flüelagranitischen Assoziation" (sog. Jüngere Orthogneise) des Silvretakristallins Graubünden – Schweiz*. Dissertation, Fribourg, 233 pp

- Liebetrau V, Nägler T (1994) Geochronologische und geochemische Diskussion der sogenannten "flüelgranitischen Assoziation" des Silvrettakristallins (Graubünden/Schweiz). *Schweiz Mineral Petr Mitt* 74:265–268
- Liñan E, Quesada C (1990) Rift phase (Cambrian). In: Dallmeyer RD, Martinez Garcia E (eds) *Pre-Mesozoic geology of Iberia*. Springer, Berlin Heidelberg New York, pp 259–266
- Linnemann U (1999) The geotectonic setting of the Ordovician of the Saxo-Thuringian Terrane (Central European Variscides, Germany). *Acta Univ Carolinae Geol* 43:97–99
- Linnemann U, Buschmann B (1995) Die cadomische Diskordanz im Saxothuringikum (oberkambrisch-tremadocische overlap-Sequenzen). *Z Geol Wiss* 23:707–727
- Linnemann U, Gehmlich M, Tichomirova M, Buschmann B (1998a) Introduction to the pre-Symposium excursion: the peri-Gondwanan basement of the Saxothuringian composite terrane. *Schr Staatl Mus Min Geol Dresden* 9:7–13
- Linnemann U, Gehmlich M, Tichomirova M, Buschmann B, Bombach K (1998b) Tectonostratigraphic events of the Peri-Gondwanan basement of the Saxo-Thuringian composite terrane (Central European Variscides). *Schr Staatl Mus Min Geol Dresden* 9:159–161
- Linnemann U, Gehmlich M, Tichomirova M, Buschmann B, Bombach K (1999) The reflection of orogenic processes within the neoproterozoic to lower Carboniferous sedimentary record of the Saxothuringian terrane. Part I. Cadomian basement to Silurian). *Terra Nostra* 99:138
- Linnemann U, Gehmlich M, Tichomirova M, Buschmann B, Nasdala L, Jonas P, Lützner H, Bombach K (2000) From Cadomian subduction to Early Palaeozoic rifting: the evolution of the Saxo-Thuringian terrane at the margin of Gondwana in the light of single zircon geochronology (Central European Variscides, Germany). *Geol Soc Lond Spec Publ* 179:131–154
- Loeschke J, Heinisch H (1993) Palaeozoic volcanism of the Eastern Alps and its palaeotectonic significance. In: von Raumer J, and Neubauer F (eds), *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 441–455
- Lombardo B, Colombo F, Compagnoni R, Ghiglione G, Rubatto D (1997) Relics of pre-Variscan events in the Malinvern-Argentera Complex, Argentera Massif, Western Alps. *Third Workshop on Alpine Geological Studies. Quad Geodin Milano* 4:66
- Loth G, Eichhorn R, Schauder P, Höll R, Kennedy A (1999) U–Pb SHRIMP data from the Wildschönau ophiolite complex: evidence for Cambro-Ordovician rifting in the Western Greywacke supergroup (Eastern Alps). *Ber Dt Min Ges Eur J Mineral* 11 (Beih):149
- MacNiocaill C, van der Pluijm BA, van der Voo R (1997) Ordovician paleogeography and the evolution of the Japetus ocean. *Geology* 25:159–162
- Maniar PD, Piccoli PM (1989) Tectonic discrimination of granitoids. *Bull Geol Soc Am* 101:635–643
- Matte P (1986) Tectonics and plate tectonics model for the Variscan belt of Europe. *Tectonophysics* 126:329–374
- McKerrow WS, Scotese CR, Brasier MD (1992) Early Cambrian continental reconstructions. *J Geol Soc Lond* 149:599–606
- Mélou M, Oulebsir L, Paris F (1999) Brachiopodes et chitinozoaires ordoviens dans le NE du Sahara Algérien: implications stratigraphiques et paléogéographiques. *Geobios* 32:823–839
- Meng Q-R, Zhang G-W (1999) Timing of collision of the north and south China blocks: controversy and reconciliation. *Geology* 27:123–126
- Ménot RP (1987) Magmatismes paléozoïques et structuration carbonifère du Massif de Belledonne (Alpes Françaises). *Mém et Doc Centre Armoricain Etudes Structurales des Socles*, vol 21, Rennes, 465 pp
- Ménot RP, Peucat JJ, Scarni D, Piboule M (1988) 496 My age of plagiogranites in the Chamrousse ophiolite complex (external crystalline massifs in the French Alps): evidence of a lower Paleozoic oceanization. *Earth Planet Sci Lett* 88:82–92
- Mingram B (1996) Geochemische Signaturen der Metasedimente des erzgebirgischen Krustenstapels. *GeoForschungsZentrum Potsdam Scientific Technical Report STR96/04:1–104*
- Mingram B (1998) The Erzgebirge a subducted part of Northern Gondwana: geochemical evidence for repetition of early Palaeozoic metasedimentary sequences in metamorphic thrust units. *Geol Mag* 135:785–801
- Mingram B, Rötzler K (1999) Geochemische, petrologische und geochronologische Untersuchungen im Erzgebirgskristallin-Rekonstruktion eines Deckenstapels. *Schriftenr Geowiss* 9:1–80
- Mingram B, Rötzler K, Tichomirova M, Haase G, Teufel S (1998) Early Palaeozoic rift-related magmatism and passive margin sediments preserved in HP units of the Erzgebirge. *Terra Nostra* 98:105–107
- Montero P, Salman K, Bea F, Azor A, Expósito Ramos I, González Lodeiro F, Martínez-Poyatos D, Simancas JF (2000) New data on the geochronology of the Ossa-Morena zone, Iberian Massif. *Basement Tectonics* 15:136–138
- Müller B, Klötzli U, Schaltegger U, Flisch M (1996) Early Cambrian oceanic plagiogranite in the Silvretta nappe. Eastern Alps: geochemical, zircon U–Pb and Rb–Sr data from garnet-hornblende-plagioclase gneisses. *Geol Rundsch* 85:822–831
- Murphy JB, Nance RD (1989) Model for the evolution of the Avalonian-Cadomian belt. *Geology* 17:735–738
- Nance R, Murphy JB (1994) Contrasting basement isotopic signatures and the palinspastic restoration of peripheral orogens: example from the Neoproterozoic Avalonian-Cadomian belt. *Geology* 22:617–620
- Nance RD, Murphy JB (1996) Basement isotopic signatures and Neoproterozoic palaeogeography of Avalonian-Cadomian and related terranes in the circum-North Atlantic. In: Nance RD, Thompson MD (eds) *Avalonian and related peri-Gondwanan terranes of the Circum-North Atlantic*. *Geol Soc Am Spec Pap* 304:333–346
- Navidad M, Carreras J (1995) Pre-Hercynian magmatism in the Eastern Pyrenees (Cap de Creus and Albera massifs) and its geodynamical setting. *Geol Mifnb* 74:65–77
- Neubauer F (1991) Late Proterozoic and Early Palaeozoic tectono-thermal evolution of the Eastern Alps. In: Dallmeyer RD, Lecorché JP (eds) *The west African orogens and circum-Atlantic correlatives*. Springer, Berlin Heidelberg New York, pp 307–314
- Neubauer F, Raumer J von (1993) The Alpine basement: linkage between Variscides and East-Mediterranean mountain belt. In: von Raumer JF, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, p 641–663
- Neubauer F, Sassi FP (1993) The Austroalpine quartzphyllites and related Palaeozoic formations. In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 423–439
- Neubauer F, Hoinkes G, Sassi FP, Handler R, Höck V, Koller F, Frank W (1999) Pre-Alpine metamorphism of the Eastern Alps. *Schweiz Mineral Petr Mitt* 79:41–62
- Noblet C, Lefort JP (1990) Sedimentological evidence for a limited separation between Armorica and Gondwana during the Early Ordovician. *Geology* 18:303–306
- Nussbaum C, Marquer D, Biino G (1998) Two subduction events in a polycyclic basement: Alpine and pre-Alpine high-pressure metamorphism in the Suretta nappe, Swiss Eastern Alps. *J Metam Geol* 16:591–605
- Oberli F, Meier M, Biino G (1994) Time constraints on the pre-Variscan magmatic/metamorphic evolution of the Gotthard and Tavetsch units derived from single-zircon U–Pb results. *Schweiz Mineral Petrogr Mitt* 74:483–488

- O'Brien PJ, Kröner A (1999) Zircon ages of meta-eclogites in the Monotonous series of the Moldanubian zone in Austria and their significance. *Ber Dtsch Min Ges Eur J Mineral* 11 (Beih):170
- O'Brien PJ, Duyster J, Grauert B, Schreyer W, Stöckhert B, Weber K (1997) Crustal evolution of the KTB drill site: from oldest relics to the late Hercynian granites. *J Geophys Res* 102:18203–18220
- Oliver GJH, Corfu F, Krogh TE (1993) U–Pb ages from SW Poland: evidence for a Caledonian suture zone between Baltica and Gondwana. *J Geol Soc Lond* 150:355–369
- Oncken O (1997) Transformation of a magmatic arc and an orogenic root during oblique collision and its consequences for the evolution of the European Variscides (Mid-German Crystalline rise). *Geol Rundsch* 86:2–20
- Özmen F, Reischmann T (1999) The age of the Sakarya continent in West Anatolia: implication for the evolution of the Aegean region. *J Conf Abstr* 4:805–806
- Ordóñez Casado B (1998) Geochronological studies of the pre-Mesozoic basement of the Iberian Massif: the Ossa Morena zone and the Allochthonous Complexes within the Central Iberian zone. Dissertation ETH no. 12.940, 235 pp
- Panwitz C, Handler R, Heinisch H, Neubauer F (2000) $^{40}\text{Ar}/^{39}\text{Ar}$ -dating on detrital white micas from Austroalpine Palaeozoic of Northern Greywacke Zone (Tyrol/Salzburg–Austria): a pilot study. *Terra Nostra* 2000:83
- Paquette JL, Ménot RP, Peucat JJ (1989) REE, Sm–Nd and U–Pb zircon study of eclogites from the Alpine External massifs (Western Alps): evidence for crustal contamination. *Earth Planet Sci Lett* 96:181–198
- Paris F (1993) Evolution paléogéographique de l'Europe au Paléozoïque inférieur: le test des Chitinozoaires. *CR Acad Sci Paris* 316:273–280
- Paris F, Robardet M (1990) Early Palaeozoic palaeobiogeography of the Variscan regions. *Tectonophysics* 177:193–213
- Paris F, Robardet M, Dabard MP, Ghienne JF, Guillocheau F, Le Hérisse A, Loi A, Mélou M, Feist R, Servais T, Shergold J, Vidal M, Vizcaïno D (1999) Ordovician sedimentary rocks of France. *Acta Univ Carolinae Geol* 43:85–88
- Pearce JA, Nigel B, Harris W, Tindle AG (1984) Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *J Petrol* 25:956–983
- Pin C (1991) Central Western Europe: major stages of development during Precambrian and Paleozoic times. In: Dallmeyer RD, Lécorché JP (eds) *The West African orogens and circum-atlantic correlatives*. Springer, Berlin Heidelberg New York, pp 295–306
- Pin C, Lancelot J (1982) U–Pb dating of an Early Palaeozoic bimodal magmatism in the French Massif Central and of its further metamorphic evolution. *Contrib Mineral Petrol* 79:1–12
- Pin C, Marini F (1993) Early Ordovician continental break-up in Variscan Europe: Nd–Sr isotope and trace element evidence from bimodal igneous associations of the Southern Massif central, France. *Lithos* 29:177–196
- Piper JDA (2000) The Neoproterozoic Supercontinent: Rodinia or Palaeopangea? *Earth Planet Sci Lett* 176:131–146
- Poller U, Nægler TF, Liebetrau V, Galetti G (1997) The Mönch-alpgneiss: geochemical characteristics and Sm–Nd data of a polymetamorphic S-type granulite (Silvretta nappe/Switzerland). *Eur J Mineral* 9:411–422
- Powell C, McElhinny MW, Meert JG, Park JK (1993) Palaeomagnetic constraints on timing of the neoproterozoic breakup of Rodinia and the Cambrian formation of Gondwana. *Geology* 21:889–892
- Prigmore JK, Butler AJ, Woodcock NJ (1997) Rifting during separation of Eastern Avalonia from Gondwana. Evidence from subsidence analyses. *Geology* 25:203–206
- Quadt A von (1992) U–Pb zircon and Sm–Nd geochronology of mafic and ultramafic rocks from the central part of the Tauern Window (Eastern Alps). *Contrib Mineral Petrol* 110:57–67
- Quadt A von, Günther D, Frischknecht R, Zimmermann R, Franz G (1997) The evolution of pre-Variscan eclogites of the Tauern Window (Eastern Alps): A Sm/Nd-, conventional and Laser ICP-MS zircon U–Pb study. *Schweiz Mineral Petrogr Mitt* 77:265–279
- Rapela CW, Coira B, Toselli A, Saavedra J (1992) El magmatismo del Paleozoico inferior en el sudoeste de Gondwana. In: Gutierrez-Marco JC, Saavedra J, Rabano J (eds) *Paleozoico inferior de Ibero-America*, Univ Extremadura, Spain, pp 21–68
- Rast N, Skehan JW (1983) The evolution of the Avalonian plate. *Tectonophysics* 100:257–286
- Raumer JF von (1998) The Palaeozoic evolution in the Alps: from Gondwana to Pangea. *Geol Rundsch* 87:407–435
- Raumer J von, Neubauer F (1993a) Late Precambrian and Palaeozoic evolution of the Alpine Basement: an overview. In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 625–639
- Raumer J von, Neubauer F (eds) (1993b) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York
- Raumer J von, Bussy F, Schaltegger U (1999) Ordovician granulites in the Alps: significance and Peri-Gondwanan correlation. *Terra Nostra* 99:165
- Reischmann T, Anthes G (1996a) Das kristalline Grundgebirge am NW Rand des Rheingrabens. In: Geiger M (ed) *Haardt und Weinstrasse: Beiträge zur Landeskunde*. Veröffentlichungen der pfälzischen Gesellschaft zur Förderung der Wissenschaften, Speyer, pp 36–59
- Reischmann T, Anthes G (1996b) Geochronology of the mid-German crystalline rise west of the River Rhine. *Geol Rundsch* 85:761–774
- Reitz E, Höll R (1991) Biostratigraphischer Nachweis von Arenin in der nördlichen Grauwackenzone. *Jahrb Geol BA* 134:329–344
- Robardet M (1996) Early paleozoic palaeogeography of North-Gondwanan Europe: Serie Correlación Geol 12:167–180
- Robardet M, Paris F, Racheboeuf PR (1990) Palaeogeographic evolution of southwestern Europe during Early Palaeozoic times. *Geol Soc Am Mem* 12:411–419
- Robardet M, Verniers J, Feist R, Paris F (1994) Le Paléozoïque de France, contexte paléogéographique et géodynamique. *Géol France* 3:3–31
- Romer R, Franz L (1998) Ordovician Barrow-type metamorphism in the Strona-Ceneri Zone (northern Italy) dated by U–Pb on staurolite. *Schweiz Mineral Petr Mitt* 78:383–395
- Romer RL, Schärer U, Steck A (1996) Alpine and pre-Alpine magmatism in the root-zone of the Western Central Alps. *Contrib Mineral Petrol* 123:138–158
- Santos Zalduegui JF, Schärer U, Gil Iburguchi JI (1995) Isotope constraints on the age and origin of magmatism and metamorphism in the Malpica-Tuy allochthon, Galicia, NW Spain. *Chem Geol* 121:91–103
- Sassi FP, Neubauer F, Mazzoli C, Sassi R, Spiess R, Zirpoli G (1994) A tentative comparison of the Palaeozoic evolution of the Austroalpine and Southalpine quartzphyllites in the Eastern Alps. *Per Mineral* 63:35–52
- Schaltegger U (1992) Die polymetamorphe Geschichte des Aarmassivs: neue U/Pb-Resultate. *Schweiz Mineral Petr Mitt* 72:148–149
- Schaltegger U (1993) The evolution of the polymetamorphic basement in the central Alps unravelled by precise U–Pb zircon dating. *Contrib Mineral Petrol* 113:466–478
- Schaltegger U, Gebauer D (1999) Pre-Alpine geochronology of the Central, Western and Southern Alps. *Schweiz Mineral Petrogr Mitt* 79:79–87
- Schaltegger U, Nægler T N, Corfu F, Maggetti M, Galetti G, Stosch H (1997) A Cambrian island arc in the Silvretta nappe: constraints from geochemistry and geochronology. *Schweiz Mineral Petrogr Mitt* 77:337–350

- Schätz M, Bachtadse V, Tait JA, Soffel HC (1997) Palaeomagnetic results from Lower Devonian sediments of the Southern Alps. *Terra Nostra* 97:152–155
- Schermaier A, Schindlmayr A, Finger F, Roberts M, Pin C (1998) Evidence for Early Palaeozoic subduction and arc-continent collision in the Oetzal-Silvretta crystalline basement (Western Eastern Alps). *Acta Univ Carolinae Geol* 42:330–331
- Schindlmayr A (1999) Granitoids and plutonic evolution of the Ötzal-Stubai Massif. Dissertation, Univ Salzburg, 287 pp
- Schmidt K (1977) Der altpaläozoische Magmatismus und seine Stellung in der tektonischen Geschichte Mittel- und Südwesteuropas. *Z Dtsch Geol Ges* 128:121–141
- Schönlaub HP (1993) Stratigraphy, biogeography and climatic relationships of the Alpine Paleozoic. In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 65–91
- Schönlaub HP (1997a) The biogeographic relationship of Ordovician strata and fossils of Austria. *Ber Geol Bundesanstalt Wien* 40:6–19
- Schönlaub HP (1997b) The Silurian of Austria. *Ber Geol Bundesanstalt Wien* 40:20–41
- Schönlaub HP, Heinisch H (1993) The classic fossiliferous palaeozoic units of the eastern and southern alps. In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 395–422
- Schönlaub HP, Histon K (2000) The palaeozoic evolution of the Southern Alps. *Mitt Österr Geol Ges* 92:15–34
- Sdzuy K, Liñan E, Gozalo R (1999) The Leonian stage (early Middle Cambrian), a unit for Cambrian correlation in the Mediterranean subprovince. *Geol Mag* 136:39–48
- Sengör AMC, Natal'in BA (1996) Palaeotectonic of Asia: fragments of a synthesis. In: Yin A, Harrison M (eds) *The tectonic evolution of Asia*. Cambridge University Press, Cambridge, pp 486–640
- Sergeev SA, Steiger RH (1993) High-precision U–Pb single zircon dating of Variscan and Caledonian magmatic cycles in the Gotthard massif, Central Swiss Alps. *Terra Nova Abstr* 5:394–395
- Skehan JW, Rast N (1983) Relationship between Precambrian and lower Palaeozoic rocks of southeastern New England and other north Atlantic Avalonian terranes. In: Schenk PE (ed) *Regional trends in the geology of the Appalachian-Caledonian-Hercynian-Mauritanide orogen*. Mathematical and physical sciences. Reidel, Dordrecht, pp 131–162
- Söllner F, Höll R, Miller H (1991) U–Pb Systematik der Zirkone in Metavulkaniten (“Porphyroiden”) aus der nördlichen Grauwackenzone und dem Tauernfenster (Ostalpen, Österreich). *Z Dtsch Geol Ges* 142:285–299
- Stampfli G (1996) The intra-Alpine terrain: a palaeotethyan remnant in the Alpine Variscides. *Eclog Geol Helv* 89:12–42
- Stampfli G (2000) Tethyan oceans. In: Boxkurt E, Winchester JA, Piper JD (eds) *Tectonics and magmatism in Turkey and the surrounding area*. *Geol Soc Lond Spec Publ* 173:1–23
- Stampfli G M, Mosar J (1996) Palaeozoic evolution of the Tethyan domain. 14. Schweiz Tektonikertreffen, abstract vol:34
- Stampfli G, Marcoux J, Baud A (1991) Tethyan margins in space and time. *Palaeogeogr Palaeoclimatol Palaeoecol* 87:373–409
- Stille P, Tatsumoto M (1985) Precambrian tholeiitic-dacitic rock-suites and Cambrian ultramafic rocks in the Pennine nappe system of the Alps: evidence from Sm–Nd isotopes and rare earth elements. *Contrib Mineral Petrol* 89:184–192
- Stipska P, Schulmann K, Kroener A (1998) Role of Cambro-Ordovician rifting in Variscan collision at the NE margin of the Bohemian Massif. *Acta Univ Carolinae Geol* 42:343–344
- Tait J, Schätz M, Bachtadse V, Soffel H (1998) Paleogeography of paleozoic terranes in the Varican and Alpine fold belts. *Schr Staatl Mus Min Geol Dresden* 9:192–193
- Tessensohn F (1999) Alpine-type orogens at the Palaeo-Pacific margin of Gondwana: the lack of Wilson cycles. *Terra Nostra* 99:198–199
- Thélin P, Sartori M, Burri M, Gouffon Y, Chessex R (1993) The pre-Alpine basement of the Briançonnais (Wallis, Switzerland). In: von Raumer J, Neubauer F (eds) *The pre-Mesozoic geology in the Alps*. Springer, Berlin Heidelberg New York, pp 297–315
- Thöni M (1999) A review of geochronological data from the Eastern Alps. *Schweiz Mineral Petrogr Mitt* 79:209–230
- Torsvik H, Eide EA (1998) Phanerozoic paleogeography and geodynamics with Atlantic details. *NGU report*, p 82
- Torsvik TH, Smethurst MA (1994) Geographic mapping and paleoreconstruction package (GMAP). Software description and examples of paleoreconstructions
- Torsvik TH, Smethurst MA (1996) Continental break-up and collision in the Neoproterozoic and Palaeozoic: a tale of Baltica and Laurentia. *Earth Sci Rev* 40:229–258
- Torsvik TH, Smethurst MA (1999) Plate tectonic modelling: virtual reality with GMAP. *Comput Geosci* 25:395–402
- Torsvik TH, Trench A, McKerrow WS (1994) Implications of paleomagnetic data from the Tortworth Silurian inlier (southern Britain) to paleogeography and Variscan tectonism. *Geophys J Int* 119:91–100
- Tremblay A, Laflèche MR, McNutt RH, Bergeron M (1994) Petrogenesis of Cambro-Ordovician subduction-related granitic magmas of the Quebec Appalachians, Canada. *Chem Geol* 113:205–220
- Ugidos JM, Valladares MI, Recio C, Rogers G, Fallick AE, Stephens WE (1997) Provenance of Upper Precambrian–Lower Cambrian shales in the Central Iberian Zone, Spain: evidence from a chemical and isotopic study. *Chem Geol* 136:55–70
- Unrug R (1997) Rodinia to Gondwana: the geodynamic map of Gondwana supercontinent assembly. *Geol Soc Am Today* 7:1–6
- Unrug R, Haranczyk C, Chosyk-Jaminska M (1999) Easternmost Avalonian and Armorican-Cadomian terranes of central Europe and Caldeonian-Variscan evolution of the polydeformed Kraków mobile belt: geological constraints. *Tectonophysics* 302:133–157
- Üstaömer PA (1999) Pre-Early Ordovician Cadomian arc-type granitoids, the Bolu Massif, West Pontides, northern Turkey: geochemical evidence. *Int J Earth Sci* 88:2–12
- Valverde-Vaquero P, Dunning GR (2000) New U–Pb ages for Early Ordovician magmatism in Central Spain. *J Geol Soc Lond* 157:15–26
- Van der Voo R (1993) Paleomagnetism of the Atlantic, Tethys and Iapetus oceans. Cambridge University Press, Cambridge
- Van Staal CR, Dewey JF, Niocail CM, Mckerrow WS (1998) The Cambrian-Silurian tectonic evolution of the Northern Appalachians and British Caledonides: history of a complex, west and southwest Pacific-type segment of Iapetus. In: Blundell DJ, Scott AC (eds) *Lyell: the past is the key to the present*. *Geol Soc Lond Spec Publ* 143:199–242
- Weger M, Loth G, Höll R, Masch L, Köhler H, Rohrmüller J (1999) Cambrian sedimentation, Ordovician and Devonian subduction, collision and uplift in the southern ZEV (Zone of Erbendorf-Vohenstraus, Western margin of Bohemian Massif). *Terra Nostra* 99:207
- Winchester JA, Floyd PA, Franke W, Zelazniewicz A, Patocka F, Kryza R, Cymerman Z, Piaseck MAJ, Seston R, Crowley QG, Kachlik V, Stedra V, Holland JG (1998) A review of recent geochemical and structural evidence for major Palaeozoic rifting and re-assembly within the Bohemian Massif. *Acta Univ Carolinae Geol* 42:359–360
- Williams KE (1997) Early Palaeozoic palaeogeography of Laurentia and western Gondwana: evidence from tectonic subsidence analysis. *Geology* 25:747–750
- Wirsing A (1997) Die Orthogneise des oberen Val Bérard (Aiguilles-Rouges-Massiv, Westalpen, Frankreich). Dissertation, Univ Fribourg

- Yin A, Nie S (1996) A Phanerozoic palinspastic reconstruction of China and its neighboring regions. In: Yin A, Harrison M (eds) *The tectonic evolution of Asia*. Cambridge Univ Press, Cambridge, pp 442–485
- Zeh A, Brätz H, Cosca M, Tichomirova M, Okrusch M (1997) $^{39}\text{Ar}/^{40}\text{Ar}$ und $^{207}\text{Pb}/^{206}\text{Pb}$ Datierungen im Ruhlaer Kristallin, Mitteldeutsche Kristallinzone. *Terra Nostra* 97:212–215
- Ziegler P (1984) Caledonian and Hercynian crustal consolidation of western and central Europe: a working hypothesis. *Geol Mijnb* 63:93–108
- Zingg MA (1989) Die Siviez-Mischabel-Decke: Entstehung und Entwicklung eines Altkristallins. *Schweiz Mineral Petrogr Mitt* 69:130
- Zonenshain LP, Kuzmin MI, Kononov MV (1985) Absolute reconstructions of the Paleozoic oceans. *Earth Planet Sci Lett* 109:103–116
- Zulauf G, Schitter F, Riegler G, Finger F, Fiala J, Vejnar Z (1999) Age constraints on the Cadomian evolution of the Teplá Barrandian unit (Bohemian Massif) through electron microprobe dating of metamorphic monazite. *Z Dtsch Geol Ges* 150:627–639
- Zurbruggen R, Franz L, Handy M (1997) Pre-Variscan deformation, metamorphism and magmatism in the Strona-Ceneri Zone (southern Alps of northern Italy and southern Switzerland). *Schweiz Mineral Petrogr Mitt* 77:361–381
- Zwart HJ, Dornsiepen UF (1978) The tectonic framework of Central and Western Europe. *Geol Mijnb* 57:627–654