

Primary cell culture of meningotheelial cells—a new model to study the arachnoid in glaucomatous optic neuropathy

Xiaorong Xin · Bin Fan · Hanspeter E. Killer ·
Albert Neutzner · Josef Flammer · Peter Meyer

Received: 14 December 2009 / Revised: 15 March 2010 / Accepted: 18 March 2010 / Published online: 8 April 2010
© Springer-Verlag 2010

Abstract

Background In a previous report, we found that the occurrence and amount of meningotheelial cell nests in the subarachnoid space are significantly increased in glaucomatous optic nerves compared to normals. In order to allow research into the role of meningotheelial cells during diseases of the optic nerve, an in vitro model is necessary. For this purpose, we developed a culture method for porcine meningotheelial cells from the arachnoid layer covering the optic nerve.

The authors have full control of all primary data, and they agree to allow Graefe's Archive for Clinical and Experimental Ophthalmology to review their data upon request.

X. Xin · B. Fan · H. E. Killer · A. Neutzner · J. Flammer ·
P. Meyer
Department of Biomedicine, University Eye Clinic,
University Hospital Basel,
Basel, Switzerland

H. E. Killer
Department of Ophthalmology, Kantonsspital Aarau,
Aarau, Switzerland

X. Xin
Department of Ophthalmology,
Qinghai Provincial People's Hospital,
Xining, Qinghai, China

B. Fan
Department of Ophthalmology,
2nd Clinical Hospital Jilin University,
Changchun, China

P. Meyer (✉)
Department of Ophthalmic Pathology,
University Eye Clinic Basel,
CH-4012 Basel, Switzerland
e-mail: peter.meyer@unibas.ch

Methods Meningotheelial cells were scraped from the arachnoid layer of porcine optic nerves and cultured for 2–3 weeks until the cells formed a monolayer. To eliminate contaminating fibroblasts from the culture, cells were negatively selected using magnetic anti-fibroblast beads after the first passage. Cells were detached using 0.05% Trypsin-EDTA, incubated with anti-fibroblast beads, separated using a magnetic column and the flow-through was collected. The purified primary meningotheelial cells were characterized by electron microscopy and immunocytochemistry using anti-glia fibrillary acidic protein (GFAP) and anti-keratan sulfate antibodies.

Results Primary cells grew out after dissection and formed a monolayer within 2–3 weeks, which was composed of two morphologically different cell types, flattened cells with round nuclei and fibroblast-like cells with long processes. The fibroblast-like cells in the culture could be labelled and selected using anti-fibroblast microbeads. The second cell type did not bind to the anti-fibroblast beads, and upon immunocytochemistry showed a marked expression of both GFAP and keratan sulphate. In addition, examination of these cells by electron microscopy revealed morphological characteristics of meningotheelial cells, including hemidesmosomes and cytoplasmic filaments.

Conclusions The technique described in this paper for the primary culture of meningotheelial cells from the subarachnoid space of the optic nerve and using magnetic beads for the removal of fibroblasts is effective in obtaining a highly enriched meningotheelial cell culture.

Keywords Meningotheelial cell · Fibroblast · Subarachnoid space · Cell culture · Optic nerve · Glaucoma · Arachnoid

Introduction

The optic nerve is a white matter tract of the central nervous system that connects the retinal axons to the brain. It is enveloped by the meninges throughout its intraorbital and intracanalicular course. The meningeal sheath of the optic nerve has the same lamellar structure as the meninges of the brain, the pia mater, the arachnoid and the dura mater. The inner layer of the arachnoid is covered with meningotheial cells. A variety of trabecules and septae covered with meningotheial cells transverse the subarachnoid space between the arachnoid and the pia. Located between the dura and the arachnoid mater is the subdural space [1, 2]. The subdural and the subarachnoid space of the optic nerve are continuous with those of the brain.

The subarachnoid space and the subdural space are occupied by the cerebrospinal fluid (CSF). CSF is mainly produced by the choroid plexus epithelium and the ependymal cells of the ventricular system, from where it flows into cisterns and the subarachnoid spaces, including the subarachnoid space of the optic nerve [3]. CSF is generally thought to communicate freely among the different CSF spaces and to have a homogeneous pressure throughout the brain and optic nerve. However, recent anatomical studies demonstrated that a complex architecture of arachnoid trabeculae, pillars, and septa in the subarachnoid space of the optic nerve might strongly influence CSF dynamics in the subarachnoid space via a dynamic valve function which might lead to a pressure gradient along the optic nerve [2, 4].

Meningotheial cells that cover the meninges of the optic nerve build a fluid barrier that prevents the outflow of CSF. Aging, and disorders such as meningitis, can influence the anatomical structure of the optic nerve sheath. A recent study demonstrated an age-dependent increase of meningotheial cell proliferation, causing the formation of cell nests, which eventually hyalinize and calcify [5]. A significant increase in occurrence and size of those meningotheial cell nests in the subarachnoid space has been observed in glaucoma compared to healthy optic nerves. The function is yet not known; however, a secondary reaction of the meningotheial cells to axon loss during glaucoma can be assumed [1]. In addition, impairment of meningotheial cell layer function might result in the failed clearance of biologically active molecules or waste products from the subdural and subarachnoidal space, and this might contribute to optic nerve diseases such as progressive anterior and posterior ischemic optic neuropathy, or optic nerve sheath compartment syndrome and even glaucoma [4, 6–8].

As the major cell population covering the orbital optic nerve sheath, the possible involvement of meningotheial

cells in optic nerve diseases warrants further investigation. The culture of primary meningotheial cells from the optic nerve sheath makes it possible for the first time to study the response of those cells to various physiological and pathological conditions, and a more detailed understanding of their function during health and disease might improve clinical intervention for optic nerve disorders.

Materials and methods


Dissection and cell culture

Porcine eyes were obtained from the local slaughter house within 4 h of death. Eyes were transported to the laboratory in Krebs-Ringer solution, and rinsed with sterile phosphate buffered saline (PBS, Sigma, Germany) containing 10,000 U/ml penicillin and 10,000 ug/ml streptomycin (Sigma, Germany). Tissue and muscles around the eye and the optic nerve were removed (Fig. 1a). The dura-arachnoid sheath was freed from the optic nerve by using two tooth-forceps, one for fixating the optic nerve, and the other for tearing the dura-arachnoid sheath towards the globe while keeping the subarachnoid side upward (Fig. 1b). The dissected layer was transferred to a sterile, collagen pre-coated, 35-mm cell culture dish (Fig. 1c) containing 1 ml Dulbecco's modified Eagle's medium (DMEM, Gibco, Germany) supplemented with 10% fetal bovine serum (FBS, Sigma, Germany) and PSA (10,000 U/ml penicillin, 10,000 ug/ml streptomycin, 25 ug/ml amphotericin B). Collagen-coating was done with 0.1% Bornstein and Traub Type I collagen in 0.1 M acetic acid (Sigma, C8919) by covering the plates with collagen followed by 1 hour air-drying. With the subarachnoid layer facing upwards, superficial cells were gently scratched from the subarachnoid side into the medium using a blade (Fig. 1d). Cells were incubated at 37°C in a 95% air/5% CO₂ atmosphere. The growth medium was changed twice a week.

Separation of meningotheial cells from contaminating fibroblasts

After outgrowth, cells were washed with PBS, trypsinized (0.05% Trypsin-EDTA) for 2 min, and centrifuged at 1100 rpm for 3 min at room temperature. The supernatant was aspirated and the cell pellet was resuspended in PBS buffer (0.5% BSA, 2 mM EDTA, pH 7.2), anti-fibroblast microbeads (Miltenyi Biotec GmbH, Germany) (20 ul/10⁷ cells) were added and incubated for 30 min at room temperature. Cells were washed with PBS buffer, centrifuged at 100×g for 10 min, and were resuspended in PBS buffer. To eliminate fibroblasts, 0.5 ml cell suspension were

Fig. 1 **a** Porcine eye with anterior part of the optic nerve surrounded by the optic nerve sheath. **b** Dura and arachnoid layers were cut and torn toward the globe by keeping the sub-arachnoid side upward. **c** The dissected layer was transferred into a dish with culture medium. **d** Superficial cells were gently scraped from subarachnoid side into the medium by using a blade



applied onto MACS column (Miltenyi Biotec GmbH, Germany); the cells flow through column were collected as meningotheial cells and analyzed for the presence of meningotheial cells by immunochemistry and electron microscopy. Fibroblasts were recovered by eluting the magnetically labeled cells from the column with a plunger.

Immunocytochemistry

After magnetic separation, the two cell populations were spun onto slides using a Cytospin III centrifuge (Shandon, Pittsburgh, PA, USA). The cells were fixed with 4% carbodiimide in PBS, incubated with GFAP antibody (Santa Cruz, CA, USA) that diluted to 1:100 in PBS. The antigen was visualized using goat anti-mouse IgG conjugated to horseradish peroxidase and 3,3-diaminobenzidine tetrahydrochloride. The sections were counterstained with hematoxylin, dehydrated and mounted. Incubation without primary antibodies served as negative control.

After separation, meningotheial cells were grown on coverslips, fixed with acetone and methanol for 3 min, permeabilized with 0.1% Triton X-100+1% BSA/FCS in PBS for 15 min at room temperature, rinsed three times with PBS and non-specific binding sites were blocked with 1% BSA in PBS/0.1% Triton X-100 for 1 h. The cells were incubated with anti-keratan sulphate antibodies (AbD Serotec, USA) for 2 h at room temperature; the concentration of antibody was diluted to a ratio of 1:100, and then washed three times with PBS. The primary antibody was visualized using Alexa Fluor 488-labeled anti-mouse secondary antibodies

(Invitrogen, USA). Cells were counterstained with DAPI for 3 min, mounted on glass slides, and observed with a fluorescence microscope (Olympus B×51, Japan).

Ultrastructural examination by electron microscopy

After magnetic separation, 4 to 5×10^6 meningotheial cells were pelleted and fixed for 180 min with cold 3% glutaraldehyde in 0.2 M sodium cacodylate phosphate buffer. After a brief rinse with PBS buffer, pellets were postfixed for 1 hr with cold 1% osmium tetroxide in phosphate buffer, pH 7.3. The pellets were cut into fragments. Both dehydration in ethanol and propylene oxide and embedding in Epon were done by routine procedures; 1.5 μm sections were mounted on slide and stained with toluidine blue. After that, 85 nm sections mounted on uncoated grids and stained with uranyl and lead were imaged using a FEI Morgagni electron microscope.

Results

Primary cell culture from the subarachnoid layer of porcine optic nerves

Cells grew out after dissection and formed a monolayer within 2–3 weeks, which was composed of two different cell types, flattened cells with round nuclei and fibroblast-like cells with long processes (Fig. 2a).




Fig. 2 a Micrograph of primary cultured cells shows two different cell types, flattened cells with round nuclei (*) and cells with long processes (**) (fibroblast-like). **b** Meningothelial cells after separation

with anti-fibroblast microbeads ($\times 10$). The results are shown for a representative cell culture out of four independent primary cell preparations

Purification of meningeal cells with magnetic anti-fibroblast microbeads

Cells were separated by magnetic fibroblast microbeads after primary cells reached confluency; positive and negative cells were collected respectively and continued to be cultured. Positive cells labelled with fibroblast microbeads showed obvious elongated fibroblast-like morphology; negative cells against fibroblast bead presented flattened cell type (Fig. 2b).

Characterization of meningeal cells with GFAP and keratan sulphate antibodies

Based on the observation of GFAP staining at the optic nerve, we found that not only astrocyte expressed GFAP, but meningeal cells also exhibited positive immunostaining for GFAP. We used GFAP as well as keratan sulphate antibody as markers for identifying meningeal cells. The result revealed strong expression of meningeal cells for both GFAP and keratan sulphate antibodies (Fig. 3a, b), while fibroblasts stained negative for GFAP (Fig. 3c). To establish the purity of the meningeal cell culture, cells after six passages were stained with anti-GFAP antibodies and counterstained using DAPI. Counting revealed less than 1% GFAP-negative cells (99.3% GFAP positive, $n=3$) confirming a pure meningeal cell culture.

Morphological analysis of meningeal cells by electron microscopy

The ultrastructural analysis revealed cells with unlabelled anti-fibroblast microbeads which contained oval nuclei (Fig. 4a) and showed slight membrane irregularity. Desmosomes were found between the cells (Fig. 4b). The

cytoplasm contained filaments arranged in aggregates (Fig. 4c).

Discussion

In order to establish a method for the culture of primary meningeal cells, we used a cut and tear method to separate the arachnoid layer from the optic nerve. We found this method to be superior to mechanical dissection of the arachnoid layer in terms of speed and the yield of meningeal cells.

Meningeal cells are the most common cell type, covering both sides of the arachnoid layer; however, there are fibroblasts present in the arachnoid trabeculae in the form of a very thin connective tissue. Therefore, they are always part of the primary culture of meningeal cells, and the removal of those contaminating fibroblasts is crucial to get pure meningeal cells. The use of magnetic anti-fibroblast microbeads proved useful to separate out fibroblasts and purify meningeal cells for further cultivation.

For identification of the purified meningeal cells, we relied on a previous observation described by Hogan [5]; in contrast to the contaminating fibroblasts, the morphology of the meningeal cells appeared to be flattened. Furthermore, immunocytochemical analysis identified the negatively selected cells as meningeal cells. We found that meningeal cells stained positive for GFAP, which is in accordance with our findings in post-mortem optic nerves (our unpublished observation) and the positive GFAP staining of ependymal cells as shown by Kivelä [9]. In addition to GFAP, keratan sulfate was previously reported to be expressed in meningeal cells [10] and, as expected, staining of the purified meningeal cells revealed the presence of this marker protein in our cultured

primary cells. Furthermore, the ultrastructural features of the primary cells we cultivated are consistent with meningeothelial cells, which presented with fine filaments extending throughout the cytoplasm, frequently joined by desmosomes with other cells.

Recent studies have brought new understanding to the pathogenesis of optic nerve diseases, in which the generally accepted concept that the CSF pressure and composition are homogeneous within small limits throughout all CSF




Fig. 3 **a** Positive immunostaining of meningeothelial cells for GFAP antibody after the separation with anti-fibroblast microbeads. **b** Marked immunostaining of meningeothelial cells for keratan sulphate antibody after separation with microbeads ($\times 40$). **c** Anti-fibroblast beads selected cells stained negative for GFAP using anti-GFPA antibodies, making GFAP a good marker to distinguish between meningeothelial cells and contaminating fibroblasts




Fig. 4 Electron micrograph of meningeothelial cells (**a**) with ovoid nuclei. Ultrastructural features of hemidesmosomes (*arrows*) (**b**), filaments (*arrowhead*), and lipid vacuoles (*arrow*) (**c**) of meningeothelial cells

compartments has been challenged by evidence suggesting that the subarachnoid space of the optic nerve can become separated from other CSF spaces in the wake of optic nerve disease. Compartmentation of the subarachnoid space of the optic nerve might result in an imbalanced composition of the CSF due to impaired removal of waste products [11, 12]. This is thought to contribute to the development of optic nerve sheath compartment syndromes, and might be related to progressive anterior and posterior ischemic optic neuropathy or chronic open-angle and normal-tension glaucoma [10, 13–15]. Meningeothelial cells are the elemental cells lining the inner surface of the dura mater, both side of arachnoid layer, and the outer surface of pia mater, and are therefore in direct contact to the CSF. They perform an essential role in keeping the CSF barrier intact, and it is conceivable that any ailment of meningeothelial cells such as inflammation, ischemia or and hypoxia possibly results in the impairment of the CSF composition. The significantly increased presence of the meningeothelial cell nests in the subarachnoid space in glaucoma has drawn our attention and interest to further investigate the pathophysiological role of meningeothelial cells in optic nerve diseases. Taken together, the primary culture of meningeothelial cells presented here is the first step in studying those cells in vitro, and might prove useful in the future to unravel the etiology of optic nerve diseases.

References

1. Pache M, Meyer P (2006) Morphological changes of the retrobulbar optic nerve and its meningeal sheaths in glaucoma. *Ophthalmologica* 220:393–396

2. Killer HE, Laeng HR, Flammer J, Groscurth P (2003) Architecture of arachnoid trabeculae, pillars, and septae in the subarachnoid space of the human optic nerve: anatomy and clinical considerations. *Br J Ophthalmol* 87:777–781
3. Wood JH (1980) *Neurobiology of cerebrospinal fluid*. Plenum Press, New York
4. Killer HE, Jaggi GP, Flammer J, Miller NR, Huber AR, Mironov A (2007) Cerebrospinal fluid dynamics between the intracranial and the subarachnoid space of the optic nerve. Is it always bidirectional? *Brain* 130:514–520
5. Hogan MJ, Alvarado JA, Weddell JE (1971) Optic nerve. *Histology of the human eye*. Saunders, Philadelphia, pp 592–606
6. Rubenstein E (1998) Relationship of senescence of cerebrospinal fluid circulatory system to dementias of the aged. *Lancet* 351:283–285
7. Silverberg GD, Mayo M, Saul T, Rubenstein E, McGuire D (2003) Alzheimer's disease, normal-pressure hydrocephalus, and senescent changes in CSF circulatory physiology: a hypothesis. *Lancet Neurol* 2:506–511
8. Killer HE, Jaggi GP, Mironov A, Flammer J, Miller NR, Huber AR (2006) The optic nerve: a new window into cerebrospinal fluid composition? *Brain* 129:1027–1030
9. Kivelä T, Uusitalo M (1998) Structure, development and function of cytoskeletal elements in non-neuronal cells of the human eye. *Prog Retin Eye Res* 17:385–428
10. Hirsch S, Bähr M (1999) Immunocytochemical characterization of reactive optic nerve astrocytes and meningeal cells. *Glia* 26:36–46
11. Reiber H (2001) Dynamics of brain-derived proteins in cerebrospinal fluid. *Clin Chim Acta* 310:173–186
12. TUMANI H, Nau R, Felgenhauer K (1998) Beta-trace protein in cerebrospinal fluid: a blood-CSF barrier-related evaluation in neurological diseases. *Ann Neurol* 44:882–889
13. Killer HE, Mironov A, Flammer J (2003) Optic neuritis with marked distension of the optic nerve sheath due to local fluid congestion. *Br J Ophthalmol* 87:249
14. Jaggi GP, Mironov A, Huber AR, Killer HE (2007) Optic nerve compartment syndrome in a patient with optic nerve sheath meningioma. *Eur J Ophthalmol* 17:454–458
15. Killer HE, Jaggi GP, Flammer J, Miller NR (2008) Is open-angle glaucoma caused by impaired cerebrospinal fluid circulation: around the optic nerve? *Clin Experiment Ophthalmol* 36:308–311