

## Microsatellite markers for the common vole (*Microtus arvalis*) and their cross-species utility

Barbara Walser · Gerald Heckel

Received: 27 April 2007 / Accepted: 1 June 2007 / Published online: 24 July 2007  
© Springer Science+Business Media B.V. 2007

**Abstract** Fragmentation of natural habitats of the common vole (*Microtus arvalis*) provides an excellent model system to study the consequences of restricted gene flow and small population sizes for isolated populations. Here we describe the isolation and characteristics of 10 autosomal and one X-linked microsatellite marker. These new markers were tested in 24 voles from a natural population in eastern Germany. Loci were highly polymorphic with numbers of alleles per locus ranging from three to 26 and expected heterozygosities from 0.51 to 0.97. All loci except for the X-linked locus Mar105 followed Hardy–Weinberg expectations. Cross-species amplifications revealed that most loci were polymorphic as well in *M. agrestis*, *M. thomasi*, and *M. pennsylvanicus*.

**Keywords** *Microtus* · Genetic markers · Microsatellites · Short tandem repeats · STR

The common vole, *Microtus arvalis*, is a small rodent, which plays due to its abundance a prominent role in many open habitats in temperate Europe. The species has a continuous range from the Atlantic coast of France to the Ukraine and Russia, and it is not yet well understood if eastern populations form separate species (Fink et al. 2004; Mitchell-Jones et al. 1999). European populations are genetically highly differentiated due to the limited dispersal abilities (Heckel et al. 2005). On a smaller scale, the fragmentation of habitats through modern agricultural practices provides an excellent model system to study the effects of restricted gene

flow, small effective sizes and inbreeding on isolated populations (see Schweizer et al. 2007). In this study, we describe the isolation of 10 autosomal and one X-linked microsatellite locus of *M. arvalis* and their utility in cross-species amplification with the European species *M. agrestis*, *M. thomasi* and the American *M. pennsylvanicus*.

Microsatellites were isolated following the enrichment protocols by Glenn and Schable (2005) with slight modifications as follows. A genomic library was constructed with DNA ( $\approx 140 \mu\text{g}/\mu\text{l}$ ) isolated from muscle tissue of a *M. arvalis* individual from a population near Belp, Switzerland ( $46^{\circ}55'N$ ,  $7^{\circ}29'E$ ). DNA was digested with Rsa I and fragments of approximately 500 bp were screened for microsatellite sequences with a mixture of the five probes (TG)<sub>12</sub>, (GA)<sub>15</sub>, (AATC)<sub>6</sub>, (AATG)<sub>6</sub> and (ATTG)<sub>6</sub>. For cloning of vole DNA, we used the Qiagen PCR Cloning<sup>plus</sup> Kit. 136 of approximately 695 colonies contained positively hybridizing clones. We used M13 primers and 1.5  $\mu\text{l}$  of bacterial pellet for PCR-amplification of the region containing the microsatellite sequences in a reaction volume of 25  $\mu\text{l}$  on a GeneAmP® PCR system 9700 (Applied Biosystems) according to Glenn and Schable (2005). PCR-products were separated on 1% agarose gel, length differences were scored by comparison with a 100 bp ladder (Invitrogen), and PCR-products were purified for sequencing with GenElute™ PCR Clean-Up-Kit (Sigma). Vole DNA inserts were sequenced with the Terminator Ready Reaction Mix ‘Big Dye’ Version 3.1 from Applied Biosystems in a reaction volume of 10  $\mu\text{l}$  following the recommendations of the manufacturer. The sequencing products were precipitated, and separated and detected on an ABI Prism 3100 Genetic Analyser from Applied Biosystems.

PCR-primers based on the clone sequences were designed using Primer3 (Rozen and Skaletsky 2000). PCR

B. Walser · G. Heckel (✉)  
Computational and Molecular Population Genetics (CMPG),  
Zoological Institute, University of Bern, Baltzerstrasse 6, Bern  
3012, Switzerland  
e-mail: gerald.heckel@zoo.unibe.ch

**Table 1** Characteristics of eleven microsatellite loci of *M. arvalis*

Locus	Primer sequence (5'-3')	Repeat motif	A	Size range (bp)	PCR set	Dye label	$H_o$	$H_e$	HWE $p=$	Accession No.
Mar03	F: GGAGATAACAAGGCCAAACAA R: TGGCATATTAGATGACCTGTGG	(TG) <sub>21</sub>	17	140–186	1	FAM	1.00	0.93	0.79	EF666981
Mar012	F: TTGCTCAATTCCTCATAAAAGG R: TGTCAATGGATTGGCATACA	(GA) <sub>23</sub>	16	92–120	1	VIC	0.88	0.92	0.15	EF666982
Mar016	F: CATCATCTCTGGGCACTG R: ACGGTCTGTGCAAACCACTT	(CA) <sub>19</sub>	7	155–167	1	VIC	0.92	0.67	0.11	EF666983
Mar049	F: GGAAAATGCGCTGTGTTATGAGA R: TGTGATGTGGGAACCGAGAGA	(GA) <sub>8</sub> AA(GA) <sub>23</sub>	21	217–284	2	NED	1.00	0.94	0.99	EF666984
Mar058	F: CAGGACCTTTGGAAAGAGCAG R: ^TGTGCACACATGAAACATAAACTTC	(TC) <sub>21</sub>	3	464–472	2	VIC	0.58	0.51	0.80	EF666985
Mar063	F: GCCTGGACACAAACCAAACCTT R: GGCTATGGCAGCTCTCTG	(AC) <sub>23</sub>	16	269–313	1	VIC	0.88	0.94	0.14	EF666986
Mar076	F: TCACCAAGGACCTACTGAGCA R: GCCAGCTTCATTCAAGAGG	(AC) <sub>16</sub>	13	108–140	1	PET	1.00	0.91	0.97	EF666987
Mar080	F: ATGGATCATTCCGCTTCTGT R: AACCTTCAGCCCCAACCAT	(TG) <sub>12</sub> (CA) <sub>2</sub> (TGCG) <sub>3</sub>	15	206–233	2	VIC	1.00	0.90	0.62	EF666988
Mar102	F: CCAGTGTGGAGGTCCCTGTAA R: TGTAAGAAAGCTGGAAAACACA	(GA) <sub>23</sub>	26	353–463	2	VIC	0.92	0.97	0.44	EF666989
Mar105	F: GCCCAGGGATAATCCAAT R: TTAATCCTCTCCCCCTCCTC	(AC) <sub>16</sub>	8	287–303	2	FAM	0.42	0.73	>0.001	EF666990
Mar113	F: AAGAGCTGCTGTGGTTGT R: TCAGCTGGGAATCAGGTCTT	(AC) <sub>12</sub>	16	100–125	2	PET	1.00	0.94	0.88	EF666991

Forward (F) primers were labeled with a fluorescent dye (FAM, VIC, NED, PET). R: reverse primer. Repeat motifs are given for the cloned allele. Number of alleles (A), size ranges, expected ( $H_e$ ) and observed ( $H_o$ ) heterozygosities and  $p$ -values of departure from Hardy–Weinberg-Equilibrium (HWE) are given for 24 individuals from a population near Gotha (Germany). Loci can be amplified in two PCR multiplex reactions (set 1 and 2)

<sup>a</sup> a 5' pig-tail extension (GTTTCTT) was used for Mar058R

**Table 2** Results of cross-species amplifications with the newly developed primers for the two European species *M. agrestis* and *M. thomasi* and the American species *M. pennsylvanicus*

Locus	<i>M. agrestis</i>	<i>M. thomasi</i>	<i>M. pennsylvanicus</i>
Mar003	p	p	–
Mar012	–	p	–
Mar016	p	m	p
Mar049	p	p	p
Mar058	–	–	–
Mar063	p	p	p
Mar076	–	p	p
Mar080	p	p	p
Mar102	–	–	–
Mar105	–	–	–
Mar113	p	–	p

p: polymorphic; m: monomorphic; –: no amplification

amplifications of microsatellite loci were carried out in two sets of five and six loci each using the Qiagen multiplex PCR kit (Table 1). Amplifications were done in 10 µl reaction volumes following otherwise the recommendations of the manufacturer. Fragment separation was done by electrophoresis on an ABI Prism 3100 Genetic Analyser (Applied Biosystems), and fragment length was determined in comparison to an internal standard (GeneScan™-500LIZ™, Applied Biosystems) using Genemapper software version 3.7 (Applied Biosystems).

Variability at the loci was tested with 24 samples from a *M. arvalis* population from Gotha, Germany (50°59'N, 10°41'E). Numbers of alleles per locus, fragment sizes, and observed and expected heterozygosities are listed in Table 1. Statistical tests with Arlequin 3.1 (Excoffier et al. 2005) provided no significant evidence for linkage disequilibrium among loci and no significant deviations from Hardy-Weinberg expectations except for locus Mar105 ( $p < 0.001$ ). Closer inspection of the data revealed that all males in the population were apparently homozygous whereas 10 out of 12 females were heterozygous at this locus which strongly suggests X-chromosomal linkage.

We evaluated cross species amplification in two European and one American *Microtus* species: *M. agrestis*

( $N = 4$ ), *M. thomasi* ( $N = 4$ ), *M. pennsylvanicus* ( $N = 2$ ). Most loci were amplified successfully in several species, and only Mar058, Mar102, and Mar105 could not be amplified in any other species than *M. arvalis* (Table 2). Polymorphisms were detected in most species even though cross species tests were performed with only a few samples. This suggests that at least some of these new markers could be applied to a wide variety of species within the genus *Microtus*. The markers are currently being employed in studies of dispersal and reproductive success in natural and experimental populations (Fink et al. 2006).

**Acknowledgements** Thanks go to Sabine Fink, Jörg Hahne and Susanne Tellenbach for technical assistance. Support by the Stiftung zur Förderung der wissenschaftlichen Forschung an der Universität Bern is gratefully acknowledged. This work was funded in part by Swiss National Science Foundation grant 3100A0-112072.

## References

- Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: An integrated software package for population genetics data analysis. Evol Bioinform 1:47–50
- Fink S, Excoffier L, Heckel G (2004) Mitochondrial gene diversity in the common vole *Microtus arvalis* shaped by historical divergence and local adaptations. Mol Ecol 13:3501–3514
- Fink S, Excoffier L, Heckel G (2006) Mammalian monogamy is not controlled by a single gene. Proc Natl Acad Sci USA 103: 10956–10960
- Glenn TC, Schable NA (2005) Isolating microsatellite DNA loci. Meth Enzymol 395:202–222
- Heckel G, Burri R, Fink S, Desmet J-F, Excoffier L (2005) Genetic structure and colonization processes in European populations of the common vole *Microtus arvalis*. Evolution 59:2231–2242
- Mitchell-Jones AJ, Amori G, Bogdanowicz W, Krystufek B, Reijnders PJH, Spitsenberger F, Stubbe M, Thissen JBM, Vohralík V, Zima J (1999) The atlas of European mammals. T. Poyser, A.D. Poyser, London
- Rozén S, Skáletsky H (2000) Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) Bioinformatics methods and protocols: methods in molecular biology. Humana Press, Totowa, NJ, pp 365–386
- Schweizer M, Excoffier L, Heckel G (2007) Fine-scale genetic structure and dispersal patterns in the common vole *Microtus arvalis*. Mol Ecol 16:2463–2473