

# Molecular and functional heterogeneity of GABAergic synapses

Jean-Marc Fritschy · Patrizia Panzanelli ·  
Shiva K. Tyagarajan

Received: 19 November 2011 / Revised: 16 January 2012 / Accepted: 19 January 2012 / Published online: 8 February 2012  
© Springer Basel AG 2012

**Abstract** Knowledge of the functional organization of the GABAergic system, the main inhibitory neurotransmitter system, in the CNS has increased remarkably in recent years. In particular, substantial progress has been made in elucidating the molecular mechanisms underlying the formation and plasticity of GABAergic synapses. Evidence available ascribes a key role to the cytoplasmic protein gephyrin to form a postsynaptic scaffold anchoring GABA<sub>A</sub> receptors along with other transmembrane proteins and signaling molecules in the postsynaptic density. However, the mechanisms of gephyrin scaffolding remain elusive, notably because gephyrin can auto-aggregate spontaneously and lacks PDZ protein interaction domains found in a majority of scaffolding proteins. In addition, the structural diversity of GABA<sub>A</sub> receptors, which are pentameric channels encoded by a large family of subunits, has been largely overlooked in these studies. Finally, the role of the dystrophin-glycoprotein complex, present in a subset of GABAergic synapses in cortical structures, remains ill-defined. In this review, we discuss recent results derived mainly from the analysis of mutant mice lacking a specific GABA<sub>A</sub> receptor subtype or a core protein of the GABAergic postsynaptic density (neuroligin-2, collybistin),

highlighting the molecular diversity of GABAergic synapses and its relevance for brain plasticity and function. In addition, we discuss the contribution of the dystrophin-glycoprotein complex to the molecular and functional heterogeneity of GABAergic synapses.

**Keywords** GABA<sub>A</sub> receptor · Gephyrin · Neuroligin, collybistin, dystrophin-glycoprotein complex · Synaptic plasticity

## Introduction

Early electron microscopy studies uncovered a fundamental dichotomy in the ultrastructure of synapses in vertebrate CNS, classified as type I (asymmetric) and type II (symmetric) synapses, based on the width of their electron-dense postsynaptic density (PSD) [1]. Subsequently, identification of glutamate (and aspartate) and GABA (and glycine) as neurotransmitters in the CNS, combined with the advent of immunohistochemistry, showed that this dichotomy corresponds to excitatory and inhibitory synapses, respectively. This structural and neurochemical distinction reflects major differences in biochemical composition of the PSD in both types of synapses. The PSD contains distinct sets of scaffolding proteins associated with neurotransmitter receptors and signaling molecules. Biochemical fractionation revealed hundreds of proteins in the PSD of glutamatergic synapses. These include numerous PDZ domain-containing scaffolding proteins, notably PSD-95 and other members of the family of membrane-associated guanylate kinase proteins (MAGUKs) [2]. The PDZ interaction motifs are a characteristic feature of many scaffolding proteins, allowing to build modular protein complexes [3], thereby ensuring rapid, efficient, and

---

J.-M. Fritschy (✉) · S. K. Tyagarajan  
Institute of Pharmacology and Toxicology, University of Zurich,  
Winterthurerstrasse 190, 8057 Zurich, Switzerland  
e-mail: fritschy@pharma.uzh.ch

J.-M. Fritschy · S. K. Tyagarajan  
Neuroscience Center Zurich (ZNZ), University of Zurich,  
Winterthurerstrasse 190, 8057 Zurich, Switzerland

P. Panzanelli  
Department of Anatomy, Pharmacology and Forensic Medicine  
and National Institute of Neuroscience-Italy, University of Turin,  
Turin, Italy

site-specific propagation of incoming synaptic inputs [4]. The phototransduction complex assembled by INAD in *Drosophila* photoreceptors is a prototypical example of such signaling scaffold organized by PDZ motifs [5].

In glutamatergic synapses, MAGUKS play a key role in assembly and regulation of the PSD [6, 7]. MAGUKS are membrane-anchored by specific lipid modifications (e.g., palmitoylation) and/or interact with components of the cytoskeleton to ensure long-term structural stability of the PSD [8, 9]. In addition, they contribute to plasticity of synaptic transmission, mainly by binding to specific effector proteins, which induce posttranslational modifications regulating the trafficking and function of neurotransmitter receptors and other signaling molecules. Given the central role of PSD scaffolding molecules, even minor alterations in their expression, trafficking, regulation, or function have been associated with brain disease [7, 10–12]. Therefore, identifying and correcting these dysfunctions might provide promising new avenues for drug therapy.

In striking contrast with type I synapses, the PSD of GABAergic and glycinergic synapses contains only a few proteins with a PDZ domain. Rather, they are organized around a core scaffolding protein, gephyrin, which forms multimeric complexes by auto-aggregation and by interacting with other postsynaptic proteins, including GABA<sub>A</sub> receptors (GABA<sub>A</sub>R), glycine receptors (GlyR), neuroligins (NL), and collybistin (CB) [13]. Models to explain the formation and maintenance of either glycinergic or GABAergic synapses are still fragmentary [14–16], notably because the biochemical composition of their PSD is poorly characterized and because gephyrin structure is only partially resolved. The paucity of PDZ domains in proteins forming GABAergic and glycinergic PSDs implies, however, that protein–protein interactions are based largely on other motifs. While the binding site of the GlyR  $\beta$  subunit intracellular loop to gephyrin has been well characterized [17, 18], much less is known about GABA<sub>A</sub>R subunits, although information available suggests direct interactions with members of the  $\alpha$  subunit family [19–21]. A major challenge for studying GABA<sub>A</sub>R-gephyrin interactions is the presence of multiple receptor subtypes differing in subunit composition, each of them likely interacting differently with gephyrin and its partners.

PSDs typically are associated with specific subcellular compartments; glutamatergic synapses being formed on dendritic spines whereas GABAergic synapses occurring mainly on neuronal somata, dendritic shafts, and axon initial segments. Therefore, association with specific cytoskeletal elements and/or transmembrane proteins linked to extracellular matrix proteins likely is of primary relevance for the formation and maintenance of PSDs. Although gephyrin was characterized initially as a tubulin-

binding protein, the role of the cytoskeleton for anchoring GABAergic and glycinergic PSDs remains largely obscure. Furthermore, PSDs interact closely with presynaptic terminals by means of trans-synaptic ligand/receptor protein complexes, such as neuroligins/NLs, integrins, ephrins, cadherins, etc., raising the possibility that the subcellular localization of synapses is determined by specific molecular components of the PSD.

The purpose of this review is to highlight current knowledge of proteins interacting with gephyrin at GABAergic synapses. We will discuss how molecular and functional heterogeneity of GABAergic synapses arises from these differential interactions. We will also present evidence to reconsider the role of gephyrin as anchoring molecule in favor of gephyrin being the hub of a signaling scaffold organized in concert with specific GABA<sub>A</sub>R subtypes to regulate GABAergic synapse function.

## Overview of major PSD proteins at inhibitory synapses

### Gephyrin

At the PSD of inhibitory synapses, anchoring GABA<sub>A</sub>R and GlyR is ensured, at least in part, by a scaffold assembled upon self-oligomerization of gephyrin. These interactions will be discussed in more detail in the following sections. Additional well-characterized proteins associated with gephyrin include the transmembrane protein NL2 and CB, a member of the Dbl family of guanine nucleotide exchange factors (GEF), discovered upon its ability to translocate gephyrin to the cell surface in non-neuronal cells (reviewed in [13]). A number of additional gephyrin-interacting proteins have been reported, which are involved mainly in regulation of the cytoskeleton and gephyrin trafficking. So far, however, proteins directly regulating gephyrin auto-aggregation have not been described, despite the fact this aggregation exclusively occurs at postsynaptic sites.

Gephyrin is a highly conserved molecule, whose primary role in the living kingdom is to catalyze the synthesis of molybdenum cofactor (Moco) [22]. In vertebrates, gephyrin acts in addition as a postsynaptic scaffolding molecule, and current evidence from rodents indicates that Moco synthesis in brain is restricted to astrocytes [23]. Gephyrin has two major functional domains, E and G, linked by an unstructured C-domain, which contains most gephyrin regulatory sites and binding sites for interacting proteins. While the crystal structure of the G and E domains has been partially determined, the instability of full-length recombinant gephyrin in solution and the lack of information about the structure of the C-domain are major obstacles for elucidating gephyrin function and the

mechanism of gephyrin auto-aggregation [13]. Current models posit either formation of hexameric gephyrin scaffolds, with six gephyrin molecules binding to three GlyR [18], or aggregation of gephyrin trimers, each of which is bound to one GlyR [13].

These models of gephyrin aggregation are based on partial structural information, localization of the GlyR binding site in the E-domain, and identification of surface-exposed loops regulating gephyrin postsynaptic aggregation [13, 18, 24]. Although available experimental evidence does not allow distinguishing between them, an important outcome of gephyrin structure is its enzymatic activity, which requires the active sites on the G and the E domains to be in close spatial proximity. The model of gephyrin trimers takes this requirement into consideration [13]. Data from GlyR single-particle tracking experiments and gephyrin fluorescence recovery after photobleaching revealed that gephyrin modulates synaptic retention time and lateral mobility of these receptors, and that GlyR diffusion properties modulate gephyrin clustering [25, 26]. Therefore, a dynamic exchange of gephyrin and receptors occurs in apparently stable postsynaptic structures. Furthermore, time-lapse recordings from neurons expressing fluorescently-tagged gephyrin revealed that gephyrin clusters are constantly remodeled and can move in dendrites over distances of several micrometers in concert with apposed presynaptic terminals [27]. Collectively, these observations are consistent with both views that gephyrin units (e.g., trimer) can be added and removed from the scaffold and that the gephyrin scaffold acts as a stable but mobile entity in dendrites.

There are multiple splice-variants of gephyrin generated by alternative splicing of cassettes localized mainly in the G- and C-domain (see [13] for nomenclature). The splice variants capable of Moco synthesis have been identified, but they do not necessarily differ from those forming postsynaptic clusters in neurons; further, insertion of the G2 splice cassette, a gephyrin isoform expressed in both neuronal and non-neuronal tissues, prevents gephyrin aggregation and Moco synthesis, and has been proposed to regulate gephyrin function, synaptic localization, and clustering [23, 25, 28]. Further characterization of gephyrin splice variants, notably in the C-domain, will be necessary to better understand the regulation of gephyrin enzymatic activity and clustering in different types of synapses.

Recent evidence points to gephyrin posttranslational modification by phosphorylation as a mechanism regulating its postsynaptic function. Although gephyrin is a known phospho-protein [29], it is only in 2007 that a first report suggested that phosphorylation-dependent recruitment of the peptidyl-prolyl isomerase Pin1 to a site located in the C-domain regulates GlyR binding [30]. We have identified a novel phosphorylation site on gephyrin,


Ser270, and shown that it selectively regulates formation of postsynaptic gephyrin clusters in vitro and in vivo under the control of GSK3 $\beta$  [31]. These findings provide a novel mechanism to regulate GABAergic synapse formation and function by a phosphorylation cascade activated by a plethora of extra- and intra-cellular signals and controlling basic cellular functions.

## Neurologin-2

Neurexins and NLs gained major interest upon discovery that they induce formation of synapses in recombinant non-neuronal cell systems. Reports of preferential distribution of NL1 in glutamatergic synapses and NL2 in GABAergic synapses [32–34] fuelled considerable efforts to elucidate their role in synapse formation and specification in the CNS. Importantly, in the context of the present review, NL2 has been shown to interact directly with gephyrin and this interaction is necessary for postsynaptic localization of gephyrin clusters at GABAergic sites [35]. Apparently, NL2 is absent from glycinergic synapses, whereas NL3 and NL4 have been reported to be present in a subset of GABAergic and glycinergic synapses, respectively [36, 37]. The significance of a segregated distribution of NLs to different types of synapses is not fully understood [38], because this apparent selectivity is lost upon overexpression and because NL3 has been reported to be present in both glutamatergic and GABAergic synapses in vivo. It is also likely that NLs are functionally interchangeable to some degree, as seen in targeted gene deletion experiments, in which the loss of a particular NL isoform appears to be compensated in part by another isoform [39]. Nevertheless, the significance of the NL2-gephyrin interaction was highlighted by a study showing that NL2 also interacts with CB, in a manner that favors formation and regulation of gephyrin clusters at postsynaptic sites (see below). Therefore, NL2 is widely considered to play a crucial role to initiate the formation of a GABAergic PSD.

## Collybistin

CB is a neuron-specific Rho-GEF encoded by a single gene (*ARHGEF9*) subject to alternative splicing. In rat, there are three main CB variants, CB1–3, differing in the C-terminal region. They all possess three major functional domains: an N-terminal SH3 domain, encoded by a spliced exon (SH3+ or SH3–), a DH (or GEF) domain selectively activating the small GTPase Cdc42, and a PH domain (Fig. 1). Available literature is somewhat confusing about the existence and structure of CB variants in other species. In human, CB is also known as hPEM2 and only one sequence, with a C terminus identical to rat CB3 has been described [40]. In mouse, four transcripts have been identified, which are



**Fig. 1** Comparison of CB isoforms in rat, human, and mouse. The three main functional domains are indicated, *different colors* denote distinct sequences in the N- or C-termini. The number of residues present in each domain or in each isoform is given on top of each variant. CB isoforms are best characterized in rat, where the existence of three splice variants (CB1–CB3) differing in the C-terminal domain is well established. In addition, each of CB1–CB3 variants is

believed to be N-terminally spliced to include or exclude the SH3 domain. The C-terminal region of CB3 is identical to hPEM2, the only CB isoform described in human. The N terminus of CB3 differs slightly from that of CB1/2. In mice, the CB1–CB3 nomenclature does not apply because the sequence closest to rat CB3 is encoded in three variants with different N-termini, whereas the sequence closest to either CB1 or CB2 has a unique C terminus

largely conserved compared to human and rat CB3, but differ in their N terminus (Fig. 1). The relative abundance of CB splice variants in the CNS and their expression pattern across different brain regions or developmental stages have not been investigated systematically. A recent report using antibodies recognizing the majority of CB isoforms (notably CB3) confirmed the localization of CB at GABAergic PSDs in most brain regions [41].

The significance of CB for glycinergic and GABAergic synaptic function is underscored by the discovery of several loss-of-function mutations in *ARGF9* underlying severe forms of X-linked mental retardation and hyperkplexia [42, 43]. In recombinant expression systems, CB is essential for cell-surface translocation of gephyrin [44]. However, this function is only assumed by CB variants lacking the SH3 domain, mainly CB2<sub>SH3-</sub>, suggesting that the SH3 domain somehow controls CB activity. A plausible model posits that, in analogy to other GEFs, the SH3 domain exerts an auto-inhibitory effect on the DH domain (and hence on activation of Cdc-42) [35]. CB binds gephyrin in the C-domain, whereas the binding site of gephyrin on CB partially overlaps with the binding of Cdc42, which led to speculation that they might be

mutually exclusive [45] (but see [46]). A recent study suggested, however, that CB-mediated gephyrin postsynaptic clustering is not dependent on a functional DH domain, but requires the PH domain [47]. Because gephyrin postsynaptic clusters were observed in the brain of conditional *Cdc42*-deficient mice, this study even concluded that *Cdc42* is dispensable for gephyrin aggregation.

In addition to gephyrin, CB interacts directly with NL2 via its SH3 domain. Based on data from recombinant systems, this interaction was suggested to cause a conformational change relieving the auto-inhibition of the DH domain, thereby facilitating cell-surface translocation of gephyrin and GABA<sub>A</sub>R [35]. In neurons, the activation of CB<sub>SH3+</sub> by NL2 was postulated to facilitate gephyrin clustering at postsynaptic sites upon binding to phosphoinositol-3-phosphate in the cell membrane via its PH domain. This model received further support in a report showing that CB binds to GABA<sub>A</sub>R  $\alpha 2$  (and  $\alpha 3$ ) subunit and that this interaction leads to CB activation and translocation of trimeric CB-gephyrin- $\alpha 2$  subunit complexes to the cell membrane [48]. However, the role of NL2 was not examined in this study. Furthermore, in view of the multiplicity of CB isoforms and of the heterogeneity of

GABA<sub>A</sub>Rs, which form multiple subtypes differing in subunit composition and cellular/subcellular localization [49], such models are necessarily fragmentary and do not take into account the diversity and exquisite functional specialization of GABAergic synapses across the CNS.

Insight into the roles of CB isoforms for gephyrin clustering was derived from over-expression studies in transfected neurons, which aimed to distinguish between CB2<sub>SH3+</sub> and CB2<sub>SH3-</sub> [46, 50], as well as to elucidate the contribution of Cdc42, the only known substrate of CB. In both studies, CB over-expression contributed to stabilize gephyrin, as evidenced by a marked increase in size and density of postsynaptic gephyrin clusters. Interestingly, CB2<sub>SH3-</sub> facilitated the formation of supernumerary postsynaptic clusters, whereas CB2<sub>SH3+</sub> favored formation of intracellular gephyrin aggregates, as illustrated in Fig. 2. Further, deletion of the PH domain of CB2 led to gephyrin retention along the dendritic cytoskeleton, whereas deletion of the DH domain strongly interfered with gephyrin postsynaptic clustering. Finally, over-expression of Cdc42, using dominant negative and constitutively active mutants for comparison, revealed that Cdc42 cooperates with CB2<sub>SH3-</sub> for postsynaptic targeting of gephyrin and that it regulates gephyrin cluster size at postsynaptic sites [46]. These data add to the model proposed by Poulopoulos et al. [35], by unraveling distinct functions of CB2 isoforms for gephyrin transport, postsynaptic targeting, and regulation of postsynaptic clustering in concert with Cdc42. However, they suggest that the postulated conformation-dependent activation of CB<sub>SH3+</sub> by NL2 might not be crucial for proper CB function at GABAergic PSDs. Further insight into the roles of CB for gephyrin trafficking and clustering will require the analysis of the other splice variants (CB1, CB3), as well as more knowledge of their expression pattern and regulation.

### Differences between GABAergic and glycinergic synapses

Because gephyrin is found in both glycinergic and GABAergic synapses, it is generally considered that the assembly of signaling scaffolds by gephyrin is similar in both types of synapses. This concept is reinforced by the homology between GABA<sub>A</sub>R and GlyR, which are members of the family of Cys-loop ligand-gated ion channels. However, there are a number of important differences between the two types of synapses, which suggest that their formation and regulation might depend on distinct mechanisms (Table 1).

First, despite being closely related, GABA<sub>A</sub>Rs and GlyRs differ fundamentally in their affinity for gephyrin. Recent reports identified the binding site of the GABA<sub>A</sub>R  $\alpha 1$ – $\alpha 3$  subunits on the gephyrin E-domain and showed substantial overlap with the GlyR  $\beta$  subunit, albeit with a


>30-fold lower affinity [19, 51]. Nevertheless, expression of a mutant GABA<sub>A</sub>R lacking the interaction site in the M3–M4 loop of the  $\alpha 3$  subunit disrupted its postsynaptic clustering in cultured hippocampal neurons. These findings extend a previous report, which identified a gephyrin-interaction motif in the M3–M4 intracellular loop of the  $\alpha 2$  subunit, sufficient to direct chimeric GABA<sub>A</sub>R to gephyrin-rich postsynaptic sites [20].

Second, as discussed in the next section, postsynaptic clustering of some GABA<sub>A</sub>R subtypes is possible in the absence of gephyrin. In contrast, GlyR postsynaptic clustering appears to be dependent on the presence of gephyrin clusters [52], mirroring the high affinity of their interaction and confirming observations that GlyR trafficking in neurons is facilitated by association with gephyrin [53].

A third major difference between GABAergic and glycinergic synapses is the dependence of gephyrin on CB for postsynaptic aggregation. Unexpectedly, CB-knockout (KO) mice exhibit no apparent morphological and functional alteration of glycinergic synapses in the brainstem and spinal cord [54]. In particular, they do not show any signs of spasticity typical of hypo-glycinergic function at birth, which are prominent in mice lacking the type 1 glycine transporter (GlyT1) [55]. These results indicated that CB either is dispensable for gephyrin/GlyR clustering at glycinergic PSDs or is functionally replaced by a homologous protein. In contrast, as detailed in the next section, CB deficiency produces marked alterations of gephyrin clustering at GABAergic postsynaptic sites.

A fourth difference is the differential targeting of NLs to GABAergic and glycinergic synapses. As noted above in the section on NL2, GABAergic, but not glycinergic synapses, selectively contain this NL isoform, whereas NL3 and NL4 have been reported in glycinergic synapses, at least in the retina [37]. Since NL2 interacts with CB and gephyrin at GABAergic synapses, this difference in localization might explain why CB deletion does not affect the PSD of glycinergic synapses.

A fifth difference is the existence of multiple, presumably dozens of GABA<sub>A</sub>R subtypes, differing in subunit composition in adult brain, compared to a few GlyR subtypes, assembled mainly from  $\alpha 1$  or  $\alpha 3$  and  $\beta$  subunit variants. The multiplicity of GABA<sub>A</sub>R underlies molecular and functional heterogeneity of GABAergic synapses, as well as differential targeting to extrasynaptic versus postsynaptic sites. There is evidence that receptors differing in  $\alpha$  subunit variant are functionally not interchangeable, and that only GABA<sub>A</sub>R targeted postsynaptically contribute to formation of gephyrin clusters [56, 57]. Therefore, interactions of GABA<sub>A</sub>R subtypes with gephyrin and other GABAergic PSD proteins are likely more heterogeneous and synapse-specific than GlyR-gephyrin interactions. These limitations are well evident in single-particle



**Fig. 2** Differential effects of  $CB2_{SH3+}$  and  $CB2_{SH3-}$  on the postsynaptic clustering of gephyrin; adapted from [46]. Primary cultures from hippocampal embryonic neurons were transfected with tagged cDNA constructs after 11 days-in vitro (DIV) and processed for immunofluorescence staining with the markers indicated 4–7 days later. **a** Control cell showing extensive co-localization of gephyrin and  $\alpha 2$  subunit immunofluorescence at postsynaptic sites. **b** Upon transfection with  $mycCB2_{SH3-}$ , there is a marked increase in the density and size of gephyrin/ $\alpha 2$  clusters in the soma and on dendrites, suggesting that

$CB2_{SH3-}$  stabilizes gephyrin and favors its postsynaptic clustering. **c,d** Similar effects were observed in neurons co-transfected with eGFP-gephyrin and  $myc-CB2_{SH3+}$  (**c**) or  $myc-CB2_{SH3-}$  (**d**); however,  $CB2_{SH3+}$  favored the formation of non-synaptic aggregates (*arrows*), whereas  $CB2_{SH3-}$  led to increased density of postsynaptic aggregates, recognized by apposition to the presynaptic marker synapsin 1 (*arrowheads*). The formation of non-synaptic aggregates suggests saturation of molecules, such as NL2, which normally ensure postsynaptic localization of  $CB2_{SH3+}$ . Scale bars 20  $\mu m$

tracking studies of GABA<sub>A</sub>R cell surface mobility, which cannot be performed on a uniquely identified receptor subtype. Nevertheless, evidence available indicates that diffusion properties and synaptic confinement of GABA<sub>A</sub>Rs containing the  $\gamma 2$  subunit are modulated by

enhanced excitatory synaptic activity, in a  $Ca^{2+}$ -dependent manner [58].

Several reports suggested that, despite the tight biochemical association between GlyRs and gephyrin, there is no direct correlation between gephyrin availability at

**Table 1** Differences between GABAergic and glycinergic synapses related to gephyrin

Feature	GABAergic	Glycinergic
1. Gephyrin–receptor interaction	Low affinity ( $\alpha 1$ , $\alpha 2$ , $\alpha 3$ subunits)	High affinity ( $\beta$ subunit); insertion of the binding motif (residues 398–410 of the $\beta$ subunit) into a recombinant protein is sufficient for interaction with gephyrin [106]
2. Receptor dependence on gephyrin for postsynaptic clustering	Receptor subtype-dependent ( $\alpha 2 \gg \alpha 1$ )	Very high (clustering of GlyRs depends on gephyrin [107])
3. Gephyrin dependence on collybistin for postsynaptic clustering	Variable (neuron and/or synapse dependent)	None
4. Neuroligin isoform present	NL2 (NL3, NL4)	NL4
5. Receptor heterogeneity	Numerous receptor subtypes (19 subunits)	Limited repertoire

See main text for further explanations and references

postsynaptic sites and synaptic retention time or cell-surface diffusion parameters of GlyRs in neurons. Therefore, the high affinity of gephyrin for the GlyR  $\beta$  loop trafficking does not imply a one-to-one relationship. It is conceivable that gephyrin scaffolds contain a limited number of binding sites for GlyR [25, 59, 60]. Further evidence for a dissociation between GlyR and gephyrin clustering dynamics was provided by analyzing the interaction between gephyrin and heat-shock cognate protein 70 (Hsc70), a chaperone modulating ubiquitination of its substrates [61]. These authors showed that Hsc70 selectively regulates gephyrin clustering in cultured spinal cord neurons, without affecting GlyR clustering and cell-surface diffusion kinetics.

### Heterogeneity of GABAergic synapses revealed in mice with targeted gene deletions

By immunofluorescence staining and immunoelectron microscopy in brain sections, gephyrin is selectively detected at postsynaptic sites of glycinergic and most GABAergic synapses, forming puncta (clusters) that are colocalized with GlyRs and GABA<sub>A</sub>Rs, respectively [62–65]. Initial observations of a functional interaction between gephyrin and GABA<sub>A</sub>R were made in  $\gamma 2$ -KO mice, in which postsynaptic clustering of GABA<sub>A</sub>R and gephyrin was strongly impaired compared to wild-type and heterozygous littermates [66–68]. Conversely, GABA<sub>A</sub>R clustering was widely disrupted in neurons from gephyrin-KO mice [69]. This interdependence was later shown not to be absolute, as GABA<sub>A</sub>R containing the  $\alpha 1$  subunit, for instance, can form postsynaptic clusters mediating synaptic transmission in the absence of gephyrin [70, 71]. However, ablation of GABA<sub>A</sub>Rs by targeted deletion of an  $\alpha$  subunit variant often results in the disruption of postsynaptic gephyrin clusters and formation of large intra-cellular gephyrin aggregates [56, 72–76], confirming the dependence of gephyrin on

GABA<sub>A</sub>Rs for postsynaptic localization. Importantly, in thalamic neurons expressing both  $\alpha 1$ -GABA<sub>A</sub>R postsynaptically and  $\alpha 4$ -GABA<sub>A</sub>R extrasynaptically, gephyrin clustering was not rescued following  $\alpha 1$  subunit deletion, indicating that  $\alpha 4$ -GABA<sub>A</sub>R cannot interact with gephyrin [56, 75].

Beyond GABA<sub>A</sub>Rs, evidence for multiple mechanisms of gephyrin postsynaptic clustering at GABAergic synapses was provided by the analysis of CB- and NL2-KO mice. In particular, in GABAergic synapses of the forebrain and cerebellum, multiple phenotypes were observed in CB-KO mice, ranging from disruption of both gephyrin and GABA<sub>A</sub>R clusters (e.g., CA1 pyramidal cells) to complete preservation of these clusters (e.g., PV<sup>+</sup> interneurons in CA1), passing by partial alteration (loss of gephyrin, but not GABA<sub>A</sub>R, clusters in Purkinje cells of the cerebellum) [54, 77]. Although preservation of gephyrin clusters might point to a functional homologue of CB, the phenotype of Purkinje cells rather indicates that clustering of gephyrin and of GABA<sub>A</sub>R depends on different mechanisms, with only the former requiring the presence of CB. In NL2-KO mice, differential loss of postsynaptic clusters was observed in different subcellular compartments of CA1 pyramidal cells and dentate gyrus granule cells, with the main disruption of gephyrin and GABA<sub>A</sub>R clusters occurring on the soma, at synapses formed by basket cells [35, 78, 79]. Thus, these reports pointed to distinct clustering mechanisms for postsynaptic proteins of perisomatic versus dendritic GABAergic synapses.


Further insight into GABAergic synapse heterogeneity was provided by the analysis of postsynaptic proteins in the hippocampal formation of  $\alpha 2$ -KO mice [74]. Focusing on CA1 pyramidal cells, where  $\alpha 2$ -GABA<sub>A</sub>R predominate in the perisomatic area and the axon-initial segment [80, 81], a pronounced reduction of gephyrin clusters was observed, notably on cell somata, on the axon-initial segment, as well as in stratum radiatum and stratum oriens (Fig. 3a). While this loss of gephyrin clusters was suggestive of a disruption

of the PSD, only moderate changes in GABAergic miniature inhibitory postsynaptic currents (mIPSC) were observed in mice lacking  $\alpha 2$ -GABA<sub>A</sub>R, with unchanged amplitude and 40% reduction in frequency. Therefore, these results indicated preservation of GABAergic function despite strongly altered gephyrin clustering at postsynaptic sites. Analysis of  $\alpha 1$ -GABA<sub>A</sub>R and NL2 revealed that their presence at perisomatic postsynaptic sites was preserved in CA1 pyramidal cells of  $\alpha 2$ -KO mice (Fig. 3b), whereas they were absent from the axon-initial segment. This striking subcellular difference was correlated with the presence of the dystrophin–glycoprotein complex (DGC) in perisomatic synapses (Fig. 3c), suggesting that this protein complex represents an alternate mechanism contributing to formation and maintenance of PSD in a subset of GABAergic synapses [74], and confirming that perisomatic and dendritic GABAergic PSDs are molecularly distinct.

### The dystrophin–glycoprotein complex

The DGC is a transmembrane signaling complex present in striate and cardiac muscle cells, kidney tubular epithelial cells, neurons, and astrocytes, linking the extracellular matrix to the actin cytoskeleton (reviewed in [82–84]). The DGC is essential for normal brain development and synaptic function and for maintaining the structural integrity of the sarcolemma. Mutations affecting the formation, post-translational modification, or function of the DGC cause multiple forms of hereditary muscle dystrophies. While the contribution of the DGC to muscle disease has been thoroughly investigated, patients with Duchenne and Becker muscular dystrophies also show signs of cognitive impairments and learning disabilities that have been related to its functions in the CNS [83, 85]. In neurons, dystrophin was shown initially to be present in PSD fraction of purified brain membranes [86]. Subsequent studies demonstrated its association with a subset of GABAergic synapses [87, 88], where it regulates GABA<sub>A</sub>R clustering and GABAergic synaptic function, as well as glutamatergic synaptic plasticity (reviewed in [82, 83]). In mdx mice, direct involvement of dystrophin in synaptic alterations has been demonstrated by dystrophin rescue experiments, which normalize mIPSC amplitude and frequency, as well as LTP in CA1 pyramidal cells [89, 90].

The molecular composition of the DGC differs between muscle sarcolemma, neuromuscular junction, brain and retina, and astrocytes [91, 92], where, for instance, the DGC is involved in proper membrane targeting of aquaporin four at end-feet in contact with brain blood vessels [93, 94]. Common components of the DGC include  $\alpha$ - and  $\beta$ -dystroglycan, full-length dystrophin (Dp427), sarcoglycans, a dystrobrevin isoform, and syntrophins. The



predominant dystrophin isoform found in astrocytes is the short N-terminal variant Dp71.

Association of the DGC with cytoplasmic signaling molecules is ensured by direct protein–protein interactions



◀ **Fig. 3** Loss of gephyrin clusters, but preservation of  $\alpha 1$ -subunit, NL2, and dystrophin clusters in perisomatic synapses of  $\alpha 2$ -KO mice; adapted from [74]. **a, a'** Distribution of gephyrin clusters in the CA1 region of the hippocampus, illustrating the strong reduction in  $\alpha 2$ -KO mice and the appearance of large intracellular gephyrin aggregates. **b, b'** Despite this effect, clustering of  $\alpha 1$  subunit and NL2 was not affected around the pyramidal cell body in mutant mice, as seen by triple immunofluorescence staining. The framed area is shown below in color-separated images. **c, c'** Similarly, the postsynaptic clustering of dystrophin and its colocalization with the  $\alpha 1$  subunit and NL2 is not affected in  $\alpha 2$ -KO mice. Scale bars **a** 50  $\mu\text{m}$ , **b, c** 10  $\mu\text{m}$

with  $\beta$ -dystroglycan or dystrophin or by means of PDZ domains present in syntrophin isoforms. Identified partners of syntrophins include nNOS, aquaporin 4, voltage-gated  $\text{Na}^+$  and  $\text{K}^+$  channels [83], and NLs [95]. Interactions with extracellular signaling molecules critically depends on  $\alpha$ -dystroglycan posttranslational modification by glycosylation. Consequently, mutations in glycosylating enzymes contribute to the pathogenesis of a subset of congenital muscle dystrophies associated with lissencephaly and mental retardation [84, 96].  $\alpha$ -dystroglycan function on glial end-feet appears to be especially crucial for normal brain development, whereas neuronal  $\alpha$ -dystroglycan glycosylation is necessary for proper synaptic plasticity [97].

Localization of the DGC by immunofluorescence has been documented in neurons, using antibodies to full-length dystrophin, Dp71, utrophin,  $\alpha$ -/ $\beta$ -dystroglycan, dystrobrevin, and syntrophins [82, 98]. These studies reveal a systematic colocalization with proteins of the GABAergic PSD in a subset of neurons of cortical areas, including the entire cerebral cortex, hippocampal formation (where Dp71 is selectively present in dentate gyrus granule cells), tectum, and cerebellum. In the hippocampus, the DGC appears to be restricted to principal cells, and is present selectively on the soma and proximal dendrites, suggesting an association with perisomatic synapses (formed by basket cells) [87, 99].

It is not known, however, what determines the presence of the DGC in specific GABAergic synapses of cortical pyramidal cells. In primary neuronal cultures containing only few interneurons, we have shown that presence of a GABAergic terminal is required for the formation of the DGC [100], whereas both gephyrin and GABA<sub>A</sub>Rs can form ectopic clusters facing glutamatergic terminals. In vivo, it is therefore conceivable that a presynaptic factor present selectively in a subset of GABAergic neurons (such as basket cells, but not chandelier cells or dendritic-targeting interneurons in the hippocampus) induces the formation of the DGC and therefore entails these GABAergic synapses with distinct properties for clustering postsynaptic proteins.

In the context of the present review, interaction of DGC members with GABAergic PSD proteins deserves

particular attention. Interaction between  $\alpha$ -dystroglycan and  $\alpha$ -/ $\beta$ -neurexins by means of laminin–neurexin–sex hormone-binding globulin (LNS)/laminin G domains has been reported several years ago [101], and was proposed to compete with  $\alpha$ -latrotoxin binding on neurexins. In addition, binding of synaptic scaffolding molecule (S-SCAM) to  $\beta$ -dystroglycan and NL2 has been characterized in studies showing the selective presence of S-SCAM at inhibitory synaptic sites [102, 103]. S-SCAM has been identified for its synaptogenic role and interactions with NMDA-receptor subunits and NLs. Its binding to  $\beta$ -dystroglycan involves one of two WW domains, whereas binding to NL2 occurs through the second of its three PDZ domains. Recently S-SCAM was shown to bind also SynArfGEF, which selectively activates Arf6, a GTP-binding protein member of the ADP ribosylation factor family. S-SCAM is able to interact also with several MAGUKs (including PSD-95, SAP97, and Homer) [103]. Further, these authors showed that SynArfGEF also interacts with dystrophin and utrophin via its PDZ domain and that it colocalizes with dystrophin and gephyrin at GABAergic synapses. Besides CB, SynArfGEF is only the second GEF identified so far in GABAergic synapses, and may represent an effector signaling molecule related to the DGC. Owing to the multiple effects of Arf6, the role of SynArfGEF is not yet determined. However, it is clear that it cannot substitute CB for gephyrin or GABA<sub>A</sub>R clustering at synapses containing the DGC, as shown in CB-KO mice.

While interactions with S-SCAM appear important for the association of the DGC with NL2 at GABAergic postsynaptic sites, they also provide for the possibility of cross-talk with glutamatergic synapses, in particular via activation of Arf6. Further, the selective association of S-SCAM with NL2 might be of relevance to explain the selective loss of inhibitory PSD proteins in perisomatic synapses of NL2-KO mice (see next section).

### Model of PSD protein clustering at GABAergic synapses

Differences in PSD protein clustering properties in GABAergic synapses in relation to the presence or absence of the DGC indicate that models of GABAergic synapse formation and maintenance need to incorporate the role DGC proteins (and possibly SynArfGEF and Arf6) to account for published results. However, since the DGC is present in a subset of GABAergic synapses only, and is dispensable for GABAergic synapse formation in cultured hippocampal neurons [104], models of clustering that do not involve the DGC are also required.

The models proposed here (Fig. 4) are elaborated on the basis of observations made in  $\alpha 1$ -KO,  $\alpha 2$ -KO, NL2-KO,

CB-KO, and mdx mice and take the following considerations into account:


1. In wild-type mice, while gephyrin, NL2, and CB likely are present in the majority of GABAergic synapses [14, 32, 41, 65], a major distinction is brought about by the presence or absence of the DGC, as exemplified by thalamic relay neurons and cerebellar neurons (Fig. 4a, b). Some cells, like hippocampal pyramidal cells, carry a mixture of both types of synapses (Fig. 4e, f).
2. Interactions between CB, NL2, gephyrin and possibly GABA<sub>A</sub>R contribute to the formation of the postsynaptic scaffold (Fig. 4a, b) [14, 15, 50].
3. In synapses containing multiple  $\alpha$  subunit variants, such as  $\alpha 1$ - and  $\alpha 2$ -subunits in perisomatic synapses of CA1 pyramidal cells (Fig. 4e) [74, 81], it is not known whether they correspond to different receptor subtypes, or whether they are intermingled within single receptors [74].
4. Gephyrin clustering, but not NL2 clustering, requires the presence of postsynaptic GABA<sub>A</sub>R (Fig. 4c, d) [105], in particular those containing the  $\alpha 2$  subunit (Fig. 4g) [74].
5. In contrast, the DGC is not altered in neurons lacking functional GABAergic transmission or gephyrin clusters (Fig. 4d, g) [74, 105], indicating that its formation and postsynaptic localization is not dependent on GABA<sub>A</sub>R or gephyrin.
6. Preservation of NL2 clustering in the absence of GABA<sub>A</sub>R and/or gephyrin suggests a mandatory association between NL2 and S-SCAM/ $\beta$ -dystroglycan complex (Fig. 4d, g) [102]. Further, this association raises the unresolved issue whether the DGC requires NL2 for localizing at GABAergic synapses.
7. Disruption of gephyrin and GABA<sub>A</sub>R clusters in perisomatic, but not dendritic synapses of hippocampal neurons of NL2-KO mice [35, 79] suggests that NL2 is the only NL isoform that interacts with S-SCAM/ $\beta$ -dystroglycan [102] and indicates that postsynaptic clustering of GABA<sub>A</sub>R requires the presence of a NL isoform. As a corollary, gephyrin cluster preservation in dendritic synapses of NL2 mice suggests compensation by another NL isoform.
8. However, these observations cannot be generalized throughout the brain, as, for example, there is no loss of  $\alpha 1$ -GABA<sub>A</sub>R and gephyrin clusters in Purkinje cells of NL2-KO mice (personal observation); the nature of the mechanisms compensating for the absence of NL2 in these cells is not known, but a compensation by NL3 or NL4 is not unlikely.

**Fig. 4** Models of GABAergic PSD scaffold structure in distinct cell types of  $\alpha 1$ -KO,  $\alpha 2$ -KO, mdx (lacking full-length dystrophin), and wild-type mice. **a, b** In wild-type mice, GABAergic PSDs contain GABA<sub>A</sub>Rs, gephyrin, and, in most cases, NL2/3, and CB2<sub>SH3+</sub> and CB2<sub>SH3-</sub>, although these have not been demonstrated, for example in the thalamus (?). In a subset of synapses, for example in Purkinje cells, the DGC is also present, along with its interacting proteins S-SCAM and SynArfGEF. The PSD is anchored to the actin cytoskeleton. Phosphorylation sites on gephyrin and GABA<sub>A</sub>Rs (-P) contribute to dynamic regulation of GABAergic synapses. **c, d** In  $\alpha 1$ -KO mice, gephyrin clusters are disrupted in neurons expressing  $\alpha 1$ -GABA<sub>A</sub>R, but NL2 clustering is preserved in the presence of the DGC, presumably via binding to S-SCAM. **e, f** In CA1 pyramidal cells, distinct GABAergic PSDs are present on the soma and dendrites, characterized by the presence or absence of the DGC, respectively. These synapses contain both  $\alpha 1$ - and  $\alpha 2$ -GABA<sub>A</sub>Rs, or possibly receptors containing both subunit variants. In these cells, CB2<sub>SH3-</sub> and Cdc42 contribute to postsynaptic targeting of gephyrin, as well as regulation of the gephyrin scaffold shape and size. CB2<sub>SH3-</sub> interacts with NL2/3 and GABA<sub>A</sub>Rs (see [46] for details). **g** In  $\alpha 2$ -KO mice, gephyrin clustering is disrupted in synapses containing  $\alpha 2$ -GABA<sub>A</sub>R. In perisomatic synapses of CA1 pyramidal cells, the DGC anchors NL2 via S-SCAM, thereby preserving NL2 and  $\alpha 1$  subunit clustering. **h** Conversely, in mdx mice, a partial reduction of GABA<sub>A</sub>R at perisomatic synapses might be due to absence of NL2 linked to the DGC. Preservation of gephyrin clustering is best explained by the interaction between CB2/NL2/ $\alpha 2$  subunit. Our model postulates a tight interaction between NL2 (*black*) and the DGC, whereas NL isoforms interacting with CB2<sub>SH3+</sub> might be either NL2 or NL3 (*dark grey*)

9. In  $\alpha 2$ -KO mice, a possible direct interaction between  $\alpha 1$ -GABA<sub>A</sub>R and NL2 might explain preservation of  $\alpha 1$ -GABA<sub>A</sub>R clusters in perisomatic synapses (Fig. 4g); whereas the loss of gephyrin clusters in both perisomatic and dendritic synapses (not shown) reveals their dependence on  $\alpha 2$ -GABA<sub>A</sub>Rs, which interact directly with both gephyrin and CB2<sub>SH3+</sub> [48].
10. Gephyrin-independent  $\alpha 1$ -GABA<sub>A</sub>R clustering has been observed in CB-KO mice [54], indicating that these GABA<sub>A</sub>R do not require CB for their proper localization.
11. Absence of dystrophin, such as occurs in mdx mice, causes a partial disruption of  $\alpha 1$ - and  $\alpha 2$ -GABA<sub>A</sub>R clustering, but not gephyrin clustering (Fig. 4h) [87]. Therefore, the mdx mutation might cause loss of postsynaptic NL2 interacting via S-SCAM, whereas the CB2<sub>SH3+</sub>/NL2/gephyrin complex is likely preserved [35, 46]. It is conceivable that the function of NL2 associated with the DGC is distinct from that of NL2/3 associated with CB and gephyrin.

These models also incorporate the postulated functions of CB2 isoforms and Cdc42 (Fig. 4e, f) [35, 46]:

- (1) The ability of CB2<sub>SH3-</sub> to form trimeric complexes with gephyrin and Cdc42 likely contributes to gephyrin postsynaptic targeting, (2) the role of membrane anchored



CB2<sub>SH3+</sub>, interacting with NL2, gephyrin, and  $\alpha 2$ -GABA<sub>A</sub>R for PSD formation and gephyrin clustering, (3) the stabilization of gephyrin clusters might be mediated by membrane-anchored CB2<sub>SH3-</sub>, (4) the regulation of gephyrin postsynaptic cluster size and shape relies on activated (GTP-bound) Cdc-42.

To explain the phenotype of CB-KO mice, in which a differential loss of gephyrin and GABA<sub>A</sub>R occurs in various cell types, the presence or absence of the DGC is not sufficient. For instance,  $\alpha 1$ -GABA<sub>A</sub>Rs are retained in Purkinje cells of CB-KO mice in the absence of gephyrin [54], suggesting their synaptic anchoring via the DGC and NL2. In contrast, in CA1 pyramidal cells, gephyrin,  $\alpha 1$ - and  $\alpha 2$ -GABA<sub>A</sub>R disappear in CB-KO mice, whereas in PV<sup>+</sup> interneurons (and in thalamic VB neurons), gephyrin and  $\alpha 1$ -GABA<sub>A</sub>R are retained [77]. However, since the  $\alpha 2$ -subunit interacts directly with CB2 [48], one might hypothesize that GABAergic synapses containing  $\alpha 2$ -GABA<sub>A</sub>R are selectively affected in CB-KO mice. Consequently, in CA1 pyramidal cells, perisomatic synapses might be made of heteromeric  $\alpha 1/\alpha 2/\beta/\gamma 2$  receptors, which would fail to cluster in the absence of CB.

## Conclusions

Taken together, the results discussed in this review indicate that GABAergic synapses are molecularly heterogeneous, and that this heterogeneity on the postsynaptic side largely arises from the multiple proteins regulating gephyrin clustering and its function as signaling scaffold at the PSD. Moreover, the view that gephyrin forms an inert structure anchoring GABA<sub>A</sub>Rs needs to be replaced by the concept that these receptors are required at postsynaptic sites to enable gephyrin clustering and to establish a signaling hub in the GABAergic PSD. Therefore, the diversity of GABA<sub>A</sub>R subunits likely reflects a diversity of signaling pathways operating at GABAergic synapses. While the precise mechanisms underlying GABA<sub>A</sub>R postsynaptic clustering remain to be elucidated, the models of PSD formation proposed here impart a major role to the neuexin/NL2 complex for specifying the location of GABAergic PSDs in relation to presynaptic terminals. The model is compatible with the proposed seeding role of CB/NL2 interaction for initiating GABA<sub>A</sub>R and gephyrin clustering and for CB2/gephyrin/Cdc-42 interactions for regulating shape and size of the PSD. Further, extending previous concepts, our model introduces the DGC and its interaction with NL2 via S-SCAM as a second molecular complex present in a subset of GABAergic synapses, possibly under the influence of presynaptic factors. This molecular complex might enable gephyrin-independent  $\alpha 1$ -GABA<sub>A</sub>R clustering, possibly via interactions between

NL2 and the  $\alpha 1$  subunit. Moreover, the differential clustering of  $\alpha 1$ - and  $\alpha 2$ -GABA<sub>A</sub>R observed in mutant mice raises the possibility that these two receptor subtypes belong to functionally distinct signaling scaffolds in GABAergic synapses. Although the DGC is present at a subset of GABAergic synapses only, it might fulfill at least three major functions: stabilization and maintenance of GABAergic synapses, anchoring signaling molecules, notably those containing a PDZ domain, such as Syn-ArfGEF or nNOS, in close vicinity to GABAergic synapses, transducing extracellular signals acting on dystroglycan to regulate the function and plasticity of GABAergic transmission.

**Acknowledgments** This work was supported by the Swiss National Science Foundation (grant 31003A\_130495 to JMF, the San Paolo Foundation (grant 2008-2254 to PP), the World Wide Style Program (University of Turin) (PP), and the Forschungskredit of the University of Zurich (SKT). We are grateful to our colleagues Uwe Rudolph (McLean Hospital) and Greg Homanics (University of Pittsburgh) for providing the mutant mice used in our studies and we thank Claire de Groot (University of Zurich) for preparing Fig. 1.

## References

1. Gray EG (1969) Electron microscopy of excitatory and inhibitory synapses: a brief review. *Prog Brain Res* 31:141–155
2. Collins MO, Husi HYL, Brandon JM, Anderson CN, Blackstock WP, Choudhary JS, Grant SG (2006) Molecular characterization and comparison of the components and multiprotein complexes in the postsynaptic proteome. *J Neurochem* 97(Suppl 1):16–23
3. Sheng M, Sala C (2001) PDZ domains and the organization of supramolecular complexes. *Annu Rev Neurosci* 24:1–29
4. Kennedy MB (1998) Signal transduction molecules at the glutamatergic postsynaptic membrane. *Brain Res Rev* 26:243–257
5. Tsunoda S, Zuker CS (1999) The organization of INAD-signaling complexes by a multivalent PDZ domain protein in *Drosophila* photoreceptor cells ensures sensitivity and speed of signaling. *Cell Calcium* 26:165–171
6. Oliva C, Escobedo P, Astorga C, Molina C, Sierralta J (2012) Role of the MAGUK protein family in synapse formation and function. *Dev Neurobiol* 72:57–72
7. Gardoni F, Marcello E, Di Luca M (2009) Postsynaptic density-membrane associated guanylate kinase proteins (PSD-MAGUKs) and their role in CNS disorders. *Neuroscience* 158:324–333
8. Ho GP, Selvakumar B, Mukai J, Hester LD, Wang Y, Gogos JA, Snyder SH (2011) S-nitrosylation and S-palmitoylation reciprocally regulate synaptic targeting of PSD-95. *Neuron* 71:131–141
9. Sturgill JF, Steiner P, Czervionke BL, Sabatini BL (2009) Distinct domains within PSD-95 mediate synaptic incorporation, stabilization, and activity-dependent trafficking. *J Neurosci* 29:12845–12854
10. Südhof TC (2008) Neuroligins and neuexins link synaptic function to cognitive disease. *Nature* 455:903–911
11. Siddiqui TJ, Craig AM (2011) Synaptic organizing complexes. *Curr Opin Neurobiol* 21:132–143
12. Lin YC, Koleske AJ (2010) Mechanisms of synapse and dendrite maintenance and their disruption in psychiatric and neurodegenerative disorders. *Annu Rev Neurosci* 33:349–378
13. Fritschy JM, Harvey RJ, Schwarz G (2008) Gephyrin, where do we stand, where do we go? *Trends Neurosci* 31:257–264

14. Papadopoulos T, Soykan T (2011) The role of collybistin in gephyrin clustering at inhibitory synapses: facts and open questions. *Front Cell Neurosci* 5:11
15. Sassoè-Pognetto M, Frola E, Pregno G, Briatore F, Patrizi A (2011) Understanding the molecular diversity of GABAergic synapses. *Front Cell Neurosci* 5:4
16. Arancibia-Cárcamo IL, Kittler JT (2009) Regulation of GABA<sub>A</sub> receptor membrane trafficking and synaptic localization. *Pharmacol Ther* 123:17–31
17. Kim EY, Schrader N, Smolinsky B, Bedet C, Vannier C, Schwarz G, Schindelin H (2006) Deciphering the structural framework of glycine receptor anchoring by gephyrin. *EMBO J* 25:1385–1395
18. Sola M, Bavro VN, Timmins J, Franz T, Ricard-Blum S, Schoehn G, Ruijgrok RW, Paarmann I, Saiyed T, O'Sullivan GA, Schmitt B, Betz H, Weissenhorn W (2004) Structural basis of dynamic glycine receptor clustering by gephyrin. *EMBO J* 23:2510–2519
19. Tretter V, Kerschner B, Milenkovic I, Ramsden SL, Ramerstorfer J, Saiepour L, Maric HM, Moss SJ, Schindelin H, Harvey RJ, Sieghart W, Harvey K (2011) Molecular basis of the GABA<sub>A</sub> receptor  $\alpha 3$  subunit interaction with gephyrin. *J Biol Chem* [Epub ahead of print]
20. Tretter V, Jacob TC, Mukherjee J, Fritschy JM, Pangalos MN, Moss SJ (2008) The clustering of GABA<sub>A</sub> receptor subtypes at inhibitory synapses is facilitated via the direct binding of receptor  $\alpha 2$  subunits to gephyrin. *J Neurosci* 28:1356–1365
21. Mukherjee J, Kretschmannova K, Gouzer G, Maric HM, Ramsden SL, Tretter V, Harvey K, Davies PA, Triller A, Schindelin H, Moss SJ (2011) The residence time of GABA<sub>A</sub>Rs at inhibitory synapses is determined by direct binding of the receptor  $\alpha 1$  subunit to gephyrin. *J Neurosci* 31:14677–14687
22. Schwarz G, Mendel RR, Ribbe MW (2009) Molybdenum cofactors, enzymes and pathways. *Nature* 460:839–847
23. Smolinsky B, Eichler SA, Buchmeier S, Meier JC, Schwarz G (2008) Splice-specific functions of gephyrin in molybdenum cofactor biosynthesis. *J Biol Chem* 283:17370–17379
24. Saiyed T, Paarmann I, Schmitt B, Haeger S, Sola M, Schmalzing G, Weissenhorn W, Betz H (2007) Molecular basis of gephyrin clustering at inhibitory synapses: role of G- and E-domain interactions. *J Biol Chem* 282:5625–5632
25. Calamai M, Specht CG, Heller J, Alcor D, Machado P, Vannier C, Triller A (2009) Gephyrin oligomerization controls GlyR mobility and synaptic clustering. *J Neurosci* 29:7639–7648
26. Bedet C, Bruusgaard JC, Vergo S, Groth-Pedersen L, Eimer S, Triller A, Vannier C (2006) Regulation of gephyrin assembly and glycine receptor synaptic stability. *J Biol Chem* 281:30046–30056
27. Dobie FA, Craig AM (2011) Inhibitory synapse dynamics: coordinated presynaptic and postsynaptic mobility and the major contribution of recycled vesicles to new synapse formation. *J Neurosci* 31:10481–10493
28. Meier J, Grantyn R (2004) A gephyrin-related mechanism restraining glycine receptor anchoring at GABAergic synapses. *J Neurosci* 24:1398–1405
29. Langosch D, Hoch W, Betz H (1992) The 93 kDa protein gephyrin and tubulin associated with the inhibitory glycine receptor are phosphorylated by an endogenous protein kinase. *FEBS Lett* 298:113–117
30. Zita MM, Marchionni I, Bottos E, Righi M, Del Sal G, Cherubini E, Zacchi P (2007) Post-phosphorylation prolyl isomerisation of gephyrin represents a mechanism to modulate glycine receptors function. *EMBO J* 26:1761–1771
31. Tyagarajan SK, Ghosh H, Yevenes GE, Nikonenko I, Ebeling C, Schwerdel C, Sidler C, Zeilhofer HU, Gerrits B, Muller D, Fritschy JM (2011) Regulation of GABAergic synapse formation and plasticity by GSK3 $\beta$ -dependent phosphorylation of gephyrin. *Proc Natl Acad Sci USA* 108:379–384
32. Varoqueaux F, Jamain S, Brose N (2004) Neuroligin 2 is exclusively localized to inhibitory synapses. *Eur J Cell Biol* 83:449–456
33. Song JY, Ichtchenko K, Sudhof TC, Brose N (1999) Neuroligin 1 is a postsynaptic cell-adhesion molecule of excitatory synapses. *Proc Natl Acad Sci USA* 96:1100–1105
34. Hoon M, Bauer G, Fritschy JM, Moser T, Falkenburger BH, Varoqueaux F (2009) Neuroligin 2 controls the maturation of GABAergic synapses and information processing in the retina. *J Neurosci* 29:8039–8050
35. Pouloupoulos A, Aramuni G, Meyer G, Soykan T, Hoon M, Papadopoulos T, Zhang M, Paarmann I, Fuchs C, Harvey K, Jedlicka P, Schwarzacher SW, Betz H, Harvey RJ, Brose N, Zhang W, Varoqueaux F (2009) Neuroligin 2 drives postsynaptic assembly at perisomatic inhibitory synapses through gephyrin and collybistin. *Neuron* 63:628–642
36. Budreck EC, Scheiffele P (2007) Neuroligin-3 is a neuronal adhesion protein at GABAergic and glutamatergic synapses. *Eur J Neurosci* 26:1738–1748
37. Hoon M, Soykan T, Falkenburger BH, Hammer M, Patrizi A, Schmidt KF, Sassoè-Pognetto M, Löwel S, Moser T, Taschenberger H, Brose N, Varoqueaux F (2011) Neuroligin-4 is localized to glycinergic postsynapses and regulates inhibition in the retina. *Proc Natl Acad Sci USA* 108:3053–3058
38. Lise MF, El-Husseini A (2006) The neuroligin and neuroligin families: from structure to function at the synapse. *Cell Mol Life Sci* 63:1833–1849
39. Varoqueaux F, Aramuni G, Rawson RL, Mohrmann R, Missler M, Gottmann K, Zhang W, Südhof TC, Brose N (2006) Neuroligins determine synapse maturation and function. *Neuron* 51:741–754
40. Harvey K, Duguid IC, Alldred MJ, Beatty SE, Ward H, Keep NH, Lingenfelter SE, Pearce BR, Lundgren J, Owen MJ, Smart TG, Luscher B, Rees MI, Harvey RJ (2004) The GDP–GTP exchange factor collybistin: an essential determinant of neuronal gephyrin clustering. *J Neurosci* 24:5816–5826
41. Patrizi A, Viltono L, Frola E, Harvey K, Harvey RJ, Sassoè-Pognetto M (2011) Selective localization of collybistin at a subset of inhibitory synapses in brain circuits. *J Comp Neurol* 520:130–141
42. Kalscheuer VM, Musante L, Fang C, Hoffmann K, Fuchs C, Carta E, Deas E, Venkateswarlu K, Menzel C, Ullmann R, Tommerup N, Dalpra L, Tzschach A, Selicorni A, Luscher B, Ropers HH, Harvey K, Harvey RJ (2009) A balanced chromosomal translocation disrupting ARHGAP9 is associated with epilepsy, anxiety, aggression, and mental retardation. *Hum Mutat* 30:61–68
43. Shimojima K, Sugawara M, Shichiji M, Mukaida S, Takayama R, Imai K, Yamamoto T (2011) Loss-of-function mutation of collybistin is responsible for X-linked mental retardation associated with epilepsy. *J Hum Genet* 56:561–565
44. Kins S, Betz H, Kirsch J (2000) Collybistin, a newly identified brain-specific GEF, induces submembrane clustering of gephyrin. *Nat Neurosci* 3:22–29
45. Xiang S, Young Kim E, Connelly JJ, Nassar N, Kirsch J, Winking J, Schwarz G, Schindelin H (2006) The crystal structure of Cdc42 in complex with collybistin II, a gephyrin-interacting guanine nucleotide exchange factor. *J Mol Biol* 359:35–46
46. Tyagarajan SK, Ghosh H, Harvey K, Fritschy JM (2011) Collybistin splice variants differentially interact with gephyrin and Cdc42 to regulate gephyrin clustering at GABAergic synapses. *J Cell Sci* 124:2786–2796
47. Reddy-Alla S, Schmitt B, Birkenfeld J, Eulenburg V, Dutertre S, Böhringer C, Götz M, Betz H, Papadopoulos T (2010) PH-

- Domain-driven targeting of collybistin but not Cdc42 activation is required for synaptic gephyrin clustering. *Eur J Neurosci* 31:1173–1184
48. Saiepour L, Fuchs C, Patrizi A, Sassoè-Pognetto M, Harvey RJ, Harvey K (2010) Complex role of collybistin and gephyrin in GABA<sub>A</sub> receptor clustering. *J Biol Chem* 285:29623–29631
  49. Fritschy JM, Brünig I (2003) Formation and plasticity of GABAergic synapses: physiological mechanisms and pathophysiological implications. *Pharmacol Ther* 98:299–323
  50. Chiou TT, Bonhomme B, Jin H, Miralles CP, Xiao H, Fu Z, Harvey RJ, Harvey K, Vicini S, De Blas AL (2011) Differential regulation of the postsynaptic clustering of  $\gamma$ -aminobutyric acid type A (GABA<sub>A</sub>) receptors by collybistin isoforms. *J Biol Chem* 286:22456–22468
  51. Maric HM, Mukherjee J, Tretter V, Moss SJ, Schindelin H (2011) Gephyrin-mediated GABA<sub>A</sub> and glycine receptor clustering relies on a common binding site. *J Biol Chem* 286:42105–42114
  52. Kirsch J, Betz H (1998) Glycine-receptor activation is required for receptor clustering in spinal neurons. *Nature* 392:717–720
  53. Hanus C, Vannier C, Triller A (2004) Intracellular association of glycine receptor with gephyrin increases its plasma membrane accumulation rate. *J Neurosci* 24:1119–1128
  54. Papadopoulos T, Korte M, Eulenburg V, Kubota H, Retiounskaia M, Harvey RJ, Harvey K, O'sullivan GA, Laube B, Hulsmann S, Geiger JR, Betz H (2007) Impaired GABAergic transmission and altered hippocampal synaptic plasticity in collybistin-deficient mice. *EMBO J* 26:3888–3899
  55. Gomeza J, Hülsmann S, Ohno K, Eulenburg V, Szöke K, Richter DW, Betz H (2003) Inactivation of the glycine transporter 1 gene discloses vital role of glial glycine uptake in glycinergic inhibition. *Neuron* 40:785–796
  56. Kralic JE, Sidler C, Parpan F, Homanics G, Morrow AL, Fritschy JM (2006) Compensatory alteration of inhibitory synaptic circuits in thalamus and cerebellum of GABA<sub>A</sub> receptor  $\alpha$ 1 subunit knockout mice. *J Comp Neurol* 495:408–421
  57. Schneider Gasser EM, Duveau V, Prenosil GA, Fritschy JM (2007) Reorganization of GABAergic circuits maintains GABA<sub>A</sub> receptor-mediated transmission onto CA1 interneurons in  $\alpha$ 1-subunit-null mice. *Eur J Neurosci* 25:3287–3304
  58. Bannai H, Lévi S, Schweizer C, Inoue T, Launey T, Racine V, Sibarita JB, Mikoshiba K, Triller A (2009) Activity-dependent tuning of inhibitory neurotransmission based on GABA<sub>A</sub>R diffusion dynamics. *Neuron* 62:670–682
  59. Specht CG, Grünwald N, Pascual O, Rostgaard N, Schwarz G, Triller A (2011) Regulation of glycine receptor diffusion properties and gephyrin interactions by protein kinase C. *EMBO J* 30:3842–3853
  60. Lévi S, Schweizer C, Bannai H, Pascual O, Charrier C, Triller A (2008) Homeostatic regulation of synaptic GlyR numbers driven by lateral diffusion. *Neuron* 59:261–273
  61. Machado P, Rostaing P, Guigonis JM, Renner M, Dumoulin A, Samson M, Vannier C, Triller A (2011) Heat shock cognate protein 70 regulates gephyrin clustering. *J Neurosci* 31:3–14
  62. Triller A, Cluzeaud F, Pfeiffer F, Betz H, Korn H (1985) Distribution of glycine receptors at central synapses: an immunoelectron microscopy study. *J Cell Biol* 101:683–688
  63. Geiman EJ, Knox MC, Alvarez FJ (2000) Postnatal maturation of gephyrin/glycine receptor clusters on developing Renshaw cells. *J Comp Neurol* 426:130–142
  64. Sassoè-Pognetto M, Kirsch J, Grünert U, Greferath U, Fritschy JM, Mohler H, Betz H, Wässle H (1995) Colocalization of gephyrin and GABA<sub>A</sub>-receptor subunits in the rat retina. *J Comp Neurol* 357:1–14
  65. Sassoè-Pognetto M, Panzanelli P, Sieghart W, Fritschy JM (2000) Co-localization of multiple GABA<sub>A</sub> receptor subtypes with gephyrin at postsynaptic sites. *J Comp Neurol* 420:481–498
  66. Gunther U, Benson J, Benke D, Fritschy JM, Reyes GH, Knoflach F, Crestani F, Aguzzi A, Arigoni M, Lang Y, Bluethmann H, Mohler H, Luscher B (1995) Benzodiazepine-insensitive mice generated by targeted disruption of the  $\gamma$ 2-subunit gene of  $\gamma$ -aminobutyric acid type A receptors. *Proc Natl Acad Sci USA* 92:7749–7753
  67. Essrich C, Lorez M, Benson JA, Fritschy JM, Luscher B (1998) Postsynaptic clustering of major GABA<sub>A</sub> receptor subtypes requires the  $\gamma$ 2 subunit and gephyrin. *Nat Neurosci* 1:563–571
  68. Schweizer C, Balsiger S, Bluethmann H, Mansuy IM, Fritschy JM, Mohler H, Luscher B (2003) The  $\gamma$ 2 subunit of GABA<sub>A</sub> receptors is required for maintenance of receptors at mature synapses. *Mol Cell Neurosci* 24:442–450
  69. Kneussel M, Brandstätter JH, Laube B, Stahl S, Müller U, Betz H (1999) Loss of postsynaptic GABA<sub>A</sub> receptor clustering in gephyrin-deficient mice. *J Neurosci* 19:9289–9297
  70. Levi S, Logan SM, Tovar KR, Craig AM (2004) Gephyrin is critical for glycine receptor clustering but not for the formation of functional GABAergic synapses in hippocampal neurons. *J Neurosci* 24:207–217
  71. Kneussel M, Brandstätter JH, Gasnier B, Feng G, Sanes JR, Betz H (2001) Gephyrin-independent clustering of postsynaptic GABA<sub>A</sub> receptor subtypes. *Mol Cell Neurosci* 17:973–982
  72. Studer R, von Boehmer L, Haenggi T, Schweizer C, Benke D, Rudolph U, Fritschy JM (2006) Alteration of GABAergic synapses and gephyrin clusters in the thalamic reticular nucleus of GABA<sub>A</sub> receptor  $\alpha$ 3 subunit-null mice. *Eur J Neurosci* 24:1307–1315
  73. Fritschy JM, Panzanelli P (2006) Molecular and synaptic organization of GABA<sub>A</sub> receptors in the cerebellum: Effects of targeted subunit gene deletions. *Cerebellum* 5:275–285
  74. Panzanelli P, Gunn BG, Schlatter MC, Benke D, Tyagarajan SK, Scheiffele P, Belelli D, Lambert JJ, Rudolph U, Fritschy JM (2011) Distinct mechanisms regulate GABA<sub>A</sub> receptor and gephyrin clustering at perisomatic and axo-axonic synapses on CA1 pyramidal cells. *J Physiol* 589:4959–4980
  75. Peden DR, Petitjean CM, Herd MB, Durakoglugil M, Rosahl JW, Wafford K, Homanics GE, Belelli D, Fritschy JM, Lambert JJ (2008) Developmental maturation of synaptic and extrasynaptic GABA<sub>A</sub> receptors in mouse thalamic ventrobasal neurones. *J Physiol* 586:965–987
  76. Fritschy JM, Panzanelli P, Kralic JE, Vogt KE, Sassoè-Pognetto M (2006) Differential dependence of axo-dendritic and axosomatic GABAergic synapses on GABA<sub>A</sub> receptors containing the  $\alpha$ 1 subunit in Purkinje cells. *J Neurosci* 26:3245–3255
  77. Papadopoulos T, Eulenburg V, Reddy-Alla S, Mansuy IM, Li Y, Betz H (2008) Collybistin is required for both the formation and maintenance of GABAergic postsynapses in the hippocampus. *Mol Cell Neurosci* 39:161–169
  78. Gibson JR, Huber KM, Südhof TC (2009) Neuroigin-2 deletion selectively decreases inhibitory synaptic transmission originating from fast-spiking but not from somatostatin-positive interneurons. *J Neurosci* 29:13883–13897
  79. Jedlicka P, Hoon M, Papadopoulos T, Vlachos A, Winkels R, Pouloupoulos A, Betz H, Deller T, Brose N, Varoqueaux F, Schwarzacher SW (2011) Increased dentate gyrus excitability in neuroigin-2-deficient mice in vivo. *Cereb Cortex* 21:357–367
  80. Prenosil GA, Schneider Gasser EM, Rudolph U, Keist R, Fritschy JM, Vogt KE (2006) Specific subtypes of GABA<sub>A</sub> receptors mediate phasic and tonic forms of inhibition in hippocampal pyramidal neurons. *J Neurophysiol* 96:846–857
  81. Kasugai Y, Swinny JD, Roberts JD, Dalezios Y, Fukazawa Y, Sieghart W, Shigemoto R, Somogyi P (2010) Quantitative localisation of synaptic and extrasynaptic GABA<sub>A</sub> receptor subunits on hippocampal pyramidal cells by freeze-fracture replica immunolabelling. *Eur J Neurosci* 32:1868–1888

82. Haenggi T, Fritschy JM (2006) Role of dystrophin and utrophin for assembly and function of the dystrophin glycoprotein complex in non-muscle tissue. *Cell Mol Life Sci* 63:1614–1631
83. Perronnet C, Vaillend C (2010) Dystrophins, utrophins, and associated scaffolding complexes: role in mammalian brain and implications for therapeutic strategies. *J Biomed Biotechnol* 2010:849426
84. Barresi R, Campbell KP (2006) Dystroglycan: from biosynthesis to pathogenesis of human disease. *J Cell Sci* 119:199–207
85. Waite A, Tinsley CL, Locke M, Blake DJ (2009) The neurobiology of the dystrophin-associated glycoprotein complex. *Ann Med* 41:344–359
86. Lidov HGW (1996) Dystrophin in the nervous system. *Brain Pathol* 6:63–77
87. Knuesel I, Mastrolola M, Zuellig RA, Bornhauser B, Schaub MC, Fritschy JM (1999) Altered synaptic clustering of GABA<sub>A</sub>-receptors in mice lacking dystrophin (mdx mice). *Eur J Neurosci* 11:4457–4462
88. Knuesel I, Zuellig RA, Schaub MC, Fritschy JM (2001) Alterations in dystrophin and utrophin expression parallel the reorganization of GABAergic synapses in a mouse model of temporal lobe epilepsy. *Eur J Neurosci* 13:1113–1124
89. Dallérac G, Perronnet C, Chagneau C, Leblanc-Veyrac P, Samson-Desvignes N, Peltekian E, Danos O, Garcia L, Laroche S, Billard J, Vaillend C (2011) Rescue of a dystrophin-like protein by exon skipping normalizes synaptic plasticity in the hippocampus of the mdx mouse. *Neurobiol Dis* 43:635–641
90. Vaillend C, Perronnet C, Ros C, Gruszczynski C, Goyenvallé A, Laroche S, Danos O, Garcia L, Peltekian E (2010) Rescue of a dystrophin-like protein by exon skipping in vivo restores GABA<sub>A</sub>-receptor clustering in the hippocampus of the mdx mouse. *Mol Ther* 18:1683–1688
91. Pilgram GS, Potikanond S, Baines RA, Fradkin LG, Noordermeer JN (2010) The roles of the dystrophin-associated glycoprotein complex at the synapse. *Mol Neurobiol* 41:1–21
92. Bragg AD, Das SS, Froehner SC (2010) Dystrophin-associated protein scaffolding in brain requires  $\alpha$ -dystrobrevin. *Neuroreport* 21:695–699
93. Adams ME, Mueller HA, Froehner SC (2001) In vivo requirement of the  $\alpha$ -syntrophin PDZ domain for the sarcolemmal localization of nNOS and aquaporin-4. *J Cell Biol* 155:113–122
94. Haenggi T, Soontornmalai A, Schaub MC, Fritschy JM (2004) The role of utrophin and Dp71 for assembly of different dystrophin-associated protein complexes (DPCs) in the choroid plexus and microvasculature of the brain. *Neuroscience* 129:403–413
95. Yamakawa H, Oyama S, Mitsuhashi H, Sasagawa N, Uchino S, Kohsaka S, Ishiura S (2007) Neuroligins 3 and 4X interact with syntrophin-gamma2, and the interactions are affected by autism-related mutations. *Biochem Biophys Res Com* 355:41–46
96. Michele DE, Barresi V, Kanagawa M, Saito F, Cohn RD, Satz JS, Dollar J, Nishino I, Kelley RI, Somer H, Straub V, Mathews KD, Moore SA, Campbell KP (2002) Post-translational disruption of dystroglycan-ligand interactions in congenital muscular dystrophies. *Nature* 418:417–422
97. Satz JS, Ostendorf AP, Hou S, Turner A, Kusano H, Lee JC, Turk R, Nguyen H, Ross-Barta SE, Westra S, Hoshi T, Moore SA, Campbell KP (2010) Distinct functions of glial and neuronal dystroglycan in the developing and adult mouse brain. *J Neurosci* 30:14560–14572
98. Grady RM, Wozniak DF, Ohlemiller KK, Sanes JR (2006) Cerebellar synaptic defects and abnormal motor behavior in mice lacking  $\alpha$ - and  $\beta$ -dystrobrevin. *J Neurosci* 26:2841–2851
99. Knuesel I, Züllig RA, Bornhauser B, Schaub MC, Fritschy JM (2000) Differential expression of utrophin and dystrophin in CNS neurons: an in situ hybridization and immunohistochemical study. *J Comp Neurol* 422:594–611
100. Brünig I, Suter A, Knuesel I, Luscher B, Fritschy JM (2002) GABAergic presynaptic terminals are required for postsynaptic clustering of dystrophin, but not of GABA<sub>A</sub> receptors and gephyrin. *J Neurosci* 22:4805–4813
101. Sugita S, Saito F, Tang J, Satz JS, Campbell KP, Südhof TC (2001) A stoichiometric complex of neurexins and dystroglycan in brain. *J Cell Biol* 154:435–445
102. Sumita K, Sato Y, Iida J, Kawata A, Hamano M, Hirabayashi S, Ohno K, Peles E, Hata Y (2007) Synaptic scaffolding molecule (S-SCAM) membrane-associated guanylate kinase with inverted organization (MAGI)-2 is associated with cell adhesion molecules at inhibitory synapses in rat hippocampal neurons. *J Neurochem* 100:154–166
103. Fukaya M, Kamata A, Hara Y, Tamaki H, Katsumata O, Ito N, Takeda S, Hata Y, Suzuki T, Watanabe M, Harvey RJ, Sakagami H (2011) SynArfGEF is a guanine nucleotide exchange factor for Arf6 and localizes preferentially at post-synaptic specializations of inhibitory synapses. *J Neurochem* 116:1122–1137
104. Levi S, Grady M, Henry M, Campbell K, Sanes J, Craig A (2002) Dystroglycan is selectively associated with inhibitory GABAergic synapses but is dispensable for their differentiation. *J Neurosci* 22:4274–4285
105. Patrizi A, Scelfo B, Viltono L, Briatore F, Fukaya M, Watanabe M, Strata P, Varoqueaux F, Brose N, Fritschy JM, Sassoè-Pognetto M (2008) Synapse formation and clustering of neuroligin-2 in the absence of GABA<sub>A</sub> receptors. *Proc Natl Acad Sci USA* 105:13151–13156
106. Kins S, Kuhse J, Laube B, Betz H, Kirsch J (1999) Incorporation of a gephyrin-binding motif targets NMDA receptors to gephyrin-rich domains in HEK 293. *Eur J Neurosci* 11:740–744
107. Kirsch J, Wolters I, Triller A, Betz H (1993) Gephyrin antisense oligonucleotides prevent glycine receptor clustering in spinal neurons. *Nature* 366:745–748