

Hydrobiologia (2009) 635:181–187
DOI 10.1007/s10750-009-9910-x

PRIMARY RESEARCH PAPER

Eutrophication and endangered aquatic plants: an experimental study on *Baldellia ranunculoides* (L.) Parl. (Alismataceae)

Gregor Kozłowski · Sophie Vallelian

Received: 12 December 2008 / Revised: 10 July 2009 / Accepted: 21 July 2009 / Published online: 5 August 2009
© Springer Science+Business Media B.V. 2009

Abstract Anthropogenic enrichment of aquatic ecosystems is an increasingly common phenomenon, resulting from human population growth and the intensification of industrial and agricultural developments. Eutrophication has been considered to be one of the main causes of the decline of many aquatic plants. These observations, however, have rarely been supported by data or explored in detail using an experimental approach. Our experimental study demonstrates, for the first time, a strong direct negative influence of eutrophication on the performance of the endangered aquatic plant *Baldellia ranunculoides* s.str. (L.) Parl. (Alismataceae). Both morphological and reproductive traits were significantly affected by this phenomenon. Plants growing on eutrophic substrates were much smaller, possessed fewer and smaller leaves, and their biomass was on average half that of plants growing on a mesotrophic substrate. Additionally, plants growing on eutrophic substrates produced fewer inflorescences and flowers, and they

had a smaller number of achenes per capitulum. Consequently, they produced only 15% of the seeds produced by plants growing in non-eutrophic conditions. Thus, the chance of long-term subsistence of *B. ranunculoides* populations in strongly eutrophic habitats will be significantly reduced, and at the same time, the success of any new establishment of viable *B. ranunculoides* populations in such habitats will be highly restricted. Conservation efforts for *B. ranunculoides*, either of existing populations or in newly selected re-introduction sites, should start, therefore, with an assessment of the trophic level of those sites. On the other hand, due to its sensitivity to eutrophication, the presence of viable populations of *B. ranunculoides* can be used as an indicator of valuable aquatic habitats for oligotrophy-dependent organisms.

Keywords Aquatic macrophytes · Endangered species · Global change · Life traits · Nitrogen deposition

Handling editor: S. M. Thomaz

G. Kozłowski (✉)
Department of Biology and Botanical Garden, University
of Fribourg, Chemin du Musée 10, 1700 Fribourg,
Switzerland
e-mail: gregor.kozlowski@unifr.ch

S. Vallelian
Department of Geosciences, University of Fribourg,
Chemin du Musée 4, 1700 Fribourg, Switzerland

Introduction

In recent decades, the human-generated modification of natural habitats has reached an unprecedented level (Jenkins, 2003; Kareiva et al., 2007). In the future, the appropriation of natural resources for human needs will expose the environment to even

more drastic changes (Sala et al., 2000; Foley et al., 2005). Aquatic organisms and their habitats are particularly affected by human-caused global change and its various facets (Carpenter et al., 1992; Sala et al., 2000; Cronk & Fennessy, 2001; Malmquist & Rundle, 2002; Egertson et al., 2004). In many regions, aquatic plants are among the most threatened groups of organisms (Preston & Croft, 2001; Moser et al., 2002). The genus *Baldellia* is not an exception, as its taxa are threatened in the majority of European countries and have strongly declined throughout their area of distribution due to habitat destruction, fragmentation and other anthropogenic changes (Roelofs, 1983; Preston & Croft, 2001; Jones, 2006; Kozłowski et al., 2008).

Baldellia species are morphologically very plastic and respond to changes in environmental conditions, and especially to nutritional and hydrological change, with adjustments to their growth and physiology (Kozłowski et al., 2008). This phenotypic plasticity is typical of many aquatic plants (Sculthorpe, 1967). *Baldellia* is one of several small genera of the exclusively aquatic plant family Alismataceae (Cook, 1990). *Baldellia* taxa are perennial, rooted water plants that grow in lakes, ponds and slow streams (Vuille, 1988; Preston & Croft, 2001). There are only two generally recognized species in the genus: *Baldellia ranunculoides* (L.) Parl. [with two subspecies: *B. ranunculoides* subsp. *ranunculoides* and subsp. *repens* (Lam.) Á. et D. Löve], which is native to Europe and the Mediterranean, and *Baldellia alpestris* (Coss.) Carv. Vasc., which is endemic to the mountains of northern Portugal and northwest Spain (Cook, 1983; Kozłowski & Matthies, 2009). The present study deals exclusively with *B. ranunculoides* s.str., an emblematic taxon in the majority of European countries and a candidate for conservation priority lists.

Baldellia ranunculoides shows a strong decline in practically all regions of its natural range and it is probably the most threatened taxon of the genus (Kozłowski et al., 2009). The last region to contain numerous populations and high abundance of this taxon is Ireland. In France, Portugal, Spain and the UK, the taxon is not threatened according to the published national red lists. It was clearly demonstrated, however, that the conservation status is underestimated in these countries. In Germany, Norway and Benelux, *B. ranunculoides* is endangered (EN). In Switzerland,

Italy and Croatia the taxon is critically endangered (CR). In certain regions of Europe the species is regionally extinct (RE, e.g. in Poland and Sicily) (IUCN, 2001; Kozłowski et al., 2009).

It is a taxon growing mainly in coastal regions of Western Europe and the Mediterranean (Meusel et al., 1992; de Bolos & Vigo, 2003; Kozłowski et al., 2008). In Central and Western Europe, *B. ranunculoides* s.str. is a characteristic species of the alliance *Hydrocotylo-Baldellion* Tx. et Dierssen apud Dierssen 1957, in the class *Littorelletea uniflorae* Br.-Bl. et Tx. 1943 ex Westhoff, Dijk et Paschier 1946 (Pott, 1995; Schubert et al., 1995; Kozłowski & Matthies, 2009). The class *L. uniflorae* includes plant communities on sites with alternating water level in meso- to oligotrophic, stagnant or slowly running water. The communities are bound to a mineral soil, which is usually not covered by organic material, and their species number is rather low (Schaminée & Arts, 1992).

Eutrophication has been generally considered to be one of the main causes of decline of many aquatic communities containing *Baldellia* sp. (Schoof-van Pelt, 1973; Roelofs et al., 1984; Arts, 2002; Kennedy & Murphy, 2004). Organic enrichment causes important changes and degradation of EN aquatic plant communities (Bobbink et al., 1998; Egertson et al., 2004). However, these observations and conclusions have rarely been supported with data or explored in detail with an experimental approach.

The main aims of our study were thus to (1) estimate the sensibility of *B. ranunculoides* s.str. to eutrophication, (2) define which morphological and/or reproductive traits are the most affected by the eutrophication and (3) provide a basis for more effective conservation measures and restoration efforts.

Materials and methods

Plant material

Seeds of *B. ranunculoides* s.str. were obtained from (1) Botanical Garden Fribourg, Switzerland; (2) Botanical Garden Berlin-Dahlem; (3) Botanical Garden Nantes, France and (4) Botanical Garden Coimbra, Portugal. The seeds were collected from natural populations from Canton Vaud (Switzerland), Nordrhein-Westphalia (Germany), Loire Atlantique (France) and Beira

Litoral (Portugal). These four regions cover the geographical range of the taxon in Western and Central Europe. For the experimental assessment, seeds of all four origins were pooled together to exclude any effect of a local adaptation.

Substrate preparation

Three types of substrates were used for the experimental assessment: (1) control substrate—representing a mesotrophic substrate composition of *B. ranunculoides* s.str. based on our successful *ex situ* cultures at the Botanical Garden Fribourg (unpublished data) and our previous work (Kozłowski & Matthies, 2009). It was obtained by mixing the four parts of topsoil with one part of coarse sand (0.5–1.0 mm of diameter). The final mixture contained 46% of dry matter, 17% of organic matter, with conductivity of 123 $\mu\text{S}/\text{cm}$ and pH 7.0; (2) Treatment 1 (T1)—addition of 5 g of Hato[®] organic NPK fertilizer (Hauert HBG Dünger AG, Grossaffoltern, Switzerland) per 1 kg of the control substrate; Treatment 2 (T2)—addition of 10 g of Hato[®] per 1 kg of the control substrate. According to the manufacturer's indication, 5 g of Hato[®] fertilizer contains 400 mg of total N, 200 mg of P_2O_5 , 400 mg of K_2O , 75 mg of Mg^{2+} and ca. 13 mg of Bo, Cu, Fe, Mn, Mo and Zn (taken together).

Experimental design

For germination tests, we sowed 25 *Baldellia* seeds into $20 \times 20 \times 15$ cm pots filled with substrate. Pots were placed in containers filled with water and, thus, maintained all the time under semi-aquatic conditions (substrate stayed constantly wet). Twenty pots were sowed for each treatment (Control, T1 and T2), giving a total of 500 seeds per treatment. After germination, 100 seedlings from each of the three different treatments were chosen randomly and grown until reaching maturity for 4 months under the same conditions in a greenhouse. After that, the following morphological and reproductive traits were assessed (see Fig. 1): the height of the plant, the number of leaves per plant, the length and width of all leaves, the number of pseudowhorls per plant, the number of flowers per pseudowhorl, and the achene number per capitulum. At the end of the 4 months, all plants were harvested (whole plant with roots), then oven-dried at 60°C for 48 h and weighed. The treatments were compared by




Fig. 1 Schematic representation of the morphological traits of *B. ranunculoides* s.str. measured during the eutrophication experiments

Kruskal–Wallis one-way analyses of variance, with *P* values calculated by the Monte-Carlo permutation test available in SPSS 12.0 (10 000 runs). Statistically significant differences were defined at the level $P < 0.05$.

Results

Figure 2 shows the effect of eutrophication on the morphology and reproductive traits of *B. ranunculoides*. For the majority of traits investigated, the difference between treatments was highly significant (the only exception was germination ratio, with $P = 0.520$). After 4 months, mature plants growing on eutrophied substrates (T2 treatment) were much smaller (mean: 2.8 cm vs. 9.7 cm for control plants), possessed fewer leaves per plant (mean: 9.6 vs. 21.0), and the leaves were narrower and shorter (means: 6.5×0.4 cm vs. 9.8×0.6 cm for control plants) (see Fig. 2a, d, e, f). Consequently, the biomass of plants


growing on eutrophic substrates was, on average, almost half that of plants grown on the control substrate (mean: 3.0 g for plants with T2 treatment vs. 5.5 g for plants on control substrate, Fig. 2c). Additionally, plants growing on T2 eutrophied substrates produced fewer pseudowhorls in their inflorescence (mean: 1.2 vs. 2.3 for control plants), which additionally developed fewer flowers (only three flowers per pseudowhorl vs. seven flowers in control plants) (Fig. 2g, h). They also produced, although less significantly ($P = 0.025$), a smaller number of achenes per capitulum (Fig. 2i). The

moderate eutrophication treatment (T1) had an intermediate effect on the growth and reproduction of *B. ranunculoides* (except for the leaf dimensions; Fig. 2e, f).

Discussion

Anthropogenic enrichment of aquatic ecosystems is an increasingly common phenomenon, resulting from human population growth and the intensification of

Fig. 2 Variation of *B. ranunculoides* s.str. morphological and reproductive traits under different eutrophication treatments. *Box plots* indicate the median and the 25th and 75th percentiles of the measured parameter. *Whiskers* indicate the upper and lower inner fence, *circles* are outliers. *P* values for the differences among the treatments are results of Kruskal–Wallis one-way-ANOVAs with Monte-Carlo permutation tests (10,000 runs). Treatments: *C* control mesotrophic conditions; *T1* eutrophication (5 g of fertilizer/kg of control substrate); *T2* strong eutrophication (10 g of fertilizer/kg of control substrate). For more details, see [Materials and methods](#) section



industrial and agricultural developments (Thomas & Daldorph, 1994). The importance of eutrophication in the decline of *Baldellia* species as well as other water plants has been stressed by several previous studies (Roelofs, 1983; Willby & Eaton, 1993; Brouwer & Roelofs, 2001; Kozłowski & Matthies, 2009). However, many of those authors proposed that the eutrophication is rather indirectly responsible for this decline by favoring fast-growing species (Bobbink et al., 1998; Greulich et al., 2000). Our experimental study demonstrates for the first time a strong direct negative influence of eutrophication on the performance of the endangered *B. ranunculoides*. Both morphological and reproductive traits were significantly affected. The only exception to this effects was the germination ratio, indicating that *B. ranunculoides* is able to germinate on a broad spectrum of substrates (Fig. 2a). However, plants germinating in water bodies with high eutrophication will stay smaller, enhancing the probability that they will be out-competed by other co-occurring plants. Additionally, the reproduction of such plants will be strongly reduced. According to our results, an average plant growing on non-eutrophic substrate produces 2.3 pseudowhorls, ca. 16.5 flowers with 32 achenes per flower, and thus will produce ca. 520 seeds. In contrast, a single plant growing on a highly eutrophic substrate produces on average only one pseudowhorl with 3 flowers and 27 achenes per flower, thus producing only ca. 80 seeds (Fig. 2g, h, i). Thus, we conclude that *B. ranunculoides* plants growing in water bodies with high eutrophication will produce ca. 15% as many seeds as plants growing in non-eutrophied conditions. In consequence, the chance of long-term subsistence of *B. ranunculoides* populations in strongly eutrophied habitats will be significantly reduced and any new establishment of viable *B. ranunculoides* populations in such habitats will be highly restricted. The potential mechanisms behind this unsuccessful performance of *B. ranunculoides* in eutrophic habitats remain unknown and need further investigations. It was demonstrated that *B. ranunculoides* reacts negatively to higher ammonium ion (NH_4^+) concentrations (Smolders et al., 2002). The nitrogen and/or ammonium toxicity might, therefore, play an important role in such high sensitivity to eutrophication.

Our results confirm the previous observations that the genus *Baldellia* is physiologically and ecologically

highly specialized for oligotrophic-exposed littoral habitats with low biomass and low concurrence, making it an extremely sensitive taxon to any anthropogenic changes (Preston & Croft, 2001; Kozłowski & Matthies, 2009). Several other taxa of the exclusively aquatic family Alismataceae, e.g. *Luronium natans* (L.) Raf., show the same specialization (Willby & Eaton, 1993; Szankowski & Klosowski, 2001). Our results also confirm our previous observations that *B. ranunculoides* s.str. prefers oligotrophic sites with lower N concentrations compared to two other *Baldellia* taxa (*B. ranunculoides* subsp. *repens* and *B. alpestris*) (Kozłowski & Matthies, 2009).

Baldellia ranunculoides belongs to the so-called Littorelletean and Isoetid species and is often accompanied by other rare and EN plants, e.g. *Littorella uniflora* (L.) Asch., *Anagallis tenella* (L.) L., *Apium inundatum* (L.) Rchb. f., *Isoëtes* spp. (Kozłowski et al., 2008). Thus, the presence of morphologically well-developed plants of *B. ranunculoides* can be used as an indicator of valuable aquatic habitats and sites with high conservation value for other oligotrophy-dependent organisms.

Conclusions

Our study clearly demonstrates that *B. ranunculoides* is adapted to low-fertility sites and is very sensitive to the eutrophication of its habitats. Although able to germinate even on heavily eutrophied substrates, it will produce in such conditions only dwarf forms with drastically reduced generative reproduction and vegetative growth. In natural conditions, such plants are not able to compete with other plants adapted to eutrophic habitats. Furthermore, in strongly eutrophic water bodies, the species will not be able to create new populations or persist for a long time, even if it still possesses shallow inundated and regularly exposed shore fragments with low plant cover. Thus, conservation efforts for *B. ranunculoides*, either of existing populations or in newly selected re-introduction sites, should start with the assessment of their trophic level. In order to maintain *B. ranunculoides* for a long time in a eutrophic water body, artificial disturbances must be performed almost every year—an extremely costly measure. On the other hand, due to its sensitivity to eutrophication, the presence of self-sustaining viable populations of *B. ranunculoides* can be used as an

indicator of the intermediate nutrient concentrations, and thus of valuable aquatic habitats for oligotrophy-dependent organisms.

Acknowledgements We would like to thank Benoît Clement and Susanne Bollinger from the Botanical Garden of the University of Fribourg (Switzerland) as well as Françoise Cudré-Mauroux for their advice and support during the experimental work and manuscript preparation. We are much indebted to the Franklinia Foundation for its engagement and valuable support. We are very grateful to Christian Clerc (GEG Champ-Pittet, Switzerland) and to the Botanical Gardens of Coimbra (Portugal), Berlin-Dahlem (Germany) and Nantes (France) for seed collections.

References

- Arts, G. H. P., 2002. Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalization. *Aquatic Botany* 73: 373–393.
- Bobbink, R., M. Hornung & J. G. M. Roelofs, 1998. The effects of air-borne nitrogen pollutants on species diversity and semi-natural European vegetation. *Journal of Ecology* 86: 717–738.
- Brouwer, E. & J. G. M. Roelofs, 2001. Degraded softwater lakes: possibilities for restoration. *Restoration Ecology* 9: 155–166.
- Carpenter, S. R., S. G. Fisher, N. B. Grimm & J. F. Kitchell, 1992. Global change and freshwater ecosystems. *Annual Review of Ecology and Systematics* 23: 119–139.
- Cook, C. D. K., 1983. Aquatic plants endemic to Europe and the Mediterranean. *Botanische Jahrbücher für Systematik* 103: 539–582.
- Cook, C. D. K., 1990. *Aquatic Plant Book*. SPB Academic Publishing, The Hague.
- Cronk, J. K. & M. S. Fennessy, 2001. *Water Plants: Biology and Ecology*. Lewis Publishers, Boca Raton.
- de Bolos, O. & J. Vigo, 2003. *Flora dels Paisos Catalans*, Vol. IV. Barcino S.A., Barcelona.
- Egerton, C. J., J. A. Kopaska & J. A. Downing, 2004. A century of change in macrophyte abundance and composition in response to agricultural eutrophication. *Hydrobiologia* 524: 145–156.
- Foley, J. A., R. DeFries, G. P. Asner, C. Barford, G. Bonan, S. R. Carpenter, F. S. Chapin, M. T. Coe, G. C. Daily, H. K. Gibbs, J. H. Helkowski, T. Holloway, E. A. Howard, C. J. Kucharik, C. Monfreda, J. A. Patz, I. C. Prentice, N. Ramankutty & P. K. Snyder, 2005. Global consequences of land use. *Science* 309: 570–573.
- Greulich, S., G. C. Bornette, C. Amoros & J. G. M. Roelofs, 2000. Investigation of fundamental niche of rare species: an experiment on establishment of *Luronium natans*. *Aquatic Botany* 66: 209–224.
- IUCN, 2001. IUCN Red List categories and criteria: version 3.1. Species Survival Commission, IUCN, Gland, Switzerland, and Cambridge, UK.
- Jenkins, M., 2003. Prospects for biodiversity. *Science* 302: 1175–1177.
- Jones, R. A., 2006. Creeping water-plantain (*Dyfr lyriad ymlusgawl*), *Baldellia ranunculooides* subsp. *repens* (Lam.) A. Löve & D. Löve in Wales. In Leach, S. J. et al. (eds), *Botanical Links in The Atlantic Arc*. BSBI Conference Report No. 24: 311–319.
- Kareiva, P., S. Watts, R. Mc Donald & T. Boucher, 2007. Domesticated nature: shaping landscape and ecosystems for human welfare. *Science* 316: 1866–1869.
- Kennedy, M. P. & K. J. Murphy, 2004. Indicators of nitrate in wetland surfaces and soil-waters: interactions of vegetation and environmental factors. *Hydrology and Earth System Sciences* 8: 663–672.
- Kozłowski, G. & D. Matthies, 2009. Habitat differentiation in the aquatic plant genus *Baldellia* sp. (Alismataceae): implications for conservation. *Aquatic Botany* 90: 110–118.
- Kozłowski, G., R. A. Jones & F.-L. Vuille-Nicholls, 2008. Biological flora of the Central Europe: *Baldellia ranunculooides* (Alismataceae). *Perspectives in Plant Ecology, Evolution and Systematics* 10: 109–142.
- Kozłowski, G., S. Rion, A. Python & S. Riedo, 2009. Global conservation status assessment of the threatened aquatic plant genus *Baldellia* (Alismataceae): challenges and limitations. *Biodiversity and Conservation* 18: 2307–2325.
- Malmquist, B. & S. Rundle, 2002. Threats to the running water ecosystems of the world. *Environmental Conservation* 29: 134–153.
- Meusel, H., E. Jäger & E. Weinert, 1992. *Vergleichende Chorologie der Zentraleuropäischen Flora*, Vol. 1–3. Gustav Fischer, Jena.
- Moser, D. M., A. Gyğax, B. Bäumler, N. Wyler & R. Palese, 2002. *Liste Rouge des fougères et plantes à fleurs menacées de Suisse*. BUWAL, Bern.
- Pott, R., 1995. *Die Pflanzengesellschaften Deutschlands*, 2nd ed. Eugen Ulmer, Stuttgart.
- Preston, C. D. & J. M. Croft, 2001. *Aquatic Plants in Britain and Ireland*. Harley Books, Colchester, Essex.
- Roelofs, J. G. M., 1983. Impact of acidification and eutrophication on macrophyte communities in soft waters in the Netherlands. I. Field observations. *Aquatic Botany* 17: 139–155.
- Roelofs, J. G. M., J. A. A. R. Schuurkes & A. J. M. Smits, 1984. Impact of acidification and eutrophication on macrophyte communities in soft waters in the Netherlands. II. Experimental studies. *Aquatic Botany* 18: 389–411.
- Sala, O. E., F. S. Chapin III, J. J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L. F. Huenneke, R. B. Jackson, A. Kinzig, R. Leemans, D. M. Lodge, H. A. Mooney, M. Oesterheld, N. LeRoy Poff, M. T. Sykes, B. H. Walker, M. Walker & D. H. Wall, 2000. Global biodiversity scenarios for the year 2100. *Science* 287: 1770–1774.
- Schaminée, J. H. J. & G. H. P. Arts, 1992. Die Strandlinggesellschaften (*Littorelletea* Br.-Bl. et Tx. 43) der Niederlande, in europäischem Rahmen gefasst. *Phytocoenologia* 20: 529–558.
- Schoof-van Pelt, M. M., 1973. *Littorelletea – A Study of the Vegetation of Some Amphiphytic Communities of Western Europe*. Stichting Studentenpers, Nijmegen.
- Schubert, R., W. Hilbig & S. Klotz, 1995. *Bestimmungsbuch der Pflanzengesellschaften Mittel- und Nordostdeutschlands*. Gustav Fischer, Stuttgart.

- Sculthorpe, C. D., 1967. *The Biology of Aquatic Vascular Plants*. Edward Arnold, London.
- Smolders, A. J. P., E. C. H. E. T. Lucassen & J. G. M. Roelofs, 2002. The isoetid environment: biogeochemistry and threats. *Aquatic Botany* 73: 325–350.
- Szankowski, M. & S. Klosowski, 2001. Habitat conditions of the phytocenoses dominated by *Luronium natans* (L.) Rafin. in Poland. *Hydrobiologia* 455: 213–222.
- Thomas, J. D. & P. W. G. Daldorph, 1994. The influence of nutrient and organic enrichment on a community dominated by macrophytes and gasteropod mollusks in a eutrophic drainage channel: relevance to snail control and conservation. *Journal of Applied Ecology* 31: 571–588.
- Vuille, F.-L., 1988. The reproductive biology of the genus *Baldellia* (Alismataceae). *Plant Systematics and Evolution* 159: 173–183.
- Willby, N. J. & J. W. Eaton, 1993. The distribution, ecology and conservation of *Luronium natans* (L.) Raf. in Britain. *Journal of Aquatic Plant Management* 31: 70–76.