

Diabetologia (2005) 48: 1523–1533
DOI 10.1007/s00125-005-1816-1

ARTICLE

Domenico Bosco · Paolo Meda · Philippe Morel ·
David Matthey-Doret · Dorothée Caille ·
Christian Toso · Leo H. Bühler · Thierry Berney

Expression and secretion of alpha1-proteinase inhibitor are regulated by proinflammatory cytokines in human pancreatic islet cells

Received: 4 November 2004 / Accepted: 6 March 2005 / Published online: 7 July 2005
© Springer-Verlag 2005

Abstract *Aims/hypothesis:* Alpha1-proteinase inhibitor (α 1-PI) has been considered a key player in inflammatory processes. In humans, the main production site of α 1-PI is the liver, but other tissues, including pancreatic islets, also synthesise this molecule. The aims of this study were to assess the islet cell types that produce α 1-PI, to determine whether α 1-PI is actually secreted by islet cells, and to assess how its production and/or secretion are regulated. *Methods:* Expression of α 1-PI in human islet cells was assessed by immunofluorescence, electron microscopy and western blotting. Release of α 1-PI was analysed by reverse haemolytic plaque assay and ELISA. The effects of cytokines on α 1-PI synthesis and secretion were tested. *Results:* Immunofluorescence showed that alpha and delta cells do express α 1-PI, whereas beta cells do not. By electron microscopy, we demonstrated a colocalisation of α 1-PI with glucagon and somatostatin within secretory granules. Immunolabelling also revealed localisation of α 1-PI within the Golgi apparatus, related vesicles and lysosomal structures. The expression of α 1-PI in islet cells was also demonstrated by western blotting and ELISA of protein extracts. ELISA and reverse haemolytic plaque

assay showed that α 1-PI is secreted into the culture medium. Treatment of islet cells with IL-1 β and oncostatin M for 4 days increased the production and release of α 1-PI. *Conclusions/interpretation:* Our results demonstrate that α 1-PI is expressed by the alpha and delta cells of human islets, and that proinflammatory cytokines enhance the production and release of this inhibitor.

Keywords Human islets · IL-1 β · Oncostatin M · α 1-Proteinase inhibitor · Reverse haemolytic plaque assay

Abbreviations OSM: oncostatin M · α 1-PI: alpha1-proteinase inhibitor · RHPA: reverse haemolytic plaque assay

Introduction

Alpha1-proteinase inhibitor (α 1-PI) is a 52-kDa acute phase protein that inhibits serine proteases, especially elastase, which is released by human neutrophils during inflammatory processes [1]. Deficiency of α 1-PI is associated with chronic liver disease, and is often associated with pulmonary emphysema. In humans, the liver is the major site of expression of α 1-PI. Hepatocytes synthesise and secrete systemic α 1-PI, and plasma concentrations of α 1-PI acutely increase three to four-fold in response to 'secondary' cytokines, such as members of the IL-6 family, including IL-6 itself, leukaemia inhibitory factor and oncostatin M (OSM) [2, 3]. It is thought that α 1-PI is involved in the tissue repair reactions that follow inflammation.

The way secreted α 1-PI acts as an anti-inflammatory agent is not well understood. The inactivation of proteolytic enzymes, such as serine proteinases of inflammatory cells is undoubtedly of key importance, but various observations indicate that other mechanisms are also involved. Thus, α 1-PI has a neutrophil chemoattractant activity [4, 5], induces changes in neutrophil shape, and increases cell-to-substrate adhesion [5] and spreading of smooth muscle cells in a fibrin gel [6]. Moreover, this molecule also acts as

D. Bosco · P. Morel · D. Matthey-Doret · C. Toso ·
L. H. Bühler · T. Berney
Cell Isolation and Transplantation Center,
Department of Surgery,
Geneva University Hospitals,
Geneva, Switzerland

P. Meda · D. Caille
Department of Cell Physiology and Metabolism,
School of Medicine,
University of Geneva,
Geneva, Switzerland

D. Bosco (✉)
Cell Isolation and Transplantation Center,
Centre Médical Universitaire,
1 rue Michel-Servet,
1211 Geneva 4, Switzerland
e-mail: domenico.bosco@medecine.unige.ch
Tel.: +41-22-3795164
Fax: +41-22-3795054

a secretagogue, since it induces degranulation of neutrophils [5] and, if complexed with elastase, increases its own synthesis by macrophages [7]. α 1-PI may also have a protective effect against factors inducing apoptosis by a mechanism unidentified to date. Thus, α 1-PI has been shown to specifically inhibit the apoptosis of hepatocytes induced by tumor necrosis factor [8], as well as that induced by prolonged ischaemia followed by reperfusion of organs in vivo [9]. It has been also suggested that α 1-PI may act as an antioxidant, given its high content in methionine [10].

Although the liver is the major site of expression of α 1-PI, many studies have shown that this molecule is also synthesised in other tissues and organs, in response to an inflammatory insult. For instance, lung epithelial cells respond dramatically to OSM, a cytokine of the IL-6 family, by synthesising and secreting α 1-PI [11–14]. In addition, local inflammatory conditions have been shown to upregulate the production of α 1-PI by monocytes [15], corneal cells [16], enterocytes and Paneth cells [17], and articular chondrocytes [18]. Taken together, these data indicate that local production of α 1-PI may contribute to the defence of tissues against different types of injuries.

Preservation of a sufficient cell mass and function by endogenous mechanisms capable of protecting against cell injury is particularly important for the pancreatic islets of Langerhans. Indeed, the integrity of these micro-organs is altered in many pathological conditions. In type 1 diabetes, activation of autoimmune events leads to a local peri-islet inflammation that is particularly aggressive toward beta cells. In type 2 diabetes, amyloid deposits alter the function and viability of beta cells [19, 20], and it is known that α 1-PI can act in vitro as an inhibitor of amyloid fibril formation, while promoting disaggregation of β -amyloid fibrils [21]. In pancreatitis, endocrine cells have to defend against the aggression of proteases released as a result of the destruction of the exocrine tissue. Finally, in islet of Langerhans transplantation for the treatment of type 1 diabetes, islets infused into the liver are subject to inflammatory insults at the site of implantation in addition to more specific insults such as immune rejection [22]. In these situations, α 1-PI could play an important protective role in the progression of damage to beta cells. Even though previous reports have documented the presence of α 1-PI within pancreatic islets [23–26], the type of islet cells producing the inhibitor is still a matter of controversy. In addition, whether islet cells secrete α 1-PI, and whether and how this secretion is regulated has not been shown.

Materials and methods

Islet isolation

Pancreases were harvested from multiorgan cadaveric donors. Islets were isolated using a previously described modification of the automated method [27, 28]. Islets were

then rinsed with CMRL 1066 medium (Sigma, St Louis, MO), and aliquots of 20,000 islets were incubated in 30 ml CMRL 1066 medium containing 5.6 mmol/l glucose and supplemented with 10% fetal calf serum, 25 mmol/l *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES), 2 mmol/l glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin (hereafter, referred to as complete CMRL). After 12 h at 37°C, culture medium was changed and islets were further cultured at 24°C for 1 to 3 days.

Islet cell preparation

Islets were washed three times with phosphate-buffered saline without Mg^{2+} and Ca^{2+} (PBS). Aliquots of 10^4 islets were resuspended in 1 ml 0.025% trypsin solution containing 1 mmol/l EDTA (Gibco, Paisley, Scotland), and incubated at 37°C for 9–10 min, with occasional pipetting through a 1-ml plastic tip. Trypsinisation was stopped by addition of 10 ml cold complete CMRL, and cells were washed twice with the same medium. Cells were then counted and aliquots of either 3×10^5 or 1×10^6 cells were incubated for 24 h in 10-cm-diameter, non-tissue culture-treated petri dishes containing 10 ml complete CMRL.

Culture conditions and cytokine treatment

To study the effect of cytokines on α 1-PI expression by western blotting, aliquots of 4,500 islets were seeded in non-tissue culture-treated petri dishes (60 mm diameter) and incubated for 4 days in the presence of 4 ml complete CMRL medium supplemented, or not, with 1 mmol/l dexamethasone, 20 ng/ml IL-1 β (Biosource International, Nivelles, Belgium), 20 ng/ml OSM (Biosource International) and 0.5 mmol/l isobutyl-methylxanthine (IBMX; Sigma). To study the effect of cytokines on α 1-PI synthesis by enzyme-linked immunosorbent assay (ELISA) and immunofluorescence, aliquots of 15×10^4 islet cells were seeded onto coverslips (15 mm diameter), and placed into a 24-well plate. Then, cells were incubated for 4 days in the presence of 0.5 ml complete CMRL, supplemented, or not, with 1 mmol/l dexamethasone, 20 ng/ml IL-1 β , 20 ng/ml OSM and 0.5 mmol/l IBMX. To analyse the secretion of α 1-PI, islet cells were incubated as described above except that a serum-free complete CMRL medium was used.

Antibodies

Goat antihuman α 1-PI polyclonal antibody (A-1845) was purchased from Sigma, and rabbit antihuman α 1-PI polyclonal antibodies were purchased from Dako (A-0012; Glostrup, Denmark), Sigma (A-0409) and Zymed (18-0002; San Francisco, CA, USA). Rabbit antiglucagon (A565), somatostatin (A0566) and pancreatic polypeptide (A0619) polyclonal antibodies were purchased from Dako.

A guinea-pig anti-insulin polyclonal antibody was developed in our laboratory, using porcine insulin as immunogen, as previously described [29].

Immunofluorescence

To analyse the expression of $\alpha 1$ -PI and islet hormones by immunofluorescence, aliquots of 10^4 islets cells were left to attach for 60 min at 37°C into Cunningham chambers. These chambers were rinsed with PBS, fixed in a 4% paraformaldehyde-PBS solution for 20 min, rinsed again with PBS and stored at 4°C. Samples of human pancreases were snap frozen and embedded in optimal cutting temperature (OCT) compound (Sakura Finetek Europe Zoeterwoude, The Netherlands). Five-micrometre sections were prepared and fixed in 4% paraformaldehyde-PBS solution for 20 min. Then, sections were rinsed and used immediately for immunofluorescence labelling.

The whole immunofluorescence procedure was carried out at room temperature and all solutions were prepared using PBS. After fixation, cells and sections were rinsed three times for 5 min in PBS, treated for 15 min with 0.1% Triton X-100 and exposed to a 0.1% bovine serum albumin (BSA) solution for 20 min. Cells were then incubated for 2 h with one, two or three of the primary antibodies, as indicated in the Results section. The dilutions were 1:1,200 for the anti-insulin antibody, and 1:400 for the both anti-glucagon and the antisomatostatin antibodies. For the anti- $\alpha 1$ -PI, dilutions were 1:1,200, 1:400 and 1:200, for the Sigma goat antibody, and the rabbit antibodies from Sigma and Dako, respectively.

Immunogold electron microscopy

Pellets of isolated islets were washed twice in 0.1 mol/l phosphate buffer, followed by 5-min fixation at room temperature in 4% paraformaldehyde and 0.1% glutaraldehyde, 60-min fixation in 4% paraformaldehyde (all fixatives diluted in 0.1 mol/l phosphate buffer, pH 7.4). The islets were washed three times in 0.1 mol/l phosphate buffer, embedded in 12% gelatin and cooled on ice. Small blocks of gelatin-embedded islets were infused with 2.3 mol/l sucrose, snap frozen in liquid nitrogen, and sectioned with an EMFCS cryoultramicrotome (Leica Microsystems, Wetzlar, Germany). Ultrathin sections were mounted on Parlodion-coated copper grids. The sections were treated as per a modification [30] of a previously described technique [31]. Briefly, sections were sequentially incubated as follows: twice for 2 min in 0.1% glycine (in PBS), once for 2 min in 1% BSA (in PBS), 1 h at room temperature in the presence of the first primary antibody, rinsed four times for 2 min in 0.1% BSA, and exposed 20 min at room temperature to either protein A-coated gold particles (15 nm diameter, for islet hormones) or to goat antimouse or antirabbit antibodies, whichever appropriate, coupled to 15-nm and to 10-nm diameter gold particles, respectively. After two 2-min rinses in 0.1% BSA, and two 2-min rinses

in PBS, the sections were exposed for 5 min to 1% glutaraldehyde (in 0.1 mol/l sodium phosphate buffer, pH 7.4), rinsed twice for 5 min in PBS, exposed for 1 h at room temperature to a second primary antibody (when required), rinsed four times for 2 min in 0.1% BSA, exposed for 20 min at room temperature to either protein A-coated gold particles or an appropriate gold-conjugated goat antibodies, rinsed twice 2 min in 0.1% BSA, rinsed twice for 2 min in PBS, exposed for 5 min to 1% glutaraldehyde (in 0.1 mol/l sodium phosphate buffer, pH 7.4), rinsed twice for 5 min in PBS, rinsed six times for 1 min in distilled water, and eventually stained for 5 min in 2% uranyl acetate oxalate (pH 7), followed by 2% methylcellulose containing 0.5% uranyl acetate. Cryosections were screened and photographed in a CM10 electron microscope (Philips, Eindhoven, The Netherlands).

Single immunolabelling for $\alpha 1$ -PI was carried using the rabbit polyclonal serum (Dako) diluted 1:40, and revealed with a goat anti rabbit IgG serum coupled to 10-nm gold particles (British Biocell International, Cardiff, Wales) and diluted 1:5. Double immunolabelling for islet hormones and $\alpha 1$ -PI was performed using the same anti $\alpha 1$ -PI serum and one of the following antibodies: a mouse monoclonal antibody against glucagon (Sigma), diluted 1:100, and revealed with a goat antimouse IgG serum coupled to 15-nm gold particles (British Biocell International) diluted 1:5; a goat antisomatostatin antibody (a kind gift of M.P. Dubois), diluted 1:40, and revealed with 15-nm protein A-coated gold particles diluted 1:150; a mouse monoclonal against insulin (a kind gift of M.J. Storch), diluted 1:100, and revealed with 15-nm protein A-coated gold particles diluted 1:150.

Controls included exposure of the sections to only the protein A-coated gold particles or to the gold-conjugated goat antibodies against either rabbit or mouse IgGs, whatever was appropriate. None of these incubations resulted in a sizeable, specific staining of the sections.

Western blot analysis

For preparation of total protein extracts, islets were incubated under the different conditions detailed in the results, rinsed three times in PBS and lysed by sonication in 80 mmol/l Tris-HCl buffer (pH 7.4), containing 5% SDS, 5 mmol/l EDTA and a cocktail of protease inhibitors (Complete Mini Protease Inhibitor Cocktail Tablets, Roche, Rotkreuz, Switzerland), which was used as per the manufacturer's recommendations. Protein content was determined by the Bio-Rad DC protein assay reagent kit (Bio-Rad, CA). Ten-micrograms aliquots of total proteins were diluted in gel loading buffer (0.025 mol/l Tris pH 6.8, 0.5% SDS, 1% 2-mercaptoethanol, 0.025% bromophenol blue, 17.5% glycerol). Aliquots were loaded on a 7.5% polyacrylamide-SDS gel. After separation, proteins were transferred to a nitrocellulose membrane (Protran; Schleicher & Schuell, Keene, NH), which was incubated overnight in Tris-buffered saline (TBS; Tris 0.01 M, pH 7.4, 0.15 mol/l NaCl) containing 5% non-fat dried milk and 0.1% Tween,

and subsequently incubated for 1 h with a polyclonal rabbit antihuman α 1-PI antibody (Dako), diluted 1:2,000 in the same buffer. After extensive washes in TBS containing 0.1% Tween (TBS-Tween), the membrane was probed for 1 h with a horseradish-peroxidase-conjugated secondary antibody against rabbit IgG (Biorad), diluted 1:6,000 in TBS-Tween, rinsed again in TBS-Tween, revealed with the SuperSignal West chemiluminescent substrate kit (Amersham-Pharmacia, Buckinghamshire, UK) according to manufacturer's instructions, and exposed for 1 min to Hyperfilm (Amersham-Pharmacia).

Expression and release of α 1-PI

The islet content in α 1-PI and the amounts of inhibitor secreted in the medium were measured by ELISA (Immundiagnostik, Bensheim, Deutschland).

A reverse haemolytic plaque assay (RHPA) was used as described [32] to attribute the secretion of α 1-PI to specific islet cell types. Briefly, isolated islet cells were suspended in a Krebs-Ringer-bicarbonate buffer containing 12.5 mmol/l HEPES and 0.1% BSA (control KRB) and mixed with 5% (v/v) packed sheep red blood cells (SRBC, Behring Institute, Marburg, Germany) that had been previously coated with protein A. Then, the cells were rinsed and incubated for 2 h at 37°C in the presence of a goat antihuman α 1-PI polyclonal antibody (Sigma), diluted 1:30 in control KRB supplemented or not with 16.7 mmol/l glucose, 10 μ mol/l forskolin and 100 nmol/l phorbol 12-myristate 13 acetate (PMA). The cells were rinsed with control KRB and incubated for 1 h at 37°C in the presence of guinea pig complement (Behring Institute), diluted 1:40 in control KRB. Chambers were filled with 0.04% (wt/vol) solution of trypan blue in control KRB, rinsed with control KRB, and filled with Bouin's fixative or 4% paraformaldehyde. After immunofluorescence for insulin or α 1-PI, the proportion of secreting cells (as determined by the presence of a surrounded haemolytic plaque that was identified as a circular area centered on an islet cell and containing the dark lysed erythrocytes) was determined under a microscope equipped with a 40 \times objective and a combined phase-contrast and fluorescence illumination. Analysis was restricted to single cells that excluded trypan blue at the end of the plaque assay. Haemolytic plaques were not detected when SRBC not coated with protein A were used as controls, nor when either the anti- α 1-PI antibody or complement were omitted.

Statistical analysis

Statistical analysis was performed by analysis of variance (ANOVA) and LSD Post Hoc tests using SPSS (Statistical Package for Social Sciences, Chicago, IL). Student's *t*-test was performed for comparison between two groups using the Statistica software package (StatSoft, Tulsa, OK). Values of $p < 0.05$ were considered significant.

Results

α 1-PI is selectively expressed in alpha and delta cells

Expression of α 1-PI was studied by immunofluorescence on dispersed islet cells prepared from seven human pancreases, and cultured under conditions favouring survival of endocrine cells. In all preparations, 13–27% of the cells stained positively for α 1-PI (means \pm SEM: 22.1 \pm 2.0%), (Fig. 1a). To identify the cell types expressing α 1-PI, double staining was performed for α 1-PI and either insulin, glucagon, somatostatin or pancreatic polypeptide (Fig. 1b). Virtually no insulin-producing beta cell and only few pancreatic polypeptide-producing cells (PP-cells) expressed α 1-PI. In contrast, most glucagon- (alpha cells) and somatostatin-producing cells (delta cells) were labelled for α 1-PI. This pattern was confirmed using triple immunolabelling. Thus, we found no beta cell co-expressing α 1-PI, but observed that most glucagon- and somatostatin-immunoreactive cells also expressed α 1-PI (Fig. 2).

On frozen sections of four normal pancreases, immunofluorescence showed α 1-PI predominantly within cells situated at the periphery of islets, most of which stained for either somatostatin or glucagon, but did not detect the inhibitor in the insulin-immunoreactive cells which occupy the core of the islets (Fig. 3).





Fig. 1 α 1-PI immunoreactive cells are abundant among islet cells and correspond to alpha and delta cells. **a** The percentages of cells labelled for insulin (β), glucagon (α), somatostatin (δ), pancreatic polypeptide (PP) and α 1-PI are shown. Values are the means \pm SEM of three independent experiments. A minimum of 200 cells was scored per condition and experiment. * $p < 0.0001$ and ** $p < 0.001$ when compared with the beta-cell group (β). **b** The percentage of cells co-expressing α 1-PI and either insulin (β), glucagon (α), somatostatin (δ) or pancreatic polypeptide (PP) are shown. Values are means \pm SEM of three independent experiments. A total of 305 insulin-, 231 glucagon-, 82 somatostatin- and 123 pancreatic-polypeptide-expressing cells was scored. * $p < 0.0001$ when compared with the beta-cell group (β).

Fig. 2 α 1-PI was detected in somatostatin- but not insulin-containing cells after triple immunofluorescence labelling. Insulin-expressing cells are stained in *blue* (a), α 1-PI-expressing cells in *green* (b) and somatostatin-expressing cells in *red* (c). *Merged images* (d) show that beta cells do not contain detectable levels of α 1-PI, whereas it is abundant in one somatostatin-expressing cell (in *yellow*). A phase-contrast view of the same cells (e). *Scale bar*: 20 μ m



α 1-PI is localised in multiple secretory compartments of glucagon- and somatostatin-containing cells

Immunolabelling of ultrathin cryosections of human islets resulted in a specific staining for α 1-PI within glucagon- and somatostatin-expressing cells, but not within insulin-expressing cells (Fig. 4a). In alpha and delta cells, α 1-PI was found within multiple secretory compartments, including the stacks and vesicles of the Golgi apparatus (Fig. 5a), the hormone-containing secretory granules (Fig. 4a–d) and a variety of lysosomal structures (Fig. 5a). In contrast, no labelling was observed within nuclei, mitochondria and cell membranes (Fig. 4a, Fig. 5a).

α 1-PI synthesis by islet cells is regulated by cytokine stimulation


Isolated islet cells were cultured for 4 days in the absence or presence of 0.5 mmol/l IBMX, 20 ng/ml IL-1 β and 20 ng/ml OSM, with, or without, 1 μ mol/l dexamethasone, before α 1-PI content of islet cells was measured by ELISA

after protein extraction. In the absence of dexamethasone, IL-1 β and OSM affected α 1-PI synthesis poorly (Fig. 6a). In contrast, in the presence of dexamethasone, treatment of islet cells by both cytokines resulted in a two- to three-fold increase in α 1-PI content (Fig. 6a). Irrespective of the presence of dexamethasone, incubation with IBMX with the aim to increase cAMP had no effect on α 1-PI synthesis (Fig. 6a).

The effect of cytokines on α 1-PI expression was further assessed by quantifying the immunofluorescence labelling of α 1-PI, related to that of the cell hormone content. In the presence of dexamethasone, IL-1 β and OSM, but not IBMX, significantly increased the labelling of α 1-PI compared with that of glucagon (Fig. 6b). This change was due to an actual increase in immunolabelling for α 1-PI, inasmuch as the ratio of glucagon to insulin staining remained unchanged (Fig. 6b).

In further experiments, cultured intact human islets were exposed for 4 days to cytokines, in the presence or absence of dexamethasone, and the expression of α 1-PI was analysed by western blotting. Increased levels of immunoreactive α 1-PI were observed in islets treated with IL-1 β and

Fig. 3 α 1-PI colocalises with glucagon, but not insulin, within intact human pancreas. Insulin-containing cells appear in blue (a), α 1-PI-expressing cells in green (b) and glucagon-expressing cells in red (c). Merged images (d) demonstrate that beta cells (blue) did not contain detectable levels of α 1-PI, whereas the inhibitor was present in several glucagon-containing cells (in yellow). Cells labelled only for α 1-PI (green in d) could be either delta cells or intra-islet endothelial cells [24]. Scale bar: 40 μ m



OSM, as compared with control islets (Fig. 6c). Addition of dexamethasone resulted in a much stronger stimulation of α 1-PI expression (Fig. 6c).

α 1-PI is secreted by islet cells

To assess whether α 1-PI is secreted by islet cells, we developed a RHPA that allows for a direct visualisation of α 1-PI release by individual islet cells. After a 2 h-incubation in the presence of antibodies against α 1-PI, and a 1-h incubation in the presence of complement, several islet cells were surrounded by a haemolytic plaque, as judged by the presence of dark, non-refracting red blood cell ghosts (Fig. 7a). Immunofluorescence for insulin showed that only non-beta cells were surrounded by a haemolytic plaque. Immunofluorescence for α 1-PI confirmed that cells surrounded by a haemolytic plaque contained α 1-PI (Fig. 7a). Quantitative analysis showed that $16.7 \pm 0.7\%$ (mean \pm SEM of three independent experiments) of the cells that did not stain for insulin were surrounded by a haemolytic plaque. This percentage did not change when cells were stimulated with glucose (16.7 ± 0.9 , $n=3$) or with 0.5 mmol/l IBMX+10 μ mol/l forskolin+100 nmol/l PMA (16.0 ± 1.0 , $n=3$).

To assess whether the release of α 1-PI was regulated by dexamethasone and cytokines, islet cells were exposed for 4 days to these agents and both the cumulated (4 day culture, Fig. 7b) and acute release (1 h in control medium, Fig. 7d) of α 1-PI was measured by ELISA, and compared

with the release of insulin (Fig. 7c). Results show that both chronic and acute releases of α 1-PI were significantly increased by IL-1 β and OSM, but only when dexamethasone was present in the medium (Fig. 7b,d). In contrast, IL-1 β and OSM did not affect insulin secretion, which was stimulated by IBMX and decreased by dexamethasone (Fig. 7c).

Discussion

Previous reports have documented that α 1-PI was preferentially expressed by peripheral islet cells, but failed to positively identify these cells [25, 26, 33]. Using different approaches, our study documents now that, in pancreases from brain-dead heart-beating donors, α 1-PI is expressed in differentiated alpha, delta and PP-cells, accounting for about 22% of the isolated islet cells.

The reason why α 1-PI is located preferentially in the alpha and delta cells of human pancreatic islets, within only a few PP-cells and not in beta cells, remains to be understood. One possibility is that it is related to lineage relationships between the different islet cell types. Indeed, it has been shown that there is a close ontogenetic relationship between pancreatic polypeptide- and insulin-producing cells, whereas adult glucagon- and insulin-expressing cells originate from different precursors [34, 35].

Our experiments show that α 1-PI is expressed in multiple secretory compartments, including the Golgi apparatus, lysosomes and secretory granules. While the




Fig. 4 α 1-PI was found in glucagon- and somatostatin-containing granules, but not in insulin-containing granules. **a** Low-power view of a cryosection immunolabelled for α 1-PI showing the inhibitor within the secretory granules (*arrows*) of glucagon- (α) and somatostatin-containing cells (δ), but not within the granules (*arrowheads*) of insulin-containing cells (β). **b–d** Double immunolabelling for α 1-PI and one islet hormone demonstrated co-expression of the inhibitor (small gold particles) and either glucagon or somatostatin (large gold particles) within individual secretory granules of (**c**) alpha and (**d**) delta cells. In contrast, no α 1-PI was detected within the insulin-containing granules of beta cells (**b**), when an antibody was used that resulted in intense labelling of the granules of an adjacent alpha cell (**b**). *Scale bar*: 1 μ m (**a**), 500 nm (**b–d**)

presence of α 1-PI in the Golgi apparatus and in lysosomes is not surprising, the finding of this inhibitor in secretory granules containing glucagon or somatostatin is unexpected, in view of previous reports considering this inhibitor as a constitutive secretory marker [36]. In secretory cells, trafficking from the Golgi apparatus onwards involves vesicles that are either directly conveyed to the cell surface (constitutive secretion), targeted to the lysosomal compartments or stored in granules belonging to the regulated secretory pathway. A genetic variant of α 1-PI with a single Arg \rightarrow Met³⁵⁸ substitution was shown to be transported into granules and to inhibit conversion of hormones during their transit through the regulated secretory pathway [37–39]. Also, the native form of α 1-PI cells was found at low levels in secretory granules after transfection of MIN6 [39], suggesting that α 1-PI could be allocated to




Fig. 5 α 1-PI was detected in multiple secretory compartments of glucagon-producing cells. **a** Immunolabelling revealed α 1-PI within the Golgi apparatus (*double arrowhead*), the secretory granules (*arrows*) and the lysosomes (*single arrowheads*) of a glucagon-producing alpha cell. Note the absence of labelling in an adjacent beta cell (β). **b** Omission of the antibody against α 1-PI resulted in no gold labelling of secretory granules of an alpha cell (α). *Scale bar*, 1 μ m (**a**), 500 nm (**b**)

the regulated pathway. However, this finding does not demonstrate that the α 1-PI released and detected in the medium comes from a regulated secretory pathway. Rather, our results showing that IBMX and PMA did not stimulate α 1-PI secretion support the hypothesis that this inhibitor is predominantly released by a constitutive pathway. Our finding that α 1-PI is localised in hormone-containing granules, argues also for a role of this inhibitor in the regulation of glucagon and somatostatin conversion within differentiated islet cells. This process starts in the Golgi apparatus and immature granules, and involves proteolytic cleavage by a family of proteases called convertases [40].

We further show here that α 1-PI expression and release by human islets was increased by cytokines. The plasma levels of α 1-PI increase during acute and chronic inflammatory processes [1–3]. Certainly, the main source of this inhibitor in humans is the liver, and the secretion of α 1-PI by hepatocytes was shown to increase in response to proinflammatory cytokines, such as IL-1 β and IL-6 [2].




Fig. 7 α 1-PI is released by islet cells. **a** After a 2-h incubation in control medium, a minority of islet cells secreted α 1-PI, and are seen surrounded by circular haemolytic plaques, containing dark ghosts of complement-lysed erythrocytes (*left*). Immunostaining for insulin (*right*) revealed that beta cells were not surrounded by haemolytic plaques, which were always centred on non-beta cells. Inserts show one cell surrounded by a haemolytic plaque, which stained positively for α 1-PI. Scale bar: 30 μ m. **b** Islet cells were cultured in serum-free medium, in the absence (*-Dex*) or presence (*+Dex*) of 1 μ mol/l dexamethasone. Standard medium (*control*) was supplemented with 0.5 mmol/l IBMX, 20 ng/ml IL-1 β or 20 ng/ml OSM. Aliquots of supernatant were taken after 4 days in culture and α 1-PI content measured by ELISA. Results of control, IL-1 β , OSM (four experiments) and IBMX (two experiments) are means \pm SEM. * p <0.001 when compared to control in the presence of dexamethasone. **c** Islet cells were cultured as described (**b**). Aliquots of

medium were taken after 4 days of culture and insulin release measured by ELISA. Data show that only IBMX affected insulin release, both in the absence (*-Dex*) and presence of dexamethasone (*+Dex*). Results are expressed as means \pm SEM of six experiments for control, IL-1 β and OSM, and four experiments for IBMX. * p <0.005 and ** p <0.05 when compared with the corresponding condition without dexamethasone. **d** Islet cells were cultured as described (**b**). After 4 days, the cells were rinsed and incubated in control medium for 1 h. Aliquots of culture supernatant were taken and α 1-PI content measured by ELISA. In the absence of dexamethasone (*-Dex*), results are means \pm SEM of two experiments. *ND*, non-determined. In the presence of dexamethasone (*+Dex*), results are means \pm SEM of three experiments for all conditions, except for IBMX where n =1. * p <0.05 compared with the control in the presence of dexamethasone

Further in-vitro experiments are needed to verify that α 1-PI is really involved in protecting islet cells from the injury caused by cytokines, such as IL-1 β , TNF- α and IFN γ [51]. In this regard, it is interesting to note that IL-6, a cytokine related to OSM, was shown to confer protection to pancreatic islet beta cells from cytokine-induced cell death

[52]. It will also be important to assess the expression of α 1-PI in islets of diabetic patients. One implication of our study is that the expression of α 1-PI should be affected in islets showing a well-characterised insulinitis and beta-cell destruction. In some diabetic patients, the plasma concentration and activity of α 1-PI was found to be decreased

compared with controls [53–55], but the pathogenesis of this finding was not investigated.

In addition to a possible role in regulating cell injury induced by inflammatory cytokines, we propose that α 1-PI, produced by the pancreatic alpha and delta cells, may also act as an antiprotease against enzymes released by the surrounding exocrine tissue. In this perspective, alpha and delta cells could protect the more centrally located islet cells, mostly beta cells. This function could be particularly important during the development of acute pancreatitis when leakage of aggressive enzymes from surrounding inflamed acinar cells is expected.

Finally, our results show that islet cells are able to release α 1-PI. We measured a higher concentration of α 1-PI in the culture supernatant after cytokine treatment. This increase is accounted for by a higher synthesis of α 1-PI after cytokine treatment, than by a major increment in the release of the inhibitor. When islet cells treated with cytokines were further incubated in control medium, they continued to release more α 1-PI than untreated cells. These observations suggest that α 1-PI in islet cells is secreted in a constitutive manner.

In summary, our results clearly show that human islet alpha and delta cells express and secrete α 1-PI, and that these events are modulated by cytokines. Taken together our data suggest that local production of α 1-PI in islets may be triggered by inflammatory processes that occur at the onset of diabetes, the development of acute pancreatitis and conditions perturbing the native islet environment, such as transplantation. Further studies should determine whether and how α 1-PI is involved in these pathological conditions.

Acknowledgements This work was supported by a grant from the Swiss National Science Foundation (3200B0-102134 to P. Morel, D. Bosco, T. Berney, and L.H. Bühler). We thank C. Sinigaglia and N. Pernin for excellent technical help. P. Meda's team was supported by grants from the Swiss National Science Foundation (31-67788.02), the Juvenile Diabetes Research Foundation International (1-2001-622 and 5-2004-255), the European Union (QLRT-2001-01777) and the National Institutes of Health (1R01 DK-63443-01).

References

- Travis J, Salvesen GS (1983) Human plasma proteinase inhibitors. *Annu Rev Biochem* 52:655–709
- Baumann H, Gaudie J (1994) The acute phase response. *Immunol Today* 15:74–80
- Moshage H (1997) Cytokines and the hepatic acute phase response. *J Pathol* 181:257–266
- Banda MJ, Rice AG, Griffin GL, Senior RM (1988) The inhibitory complex of human alpha 1-proteinase inhibitor and human leukocyte elastase is a neutrophil chemoattractant. *J Exp Med* 167:1608–1615
- Parmar JS, Mahadeva R, Reed BJ et al (2002) Polymers of alpha(1)-antitrypsin are chemotactic for human neutrophils: a new paradigm for the pathogenesis of emphysema. *Am J Respir Cell Mol Biol* 26:723–730
- Ikari Y, Fujikawa K, Yee KO, Schwartz SM (2000) Alpha(1)-proteinase inhibitor, alpha(1)-antichymotrypsin, or alpha(2)-macroglobulin is required for vascular smooth muscle cell spreading in three-dimensional fibrin gel. *J Biol Chem* 275:12799–12805
- Joslin G, Krause JE, Hershey AD, Adams SP, Fallon RJ, Perlmutter DH (1991) Amyloid-beta peptide, substance P, and bombesin bind to the serpin-enzyme complex receptor. *J Biol Chem* 266:21897–21902
- Van Molle W, Libert C, Fiers W, Brouckaert P (1997) Alpha 1-acid glycoprotein and alpha 1-antitrypsin inhibit TNF-induced but not anti-fas-induced apoptosis of hepatocytes in mice. *J Immunol* 159:3555–3564
- Daemen MA, Heemskerk VH, van't Veer C et al (2000) Functional protection by acute phase proteins alpha(1)-acid glycoprotein and alpha(1)-antitrypsin against ischemia/reperfusion injury by preventing apoptosis and inflammation. *Circulation* 102:1420–1426
- Brantly M (2002) Alpha1-antitrypsin: not just an antiprotease: extending the half-life of a natural anti-inflammatory molecule by conjugation with polyethylene glycol. *Am J Respir Cell Mol Biol* 27:652–654
- Cichy J, Potempa J, Chawla RK, Travis J (1995) Stimulatory effect of inflammatory cytokines on alpha 1-antichymotrypsin expression in human lung-derived epithelial cells. *J Clin Invest* 95:2729–2733
- Cichy J, Potempa J, Travis J (1997) Biosynthesis of alpha1-proteinase inhibitor by human lung-derived epithelial cells. *J Biol Chem* 272:8250–8255
- Sallenave JM, Tremblay GM, Gaudie J, Richards CD (1997) Oncostatin M, but not interleukin-6 or leukemia inhibitory factor, stimulates expression of alpha1-proteinase inhibitor in A549 human alveolar epithelial cells. *J Interferon Cytokine Res* 17:337–346
- Boutten A, Venembre P, Seta N et al (1998) Oncostatin M is a potent stimulator of alpha1-antitrypsin secretion in lung epithelial cells: modulation by transforming growth factor-beta and interferon-gamma. *Am J Respir Cell Mol Biol* 18:511–520
- Knoell DL, Ralston DR, Coulter KR, Wewers MD (1998) Alpha 1-antitrypsin and protease complexation is induced by lipopolysaccharide, interleukin-1beta, and tumor necrosis factor-alpha in monocytes. *Am J Respir Crit Care Med* 157:246–255
- Boskovic G, Twining SS (1998) Local control of alpha1-proteinase inhibitor levels: regulation of alpha1-proteinase inhibitor in the human cornea by growth factors and cytokines. *Biochim Biophys Acta* 1403:37–46
- Molmenti EP, Perlmutter DH, Rubin DC (1993) Cell-specific expression of alpha 1-antitrypsin in human intestinal epithelium. *J Clin Invest* 92:2022–34
- Fischer DC, Siebertz B, van de Leur E et al (1999) Induction of alpha1-antitrypsin synthesis in human articular chondrocytes by interleukin-6-type cytokines: evidence for a local acute-phase response in the joint. *Arthritis Rheum* 42:1936–1945
- Zhang S, Liu J, Dragunow M, Cooper GJ (2003) Fibrillogenic amylin evokes islet beta-cell apoptosis through linked activation of a caspase cascade and JNK1. *J Biol Chem* 278:52810–52819
- Jaikaran ET, Clark A (2001) Islet amyloid and type 2 diabetes: from molecular misfolding to islet pathophysiology. *Biochim Biophys Acta* 1537:179–203
- Eriksson S, Janciauskiene S, Lannfelt L (1995) Alpha 1-antichymotrypsin regulates Alzheimer beta-amyloid peptide fibril formation. *Proc Natl Acad Sci U S A* 92:2313–2317

22. Bottino R, Fernandez LA, Ricordi C et al (1998) Transplantation of allogeneic islets of Langerhans in the rat liver: effects of macrophage depletion on graft survival and microenvironment activation. *Diabetes* 47:316–323
23. Papaccio G, Pedulla M, Ammendola E, Todaro M (2002) Cytokine regulatory effects on alpha-1 proteinase inhibitor expression in NOD mouse islet endothelial cells. *J Cell Biochem* 85:123–130
24. Lou J, Triponez F, Oberholzer J et al (1999) Expression of alpha-1 proteinase inhibitor in human islet microvascular endothelial cells. *Diabetes* 48:1773–1778
25. Ray MB, Zumwalt RE (1986) Identification of alpha-1-proteinase inhibitor-containing cells in pancreatic islets. *Cell Tissue Res* 243:677–680
26. Ray MB, Desmet VJ (1978) Immunohistochemical demonstration of alpha-1-antitrypsin in the islet cells of human pancreas. *Cell Tissue Res* 187:69–77
27. Ricordi C, Lacy PE, Finke EH, Olack BJ, Scharp DW (1988) Automated method for isolation of human pancreatic islets. *Diabetes* 37:413–420
28. Ris F, Hammar E, Bosco D et al (2002) Impact of integrin-matrix matching and inhibition of apoptosis on the survival of purified human beta-cells in vitro. *Diabetologia* 45:841–850
29. Wright PH, Makulu DR, Posey IJ (1968) Guinea pig anti-insulin serum. Adjuvant effect of *H. pertussis* vaccine. *Diabetes* 17:513–516
30. Liou W, Geuze HJ, Slot JW (1996) Improving structural integrity of cryosections for immunogold labeling. *Histochem Cell Biol* 106:41–58
31. Tokuyasu K (1997) Immuno-cytochemistry on ultrathin cryosections. In: Spector, DL, Goodman, RD, Leinwand LA (eds) *Cells, a laboratory manual*. Laboratory Press, Cold Spring Harbor, pp 131.1–131.27
32. Salomon D, Meda P (1986) Heterogeneity and contact-dependent regulation of hormone secretion by individual B cells. *Exp Cell Res* 162:507–520
33. Ray MB, Desmet VJ, Gepts W (1977) Alpha-1-Antitrypsin immunoreactivity in islet cells of adult human pancreas. *Cell Tissue Res* 185:63–68
34. Herrera PL (2000) Adult insulin- and glucagon-producing cells differentiate from two independent cell lineages. *Development* 127:2317–2322
35. Herrera PL (2002) Defining the cell lineages of the islets of Langerhans using transgenic mice. *Int J Dev Biol* 46:97–103
36. Glombik MM, Kromer A, Salm T, Huttner WB, Gerdes HH (1999) The disulfide-bonded loop of chromogranin B mediates membrane binding and directs sorting from the trans-Golgi network to secretory granules. *EMBO J* 18:1059–1070
37. Misumi Y, Oda K, Fujiwara T, Takami N, Tashiro K, Ikehara Y (1991) Functional expression of furin demonstrating its intracellular localization and endoprotease activity for processing of proalbumin and complement pro-C3. *J Biol Chem* 266:16954–16959
38. Rhodes CJ, Brennan SO, Hutton JC (1989) Proalbumin to albumin conversion by a proinsulin processing endopeptidase of insulin secretory granules. *J Biol Chem* 264:14240–14245
39. Ohkubo K, Naito Y, Fujiwara T, Miyazaki J, Ikehara Y, Ono J (2003) Inhibitory effect of the alpha-1-antitrypsin Pittsburgh type-mutant (alpha1-PIM/R) on proinsulin processing in the regulated secretory pathway of the pancreatic beta-cell line MIN6. *Endocr J* 50:9–20
40. Steiner DF (1998) The proprotein convertases. *Curr Opin Chem Biol* 2:31–39
41. Faust D, Raschke K, Hormann S, Milovic V, Stein J (2002) Regulation of alpha1-proteinase inhibitor release by proinflammatory cytokines in human intestinal epithelial cells. *Clin Exp Immunol* 128:279–284
42. Amsterdam A, Sasson R (2002) The anti-inflammatory action of glucocorticoids is mediated by cell type specific regulation of apoptosis. *Mol Cell Endocrinol* 189:1–9
43. Lambillotte C, Gilon P, Henquin JC (1997) Direct glucocorticoid inhibition of insulin secretion: an in vitro study of dexamethasone effects in mouse islets. *J Clin Invest* 99:414–423
44. Corbett JA, Wang JL, Misko TP, Zhao W, Hickey WF, McDaniel ML (1993) Nitric oxide mediates IL-1 beta-induced islet dysfunction and destruction: prevention by dexamethasone. *Autoimmunity* 15:145–153
45. Mauricio D, Andersen HU, Larsen CM, Karlens AE, Mandrup-Poulsen T, Nerup J (1997) Dexamethasone prevents interleukin-1beta-mediated inhibition of rat islet insulin secretion without decreasing nitric oxide production. *Cytokine* 9:563–569
46. Akiyama T, Takasawa S, Nata K et al (2001) Activation of *Reg* gene, a gene for insulin-producing beta-cell regeneration: poly (ADP-ribose) polymerase binds *Reg* promoter and regulates the transcription by autopoly(ADP-ribosylation). *Proc Natl Acad Sci U S A* 98:48–53
47. Fox CJ, Danska JS (1997) IL-4 expression at the onset of islet inflammation predicts nondestructive insulinitis in nonobese diabetic mice. *J Immunol* 158:2414–2424
48. Bolaffi JL, Rodd GG, Wang J, Grodsky GM (1994) Interrelationship of changes in islet nicotine adenine dinucleotide, insulin secretion, and cell viability induced by interleukin-1 beta. *Endocrinology* 134:537–542
49. Delaney CA, Green MH, Lowe JE, Green IC (1993) Endogenous nitric oxide induced by interleukin-1 beta in rat islets of Langerhans and HIT-T15 cells causes significant DNA damage as measured by the 'comet' assay. *FEBS Lett* 333:291–295
50. Mandrup-Poulsen T, Bendtzen K, Nerup J, Dinarello CA, Svenson M, Nielsen JH (1986) Affinity-purified human interleukin 1 is cytotoxic to isolated islets of Langerhans. *Diabetologia* 29:63–67
51. Hohmeier HE, Tran VV, Chen G, Gasa R, Newgard CB (2003) Inflammatory mechanisms in diabetes: lessons from the beta-cell. *Int J Obes Relat Metab Disord* 27(Suppl 3):S12–S16
52. Choi SE, Choi KM, Yoon IH et al (2004) IL-6 protects pancreatic islet beta cells from pro-inflammatory cytokines-induced cell death and functional impairment in vitro and in vivo. *Transpl Immunol* 13:43–53
53. Hall P, Tryon E, Nikolai TF, Roberts RC (1986) Functional activities and nonenzymatic glycosylation of plasma proteinase inhibitors in diabetes. *Clin Chim Acta* 160:55–62
54. Finotti P, Piccoli A, Carraro P (1992) Alteration of plasma proteinase-antiproteinase system in type 1 diabetic patients. Influence of sex and relationship with metabolic control. *Diabetes Res Clin Pract* 18:35–42
55. Sandler M, Gemperli BM, Hanekom C, Kuhn SH (1988) Serum alpha 1-protease inhibitor in diabetes mellitus: reduced concentration and impaired activity. *Diabetes Res Clin Pract* 5:249–255