

CMS 2: 87–106 (2005)

DOI: 10.1007/s10287-004-0028-3

A multivariate FGD technique to improve VaR computation in equity markets

Francesco Audrino, Giovanni Barone-Adesi*

University of Lugano, USI, Institute of Finance, Via G. Buffi 13, 6900 Lugano, Switzerland

Abstract. It is difficult to compute Value-at-Risk (VaR) using multivariate models able to take into account the dependence structure between large numbers of assets and being still computationally feasible. A possible procedure is based on functional gradient descent (FGD) estimation for the volatility matrix in connection with asset historical simulation. Backtest analysis on simulated and real data provides strong empirical evidence of the better predictive ability of the proposed procedure over classical filtered historical simulation, with a resulting significant improvement in the measurement of risk.

1 Introduction

The measurement of market risk (the risk that a financial institution incurs losses on its trading book due to unexpected changes in prices or rates) has assumed a primary importance for regulators and for internal risk control, because of the growth in trading in most financial institutions. One of the most widely used risk measures is Value-at-Risk, or VaR (see [12], for a review of the early literature on VaR). A portfolio's (or an asset's) VaR is commonly defined as the maximum loss that will be incurred on the portfolio with a given level of confidence over a specified holding period, based on the distribution of price changes over a given observation period. Or, in other words, a VaR calculation amounts to a simple quantile estimation of the Profit-and-Loss distribution of a given portfolio over a prescribed holding period.

The main advantage of using VaR as a risk measure is that it is very simple and can also be used to summarize the risk of individual positions. Because of this, it has been adopted for regulatory purposes. More specifically, the BIS has stipulated

* The authors would like to thank Peter Bühlmann and two anonymous referees for some helpful comments. Financial support by the National Centre of Competence in Research "Financial Valuation and Risk Management" (NCCR FINRISK) is gratefully acknowledged.

that the minimum capital requirement for market risk should be based on a 10-day VaR at 99% confidence level.

A lot of different ways have been proposed so far to compute VaR with univariate methods: see, for example, [11] or [20]. In this paper, we study whether the accuracy of VaR predictions for individual positions estimated with univariate techniques can be significantly improved using multivariate methods, which can also take into account the predictive contributions and interactions of other positions belonging to a common market segment. We present some simulations and results for a real data example: the U.S. biotechnological equity market segment.

Although VaR is conceptually a simple measure of risk, computing it in practice using multivariate methods which allow to take into account the possible non-linear dependence structure across different assets in large equity markets can be very difficult, due to the well known curse of dimensionality when estimating high-dimensional conditional covariance matrices. Previous work on multivariate volatility models has been done in [7, 16, 21, 15, 2] and [14] in the framework of GARCH-type models, and in [19, 10] and [1] within the stochastic volatility (state space model) framework. In the GARCH-type framework, only very simple models (see [7, 14]) are feasible in high dimensions without resorting to variance reduction techniques, whereas with stochastic volatility models, only in [10] is presented an example with dimensionality as large as 40 which is still far lower than the number of assets which often occurs in practice.

We present here a non-parametric technique for constructing accurate daily VaR estimates for individual positions which is able to take into account all the possible non-linear dependence structure across different assets and which is still computationally feasible for multivariate problems in large dimensions. Our strategy is based on functional gradient descent (FGD) estimation for the multivariate conditional covariance matrix in connection with historical simulation. FGD is a recent technique from the area of machine learning introduced to solve the classification problem; cf. [22, 8, 18, 17]. The FGD algorithm that we propose is the same as in [3], who have studied the statistical performance of FGD in the financial field. It is very general and can be further adapted to solve other multivariate problems dealing with high-dimensional (volatility) function estimation, such as asset allocation problems, involving the allocation of assets among several stocks whose returns are correlated, or risk management for large global trading portfolios with time-dependent weights.

The main advantage of our technique is its ability to construct reliable and powerful VaR predictions in a high-dimensional multivariate GARCH-type set-up. As we have already said above, it is not possible to use standard multivariate GARCH-type models, such as for example the BEKK models, to estimate the conditional covariance matrix in large dimensions, because we would have to face an intractable model-selection problem and most parameters would have to be set to zero in order to avoid overfitting. Using FGD this problem can be overcome:

this technique can also be used in situations where we deal with more parameters than observations. Choosing reasonable starting functions (for example estimated by a very simple multivariate GARCH-type model), FGD tries to improve, often successfully, those components where the initial predictions are poorest. Clearly, as in [3] has been already shown, we can not expect to learn in all d dimensions when increasing d and keeping a fixed sample size. However, although the gain on average will generally decrease with the number of fitted assets, FGD still improves the worst cases.

Once that FGD yields accurate predictions for the conditional covariance matrix, we can use a model-based bootstrap (see [13]) to generate recursively pathways for future returns. This methodology can also be seen as a multivariate extension of the method proposed and backtested in [4, 5] and [6] based on filtered historical simulation. The main difference here is that we use a multivariate GARCH-type model in connection with FGD for filtering the residuals.

Our strategy contrasts well several critiques made about the use of filtered historical simulation for estimating VaR (see in particular [25]). First of all, our FGD technique allows for the use of cross-terms as predictor variables. This is a reasonable assumption if we consider that assets belonging to a common market segment show some dependence structure and it is conceivable that one asset can be influenced and predicted by past values of some other. This possibility has not been considered in the filtered historical simulation method proposed in [4] and [5], where the volatility of an asset depends only on its own past lagged values and volatilities.

A second critique is related to the assumption of independent identically distributed (i.i.d.) innovations, which implies fixed conditional correlations in a multivariate setting. In our procedure, we only assume constant conditional correlations in a rolling (i.e. not fixed) time-window of about three years of data, using to model the dynamics of the multivariate return series the constant conditional correlation (CCC) model firstly proposed in [7]. Our method can be perhaps further improved by assuming dynamic conditional correlations (see for example [14]), but this is not in the spirit of this paper and it is left to future research.

Using different statistical and economical backtests, we collect empirical evidence of the better predictive power of our multivariate procedure over other univariate techniques, and in particular over the filtered historical simulation method. Through a simulation exercise and a 13-dimensional real data example for individual assets belonging to a common equity market segment, we found that VaR estimates from our technique are more accurate, with a resulting improvement in the measurement of market risk.

The paper is organized as follows. We present and discuss our FGD algorithm in Sect. 2. Section 3 is concerned with the description of the model-based bootstrap method used for the construction of daily VaR estimates. In Sect. 4 we propose two simulation exercises to test the goodness of our multivariate procedure, also in

comparison to the univariate method proposed in [5]. The results of the real data backtest analysis are summarized in Sect. 5. Section 6 concludes the paper.

2 Volatility estimation with Functional Gradient Descent

2.1 Starting point

The multivariate real data of interest are in our case time series of asset prices $\{P_{t,i}; t = 0, 1, \dots, T, i = 1, \dots, d\}$. Their (log)-returns (in percentages) are defined as the change in the logarithms of the individual prices

$$X_{t,i} = 100 \cdot (\log(P_{t,i}) - \log(P_{t-1,i})), \quad t = 1, \dots, n.$$

We assume stationarity of the returns (at least within a suitable time-window). In the empirical investigations of Sect. 5, results are based on a rolling time-window of about three years, which seems to be consistent with the assumption of stationarity (see [24]).

As [3] have already shown, the Functional Gradient Descent (FGD) technique (see [18] or [17]) is a powerful strategy to construct accurate predictions for the multivariate volatility matrix

$$V_t = \text{Cov}_{d \times d}(\mathbf{X}_t | \mathcal{F}_{t-1}), \quad \mathbf{X}_t = (X_{t,1}, \dots, X_{t,d})^T, \quad (1)$$

where \mathcal{F}_{t-1} denotes the information available up to time $t - 1$, i.e. the σ -algebra generated by $\{\mathbf{X}_s; s \leq t - 1\}$. As already mentioned in Sect. 1, the importance of FGD is revealed particularly in large dimensions (for example d in the dozens or hundreds) where predicting the multivariate volatility matrix raises huge challenges because of the well-known curse of dimensionality. In such a case FGD is one of the few feasible non-parametric techniques (if not the only one so far).

Our working model is a generalization of the classical constant conditional correlation (CCC) GARCH model firstly introduced in [7],

$$\mathbf{X}_t = \mu_t + \Sigma_t \mathbf{Z}_t, \quad (2)$$

where we assume the following:

- (A1) (innovations) $\{\mathbf{Z}_t\}_{t \in \mathbb{Z}}$ is a sequence of i.i.d. multivariate innovations with spherical distribution (e.g. multivariate normal) having mean zero and covariance matrix $\text{Cov}(\mathbf{Z}_t) = I_d$. Moreover, \mathbf{Z}_t is independent from $\mathcal{F}_{t-1} = \{\mathbf{X}_s; s \leq t - 1\}$.
- (A2) (CCC construction) The conditional covariance matrix $V_t = \Sigma_t \Sigma_t^T$ is almost surely positive definite for all t . The typical element of V_t is $v_{t,ij} = \rho_{ij}(v_{t,ii}v_{t,jj})^{1/2}$. The parameter $\rho_{ij} = \text{Corr}(X_{t,i}, X_{t,j} | \mathcal{F}_{t-1})$ equals the constant conditional correlation and hence $-1 \leq \rho_{ij} \leq 1$, $\rho_{ii} = 1$.

(A3) (functional form) The conditional variances are functions of the form

$$v_{t,ii} = \sigma_{t,i}^2 = \text{Var}(X_{t,i} | \mathcal{F}_{t-1}) = F_i(\{X_{t-j,k}; j = 1, 2, \dots, k = 1, \dots, d\})$$

where F_i takes values in \mathbb{R}^+ .

(A4) (conditional mean) The conditional mean μ_t is of the form

$$\mu_t = (\mu_{t,1}, \dots, \mu_{t,d})^T = A\mathbf{X}_{t-1}$$

with A a diagonal $d \times d$ matrix (parametric vector AR(1) in mean).

Note that (A2) can be represented in matrix form as

$$V_t = \Sigma_t \Sigma_t^T = D_t R D_t,$$

$$D_t = \text{diag}(\sigma_{t,1}, \dots, \sigma_{t,d}), \quad R = [\rho_{ij}]_{i,j=1}^d.$$

The functional form (A3) allows clearly for cross-terms, since the conditional variances of one series depends on past multivariate observations. This is one of the nice features of such a multivariate GARCH-type model and is motivated from the fact that in reality some instruments can be influenced and better predicted using past information from other risk factors.

2.2 Volatility estimation

FGD estimates the (squared) individual volatility functions $F_i(\cdot)$ in (A3), where $F_i(\cdot) : \mathbb{R}^{pd} \rightarrow \mathbb{R}^+$ is restricted to be a function of the last p lagged multivariate observations, with p finite. The main idea of FGD is to find the estimates for the functions $F_i(\cdot)$ which minimize a suitable loss function λ , under the constraint that the solutions $\hat{F}_i(\cdot)$ are additive expansions of “simple estimates”. These “simple estimates” are given from a statistical procedure \mathcal{S} , called the base learner, which is often constructed from a (constrained or penalized) least squares fitting; common examples of base learners are regression or decision trees, projection pursuit regressors, neural nets and splines. For more details, we remand to [18, 17] and, for a first application in the financial field, [3].

To proceed with the FGD technique, we have therefore to specify a suitable loss function which has to be minimized during the estimation. Assuming multivariate normality of the innovation variables \mathbf{Z}_t in (2), the (multivariate) negative log-likelihood (conditional on the first p observations) is given by

$$- \sum_{t=p+1}^n \log \left((2\pi)^{-d/2} |V_t|^{-1/2} \exp(-\xi_t^T V_t^{-1} \xi_t / 2) \right)$$

$$= \sum_{t=p+1}^n \left(\log |D_t| + \frac{1}{2} (D_t^{-1} \xi_t)^T R^{-1} (D_t^{-1} \xi_t) + d \log(2\pi) / 2 + \log |R| / 2 \right)$$

where $\xi_t = \mathbf{X}_t - \mu_t$, D_t is diagonal with elements $\sqrt{F_i(\mathbf{X}_{t-p}^{t-1})}$. For this reason a natural loss function is

$$\begin{aligned} \lambda_R(\mathbf{Y}, \mathbf{f}) &= \log(\det(D(\mathbf{f}))) + \frac{1}{2} (D(\mathbf{f})^{-1} \mathbf{Y})^T R^{-1} (D(\mathbf{f})^{-1} \mathbf{Y}) \\ D(\mathbf{f}) &= \text{diag}(f_1, \dots, f_d), \end{aligned} \quad (3)$$

since the terms $d \log(2\pi)/2$ and $\log(\det(R))/2$ are constants and may be dropped. As pointed out with the subscript, the loss function λ_R depends on the unknown correlation matrix R . The FGD algorithm will be constructed iteratively by estimating R and using the loss function with the estimated R to get an estimate for all F_i 's.

Estimation of the correlation matrix R can be easily done via empirical moments of residuals. Having (previous) estimates $\hat{\mathbf{F}} = (\hat{F}_1, \dots, \hat{F}_d)$, we build the residuals

$$\hat{\varepsilon}_{t,i} = (X_{t,i} - \hat{\mu}_{t,i}) / \hat{F}_i(\mathbf{X}_{t-1}, \dots)^{1/2}, \quad t = p+1, \dots, n$$

and define

$$\hat{R} = (n-p)^{-1} \sum_{t=p+1}^n \hat{\varepsilon}_t \hat{\varepsilon}_t^T, \quad \hat{\varepsilon}_t = (\hat{\varepsilon}_{t,1}, \dots, \hat{\varepsilon}_{t,d})^T. \quad (4)$$

As the name ‘‘functional gradient descent’’ suggests, we need to calculate the partial derivatives of the loss function λ_R . They are given (in the case of normality of the innovations \mathbf{Z}_t) by

$$\frac{\partial \lambda_R(\mathbf{Y}, \mathbf{f})}{\partial f_i} = \left(\frac{1}{f_i} - \sum_{j=1}^d \frac{\gamma_{ij} y_i y_j}{f_i^{3/2} f_j^{1/2}} \right) / 2, \quad i = 1, \dots, d, \quad (5)$$

where $[\gamma_{ij}]_{i,j=1}^d = R^{-1}$. This will be used when computing negative gradients (see Step 2 in the following FGD algorithm) for every component $i = 1, \dots, d$.

If the assumption of normality of the innovations \mathbf{Z}_t in (2) is violated, the estimates may be consistent but inefficient and this can result in poor performance. As it is shown in the empirical investigations of Sect. 5, an alternative can be to assume a fat-tailed distribution for the innovations (such as for example a scaled t_ν distribution with a fixed number of degrees of freedom ν), that is consistent with the belief that financial (log-) returns are leptokurtic. Another possibility could be to assume a normal inverse gaussian distribution, which seems to work quite well (see [26]).

In analogy to [3], the FGD algorithm for multivariate volatility looks as follows.

FGD algorithm

Step 1 (initialization). Choose the starting function $\hat{F}_{i,0}(\cdot)$ and denote by

$$\hat{F}_{i,0}(t) = \hat{F}_{i,0}(\mathbf{X}_{t-1}, \mathbf{X}_{t-2}, \dots), \quad i = 1, \dots, d.$$

Construct estimates $\hat{\mu}_t$ for the conditional mean from a starting model and compute \hat{R}_0 as in (4) using $\hat{\mathbf{F}}_0$. Set $m = 1$.

For every component $i = 1, \dots, d$, do the following.

Step 2_i (projection of component gradients to base learner). Compute the negative gradient

$$U_{t,i} = - \frac{\partial \lambda_{\hat{R}_{m-1}}(\mathbf{X}_t - \hat{\mu}_t, \mathbf{F})}{\partial F_i} \Big|_{\mathbf{F}=\hat{\mathbf{F}}_{m-1}(t)}, \quad t = p+1, \dots, n.$$

This is explicitly given in (5). Then, fit the negative gradient vector $U_i = (U_{p+1,i}, \dots, U_{n,i})^T$ with a base learner, using always the first p time-lagged predictor variables (i.e. \mathbf{X}_{t-p}^{t-1} is the predictor for $U_{t,i}$)

$$\hat{f}_{m,i}(\cdot) = \mathcal{S}_X(U_i)(\cdot),$$

where $\mathcal{S}_X(U_i)(x)$ denotes the predicted value at x from the base learner \mathcal{S} using the response vector U_i and predictor variables X .

Step 3_i (line search). Perform a one-dimensional optimization for the step-length,

$$\hat{w}_{m,i} = \operatorname{argmin} \sum_{t=p+1}^n \lambda_{\hat{R}_{m-1}}(\mathbf{X}_t - \hat{\mu}_t, \hat{\mathbf{F}}_{m-1}(t) + w \hat{f}_{m,i}(\mathbf{X}_{t-p}^{t-1})).$$

$(\hat{\mathbf{F}}_{m-1}(t) + w \hat{f}_{m,i}(\cdot))$ is defined as the function which is constructed by adding in the i th component only). This can be expressed more explicitly by using (3).

(Note that the line search guarantees that the negative log-likelihood is monotonely decreasing with every iteration.)

Step 4 (up-date). Select the “best component” as

$$i_m^* = \operatorname{argmin}_i \sum_{t=p+1}^n \lambda_{\hat{R}_{m-1}}(\mathbf{X}_t - \hat{\mu}_t, \hat{\mathbf{F}}_{m-1}(t) + \hat{w}_{m,i} \hat{f}_{m,i}(\mathbf{X}_{t-p}^{t-1})).$$

Up-date

$$\hat{\mathbf{F}}_m(\cdot) = \hat{\mathbf{F}}_{m-1}(\cdot) + \hat{w}_{m,i_m^*} \hat{f}_{m,i_m^*}(\cdot).$$

Then, compute the new estimate \hat{R}_m according to (4) using $\hat{\mathbf{F}}_m$.

Step 5 (iteration). Increase m by one and iterate Steps 2–4 until stopping with $m = M$. This produces the FGD estimate

$$\hat{\mathbf{F}}_M(\cdot) = \hat{\mathbf{F}}_0(\cdot) + \sum_{m=1}^M \hat{w}_{m,i_m^*} \hat{f}_{m,i_m^*}(\cdot).$$

The stopping value M is chosen by the following scheme: split the (in-sample) estimation period into two sets, the first of size $0.7 \cdot n$ used as training set and the second of size $0.3 \cdot n$ used as test set (this can also be used when the data are dependent). The optimal value of M is then chosen to optimize the cross-validated log-likelihood.

Remark 1. Initialization in Step 1 is very important here to achieve good estimates. As a starting function, we propose to use the fit from a AR(1)-CCC-GARCH(1,1) model, which is of the form (2) with (A3) specified to

$$F_i(X_{t-1}, X_{t-2}, \dots) = \sigma_{t,i}^2 = \alpha_{0,i} + \alpha_{1,i} \xi_{t-1,i}^2 + \beta_{0,i} \sigma_{t-1,i}^2, \quad i = 1, \dots, d. \quad (6)$$

We construct the estimates by maximum likelihood from the d individual series, ignoring the more general correlation structure in R . This causes some statistical decrease in efficiency, but it gains the advantage that the estimates remain quickly computable in high dimensions d .

Note that the starting estimates $\hat{\mu}_t$ for the conditional mean are kept fixed during the FGD estimation of the volatility functions.

Remark 2. The base learner in Step 2 obviously determines the FGD estimate $\hat{\mathbf{F}}_M(\cdot)$. This should be “weak” (not involving too many parameters to be estimated) enough not to immediately produce an overfitted estimate at the first iteration. The complexity of the FGD estimate $\hat{\mathbf{F}}_M(\cdot)$ is increased by adding further terms with every iteration (see [9]). We choose regression trees as base learners, since particularly in high dimensions, they have the ability to select variables by choosing few of the explanatory variables for prediction. This choice should not be regarded as exclusive: other base learners could be tried out and compared using some form of cross-validation.

As stated above, it is often desirable to make a base learner sufficiently “weak”. A simple but effective way to reduce the complexity of the base learner is via shrinkage towards zero. The up-date in Step 4 of the FGD algorithm is then replaced by

$$\hat{\mathbf{F}}_m(\cdot) = \hat{\mathbf{F}}_{m-1}(\cdot) + \nu \cdot \hat{w}_{m,i_m^*} \hat{f}_{m,i_m^*}(\cdot), \quad 0 < \nu \leq 1. \quad (7)$$

Obviously, this reduces the variance of the base learner by the factor ν^2 .

Remark 3. Stopping in Step 4 is important. It can be viewed as a regularization device which is very effective in complex model fitting. We find empirically that estimating M by the simple 70%-30% cross-validation scheme works well.

Summarizing the above three remarks, the functional form that the individual (squared) volatility functions in (A3) can take in our simulations and real data examples is

$$\mathbf{F}(t) = \mathbf{F}_0(\mathbf{X}_{t-1}, \mathbf{X}_{t-2}, \dots) + \nu \sum_{m=1}^M \sum_{k=1}^L \gamma_{i_m^*, k}^{(m)} I_{[\mathbf{X}_{t-p}^{-1} \in \mathcal{R}_{i_m^*, k}^{(m)}]}, \quad (8)$$

where the starting functions $\mathbf{F}_0(\cdot)$ are given by (6) and the cells $\mathcal{R}_{i_m^*, k}^{(m)}$ are constructed when fitting by least squares a regression tree to the negative gradient vector U (see step 2 of the above FGD Algorithm). This produces partitions $\{\mathcal{R}_{i,1}^{(m)}, \dots, \mathcal{R}_{i,L}^{(m)}\}$ of the predictor space \mathbb{R}^{pd}

$$\cup_{k=1}^L \mathcal{R}_{i,k}^{(m)} = \mathbb{R}^{pd}, \quad \mathcal{R}_{i,k}^{(m)} \cap \mathcal{R}_{i,h}^{(m)} = \emptyset \quad (k \neq h), \quad i = 1, \dots, d.$$

Standard optimal volatility parameters in (8) chosen by the FGD Algorithm are $L \in \{2, 3, 5\}$ (number of end nodes), $\nu \in [0.1, 0.5]$ (shrinkage factor) and $p \in \{1, 2\}$ (number of past lags used as predictors). In general, the estimated optimal location parameters $\gamma_{i_m^*, k}^{(m)}$ are very small (i.e. $|\gamma_{i_m^*, k}^{(m)}| \leq 0.2$), not to have overfitting. Moreover, elements of the constant conditional correlation matrix assumed in (2) are in the most cases higher (in absolute terms) than the ones from a classical filtered historical simulation approach.

Past multivariate returns are used to estimate the optimal partition cells in (8) for the chosen asset i_m^* . We have observed in the real data example of Sect. 5 that most of the predictors chosen by the Algorithm to estimate the optimal partition and therefore the individual (squared) volatility functions *are not* past returns from the same series. For example, we find that in the estimation of our AMEX data *more than 85%* of the optimal predictors are other firms' past lagged returns. This result supports our assumption that there is information included in past lagged observation of other firms that can be used for a better estimation and forecast of risk.

A good feature of such a FGD procedure, particularly in connection with tree-structured base learners (see Remark 2), is that it is a computationally feasible, simple method aiming to improve the initial estimates. FGD traces out a one-dimensional sequence of estimated predictions, which is feasible to optimize via choosing a stopping value M . One can alternatively try to estimate predictions for the volatility matrix V_t in (1) with more complex multivariate GARCH models, but this becomes quickly an intractable model-selection problem in large dimensions d . For example, if we wish to fit a multivariate BEKK model (see [15]) with $d = 10$ individual series, many of the hundreds of parameters would have to be set equal to zero in order to avoid overfitting and more than 10^{73} models would have to be fitted and checked when using a classical strategy for selecting the best subset of non-zero parameters with a model-fitting criterion, such as the Akaike's information criterion (AIC).

A feasible extension, left to future research, of our FGD algorithm could be the use, instead of our working model (2), of a generalization of the dynamic conditional

correlation (DCC) model, recently proposed in [14], using FGD for estimating the individual conditional variances. Note that in our model (2) we assume constant conditional correlations only in a rolling (i.e. not fixed) time-window consisting of the last 800 observations (i.e. at time t , the time-window contains the multivariate observations \mathbf{X}_{t-800}^{t-1} , about three years of data), and not in the full period.

3 VaR estimation

Our VaR estimation is based on a multivariate generalization of the filtered historical simulation procedure proposed in [4] and [5]. Our simulation is based on the combination of multivariate GARCH modelling, using the FGD technique introduced in Sect. 2, and historical asset returns. As we have already explained in Sect. 1, the use of a multivariate GARCH model (2) in connection with FGD as a filter for the estimation of the standardized residuals is needed to remedy the main criticisms made about the use of standard filtered historical simulation for estimating VaR (see [25]). For example, the working model (2) allows clearly for cross-terms and consequently the (squared) volatility function $F_i(\cdot)$ of an asset i can be influenced and predicted by *all* the p -past lagged multivariate observations. This is a realistic assumption if we consider (log-) returns of different assets belonging to a common market segment (in our empirical cases the chemical or the biotechnological one).

The complete methodology stands as follows. In a first step, we filter the multivariate standardized innovations \mathbf{Z}_t using our model (2)

$$\begin{aligned}\mathbf{Z}_t &= (\Sigma_t)^{-1}(\mathbf{X}_t - \mu_t), \\ V_t &= \Sigma_t \Sigma_t^T = D_t R D_t, \quad t = 1, \dots, n,\end{aligned}$$

where the individual (squared) volatility functions $\sigma_{t,i}^2 = F_i(\cdot)$, $i = 1, \dots, d$ are estimated using the FGD technique presented in the algorithm of Sect. 2. Under the assumption (A1), the standardized innovations are i.i.d. and independent from the past.

Now, the historical standardized residuals can be drawn randomly (with replacement) and may be used to generate pathways for future returns. In other words, we use a model-based bootstrap (see [13]): from an i.i.d. resampling of the standardized residuals we recursively generate a time series using the structure and the fitted parameters of the estimated optimal model (2). Thus, we choose randomly dates with corresponding standardized innovations

$$\mathbf{Z}_1^*, \mathbf{Z}_2^*, \dots, \mathbf{Z}_x^*, \quad (9)$$

where x is the time horizon at which we want to estimate the VaR (in general from 1 up to 10 days), and we construct for each asset i pathways for (squared) volatilities and returns from $t + 1$ up to $t + x$ using (2):

$$\widehat{v}_{t+b,ii}^* = (\widehat{\sigma}_{t+b,i}^*)^2 = \widehat{F}_i(\{X_{t+b-s,k}^*; s = 1, 2, \dots, p, k = 1, \dots, d\})$$

$$\widehat{v}_{t+b,ij}^* = \widehat{\rho}_{ij} \sqrt{\widehat{v}_{t+b,ii}^* \widehat{v}_{t+b,jj}^*}$$

$$X_{t+b,i}^* = \mu_{t+b,i}^* + (\widehat{\Sigma}_{t+b}^* \widehat{\mathbf{Z}}_b^*)_i, \quad b = 1, \dots, x, \quad i, j = 1, \dots, d. \quad (10)$$

Note that all quantities denoted by “ $\widehat{\cdot}$ ” use the fitted structure and parameters from the FGD algorithm of Sect. 2.

The “empirical” distribution of simulated, model-based returns at the chosen time horizon x for each asset i , $i = 1, \dots, d$, is obtained replicating the above procedure a large number of times, e.g. 2000. An estimate of the VaR at time horizon x and at level q (q in general $\in \{0.05, 0.01, 0.005\}$) is given by the corresponding q -quantile of the “empirical” returns distribution.

An alternative way to calculate VaR could be the use of extreme value theory (EVT) in connection with the popular peaks over the threshold (POT) method. Such a strategy is well illustrated in [23]. If the assumption made in the FGD estimation of Sect. 2 (i.e. normal or scaled t_ν distributed innovations) is violated, EVT can sometimes yield better VaR predictions than the simpler empirical quantiles.

4 Simulation results

In this section, we present a simulation exercise to study the accuracy of daily VaR predictions estimated with our FGD procedure. We compare our predictions with the ones from the classical filtered historical simulation method in [5] (from now we will denote BAGV) for a normal data generating process. We focus our analysis on the case of 1-day and 10-day VaR estimates for 99% confidence level. This is of particular interest since the BIS capital requirements for market risk are based on VaR at these time horizons and confidence level.

We simulate series of sample size 1500 for 12 assets from the model (2) with standard normally distributed innovations and various individual (squared) volatility functions F_i in (2, A3). One such function is the classical GARCH(1,1) volatility

$$\sigma_{t,i}^2 = F_i(X_{t-1,i}, \sigma_{t-1,i}^2) \text{ where}$$

$$F_i(x, \sigma^2) = \alpha_0 + \alpha_1 x^2 + \beta \sigma^2, \text{ where}$$

$$\alpha_0 \sim \text{Unif}([0, 0.2]), \quad \alpha_1 \sim \text{Unif}([0.05, 0.15]), \quad \beta \sim \text{Unif}([0.8, 0.84]) \quad (11)$$

and $\alpha_0, \alpha_1, \beta$ mutually independent. Another function is from a threshold model

$$\sigma_{t,i}^2 = F_i(X_{t-1,i}, \sigma_{t-1,i}^2) \text{ where}$$

$$F_i(x, \sigma^2) = \begin{cases} \alpha_1 + \alpha_2 x^2, & \text{if } x \leq d_1 = 0, \\ 0.2 + \alpha_3 x^2 + \alpha_4 \sigma^2, & \text{if } x > d_1 = 0 \text{ and } \sigma^2 \leq d_2 = 0.5, \\ 0.8 + \alpha_5 \sigma^2, & \text{if } x > d_1 = 0 \text{ and } \sigma^2 > d_2 = 0.5, \end{cases}$$

where $\alpha_1 \sim \text{Unif}([0, 0.3]), \quad \alpha_2 \sim \text{Unif}([0.4, 0.6]), \quad \alpha_3 \sim \text{Unif}([0.1, 0.3]),$
 $\alpha_4 \sim \text{Unif}([0.6, 0.8]), \quad \alpha_5 \sim \text{Unif}([0.4, 0.6]) \quad (12)$

($\alpha_1, \dots, \alpha_5$ mutually independent). A third function, in which we also allow for one cross-term, is

$$\begin{aligned} \sigma_{t,i}^2 &= F_i(X_{t-1,i}, X_{t-1,j}, \sigma_{t-1,i}^2) \text{ where} \\ F_i(x, y, \sigma^2) &= (\alpha_1 + 0.2 |y| + \alpha_2 x^2) \cdot (0.8 \exp(\alpha_3 |x| |\sigma|)) \\ &\quad + (0.4x^2 + \alpha_4 \sigma^2)^{3/4}, \\ \alpha_1 &\sim \text{Unif}([0.05, 0.15]), \alpha_2 \sim \text{Unif}([0.8, 0.95]), \\ \alpha_3 &\sim \text{Unif}([-1.6, -1.4]), \alpha_4 \sim \text{Unif}([0.4, 0.6]) \end{aligned} \quad (13)$$

($\alpha_1, \dots, \alpha_4$ mutually independent), where the component $j \in \{1, \dots, d\} \setminus i$ is chosen randomly. We choose the simple GARCH(1,1) volatility function (11) for 4 assets, the threshold function (12) for 3 assets and the general function (13) allowing also for cross-terms for the remaining 5 assets. Note that also the coefficients in these functions are randomly chosen. The constant conditional correlation matrix R is chosen to mimic the one of real log-returns. This model is “fairly close” to a CCC-GARCH(1,1) model since more than half of the volatility functions involve only auto-dependence (no dependence on a cross-series in (11) and (12)), a third of them actually being linear GARCH-type where the BAGV approach is correctly specified.

We backtest the accuracy of the VaR predictions on the last 500 observations using a rolling time-window of size 1000 to estimate the parameters for the simulated 12-dimensional system. We report results with the use of $p = 2$ (number of multivariate lagged returns used as predictors), $L = 3$ (number of end-nodes in the regression trees) and $\nu = 0.5$ (shrinkage factor) in the FGD Algorithm.

Analyzing as a first step the individual (squared) volatility functions chosen by our FGD procedure, we find the following. As expected, most of the times (13 out of 14 total FGD iterations, i.e. $M = 14$ in the FGD algorithm) the “best component” chosen by the FGD Algorithm in Step 4 corresponds to individual series generated with volatility functions (12) and (13), i.e. not being of a GARCH(1,1) type. This is not surprising, since the initial starting functions used in the FGD Algorithm (i.e. CCC-GARCH(1,1) estimates) have already the correct volatility structure for individual series generated with volatility function (11) and they can not be improved by FGD. However, in the only exception we get, a truly univariate (i.e. depending only on the same asset’s past returns) volatility function is chosen.

About 60% of the times (8 out of 14 total iterations) FGD optimize individual assets simulated using the volatility function (12). This is also reasonable, since it has already been shown that the starting GARCH(1,1) functions yield poor performance when the true data generating process is of a threshold type. In such cases, our FGD estimates (8) improve the accuracy of the initial predictions. Moreover, most of the times (about 85%) FGD chooses the correct univariate volatility functions. Similarly, when the best components correspond to series simulated using the volatility function (13) also allowing for a (randomly chosen) cross-term, in most

Table 1. Mean absolute errors (MAE) and mean squared errors (MSE) (averaged across assets simulated with each specific volatility function a first time, and averaged across all assets a second time) of VaR predictions at 99% confidence level for 1-day time horizon. Results are computed using the FGD VaR procedure and the BAGV method for a 12-dimensional data set simulated using different individual (squared) volatility functions. Improvements over the classical BAGV method are given between parenthesis. “Assets” denotes the total number of assets for which we choose each specific volatility function to simulate the data.

Function	Assets	Model	MAE	MSE
Function (11)	4	BAGV	0.2471	0.0954
		FGD VaR	0.2409 (2.5%)	0.0907 (4.9%)
Function (12)	3	BAGV	0.5292	0.4560
		FGD VaR	0.3580 (32.4%)	0.2402 (47.3%)
Function (13)	5	BAGV	0.5009	0.4398
		FGD VaR	0.4231 (15.5%)	0.3112 (29.2%)
Global	2	BAGV	0.4234	0.3290
		FGD VaR	0.3461 (18.3%)	0.2199 (33.2%)

cases (about 60%) the final FGD volatility functions have the correct multivariate volatility structure.

To verify the accuracy of VaR predictions using the different approaches, we compute classical mean absolute errors (MAE) and mean squared errors (MSE) (averaged across assets simulated with each specific volatility function in a first time, and averaged across all assets in a second time) by comparing them with “true VaR” predictions based on a full Monte-Carlo simulation. Goodness-of-fit results for 99%-VaR predictions at 1-day time horizon for the BAGV method and our FGD VaR procedure are summarized in Table 1. Similar results also hold for VaR predictions at 10-day time horizon, although with a reduction of the improvements to about one third (i.e. more or less 10% better when using our approach).

Table 1 clearly shows that the accuracy of VaR estimates computed using our FGD procedure is better globally. When considering the average gains for each different individual (squared) volatility function used in the simulation, we can see that, as expected, the largest improvements are realized in the cases where both approaches are mis-specified (functions (12) and (13)), meaning that our FGD procedure, allowing for a general and more flexible functional form given by (8) works better than the BAGV approach. These results confirm the ones found in [3].

To end this simulation exercise, we also computed correlations of the levels and changes of 1-day VaR predictions at 99% confidence level with true VaR’s for the FGD VaR procedure and the BAGV method. We found that average correlation of the VaR estimates with “true VaR” simulated values is significantly higher using our FGD VaR procedure for all type of functions used to model the individual (squared) volatilities (0.6956 vs. 0.6026). Average correlations of changes in the VaR estimates with changes in true VaR are for both approaches lower (0.5367 vs. 0.4165). However, our strategy responds faster to changes in risk than the BAGV method.

5 Backtesting VaR for a real data example

We backtest here the non-parametric procedure for the estimation of VaR proposed in Sects. 2-3 on a real data example. We use parameters $L = 3$, $p = 1$ and $\nu = 0.5$ in the FGD Algorithm. Our tests are essentially the same as in [6].

The analysis is based on two criteria: statistical and economical. The former investigate the frequency and the losses exceeding the VaR predicted by our strategy (violations); the latter examine the implications of these violations (or breaks) and of the structure of the estimated VaR in economic terms.

As VaR's and asset gains and losses are calculated consistently, they can be compared directly to each other, for the corresponding number of days ahead in the holding period. We define the following:

a *violation* (or a *break*) has occurred when ($\text{VaR} > \text{actual asset value}$). (14)

If the model to compute VaR is correct, the actual asset losses should exceed VaR a certain number of times which corresponds to the total number of testing days multiplied by the confidence level used. This means that sometimes the VaR estimated is not sufficient to cover the actual loss. For example, for 95% confidence and 1500 testing days, we should have 75 violations (or breaks). In the following backtests, we stored the risk measures for five different VaR horizons ($x = 1, 2, 3, 5, 10$ days) and three different probability levels ($q \in \{0.95, 0.99, 0.995\}$).

We focus our empirical analysis on the U.S. biotechnological market segment. We consider all 13 assets with enough liquidity belonging to the US AMEX Biotechnology Index with 1100 daily (log-) returns (in percentages): from the Affymetrix Inc., the Amgen Inc., the Biogen Inc., the Cephalon Inc., the Chiron Corporation, the Genzyme Corporation, the Gilead Sciences Inc., the Human Genome Sciences Inc., the IDEC Pharmaceuticals Corporation, the Medimmune Inc., the Millenium Pharmaceuticals Inc., the Protein Design Labs Inc. and the Vertex Pharmaceuticals Inc. The data are from the time period between June 7, 1996 and August 24, 2000. The analysis is made using for prediction a rolling time-window of 800 days and the parameters are re-calculated every 10 days.

We estimate daily VaR for each of these thirteen companies for a backtesting period of 300 days using our FGD algorithm with normal distributed innovations and the standard BAGV method (see [5]). The estimates are then compared to the actual values and the number of violations is recorded.

5.1 Statistical backtests

The first tests are classical overall frequency tests. In Table 2, we show the number of violations of each individual asset for our backtesting period (total of 300 days). The number of violations recorded for the entire backtesting period are reported in each column, where 1-Day up to 10-Day are the 1,2,3,5 and 10-day VaR horizons. We summarize the results recorded for violations at the 99% confidence level.

Table 2. Overall frequency tests: violations for the thirteen assets belonging to the AMEX Biotechnology Index recorded during the backtesting period between July 1, 1999 and August 24, 2000 (for a total of 300 trading days), at the 99% confidence level. Expected are 3 violations. The values marked with an asterisk lead to a rejection of the null-hypothesis of unconditional unbiasedness.

Asset	Model	1-Day	2-day	3-Day	5-Day	10-Day
Affymetrix	BAGV	3	4	4	5	5
	FGD VaR	4	5	5	6	5
Amgen	BAGV	6	6	4	6	9*
	FGD VaR	6	3	6	6	4
Biogen	BAGV	3	3	3	3	3
	FGD VaR	3	3	3	4	3
Cephalon	BAGV	4	5	3	5	3
	FGD VaR	4	6	3	3	4
Chiron	BAGV	8*	7*	7*	7*	6
	FGD VaR	6	5	6	6	3
Genzyme	BAGV	6	8*	6	6	7*
	FGD VaR	6	6	6	5	7*
Gilead	BAGV	5	5	5	4	5
	FGD VaR	5	4	3	3	3
H. Genome	BAGV	5	6	6	7*	6
	FGD VaR	6	6	6	7*	6
IDEC	BAGV	7*	5	7*	5	8*
	FGD VaR	5	5	6	5	7*
Medimmune	BAGV	9*	8*	8*	10*	10*
	FGD VaR	9*	9*	5	7*	8*
Millenium	BAGV	4	4	6	6	6
	FGD VaR	5	6	6	5	6
P. Design	BAGV	5	9*	5	5	5
	FGD VaR	5	6	8*	6	7*
Vertex	BAGV	4	4	5	2	5
	FGD VaR	4	4	3	3	4

Similar results also hold for other confidence levels (i.e. 95% and 99.5%) and are available upon request. The backtest results marked with an asterisk show some significant difference from the following success criterium. Under the hypothesis of unconditional unbiasedness of the VaR estimates, the numbers of violations are binomially distributed around their expected values. A two-standard deviation interval can be used as tolerance for testing the null hypothesis of unconditional unbiasedness.

As we can see, a relevant number of times, especially for high confidence levels, test values are significantly different from our success criterium, i.e we get 9 and 17 (out of 65 tests) rejections of the null hypothesis of unconditional unbiasedness using our FGD VaR strategy and the classical BAGV method, respectively. The two methods seem to underestimate risk, although our procedure is in general better for estimating VaR and lead to a consistently smaller number of rejections.

We try now with some other tests to understand the reason why our procedure and the BAGV method yield for some assets such poor daily VaR predictions. We perform individual firm tests to determine whether violations cluster for one or two companies for which risk may be miss-specified. Under the hypothesis of

Table 3. Time clustering effect: Ljung-Box tests for the number of aggregated violations for 99% confidence level and at 1-day time horizon for thirteen companies belonging to the AMEX Biotechnology Index recorded during the entire backtesting period between July 1, 1999 and August 24, 2000 (Panel A) and during the backtesting period without the dates between March 9, 2000 and March 22, 2000 (Panel B). The values marked with an asterisk lead to a rejection of the null hypothesis of no autocorrelations in the time series of number of aggregated violations at the 1% confidence level or better.

Time clustering effect: A						
	Q_1	Q_2	Q_3	Q_4	Q_5	Q_6
FGD VaR	3.799	6.065*	6.950	22.57*	23.08*	23.49*
BAGV	3.823	5.611	8.116*	20.61*	20.62*	20.88*
Time clustering effect: B						
	Q_1	Q_2	Q_3	Q_4	Q_5	Q_6
FGD VaR	0.143	1.797	3.663	7.131	7.357	7.779
BAGV	0.003	2.199	3.764	7.681	8.602	8.961

randomness the number of violations in the two halves of our backtesting period are independent. Therefore a cross-sectional regression of the violations which each asset reports in the first half on the number of violations recorded in the second half, should have zero slope. The values of these tests are for all confidence levels and at all time horizons not significant and therefore we do not report them here.

We also search for a time clustering effect applying the Ljung-Box test to the time series of the number of aggregated violations across all companies occurring each day. The autocorrelations in this series detect whether days with large number of violations across all assets tend to be followed by other days with large number of violations, pointing to a miss-specification of the time series model of volatility. The resulting values for 1-day VaR at 99% confidence level are summarized in Table 3, Panel A.

Despite the low power of such tests in detecting errors in the VaR estimates (note that the serial correlation in the VaR violations can be very low even if the VaR model is not correctly specified; see [25]), the results reject clearly for both methods the assumption of no autocorrelations in the time series of number of aggregated violations for orders bigger than 3. This result is true for VaR predictions at all confidence levels. A detailed analysis of the time series of aggregate violations show that breaks tend to cluster for a short period in March 2000 (10 business days) in relation with the well-known US technological market crash, where all companies registered several consecutive big losses. This can be the reason why daily VaR predictions using both strategies for these days are poor and the risk tends to be underestimated.

Table 4. Overall frequency tests: violations for the thirteen assets belonging to the AMEX Biotechnology Index recorded during the entire backtesting period without the dates between March 9, 2000 and March 22, 2000 (10 days), at the 99% confidence level. Expected are about 3 violations. The values marked with an asterisk lead to a rejection of the null hypothesis of unconditional unbiasedness of VaR estimates.

Asset	Model	1-Day	2-day	3-Day	5-Day	10-Day
Affymetrix	BAGV	1	2	2	2	3
	FGD VaR	3	4	3	3	4
Amgen	BAGV	5	5	3	3	7*
	FGD VaR	5	2	5	4	4
Biogen	BAGV	3	3	3	2	3
	FGD VaR	3	3	3	3	3
Cephalon	BAGV	3	4	2	2	3
	FGD VaR	3	5	2	1	3
Chiron	BAGV	7*	6	6	7*	5
	FGD VaR	5	4	5	5	2
Genzyme	BAGV	4	6	4	4	7*
	FGD VaR	4	4	4	3	5
Gilead	BAGV	5	5	5	3	3
	FGD VaR	5	4	3	3	3
H. Genome	BAGV	3	4	4	7*	4
	FGD VaR	4	4	4	5	4
IDEC	BAGV	7*	5	7*	5	7*
	FGD VaR	5	5	6	5	5
Medimmune	BAGV	7*	6	6	8*	8*
	FGD VaR	6	6	4	5	4
Millenium	BAGV	3	3	5	4	4
	FGD VaR	3	4	4	3	6
P. Design	BAGV	3	5	2	2	5
	FGD VaR	4	4	4	3	4
Vertex	BAGV	3	4	4	1	3
	FGD VaR	3	3	3	2	3

The values for the same overall frequency tests at the 99% confidence level and clustering tests (1-day time horizon, 99% confidence level) on the violations recorded during the backtesting period without the dates between March 3, 2000 and March 21, 2000 are summarized in Table 4 and Table 3 Panel B, respectively. Similar results also hold for the other confidence levels.

Without this short period in March 2000 there is no significant serial correlation's (order 1 to 6) for any confidence level at 1-day time horizon for the remaining dates. Moreover, the most values of the overall frequency tests are now turned to be not significant. The better potential in predicting daily VaR of our FGD procedure over the BAGV method is clearly shown by the results of Table 4. Particularly when considering daily VaR predictions at long-time horizons (5, 10 days) for all confidence levels, our strategy results to be more attractive for risk management than the standard filtered historical simulation BAGV method. The reason of this result can be explained with the fact that our FGD VaR procedure uses a larger number of predictor variables and the volatility estimates condition on more information than the ones from the BAGV method.

5.2 Economical backtests

To end this section, we focus our backtest analysis on some economical criteria. When focusing the attention a little more on the (absolute) size of the VaR estimates obtained using the different methods, we observed some interesting differences.

The first one is that the BAGV method particularly in the periods of low returns (in absolute values) yields too conservative VaR predictions and tends to overestimate the risk. In contrast, our approach is less conservative, capturing better the passage from stressed, high volatility periods to more stable periods and vice versa. Moreover, the mean absolute difference of consecutive VaR estimates is in the most cases also smaller using our FGD VaR method, indicating that VaR estimates from our procedure change more slowly.

We may interpret the average estimated individual VaR as the average necessary risk capital. Our results show that it is lower for all assets at each time horizon using our procedure. This is a consequence of our assumptions in (2), which allow for cross-terms and the use of more conditioning information than the BAGV method. Information can then flow from one asset to an other causing a better reaction to changes in market conditions and a further reduction of the VaR predictions (in absolute terms) during the periods characterized by small returns. Therefore, our FGD VaR procedure can achieve the same VaR coverage with less capital on average.

A second difference appears clearly when we consider the largest daily violation recorded during our backtesting period. One illustrative example for the largest daily violations at 10-day time horizon and for 99.5% confidence level is shown in Fig. 1.

Using our procedure to estimate daily VaR reduces some peaks with large (aggregate) violations when compared to the ones from the BAGV method. As expected, the period of time with the largest aggregate daily violation is March 2000, where we have seen that violations tend to cluster. In particular, for the 99.5% confidence level and at the 10-day time horizon, the maximal aggregate violation and the mean size of violations are significantly larger (17.21% vs. 14.88% and 3.56% vs. 3.24%) using the BAGV method.

6 Conclusions

We have presented a non-parametric technique to construct daily VaR estimates. Our strategy is based on a multivariate FGD algorithm, which is a method for estimating the conditional covariance matrix in (1), in connection with historical simulation. The use of multivariate GARCH-type models as a filter for historical simulation improves the BAGV method, based on filtered historical simulation (see [4, 5]). For example, our technique allows for cross-terms and the conditional correlation matrix is assumed to be constant only in a rolling (i.e. not fixed) time-window. So far, the use of multivariate GARCH-type models (for example BEKK models) for the estimation and the prediction of the conditional covariance matrix (1) for

Largest Daily Violation for 10-Day VaR at 99.5%




Fig. 1. Backtest analysis for thirteen assets belonging to the AMEX Biotechnology Index: largest daily violations (equally weighted, in %) for a 10-day VaR horizon and at 99.5% confidence level. They are obtained aggregating individual asset violations. Results from our FGD VaR procedure and from the BAGV method are shown by solid lines and dotted lines, respectively. The backtesting period goes from July 1, 1999 to August 24, 2000 (for a total of 300 trading days).

large dimensions was a huge challenge computationally and in most cases led to an intractable model-selection problem. Our FGD algorithm solves these problems: it is computationally feasible in multivariate set-ups with dozens up to hundreds of return series. This is the most attractive feature of FGD.

Our simulation exercise and our tests on a real 13-dimensional data-set belonging to the U.S. biotechnological market segment show that our technique produces accurate and powerful daily VaR estimates, significantly outperforming the VaR predictions from the BAGV method. The results of the backtests show that our multivariate FGD VaR technique, conditioning on more information, has the ability to correct the inaccuracies, which sometimes occur using the BAGV method, yielding better risk estimates. Moreover, we found that BAGV tends to overestimate risk during the periods of low volatility. Summarizing, there is empirical evidence that our procedure can achieve the same VaR coverage with less capital on average than the BAGV method.

When the daily VaR predictions from the standard FGD algorithm are not satisfactory, our procedure can be further improved, for example with a modification of the assumption about the distribution of the innovations in (2) or allowing for

more complex, time-varying conditional correlation dynamics). These extensions are left to future research.

References

- [1] Aguilar, O., West, M. (2000) Bayesian dynamic factor models and variance matrix discounting for portfolio allocation. *Journal of Business and Economic Statistics* **18**: 338–357
- [2] Alexander, C. (2001) A primer on the orthogonal GARCH model. ISMA Centre, Mimeo
- [3] Audrino, F., Bühlmann, P. (2003) Volatility Estimation with Functional Gradient Descent for Very High-Dimensional Financial Time Series. *Journal of Computational Finance* **6**(3): 65–89
- [4] Barone-Adesi, G., Bourgoin, F., Giannopoulos, K. (1998) Don't Look Back. *Risk* **11**: 100–104
- [5] Barone-Adesi, G., Giannopoulos, K., Vosper, L. (1999) VaR Without Correlations for Portfolio of Derivative Securities. *Journal of Futures Markets* **19**: 583–602
- [6] Barone-Adesi, G., Giannopoulos, K., Vosper, L. (2002) Backtesting Derivative Portfolios with FHS. *European Financial Management* **8**: 31–58
- [7] Bollerslev, T. (1990) Modelling the coherence in short-run nominal exchange rates: a multivariate generalized ARCH model. *The Review of Economics and Statistics* **72**: 498–505
- [8] Breiman, L. (1999) Prediction games & arcing algorithms. *Neural Computation* **11**: 1493–1517
- [9] Bühlmann, P., Yu, B. (2003) Boosting with the L_2 -loss: regression and classification. *Journal of the American Statistical Association* (forthcoming)
- [10] Chib, S., Nardari, F., Shephard, N. (1999) Analysis of high dimensional multivariate stochastic volatility models. Preprint, University of Oxford
- [11] Dowd, K. (1998) *Beyond value-at-Risk: The New Science of Risk Management*. Wiley, New York
- [12] Duffie, D., Pan, J. (1997) An Overview of Value at Risk. *Journal of Derivatives* **4**: 7–49
- [13] Efron, B., Tibshirani, R.J. (1993) *An Introduction to the Bootstrap*. Chapman & Hall, London
- [14] Engle, R.F. (2002) Dynamic Conditional Correlation: a simple class of multivariate GARCH models. *Journal of Business and Economic Statistics* (to appear)
- [15] Engle, R.F., Kroner, K.F. (1995) Multivariate simultaneous generalized ARCH. *Econometric Theory* **11**: 122–150
- [16] Engle, R.F., Ng, V.K., Rothschild, M. (1990) Asset pricing with a factor ARCH covariance structure: empirical estimates for treasury bills. *Journal of Econometrics* **45**: 231–238
- [17] Friedman, J.H. (2001) Greedy function approximation: a gradient boosting machine. *Annals of Statistics* **29**, 1189–1232
- [18] Friedman, J.H., Hastie, T., Tibshirani, R. (2000) Additive logistic regression: a statistical view of boosting. *Annals of Statistics* **28**, 337–407 (with discussion)
- [19] Harvey, A.C., Ruiz, E., Shephard, N. (1994) Multivariate stochastic variance models. *Rev. Economic Studies* **61**, 247–264
- [20] Jorion, P. (2001) *Value-at-Risk: the New Benchmark for Controlling Market Risk*. McGraw-Hill, Chicago
- [21] Lin, W.-L. (1992) Alternative estimators for factor GARCH models – a Monte Carlo comparison. *Journal of Applied Econometrics* **7**: 259–279
- [22] Mason, L., Baxter, J., Bartlett, P., Frean, M. (1999) Functional gradient techniques for combining hypotheses. In *Advances in Large Margin Classifiers*. MIT Press, Cambridge, MA
- [23] McNeil, A.J., Frey, R. (2000) Estimation of tail-related risk measures for heteroscedastic financial time series: an extreme value approach. *Journal of Empirical Finance* **7**: 271–300
- [24] Mikosch, T., Starica, C. (1999) Change of structure in financial data, long-range dependence and GARCH modelling. Technical report. University of Groningen
- [25] Pritsker, M. (2001) The Hidden Danger of Historical Simulation. Working Paper (April), University of California, Berkeley
- [26] Venter, J.H., de Jongh, P.L. (2001) Risk Estimation using the Normal Inverse Gaussian Distribution. *Journal of Risk* **4**(2): 1–23