

Int Ophthalmol (2008) 28:439–440  
DOI 10.1007/s10792-007-9157-3

CASE REPORT

# Intraocular lymphoma following a primary testicular lymphoma in remission for 10 years

K. Aliferis · C. C. Chan · G. Donati · E. Baglivo

Received: 27 September 2007 / Accepted: 2 October 2007 / Published online: 6 November 2007  
© Springer Science+Business Media B.V. 2007

**Keywords** Testicular lymphoma · Vitritis · Intraocular lymphoma · Uveitis · IL-6 · IL-10

## Case report

An 83-year-old Caucasian patient was referred with bilateral vitritis treated with topical corticosteroids for 6 months. The patient had had a history of right testicular B-cell lymphoma 10 years ago, which was treated with an orchidectomy followed by six rounds of CHOP-type chemotherapy.

Ophthalmologic examination revealed a best corrected visual acuity of 50/100 in the right eye (RE) and 30/100 in the left eye (LE). Anterior segment examination was normal. Intra-ocular pressure was 15 mmHg in both eyes.

Fundus examination disclosed mild vitritis and multiple pin-like creamy lesions in the deep retina of both eyes. A fluorescein angiography disclosed multiple pinpointed hyperfluorescent lesions (Fig. 1).

An anterior chamber aspiration revealed an interleukin-10 (IL-10) of 314 pg/ml (normal <8 pg/ml) and 150 pg/ml in the aqueous of the LE and RE, respectively.

A diagnostic vitrectomy of the LE confirmed the presence of atypical lymphoid B-cells with an *IgH* gene rearrangement detected using microdissection and PCR. An IL-10 level of 624 pg/ml was measured with ELISA in the vitreous. An intraocular B-cell lymphoma was diagnosed.

The patient was treated with a bilateral local radiotherapy. In the follow-up at 6 months after treatment his best corrected visual acuity was 10/10 in both eyes. A cicatrization of the pin-like lesions was observed in the fundus examination.

He did not develop any local or systemic adverse effects.


## Discussion

Secondary or metastatic intraocular lymphoma is usually located in the uvea without involvement of the neurosensory retina, although a few cases of predominantly retinal disease have been reported [1]. A recurrent, chronic, resistant bilateral uveitis in elderly patients is the typical presentation of primary

---

K. Aliferis · G. Donati · E. Baglivo (✉)  
Clinique d'Ophtalmologie de Genève—Hôpitaux  
Universitaires de Genève, Rue Alcide-Jentzer,  
22, Bethesda, 1205 Geneva, Switzerland  
e-mail: Edoardo.Baglivo@hcuge.ch

C. C. Chan  
Laboratory of Immunology, National Eye Institute,  
Bethesda, MD, USA



**Fig. 1** A fluorescein angiography of the LE discloses multiple pinpointed hyperfluorescent lesions

CNS/intraocular lymphoma. Rarely, a pseudohypopyon may be present [2]. Other unusual manifestations include optic disk swelling [3], serous macular detachments [4] and lymphoma—associated retinopathy [5].

The most common lymphoma subtype involving the eye is B cell lymphoma [1]. Morphologically, it may be difficult to differentiate whether an ocular diffuse large B-cell lymphoma (DLBCL) is a primary or secondary tumor. Coupland et al. [6] have reported that the expression of immunoglobulin transcription factors is different in systemic DLBCL and primary retinal and CNS lymphoma. Clonal analysis studies [7, 8] with sequencing of the polymerase chain reaction products can indicate if the clones have originated from the same tumor or not.

Up to now there have been seven documented cases of primary testicular lymphoma metastases to the eye [9]. Both the testis and eye, as well as the

brain, are immunoprivileged organs, with strong blood-tissue barriers and altered immune response [10]. The presence of a bilateral vitritis in a patient with a history of a testicular lymphoma should evoke a secondary intraocular involvement. A diagnostic vitrectomy using PCR is indicated.

## References

1. Parikh AH, Sameer HK, Wright JD Jr, Kean TO (2005) Systemic non-Hodgkin's lymphoma simulating primary intraocular lymphoma. *Am J Ophthalmol* 139:573–574
2. Pilon AF, Rhee P, Messner LV (2005) Bilateral, persistent serous macular detachments with Waldenstrom's macroglobulinemia. *Optom Vis Sci* 82:573–578
3. Pinna A, Dore S, Dore F et al (2003) Bilateral optic disc swelling as the presenting sign of Waldenstrom's macroglobulinemia. *Acta Ophthalmol Scand* 81:413–415
4. Sen NH, Chan CC, Caruso RC et al (2004) Waldenstrom's macroglobulinemia-associated retinopathy. *Ophthalmology* 111:535–539
5. Tranos PG, Andreou PG, Wickremasinghe SS, Brazier JD (2002) Pseudo-hypopyon as a feature of multiple myeloma. *Arch Ophthalmol* 120:87–88
6. Coupland SE, Damato B (2006) Lymphoma involving the eye and the ocular adnexa. *Curr Opin Ophthalmol* 17: 523–531
7. Coupland SE, Hummel M, Stein H (2005) Molecular analysis of immunoglobulin genes in primary intraocular lymphoma. *Invest Ophthalmol Vis Sci* 46:3507–3514
8. Coupland SE, Willerding G, Jahnke K et al (2005) Demonstration of identical clonal derivation in a case of ocular lymphoma. *Br J Ophthalmol* 89:238–239
9. Wallace DJ, Altemare C, Shen DF, Desmet M, Buggage RR, Nussenblatt RB, Chan CC (2006) Primary testicular and intraocular lymphomas: two case reports and a review of the literature. *Surv Ophthalmol* 51(1):41–50
10. Filippini A, Riccioli A, Padula F et al (2001) Control and impairment of immune privilege in the testis and in semen. *Hum Reprod Update* 7:444–449