

SHORT REPORT

Walter Knirsch · Nikolaus A. Haas
Martin A. G. Lewin · Ingo Dähnert · Deniz Kececioglu
Felix Berger · Frank Uhlemann

Percutaneous closure of patent ductus arteriosus in small infants of less than 8 kg body weight using different devices

Received: 8 October 2003 / Accepted: 13 May 2004 / Published online: 28 July 2004
© Springer-Verlag 2004

Therapeutic closure of patent ductus arteriosus (PDA) may become necessary in small infants with congestive heart failure refractory to medical management, regardless of the age or weight of the patients. Percutaneous closure of PDA offers an alternative to surgical treatment [3,4]. The aim of this retrospective, multicentre observational trial was to evaluate the immediate results and mid-term clinical and echocardiographic follow-up of percutaneous closure of PDA in young infants weighing less than 8 kg. Four centres were involved.

Percutaneous PDA closure was performed in 25 patients (16 female, 9 male) with a body weight of less than 8 kg (mean age 9.2 months, mean body weight 6.3 kg). The mean dimensions of the PDA were 5.9 mm at the aortic end and 2.7 mm at the pulmonary end with a mean of 2.2 mm at the narrowest part. Mean pulmonary artery pressure was 22.6 mm Hg, Qp:Qs = 2.2 to 1 and Rp:Rs = 0.13 to 1. Two patients had an additional atrial septal defect and one patient had an aorto-pulmonary collateral, which were closed during the same cardiac catheterisation (Table 1). Mean fluoroscopy time was 15.0 min. For a tubular shaped PDA, the preferred device was the pfm Ductocclud coil (pfm-AG, Cologne, Germany) or Cook detachable coils (Cook

Inc., Bloomington, IN, USA). For a fenestrated PDA, the Amplatzer Duct Occluder (AGA Medical Corp., Golden Valley, MN, USA) was used. The PDA was closed according to standardised procedures [5].

The Amplatzer Duct Occluder (ADO) was used in 17 patients, Cook detachable coils in six patients, and the pfm Ductocclud coil in two patients (Table 1). In two patients (one with a body weight of 3.3 kg), percutaneous PDA closure failed due to protrusion of the device into the aorta or left pulmonary artery; these infants successfully underwent surgical PDA closure (Table 1). Complete closure was achieved in all patients: immediate occlusion in 13 patients, the following day in four patients, and after mid-term follow-up (6 months) in six patients. In one patient, the Cook detachable coil migrated to the right pulmonary artery 1 day after the intervention; the coil was retrieved and the PDA was closed with an ADO (Fig. 1). In all patients, symptoms of heart failure resolved and medical treatment could be terminated. At mid-term follow-up (mean 18.8 months) Doppler echocardiography revealed mild flow velocity acceleration in the left pulmonary artery in two patients and in the descending aorta in one patient. Special attention was given to monitoring femoral vessel patency after catheterisation. There were no other complications (haematoma, femoral artery thrombosis, haemolysis, infection, haemorrhage or recanalisation).

The rate of percutaneous closure of PDA was comparable to other studies in small children of less than 10 kg body weight described by Aydogan et al. [1] (94%) or by Dalvi et al. [2] (90%), although we used other device types, depending on the morphology of the PDA.

We conclude that percutaneous closure of PDA is a feasible, safe and effective method in infants of between 4 and 8 kg body weight.

W. Knirsch (✉) · F. Berger
Department of Paediatric Cardiology,
University Children's Hospital, Steinwiesstrasse 75,
8032 Zürich, Switzerland
E-mail: walter.knirsch@kispi.unizh.ch
Tel.: +41-1-2667617
Fax: +41-1-2667981

N. A. Haas · M. A. G. Lewin · F. Uhlemann
Department of Paediatric Cardiology and Paediatric Intensive
Care, Olgahospital, Stuttgart, Germany

D. Kececioglu
Department of Paediatric Cardiology,
University Children's Hospital, Freiburg,
Germany

I. Dähnert
Department of Paediatric Cardiology, Heart Centre,
University of Leipzig, Leipzig, Germany

References

1. Aydogan Ü (2002) Arterial duct closure with detachable coils: application in the small child. *Asian Cardiovasc Thorac Ann* 10: 124–128

Table 1 Overview of the 25 patients with a body weight of less than 8 kg with percutaneous PDA closure using different devices

| Patient | Sex | Age (years) | Body weight (kg) | Venous sheath | Arterial sheath | Devices | X-ray time (mins) | Occlusion | Special aspects |
|---------|-----|-------------|------------------|---------------|-----------------|---|-------------------|----------------|--|
| 1 | F | 1.43 | 7.83 | 5 | 4 | Amplatzer Duct Occluder | 29.8 | During 48 h | Aorto-pulmonary collateral closure during same procedure |
| 2 | Ff | 0.19 | 4.20 | 5 | 4 | Amplatzer Duct Occluder | 14.3 | Immediately | |
| 3 | M | 0.91 | 6.74 | 5 | 4 | Pfm Ductocclud coil | 10.8 | After 6 months | |
| 4 | Ff | 1.43 | 7.90 | 5 | 5 | Cook detachable coil | 17.5 | Immediately | |
| 5 | Ff | 1.06 | 8.00 | 6 | 4 | Amplatzer Duct Occluder | 13.0 | Immediately | |
| 6 | M | 0.14 | 4.70 | 5 | 4 | Amplatzer Duct Occluder | 11.8 | Immediately | |
| 7 | M | 0.76 | 7.50 | 5 | 4 | Amplatzer Duct Occluder | 14.9 | Immediately | |
| 8 | F | 0.45 | 6.90 | 4 | 4 | Pfm Ductocclud coil | 18.6 | Immediately | |
| 9 | M | 0.28 | 3.33 | 6 | 3 | Amplatzer Duct Occluder; pfm Ductocclud coil | 10.4 | Not possible | |
| 10 | F | 1.91 | 7.20 | 6 | 4 | Amplatzer Duct Occluder | 14.0 | During 48 h | |
| 11 | F | 0.69 | 7.60 | 6 | 4 | Amplatzer Duct Occluder | 7.5 | After 6 months | |
| 12 | F | 0.56 | 5.80 | 6 | 4 | Amplatzer Duct Occluder | 20.2 | During 48 h | Left pulmonary artery flow velocity < 2 m/s |
| 13 | F | 0.57 | 5.55 | 6 | 4 | Amplatzer Duct Occluder | 13.1 | After 6 months | Left pulmonary artery flow velocity < 2 m/s |
| 14 | M | 0.36 | 3.50 | 6 | 4 | Cook detachable coil | 10.1 | During 48 h | |
| 15 | F | 1.65 | 8.00 | - | 4 | Cook detachable coil | 4.9 | After 6 months | |
| 16 | F | 1.13 | 8.00 | 6 | 4 | Amplatzer Duct Occluder | 20.3 | Immediately | |
| 17 | F | 0.88 | 5.30 | 6 | 4 | Amplatzer Duct Occluder | 13.2 | Immediately | Atrial septal defect closure during same procedure |
| 18 | M | 0.57 | 7.60 | 6 | 4 | Amplatzer Duct Occluder | 9.2 | Immediately | Reduced arterial perfusion for 72 h |
| 19 | F | 0.56 | 7.10 | 6 | 4 | Amplatzer Duct Occluder | 9.4 | Immediately | |
| 20 | F | 0.61 | 6.80 | 6 | 4 | Cook detachable coil; Amplatzer Duct Occluder | 32 | Immediately | Embolisation |
| 21 | F | 0.76 | 6.90 | 6 | 4 | Amplatzer Duct Occluder | 20.5 | Immediately | Descending aorta flow velocity acceleration 2.5 m/s |
| 22 | M | 1.43 | 6.40 | 6 | 4 | Amplatzer Duct Occluder | 7.8 | Immediately | |
| 23 | M | 0.27 | 4.00 | 6 | - | Cook detachable coil | 21.4 | After 6 months | Atrial septal defect closure during same procedure |
| 24 | M | 0.11 | 4.10 | 4 | - | Cook detachable coil | 19.3 | After 6 months | Right-sided PDA |
| 25 | F | 0.54 | 6.20 | 6 | 4 | Amplatzer Duct Occluder | 11.3 | Not possible | |




Fig. 1 Posterior-anterior chest X-ray film showing the embolised Cook detachable coil in the right pulmonary artery. A 6 F long sheath was introduced into the right pulmonary artery and the Cook detachable coil was retrieved into the sheath using a snare catheter. Afterwards the sheath was used to implant the ADO

2. Dalvi B, Nabar A, Goyal V, Naik A, Kulkarni H, Ramakanthan R (1998) Transcatheter closure of patent ductus arteriosus in children weighing <10 kg with Gianturco coils using the balloon occlusion technique. *Cathet Cardiovasc Diagn* 44: 303–308

3. Masura J, Walsh KP, Thanopoulous B, Chan C, Bass J, Goussous Y, Gavora P, Hijazi ZM (1998) Catheter closure of moderate-to-large-sized patent ductus arteriosus using the new Amplatzer duct occluder: immediate- and short-term results. *J Am Coll Cardiol* 31: 878–882
4. O'Donnell C, Neutze JM, Skinner JR, Wilson NJ (2001) Transcatheter patent ductus arteriosus occlusion: evolution of techniques and results from the 1990s. *J Paediatr Child Health* 37: 451–455
5. Rosenthal E, Qureshi SA, Reidy J, Baker EJ, Tynan M (1996) Evolving use of embolisation coils for occlusion of the arterial duct. *Heart* 76: 525–530