

# Dephasing in the Semiclassical Limit is System-Dependent<sup>¶</sup>

C. Petitjean<sup>a</sup>, P. Jacquod<sup>b</sup>, and R. S. Whitney<sup>c</sup>

<sup>a</sup> *Département de Physique Théorique, Université de Genève, CH-1211 Genève 4, Switzerland*

<sup>b</sup> *Physics Department, University of Arizona, Tucson, AZ 85721, USA*

<sup>c</sup> *Institut Laue–Langevin, Grenoble, 38042 France*

Received October 8, 2007

Dephasing in open quantum chaotic systems has been investigated in the limit of large system sizes to the Fermi wavelength ratio,  $L/\lambda_F \gg 1$ . The weak localization correction  $g^{wl}$  to the conductance for a quantum dot coupled to (i) an external closed dot and (ii) a dephasing voltage probe is calculated in the semiclassical approximation. In addition to the universal algebraic suppression  $g^{wl} \propto (1 + \tau_D/\tau_\phi)^{-1}$  with the dwell time  $\tau_D$  through the cavity and the dephasing rate  $\tau_\phi^{-1}$ , we find an exponential suppression of weak localization by a factor of  $\propto \exp[-\tilde{\tau}/\tau_\phi]$ , where  $\tilde{\tau}$  is the system-dependent parameter. In the dephasing probe model,  $\tilde{\tau}$  coincides with the Ehrenfest time,  $\tilde{\tau} \propto \ln[L/\lambda_F]$ , for both perfectly and partially transparent dot-lead couplings. In contrast, when dephasing occurs due to the coupling to an external dot,  $\tilde{\tau} \propto \ln[L/\xi]$  depends on the correlation length  $\xi$  of the coupling potential instead of  $\lambda_F$ .

PACS numbers: 03.65.Yz, 05.45.Mt, 73.23.-b, 74.40.+k

DOI: 10.1134/S0021364007220079

## INTRODUCTION

Electronic transport in mesoscopic systems exhibits a range of quantum coherent effects such as weak localization, universal conductance fluctuations, and Aharonov–Bohm effects [1, 2]. Being intermediate in size between micro- and macroscopic systems, these systems are ideal playgrounds to investigate the quantum-to-classical transition from a microscopic coherent world, where quantum interference effects prevail, to a macroscopic classical world [3]. Indeed, the disappearance of quantum coherence in mesoscopic systems as dephasing processes set in has been the subject of intensive theoretical [4–7] and experimental [8–10] studies. When the temperature is sufficiently low, it is accepted that the dominant processes of dephasing are electronic interactions. In disordered systems, dephasing due to electron–electron interactions is known to be well modeled by the classical noise potential [4], which gives an algebraic suppression of the weak localization correction to conductance through a diffusive quantum dot:

$$g^{wl} = g_0^{wl}/(1 + \tau_D/\tau_\phi). \quad (1)$$


Here,  $g_0^{wl}$  is the weak localization correction (in units of  $2e^2/h$ ) without dephasing, the dephasing time  $\tau_\phi$  is given by the noise power, and  $\tau_D$  is the electronic dwell time in the dot. Equation (1) is insensitive to most noise-spectrum details and holds for other noise

sources such as electron–phonon interactions or external microwave fields.

Other, mostly phenomenological models of dephasing have been proposed to study dephasing in ballistic systems [5–7], the most popular of which is perhaps the dephasing lead model [5, 6]. A cavity is connected to two external L (left) and R (right) leads of widths  $W_L$ ,  $W_R$ . A third lead of width  $W_3$  is connected to the system via a tunnel-barrier of transparency  $\rho$ . A voltage is applied to the third lead to ensure that no current flows through it on average. A random matrix theory (RMT) treatment of the dephasing lead model leads to Eq. (1) with  $\tau_D = \tau_0 L/(W_L + W_R)$  and  $\tau_\phi = \tau_0 L/\rho W_3$ , in term of the dot’s time of flight  $\tau_0$  [6, 11]. Thus, it is commonly assumed that dephasing is system-independent. The dephasing lead model is often used phenomenologically in contexts where the source of dephasing is unknown.

Our purpose in this article is to revisit dephasing in open chaotic ballistic systems with a focus on whether dephasing remains system-independent in the semiclassical limit of large ratio  $L/\lambda_F$  of the system size to the Fermi wavelength. This regime sees the emergence of a finite Ehrenfest time scale,  $\tau_E^{cl} = \lambda^{-1} \ln[L/\lambda_F]$  ( $\lambda$  is the Lyapunov exponent), in which case dephasing can lead to an exponential suppression of weak localization,  $\propto \exp[-\tau_E^{cl}/\tau_\phi]/(1 + \tau_D/\tau_\phi)$  [12]. Subsequent numerical investigations of the dephasing lead model support this prediction [13]. Here, we analytically investigate

<sup>¶</sup> The text was submitted by the authors in English.



**Fig. 1.** (Left panel a) Scheme of the system-environment model. The system is an open quantum dot that is coupled to an environment in the shape of a second, closed quantum dot. (Right panel b) Scheme of the dephasing lead model.

two different models of dephasing and show that the suppression of weak localization is strongly system-dependent. First, we construct a new formalism that incorporates the coupling to external degrees of freedom into the scattering approach to transport. This approach is illustrated by a semiclassical calculation of weak localization in the case of an environment modeled by a capacitively coupled, closed quantum dot. We restrict ourselves to the regime of pure dephasing, where the environment does not alter the classical dynamics of the system. Second, we provide the first semiclassical treatment of transport in the dephasing lead model. We show that, in both cases, the weak localization correction to the conductance is

$$g^{wl} = g_0^{wl} \exp[-\tilde{\tau}/\tau_\phi]/(1 + \tau_D/\tau_\phi), \quad (2)$$

where  $g_0^{wl}$  is the finite- $\tau_E^{cl}$  correction in absence of dephasing. The time scale  $\tilde{\tau}$  is system-dependent. For the dephasing lead model,  $\tilde{\tau} = \tau_E^{cl} + (1 - \rho)\tau_E^{op}$  in terms of the transparency  $\rho$  of the contacts to the leads, and the open system Ehrenfest time  $\tau_E^{op} = \lambda^{-1} \ln[W^2/\lambda_F L]$ . This analytic result fits the numerics of [13], and (up to logarithmic corrections) is in agreement with [12]. Yet, for the system-environment model,  $\tilde{\tau} = \lambda^{-1} \ln[(L/\xi)^2]$  depends on the correlation length  $\xi$  of the interdot coupling potential. We thus conclude that dephasing in the semiclassical limit is system-dependent.

### TRANSPORT THEORY FOR A SYSTEM-ENVIRONMENT MODEL

In the standard theory of decoherence, one starts with the total density matrix  $\eta_{tot}$  including both system and environment degrees of freedom [3]. The time-evolution of  $\eta_{tot}$  is unitary. The observed properties of the system alone are given by the reduced density matrix  $\eta_{sys}$  obtained by tracing  $\eta_{tot}$  over the environment degrees of freedom. This is probability conserving,  $\text{Tr}\eta_{sys} = 1$ , but renders the time-evolution of  $\eta_{sys}$  non-unitary. The decoherence time is inferred from the decay rate of its off-diagonal matrix elements [3]. We

generalize this approach to the scattering theory of transport.

To this end, we consider two coupled chaotic cavities as sketched in Fig. 1a. Few-electron double-dot systems similar to the one considered here have recently been the focus of intense experimental efforts [14]. One of them (the system) is an open quantum dot connected to two external leads. The other one (the environment) is a closed quantum dot, which we model using RMT. The two dots are chaotic and capacitively coupled. In particular, they do not exchange particles. We require that  $\lambda_F \ll W_{L,R} \ll L$ , so that the number of transport channels satisfies  $1 \ll N_{L,R} \ll L/\lambda_F$  and the chaotic dynamics inside the dot have enough time to develop,  $\lambda\tau_D \gg 1$ . Electrons in the leads do not interact with the second dot. Inside each cavity, the dynamics are generated by chaotic Hamiltonians  $H_{sys}$  and  $H_{env}$ . We only specify that the capacitive coupling potential,  $\mathcal{U}$ , is smooth and has magnitude  $\mathcal{U}$  and correlation length  $\xi$ .

The environment coupling can be straightforwardly included in the scattering approach by writing the scattering matrix,  $\mathbb{S}$ , as an integral over time-evolution operators. We then use a bipartite semiclassical propagator to write the matrix elements of  $\mathbb{S}$  for the given initial and final environment positions,  $(\mathbf{q}_0, \mathbf{q})$ , as

$$\begin{aligned} \mathbb{S}_{mn}(\mathbf{q}_0, \mathbf{q}) &= (2\pi)^{-1} \int_0^\infty dt \int_L dy_0 \int_R d\mathbf{y} \langle m|\mathbf{y} \rangle \langle \mathbf{y}_0|n \rangle \\ &\times \sum_{\gamma, \Gamma} (C_\gamma C_\Gamma)^{1/2} \exp[i\{S_\gamma + S_\Gamma + \mathcal{S}_{\gamma, \Gamma}\}]. \end{aligned} \quad (3)$$

This is a double sum over classical paths, labeled  $\gamma$  for the system and  $\Gamma$  for the environment. For pure dephasing, the classical path  $\gamma(\Gamma)$  connecting  $\mathbf{y}_0(\mathbf{q}_0)$  to  $\mathbf{y}(\mathbf{q})$  in the time  $t$  is solely determined by  $H_{sys}$  ( $H_{env}$ ). The prefactor  $C_\gamma C_\Gamma$  is the inverse determinant of the stability matrix, and the exponent contains the noninteracting action integrals,  $S_\gamma$ ,  $S_\Gamma$ , accumulated along  $\gamma$  and  $\Gamma$ , and the interaction term,  $\mathcal{S}_{\gamma, \Gamma} = \int_0^t d\tau \mathcal{U}[\mathbf{y}_\gamma(\tau), \mathbf{q}_\Gamma(\tau)]$ .

Since we assume that the particles in the leads do not interact with the second cavity, we can write the initial total density matrix as  $\eta^{(n)} = \eta_{sys}^{(n)} \otimes \eta_{env}$ , with  $\eta_{sys}^{(n)} = |n\rangle\langle n|$ ,  $n = 1, 2, \dots, N_L$ . We take  $\eta_{env}$  as a random matrix, though our approach is not restricted to that particular choice. We define the conductance matrix as the following trace over the environment degrees of freedom:

$$g_{mn}^{(r)} = \langle m|\text{Tr}_{env}[\mathbb{S}\eta^{(n)}\mathbb{S}^\dagger]|m \rangle. \quad (4)$$

The conductance is then given by  $g = \sum_{m,n} g_{mn}^{(r)}$ . This construction is current conserving; however, the envi-

ronment-coupling generates decoherence and the suppression of coherent contributions to transport. To see this, we now calculate the conductance to the leading order in the weak localization correction.

We insert Eq. (3) into Eq. (4), perform the sum over channel indices with the semiclassical approximation  $\sum_m^{N_L} \langle \mathbf{y}_0 | n \rangle \langle n | \mathbf{y}'_0 \rangle \approx \delta(\mathbf{y}'_0 - \mathbf{y}_0)$  [15], and use the RMT result  $\overline{\langle \mathbf{q}_0 | \eta_{\text{env}} | \mathbf{q}'_0 \rangle} \approx \Omega_{\text{env}}^{-1} \delta(\mathbf{q}'_0 - \mathbf{q}_0)$ , where  $\Omega_{\text{env}}$  is the environment volume [16]. The conductance then reads

$$g = (4\pi^2 \Omega_{\text{env}})^{-1} \int_0^\infty dt dt' \int_{\Omega_{\text{env}}} d\mathbf{q}_0 d\mathbf{q} \int_L d\mathbf{y}_0 \int_R d\mathbf{y} \quad (5)$$


$$\times \sum_{\gamma, \Gamma; \gamma', \Gamma'} (C_\gamma C_\Gamma C_{\gamma'} C_{\Gamma'})^{1/2} e^{i(\Phi_{\text{sys}} + \Phi_{\text{env}} + \Phi_{\mathcal{U}})}.$$

This is a quadruple sum over classical paths of the system ( $\gamma$  and  $\gamma'$ , going from  $\mathbf{y}_0$  to  $\mathbf{y}$ ) and the environment ( $\Gamma$  and  $\Gamma'$ , going from  $\mathbf{q}_0$  to  $\mathbf{q}$ ), with action phases  $\Phi_{\text{sys}} = S_\gamma(\mathbf{y}_0, \mathbf{y}; t) - S_\gamma(\mathbf{y}_0, \mathbf{y}; t')$ ,  $\Phi_{\text{env}} = S_\Gamma(\mathbf{q}_0, \mathbf{q}; t) - S_\Gamma(\mathbf{q}_0, \mathbf{q}; t')$  and  $\Phi_{\mathcal{U}} = \mathcal{P}_{\gamma, \Gamma}(\mathbf{y}_0, \mathbf{y}; \mathbf{q}_0, \mathbf{q}; t) - \mathcal{P}_{\gamma, \Gamma}(\mathbf{y}_0, \mathbf{y}; \mathbf{q}_0, \mathbf{q}; t')$ . We are interested in the conductance averaged over energy variations, and, hence, look for contributions to Eq. (5) with stationary  $\Phi_{\text{sys}}$ ,  $\Phi_{\text{env}}$ . The first such contributions are the diagonal ones with  $\gamma = \gamma'$  and  $\Gamma = \Gamma'$ , for which  $\Phi_{\mathcal{U}} = 0$ . They are  $\mathcal{U}$ -independent and give the classical, Drude conductance:  $g^{\text{D}} = N_L N_R / (N_L + N_R)$ . The leading order correction to this comes from weak-localization paths  $\gamma$  and  $\gamma'$  [12, 15, 17, 18] (see Fig. 2), with  $\Gamma' = \Gamma$  for the environment paths [19]. In the absence of dephasing, these contributions accumulate the phase difference  $\delta\Phi_{\text{sys}}$ . Semiclassically, these contributions give  $g_0^{\text{wl}} = -\exp[-\tau_E^{\text{cl}}/\tau_D] N_L N_R / (N_L + N_R)^2$  [15, 17, 18].

In the presence of an environment, each weak localization pair of paths accumulates an additional action phase difference  $\delta\Phi_{\mathcal{U}}$ , which is averaged over. Dephasing occurs mostly in the loop, when the paths are more than the correlation length  $\xi$  apart (see Fig. 2). Thus, we can define  $T_\xi = \lambda^{-1} \ln[(\xi/\epsilon L)^2]$  as twice the time between the encounter and the start of dephasing. If  $\xi < \epsilon L$ , dephasing starts before the paths reach the encounter,  $T_\xi < 0$  [20]. Using the central limit theorem and assuming a fast decaying interaction correlator,  $\langle \mathcal{U}[\mathbf{y}_\gamma(\tau), \mathbf{q}_\Gamma(\tau)] \mathcal{U}[\mathbf{y}_{\gamma'}(\tau'), \mathbf{q}_{\Gamma'}(\tau')] \rangle_\Gamma \propto \tau_\phi^{-1} \exp[-\{\mathbf{y}_\gamma(\tau) - \mathbf{y}_{\gamma'}(\tau')\}^2 / \xi^2]$ , the average phase difference due to  $\mathcal{U}$  reads

$$\langle e^{i\delta\Phi_{\mathcal{U}}} \rangle = e^{-\frac{1}{2} \langle \delta\Phi_{\mathcal{U}}^2 \rangle} = \exp[-(t_2 - t_1 - T_\xi) / \tau_\phi], \quad (6)$$

where  $t_1$  ( $t_2$ ) gives the start (end) of the loop. The derivation then proceeds as for  $\mathcal{U} = 0$  [15], except that, during the time  $(t_2 - t_1 - T_\xi)$ , the dwell time is effectively divided by  $[1 + \tau_D/\tau_\phi]$ , so the  $(t_2 - t_1)$ -integral generates



**Fig. 2.** (Color online) Semiclassical contribution to weak localization for the system-environment model. The paths are paired everywhere except at the encounter. There, one crosses itself at the angle  $\epsilon$ , while the other does not (going the opposite way around the loop). Here, we show  $\xi > \epsilon L$ , so the dephasing (dotted path segment) starts in the loop ( $T_\xi > 0$ ).


an extra prefactor of  $[1 + \tau_D/\tau_\phi]^{-1} \exp[-\tau_\xi/\tau_\phi]$ , where  $\tau_\xi = \lambda^{-1} \ln[(L/\xi)^2]$ . Thus, the weak localization correction is as in Eq. (2) with  $\tilde{\tau}$  given by  $\tau_\xi$ . In contrast to [12], the exponent depends on  $\xi$  not  $\lambda_F$ .

A calculation of coherent-backscattering with dephasing to be presented elsewhere enables us to show that our approach is probability- and, thus, current-conserving. We also point out that (for  $\tau_\phi \sim \tau_D$ ) one can ignore the modifications of the classical paths due to the coupling to the environment as long as  $\xi \gg [\lambda_F L / \lambda \tau_D]^{1/2}$ . Thus, our method is applicable for  $\xi$  smaller (as well as larger) than the encounter size  $[\lambda_F L]^{1/2}$  but not for  $\xi \sim \lambda_F$ .

## DEPHASING LEAD MODEL

We next add a third lead to an otherwise closed dot (as in Fig. 1b) and tune the potential on this lead such that the net current through it is zero. Thus, every electron that leaves through lead 3 is replaced by one with an unrelated phase, leading to a loss of phase information without loss of particles. In this situation, the conductance from L to R is given by  $g = T_{LR} + T_{L3} T_{R3} (T_{L3} + T_{R3})^{-1}$  [5], where  $T_{nm}$  is the conductance from lead  $m$  to lead  $n$  when we do not tune the potential on lead 3 to ensure zero current. We next note that  $T_{nm} = T_{nm}^{\text{D}} + \delta T_{nm} + \mathcal{O}[N^{-1}]$ , where the Drude contribution,  $T_{nm}^{\text{D}}$ , is  $\mathcal{O}[N]$  and the weak localization contribution,  $\delta T_{nm}$ , is  $\mathcal{O}[1]$ . If we now expand  $g$  for large  $N$  and collect all the  $\mathcal{O}[1]$ -terms, we get

$$g^{\text{wl}} = \delta T_{LR} + \frac{(T_{L3}^{\text{D}})^2 \delta T_{R3} + (T_{R3}^{\text{D}})^2 \delta T_{L3}}{(T_{L3}^{\text{D}} + T_{R3}^{\text{D}})}. \quad (7)$$



**Fig. 3.** (Color online) Failed coherent-backscattering contribution to conductance:  $\delta T_{nm}^{\text{cbs}}$ . It involves paths that return close but antiparallel to themselves at lead  $m$  but reflect off the tunnel-barrier, remaining in the cavity to finally escape via lead  $n$ . The cross-hatched region is when the two solid paths are paired (within  $W$  of each other).

For a cavity perfectly connected to all three leads (with  $W_{L,R,3} \gg \lambda_F$ ), the Drude and weak localization results for  $\mathcal{U} = 0$  (at finite  $\tau_E^{\text{cl}}$ ) [15] can be substituted into Eq. (7), immediately giving Eq. (2) with  $\tilde{\tau}$  given by  $\tau_E^{\text{cl}}$ .

To connect with the numerics of [13], we now consider a tunnel barrier with finite transparency  $0 \leq \rho_3 \leq 1$  between the cavity and the dephasing lead. Introducing tunnel barriers into the trajectory-based theory of weak localization is detailed in [21]. It requires three main changes to the theory in [15]: (i) The dwell time (single path survival time) becomes  $\tau_{D1}^{-1} = (\tau_0 L)^{-1} \sum_m \rho_m W_m$ . (ii) The paired path survival time (for two paths closer than the lead width) is no longer equal to the dwell time, instead it is  $\tau_{D2}^{-1} = (\tau_0 L)^{-1} \sum_m \rho_m (2 - \rho_m) W_m$ , because survival requires that both paths hitting a tunnel barrier are reflected [21]. (iii) The coherent-backscattering peak contributes to transmission as well as reflection (see Fig. 3).

For the Drude conductance, we need only (i) above, giving us  $T_{nm}^{\text{D}} = \rho_m \rho_n N_m N_n / \mathcal{N}$ , where  $\mathcal{N} = \sum_k \rho_k N_k$ . For the conventional weak localization correction, we need (i) and (ii). The contribution's classical path stays within  $W$  of itself for the time  $T_W(\epsilon)/2$  on either side of the encounter (the dashed region in Figs. 2 and 3); thus, we must use the paired-paths survival time,  $\tau_{D2}$ , for these parts of the path. Elsewhere, the survival time is given by  $\tau_{D1}$ . We follow the derivation in [15] with these new ingredients, and the conventional weak localization correction becomes  $\delta T_{nm}^{(0)} = -(\rho_m \rho_n N_m N_n / \mathcal{N}^2) (\tau_{D1} / \tau_{D2}) \exp[-\Theta]$ . The exponential with  $\Theta = \tau_E^{\text{op}} / \tau_{D2} + (\tau_E^{\text{cl}} - \tau_E^{\text{op}}) / \tau_{D1}$  is the probability that the path segments survive the time  $\tau_E^{\text{op}}$  as a pair ( $\tau_E^{\text{op}}/2$  either side of the encounter) and survive an additional time  $(\tau_E^{\text{cl}} - \tau_E^{\text{op}})$  unpaired (to form a loop of length  $\tau_E^{\text{cl}}$ ). However, we must include point (iii) above

and consider the failed coherent backscattering (shown in Fig. 3). We perform the backscattering calculation following [15] (see also [18]) but using  $\tau_{D2}$  when the paths are closer than  $W$  and  $\tau_{D1}$  elsewhere. We then multiply the result by the probability that the path reflects off lead  $m$  and then escapes through lead  $n$ . This gives the contribution  $\delta T_{nm}^{\text{cbs}} = -(\rho_m (1 - \rho_m) \rho_n N_m N_n / \mathcal{N}^2) \exp[-\Theta]$  assuming  $n \neq m$ . There is a second such contribution with  $m \leftrightarrow n$ . Summing the contributions for  $m \neq n$ ,

$$\delta T_{nm} = (\rho_m \rho_n N_m N_n / \mathcal{N}^2) (\rho_m + \rho_n - \tilde{\mathcal{N}} / \mathcal{N}) e^{-\Theta}, \quad (8)$$

where  $\tilde{\mathcal{N}} = \sum_k \rho_k^2 N_k$ . If only the dephasing lead has a tunnel barrier, substituting the Drude and weak localization results into Eq. (7), we obtain Eq. (2) with  $\tilde{\tau} = (1 - \rho) \tau_E^{\text{op}} + \tau_E^{\text{cl}}$ . In this case, the exponential in Eq. (2) is the probability that a path does not escape into the dephasing lead in either the paired region or the extra time  $(\tau_E^{\text{cl}} - \tau_E^{\text{op}})$  unpaired (for the loop to form).

To generalize our results to  $j$  dephasing leads, we expand the relevant conductance formula in powers of  $N$  and collect the  $\mathcal{O}[1]$  terms. Then,  $g^{\text{wl}} = \delta T_{\text{LR}} + \sum_{m=1}^j (A_m \delta T_{Lm} + B_m \delta T_{Rm})$ , where the sum is over all the dephasing leads. The prefactors  $A_m, B_m$  are combinations of Drude conductances and thus independent of  $\tau_E^{\text{cl}}, \tau_E^{\text{op}}$ ; we need them to get power-law dephasing. However, we can already see that there must be exponential decay with  $[(1 - \rho) \tau_E^{\text{op}} + \tau_E^{\text{cl}}] / \tau_\phi$ , as for  $j = 1$ , where now  $\tau_\phi^{-1} = (\tau_0 L)^{-1} \sum_m \rho_m W_m$  and  $\rho \tau_\phi^{-1} = (\tau_0 L)^{-1} \sum_m \rho_m^2 W_m$ .

## CONCLUSIONS

We first observe that the dephasing-lead model has no independent parameter  $\xi$ . To our surprise, it is the Fermi wavelength, not the dephasing-lead's width, which plays a role similar to  $\xi$ . Thus, a dephasing-lead model cannot mimic a system-environment model with  $\xi \sim L$  at finite  $\tau_E^{\text{cl}}$ . Our second observation is that Eq. (2) with  $\tilde{\tau} = \tau_\xi$  is for a regime where  $\mathcal{U}$  does not affect the momentum/energy of classical paths. Therefore, it does not contradict the result with  $\tilde{\tau} = \tau_E^{\text{cl}}$  in [12] valid for  $\xi$  so small that dephasing occurs via a ‘‘single inelastic process with large energy transfer’’ [22]. Intriguingly, their result is similar to ours for the dephasing-lead model. Could this be due to the destruction of classical determinism by the dephasing process in both cases?

We finally note that conductance fluctuations in the dephasing-lead model exhibit an exponential depen-

dence  $\propto \exp[-2\tau_E/\tau_0]$  for  $\rho_3 \ll 1$  [13] but recover the universal behavior of Eq. (1) for  $\rho_3 = 1$  [23]. However, external noise can lead to dephasing of conductance fluctuations with an  $\tau_E$ -independent exponential term [23], similar to the one found above for weak-localization in the system-environment model. Thus, our conclusion that dephasing is system-dependent in the deep semiclassical limit also applies to conductance fluctuations.

We are grateful to I. Aleiner, P. Brouwer, M. Büttiker, and M. Polianski for useful and stimulating discussions. This work was supported by the Swiss National Science Foundation. A part of this work was done at the Aspen Center for Physics.

### REFERENCES

1. A. D. Stone, in *Physics of Nanostructures*, Ed. by J. H. Davies and A. R. Long (Inst. of Physics and Physical Society, London, 1992).
2. Y. Imry, *Introduction to Mesoscopic Physics* (Oxford Univ. Press, New York, 1997).
3. E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu, *Decoherence and the Appearance of a Classical World in Quantum Theory*, 2nd ed. (Springer, Berlin, 2003).
4. B. L. Altshuler, A. G. Aronov, and D. Khmel'nitsky, *J. Phys. C* **15**, 7367 (1982).
5. M. Büttiker, *Phys. Rev. B* **33**, 3020 (1986).
6. H. U. Baranger and P. A. Mello, *Waves Random Media* **9**, 105 (1999).
7. M. G. Vavilov and I. L. Aleiner, *Phys. Rev. B* **60**, R16311 (1999); G. Seelig and M. Büttiker, *Phys. Rev. B* **64**, 245313 (2001); M. L. Polianski and P. W. Brouwer, *J. Phys. A: Math. Gen.* **36**, 3215 (2003).
8. A. G. Huibers et al., *Phys. Rev. Lett.* **81**, 200 (1998).
9. A. E. Hansen et al., *Phys. Rev. B* **64**, 045327 (2001); K. Kobayashi, H. Aikawa, S. Katsumoto, and Y. Iye, *J. Phys. Soc. Jpn.* **71**, 2094 (2002).
10. F. Pierre et al., *Phys. Rev. B* **68**, 085413 (2003).
11. C. W. J. Beenakker, *Rev. Mod. Phys.* **69**, 731 (1997).
12. I. L. Aleiner and A. I. Larkin, *Phys. Rev. B* **54**, 14423 (1996).
13. J. Tworzydło et al., *Phys. Rev. Lett.* **93**, 186806 (2004).
14. W. G. van der Wiel et al., *Rev. Mod. Phys.* **75**, 1 (2003).
15. Ph. Jacquod and R. S. Whitney, *Phys. Rev. B* **73**, 195115 (2006); see in particular Sect. 5.
16. For a many-particle environment, the coordinate vectors  $\mathbf{q}_0$  and  $\mathbf{q}'_0$  are  $N \times d$  dimensional vectors and  $\Omega_{\text{env}} = V_{\text{env}}^N$ , where  $N$  is the number of particles in the environment, and  $V_{\text{env}} = L_{\text{env}}^d$  is its spatial volume.
17. K. Richter and M. Sieber, *Phys. Rev. Lett.* **89**, 206801 (2002); I. Adagideli, *Phys. Rev. B* **68**, 233308 (2003); S. Heusler, S. Müller, P. Braun, and F. Haake, *Phys. Rev. Lett.* **96**, 066804 (2006).
18. S. Rahav and P. W. Brouwer, *Phys. Rev. Lett.* **96**, 196804 (2006).
19. The conductance of the system is independent of environment contributions, because their integral with respect to  $q$  is zero.
20. Considering a sharp onset of dephasing at  $\xi$  is justified by the local exponential instability of classical trajectories and gives the right answer to the zeroth order in  $(\lambda\tau_D)^{-1}$ .
21. R. S. Whitney, *Phys. Rev. B* **75**, 235404 (2007).
22. Aleiner and Larkin [12] mention that they do not expect their result to hold when  $\xi$  is a classical scale.
23. A. Altland, P. W. Brouwer, and C. Tian, *Phys. Rev. Lett.* **99**, 036804 (2007); cond-mat/0605051v1.