

Eur J Pediatr (2012) 171:603  
DOI 10.1007/s00431-011-1601-8

## ERRATUM

## Erratum to: Chest physiotherapy using passive expiratory techniques does not reduce bronchiolitis severity: a randomised controlled trial

Isabelle Rochat · Patricia Leis · Marie Bouchardy · Christine Oberli ·  
Hendrika Sourial · Margrit Friedli-Burri · Thomas Perneger ·  
Constance Barazzone Argiroffo

Published online: 21 October 2011  
© Springer-Verlag 2011

**Erratum to: Eur J Pediatr**  
DOI 10.1007/s00431-011-1562-y

The author wishes to apologize for the missing figure 1 in the original article. Here is the figure and its caption:


The online version of the original article can be found at <http://dx.doi.org/10.1007/s00431-011-1562-y>.

I. Rochat · C. Barazzone Argiroffo  
Pediatric Pulmonology Unit, Children's Hospital,  
University Hospitals,  
Geneva, Switzerland

P. Leis · M. Bouchardy · C. Oberli · H. Sourial · M. Friedli-Burri  
Pediatric Physiotherapy, Children's Hospital,  
University Hospitals,  
Geneva, Switzerland

T. Perneger  
Division of Clinical Epidemiology, University Hospitals,  
Geneva, Switzerland

I. Rochat (✉)  
Pediatric Pulmonology Unit, CHUV,  
Rue du Bugnon 46,  
1011 Lausanne, Switzerland  
e-mail: [isabelle.rochat@chuv.ch](mailto:isabelle.rochat@chuv.ch)



**Fig. 1** Comparison of the time to clinical stability in children hospitalised with bronchiolitis randomised to chest physiotherapy (grey line) or control group (black line). The slight difference at day 5 was not significant ( $P=0.35$ )