

MHC presentation via autophagy and how viruses escape from it

Monique Gannage · Christian Münz

Received: 18 May 2010 / Accepted: 29 August 2010 / Published online: 21 September 2010
© Springer-Verlag 2010

Abstract T cells detect infected and transformed cells via antigen presentation by major histocompatibility complex (MHC) molecules on the cell surface. For T cell stimulation, these MHC molecules present fragments of proteins that are expressed or taken up by the cell. These fragments are generated by distinct proteolytic mechanisms for presentation on MHC class I molecules to cytotoxic CD8⁺ and on MHC class II molecules to helper CD4⁺ T cells. Proteasomes are primarily involved in MHC class I ligand and lysosomes, in MHC class II ligand generation. Autophagy delivers cytoplasmic material to lysosomes and, therefore, contributes to cytoplasmic antigen presentation by MHC class II molecules. In addition, it has been recently realized that this process also supports extracellular antigen processing for MHC class II presentation and cross-presentation on MHC class I molecules. Although the exact mechanisms for the regulation of these antigen processing pathways by autophagy are still unknown, recent studies, summarized in this review, suggest that they contribute to immune responses against infections and to maintain tolerance. Moreover, they are targeted by viruses for immune escape and could maybe be harnessed for immunotherapy.

Keywords MHC class I · MHC class II · CD8⁺ T cell · CD4⁺ T cell · Cross-presentation · Herpesvirus · Endocytosis

Introduction

T cells detect infected or transformed cells via antigen presentation by major histocompatibility complex (MHC) molecules on the cell surface. Helper CD4⁺ T cells and cytotoxic CD8⁺ T cells are restricted by MHC class I and II molecules, respectively. These molecules display antigens in the form of peptides on the cell surface. These peptides are generated via distinct proteolytic processes and loaded in different cellular compartments on MHC class I and II molecules. Some antigen processing for peptide presentation by MHC molecules is assisted by autophagy.

CD8⁺ T cells, which rapidly expand during immune responses and mediate cytotoxicity against infected and tumor cells, recognize octamer or nonamer peptides on MHC class I molecules [1]. These peptides are thought to originate primarily from protein degradation by proteasomes, large multicatalytic proteases in cytosol and nucleus [2]. These peptides are then imported into the endoplasmic reticulum (ER) via the transporter associated with antigen processing (TAP), and can be further trimmed on their N terminus via ER aminopeptidases associated with antigen processing (ERAAPs) [3]. Peptides that fulfill the binding requirements for the expressed MHC class I molecules, so-called binding motifs, are then loaded into the peptide binding groove of these molecules in the MHC class I loading complex. Upon binding of high-affinity ligands, MHC class I molecules and their peptide cargo are then released from the ER to travel to the cell surface for immune surveillance by CD8⁺ T cells. This model of antigen loading onto MHC class I molecules predicts that proteasome substrates are the main source of MHC class I ligands. Originally, it was assumed that MHC class I molecules would, therefore, primarily present cytosolic and nuclear antigens. However, it was noted that certain cell

This article is published as part of the Special Issue on Autophagy.

M. Gannage · C. Münz (✉)
Viral Immunobiology, Institute of Experimental Immunology,
University of Zürich,
Winterthurerstrasse 190,
CH-8057 Zürich, Switzerland
e-mail: christian.muenz@usz.ch

types, primarily professional antigen-presenting cells like dendritic cells (DCs) are capable to display extracellular antigens on MHC class I molecules by antigen processing via cross-presentation [4, 5]. During cross-presentation, endocytosed antigen is thought to escape into the cytosol by so far poorly defined mechanisms. Whole proteins or preprocessed protein fragments might be delivered to the cytosol, and at least for some antigens, vesicular degradation by insulin regulated aminopeptidase is required for efficient cross-presentation [6]. In the cytosol, cross-presented antigens can access the proteasome and follow the MHC class I antigen processing and presentation pathway. The stage of endosomal maturation, from which cross-presented antigen is exported, is heavily debated, and early endosomes as well as late endosomes, including some that might even fuse with the ER, have been implicated [7–10]. Thus, different antigen formulations might be cross-presented from different endosomes, but, as we will discuss later, antigen assisted in its exocytosis by macroautophagy might access one of these cross-presenting compartments very efficiently.

In contrast to MHC class I molecules, MHC class II molecules present longer peptides of heterogenic size (≥ 9 amino acids long) to CD4⁺ T cells, which orchestrate humoral and cellular immune responses by virtue of their cytokine production. MHC class II ligands are primarily products of lysosomal proteolysis [11] and are loaded with these in late endosomal compartments, called MHC class II loading compartment (MIIC). MHC class II molecules reach these late endosomes with the help of a chaperone called invariant chain (Ii), which prevents premature peptide binding of MHC class II molecules in ER and Golgi apparatus, as well as guides MHC class II molecules via its cytosolic domain to MIICs. In these vesicles, Ii is degraded by lysosomal proteolysis, and the remnant peptide occupying the peptide binding groove, class II-associated invariant chain peptide, is expelled with the help of the chaperones H2-M or HLA-DM in mouse and man, respectively. These chaperones also ensure high-affinity peptide ligand binding to MHC class II molecules. Thus, MHC class II presented antigens are in their majority proteins that can gain access to the MIIC. In the classical paradigm of MHC class II antigen processing, these were thought to be exclusively extracellular proteins that reach the MIIC after endocytosis. As discussed later, it has now been realized that autophagy can permit cytoplasmic antigen to also access this pathway.

With respect to autophagic pathways that can be involved in antigen processing for MHC presentation, only macroautophagy and chaperone-mediated autophagy have so far been investigated, and we want to briefly introduce the processes and molecules that have been used by immunologists to elucidate a role for autophagy in antigen presentation to T cells. During macroautophagy more than 30 autophagy-related gene (*atg*) products are involved in

the generation and lysosome fusion of cytosolic double-membrane surrounded vesicles, called autophagosomes [12]. The sites of autophagosome generation are characterized by the assembly of type III phosphoinositide-3 (PI3) kinase complexes, incorporating Atg6/Beclin-1. Both rough ER and the outer nuclear leaflet have been identified as autophagosome formation sites [13–15], but these vesicles might also be generated at other places. Two ubiquitin-like systems are involved in autophagosome formation at these sites with Atg8 and Atg12 as the ubiquitin-like molecules at their center [16]. In one ubiquitin-like system Atg12 gets coupled to Atg5 with the assistance of the E1- and E2-like enzymes Atg7 and Atg10. The Atg12–Atg5 heterodimer then associates with Atg16L1 and coats the outer autophagosome membrane. It dissociates from the outer membrane, once the autophagosome is completed, and has been shown to direct the conjugation of the other ubiquitin-like molecule Atg8. Atg8 and its most studied mammalian homologue LC3 are coupled to the lipid phosphatidylethanolamine at the outer and inner autophagosomal membrane with the help of the E1- and E2-like enzymes Atg7 and Atg3. While it is recycled from the outer membrane upon autophagosome completion, it stays associated with the inner membrane and is degraded with the autophagosome cargo in lysosomes. Atg8/LC3 catalyses hemifusion of membranes and might therefore support the elongation of the autophagosome membrane [17]. In addition, it is used to recruit autophagic cargo [18–20]. In addition to its role in autophagosome formation, Atg6/Beclin-1-containing PI3 kinase complexes are then once more involved in macroautophagy at the stage of autophagosome fusion with the lysosome [21–24]. In addition to macroautophagy, chaperone-mediated autophagy also imports cytosolic proteins into lysosomes for degradation [25]. Its substrates carry a KFERQ-like signal sequence, which targets them for direct transport across the lysosomal membrane. This transport is assisted by cytosolic and vesicular chaperones of the heat-shock protein family and the transmembrane protein LAMP2A. Thus, both macroautophagy and chaperone-mediated autophagy deliver cytoplasmic constituents to lysosomes for degradation. Since T cells monitor the proteolytic waste of cells for signs of infection and transformation, it is maybe not surprising that they also survey substrates of autophagy. The evidence for this will be discussed below.

Antigen processing for MHC class II presentation via autophagy

Endogenous pathway

Different studies have now demonstrated that autophagosomes can deliver cytosolic and nuclear proteins to MHC


class II loading compartments (Fig. 1), and this phenomenon occurs in different cell types: in epithelial cells, in B cells, but also in dendritic cells as the most professional antigen-presenting cells. Morphological analysis of MHC class II compartment have revealed that between 30% and 50% of MHC class II compartment co-stain with the autophagosome marker Atg8/LC3 [26]. In parallel, using different model antigens, for the generation of which proteins were coupled to Atg8/LC3, our group demonstrated that targeting an antigen to macroautophagy significantly enhances its processing and presentation on MHC class II molecules [26] and (Gannage et al., unpublished observation). In this section, we discuss different examples of intracellular antigens from self, viral, and tumor origin that have been shown to be processed for MHC class II presentation via macroautophagy.

Evidence that self-antigens can be delivered to MHC class II loading compartment via macroautophagy came first from the analysis of MHC class II bound epitopes. In human B lymphoblastoid cell lines, mass spectrometrical analysis of the MHC class II ligandome showed that 20% to 30% of self-class II epitopes are derived from cytosolic and nuclear proteins. A significant change in the MHC presented amount of these peptides occurred upon starvation-induced macroautophagy [27]. This self-antigen presentation via macroautophagy might be involved in autoimmune diseases. Along these lines, the group of Ludger Klein has implicated macroautophagy in the generation of central tolerance by MHC class II antigen presentation of self-antigens on thymic epithelial cells (TECs). Using a model of transplantation of $Atg5^{-/-}$ thymi into wild-type or TCR transgenic mice, Klein et al. demonstrated that thymic macroautophagy shapes the repertoire of $CD4^+$ T cells [28]. Macroautophagy in cortical TECs seemed to be required for positive selection of certain T cell receptor specificities, while in medullary TECs, it was

required for negative selection. Without macroautophagy-dependent negative selection, immunopathology was observed in several tissues including the gut, and the symptoms could be transferred by T cells. Similarly, Kasai and colleagues demonstrated by immunofluorescence analysis the colocalization of Atg8/LC3 with MHC class II loading compartment in TECs, both in vitro (in cTEC and mTEC cell lines) and in vivo (in thymic cryosection) [29]. In addition, Western blot analysis of H2-M vesicles isolated from TECs lines detected the lipidated form of Atg8/LC3 (LC3-II), suggesting fusion of autophagosomes with these MHC class II loading compartments. It is tempting to speculate that macroautophagy allows intracellular antigens, including peripheral tissue-derived antigen that are transcribed due to the expression of the transcription factor autoimmune regulator element, to be presented onto MHC class II molecules for central tolerance induction by TECs. However, this self-antigen presentation might also result in autoimmunity, and Janice Blum's group has indeed shown that the autoantigens GAD65 and SMA can be better presented to $CD4^+$ T cells when chaperone-mediated autophagy is enhanced [30]. These data suggest that both macroautophagy and chaperone-mediated autophagy transport self-proteins for MHC class II presentation to $CD4^+$ T cells during tolerance induction and possibly also autoimmune disease.

Apart from self-antigens, macroautophagy can also deliver pathogen-derived antigens to MHC class II loading compartments. This is exemplified by viral antigens in Epstein Barr virus (EBV) infected B cells. The pathway has been implicated in the degradation of two latent EBV antigens, EBV nuclear antigen 1 (EBNA1) and latent membrane protein 1 (LMP1). LMP1, which is considered the major oncogene of EBV, induces macroautophagy [31] and accumulates upon silencing of the two essential macroautophagy gene products Atg6/Beclin 1 and Atg7. Therefore, LMP1 seems to regulate its own clearance by

Fig. 1 Macroautophagy assists in antigen processing for MHC class I and II presentation. Cytosolic antigen can be directly transported to MHC class II loading compartments (MIIC) via autophagosomes (lower left). In addition, it might escape lysosomal degradation and get exocytosed from multivesicular bodies (MVB) for uptake by bystander cells (right and upper left). The endocytosed antigen might get loaded onto MHC class II molecules or escape to the cytosol for proteasomal degradation and presentation on MHC class I molecules



macroautophagy, but further studies will have to be performed to demonstrate if this results in CD4⁺ T cell epitope generation from LMP1. For the second latent EBV antigen that is at least partially turned over by macroautophagy, EBNA1, antigen processing for MHC class II presentation via macroautophagy has been demonstrated [32, 33]. Leung et al. identified two epitopes from EBNA1 that are processed via endogeneous pathways for MHC class II presentation by EBV-infected B cells. In line with previously published results from our group [32], one of these epitopes is delivered to MHC class II loading compartments via macroautophagy. Interestingly, preventing nuclear import of EBNA1 by mutating its nuclear localization sequence strongly enhanced the presentation of both this and the other CD4⁺ T cell epitopes. Both of them were now delivered to MHC class II loading compartment via macroautophagy. Thus, cytosolic relocation of EBNA1 resulted in broadening the range of CD4⁺ T cell epitopes, displayed on MHC class II molecules after macroautophagy. How the relocation of an antigen from the nucleus to the cytosol can enhance its endogeneous delivery to MHC class II compartment by macroautophagy remains unclear. In contrast to this finding, the study by Riedel et al. [34] suggested that for endogenous MHC class II presentation of the cytosolic bacterial antigen neomycin phosphotransferase II via macroautophagy, the relocation of the antigen to the nucleus does not affect its antigen recognition. The differences in these two studies might be related to the nature and expression level of the antigen itself, allowing them to form cytosolic malformed protein pools to different degrees. In another viral infection, herpes simplex virus type 1 (HSV-1) induces macroautophagy, but this response is antagonized by the HSV-1 neurovirulence gene product, infected cell protein (ICP) 34.5, which interacts with the essential autophagy protein Atg6/Beclin-1. Interestingly, a mutant virus unable to bind to the Atg6/Beclin-1 protein induces a significantly stronger CD4⁺ T cell response in infected mice [35]. This might imply an essential role of macroautophagy-mediated MHC class II presentation during HSV-1 infection. Indeed, Akiko Iwasaki and colleagues have recently demonstrated that HSV infection of mice, lacking macroautophagy in DCs, was compromised in raising virus-specific CD4⁺ T cell responses [36]. Finally, in macrophages and DCs infected with mycobacterium tuberculosis (Mtb), induction of macroautophagy by rapamycin, IFN- γ , or starvation significantly enhanced the processing of immunodominant CD4⁺ T cell epitopes from the mycobacterial antigen Ag85B [37]. This mechanism is an active process, occurring only in live Mtb infection, and is specifically inhibited by RNA silencing of Atg6/Beclin-1. In addition, vaccination of mice with rapamycin-pretreated Mtb-infected DCs significantly improved the efficiency of T cell responses prior to Mtb challenge. This work described

for the first time how the manipulation of macroautophagy could improve a vaccine strategy. Thus, in the MHC class II presentation of individual viral antigens via macroautophagy, a role for this pathway in vivo and even immune evasion by viral inhibition of macroautophagy has been demonstrated.

A third group of antigens that might fall prey to macroautophagy are tumor antigens. The implications of macroautophagy during tumorigenesis are still unclear. Among others, macroautophagy has been shown to promote cell survival of cancer cells, but also in parallel to function as a tumor suppressor pathway in some models. Nevertheless, the pathway is active in cancer cells and might generate tumor-derived CD4⁺ T cell epitopes. As an example, the protein encoded by the mucin gene 1 (MUC1) has been described to be processed by macroautophagy after expression in DCs [38]. In MUC1-transfected DCs, the processing of MUC1 for MHC class II presentation was found to be dependent on macroautophagy. This was documented by significant inhibition of MUC1-specific CD4⁺ T cell proliferation during coculture with antigen-expressing DCs in the presence of PI3 kinase inhibitors (3-methyladenine or wortmannin), which compromise autophagosome generation. In contrast, MUC1-specific CD8⁺ T cell responses were unaffected by this pathway. Thus, self-derived, pathogen-derived, and tumor antigens have now been demonstrated to follow macroautophagy for MHC class II presentation to CD4⁺ T cells. This pathway of immune surveillance should be harnessed during vaccination and for the development of novel immunotherapies.

Exogenous pathway

In addition to its role in MHC class II presentation of cytosolic antigens, macroautophagy has recently been shown to participate in the processing of exogenous antigens and their delivery to MHC class II compartment (Fig. 1). Evidence for this pathway has been gathered primarily in macrophages and DCs. Indeed, three recent studies have established a role for macroautophagy in the processing of phagocytosed antigens for MHC class II presentation by DCs. Particularly, enhanced antigen processing after DC activation by pathogen associated molecular patterns, as recognized by toll-like receptors (TLRs) and NOD-like receptors (NLRs), was supported by macroautophagy.

The first study by the group of Akiko Iwasaki highlighted the importance of macroautophagy in DCs for CD4⁺ T cell priming in vivo [36]. Interestingly, during HSV infection, macroautophagy seems to be important not only for the processing of cytosolic endogeneous antigens and their delivery to MHC class II compartment, but also for the processing of exogenous phagocytosed antigens. Using chimeric mice reconstituted with a hematopoietic system

deficient for Atg5, the authors could demonstrate a defect in CD4⁺ T cell responses in vivo upon HSV infection. In addition, in these chimeric mice, adoptively transferred ovalbumin (OVA)-specific CD4⁺ T cells showed an impairment of proliferation after infection with OVA-expressing *Listeria monocytogenes* or OVA-expressing HSV. A role of macroautophagy for MHC class II antigen processing and presentation in vivo in professional antigen-presenting cells was further supported by mice with a conditional deletion of Atg5 in their DCs. Indeed in these mice, HSV infection failed to prime CD4⁺ T cell responses in vivo, and the mice succumbed to a more severe disease compared with wild-type mice and developed a higher clinical score upon infection. In the same study, the authors ruled out a general defect in macroautophagy-deficient DCs, by demonstrating that they had no defect in their migration capability, in their phenotypic maturation (CD86, CD40, and MHC class II expression), and in their cytokines production, and were not impaired in endocytosis or phagocytosis compared with wild-type DCs. Instead, the defect in extracellular antigen presentation on MHC class II molecules was rather due to an impairment of processing of phagocytosed antigens that trigger TLR stimulation. Using lipopolysaccharide-coated OVA beads, an impairment of phagosome to lysosome fusion was observed in Atg5^{-/-} DCs, resulting in a reduction of the processing of phagocytosed antigen for MHC class II presentation. This phenotype was the consequence of a defect in the delivery of lysosomal proteases to phagosomes, as well as a delay of fusion of phagosomes with lysosomes. The precise mechanism of how the macroautophagy machinery is recruited to phagosomes and might enhance fusion with lysosomes remains unclear, but these findings are in line with previous studies, documenting enhanced antigen presentation of TLR ligand-coated antigen and Atg-supported fusion of phagosomes containing such antigen with lysosomes [39, 40]. In one of these studies, the authors described that TLR engagement induces “LC3-associated phagocytosis” (LAP). Upon TLR2 stimulation, a recruitment of Atg8/LC3 to the phagosomal membrane was observed, resulting in an enhanced maturation of the phagosome. These studies suggest that macroautophagy facilitates endocytosed antigen delivery to lysosomes and MHC class II presentation of this antigen.

Along the same lines, Alison Simmons and colleagues demonstrated that upon nucleotide-binding oligomerization domain-containing-2 (NOD2) activation, macroautophagy is upregulated and involved in MHC class II processing of phagocytosed antigens [41]. This study provided for the first time evidence that a defect in macroautophagy mediated MHC class II antigen presentation could result in defective CD4⁺ T cell responses, which might then not be able to control the gut commensals in Crohn disease (CD). This defect in macroautophagy-mediated MHC class II antigen presentation could be caused by mutations in

both the Atg16L1 and NOD2 gene, which had been identified as major risk factors for the occurrence of the disease. The authors demonstrated that a bacterial peptidoglycan muramyl dipeptide (MDP) induces macroautophagy upon NOD2 activation in human DCs. MHC class II expression on the cell surface of MDP-activated DCs was upregulated and this upregulation was dependent on macroautophagy. In parallel, a colocalization of Atg8/LC3 with MHC class II loading compartments was observed. Moreover, DC-mediated priming of CD4⁺ T lymphocytes after *Salmonella enterica* infection was dependent on NOD2 and Atg16L1 because CD4⁺ T cell proliferation was compromised by siRNA-mediated silencing of these two genes. The second part of this study analyzed the function of DCs from CD patients, carrying either the NOD (3020insC) or the Atg16L1 (T300A) mutation, which are associated with the familial form of the disease. In both cases, the cells demonstrated a defect in macroautophagy upregulation upon MDP stimulation or *S. enterica* infection. As a consequence, MHC class II upregulation and CD4⁺ T cell proliferation were compromised. Confirming these findings, the group of Dana Philpott could demonstrate that NOD1 and 2 stimulation upregulated macroautophagy and led to autophagosome recruitment to the plasma membrane [42]. This regulation was independent of the classical pathway of NLR activation, involving usually the RIP2 adaptor and NF- κ B transcription factor. These data suggested that also NLR-ligand stimulated MHC class II upregulation and antigen presentation requires macroautophagy.

Finally, a third recent study describes a new mechanism of immune escape of human immunodeficiency virus (HIV) 1 virus in DCs [43]. In this work, HIV-1 was shown to inhibit macroautophagy initiation by activating the mTor pathway. This inhibition resulted in an impairment of TLR4 and TLR8 activation by viral replication intermediates and of antigen presentation to CD4⁺ T cells. Indeed, in an in vitro assay of DC capture of inactivated HIV virus, macroautophagy was shown to be important for MHC class II antigen processing of a HIV gag-derived CD4⁺ T cell epitope. TNF- α secretion of an HIV gag-specific CD4⁺ T cell clone was significantly reduced upon stimulation with macroautophagy-deficient DCs, pulsed with inactivated HIV virus. In parallel, CD8⁺ T cell activation was not compromised by macroautophagy inhibition. This study suggests that efficient HIV antigen processing for MHC class II presentation by DCs requires macroautophagy.

Macroautophagy in antigen processing for MHC class I presentation

In contrast to a quite robust role for macroautophagy in MHC class II antigen presentation, much less is known

about how this pathway influences antigen processing for MHC class I presentation. Indeed, in several studies, MHC class I-restricted antigen recognition was actually used as a control, and no effect of macroautophagy inhibition on CD8⁺ T cell recognition was observed [26, 36, 43]. However, under certain circumstances, macroautophagy might enhance antigen presentation on MHC class I molecules to CD8⁺ T cells, and we will summarize the so far sparse evidence for this here.

The only study so far that argues for a role of macroautophagy in endogenous antigen processing for MHC class I presentation by the group of Michel Desjardins suggested that late during HSV infection macroautophagy supports MHC class I presentation of viral antigens [15]. This *in vitro* finding was restricted to an immunodominant CD8⁺ T cell epitope of the glycoprotein B of the virus, and to its processing during the late stage of the infection. Interestingly, early after infection (6–8 h), the presentation of this epitope followed the classical MHC class I pathway, but later on (8–12 h), macroautophagy inhibition compromised presentation to CD8⁺ T cells; even so, processing of this antigen was still dependent on components of the classical MHC class I antigen processing pathway like the proteasome and TAP. In infected macrophages, the authors demonstrated that silencing the essential autophagy gene *atg5* resulted in a decrease of HSV gB-specific CD8⁺ T cell recognition. Electron microscopy analysis revealed that the formation of autophagosomes after HSV-1 infection involved the outer nuclear membrane. However, it was not clarified by which mechanism autophagosome cargo can escape to the cytosol to be further processed by proteasomes. Irrespective of the mechanism, this study suggests that macroautophagy might intersect with the cross-presentation pathway of MHC class I antigen processing.

A similar intersection with endosomes capable of cross-presentation was proposed for macroautophagy substrates of antigen donor cells (Fig. 1). The first report showing a direct involvement of macroautophagy in cross-presentation was by Li and colleagues [44]. In this study, 293T cells expressing the model antigen OVA *in vitro*, or melanoma cells expressing the gp100 melanoma antigen *in vivo*, were used as donor cells to access cross-presentation. In both cases, siRNA-mediated silencing of macroautophagy in antigen donor cells resulted in a significant reduction of CD8⁺ T cell activation after cross-presentation by DCs. Interestingly, purified autophagosomes from antigen donor cells were also efficiently cross-presented to CD8⁺ T cells, probably acting like exosomes. Therefore, antigen wrapped in macroautophagic membranes might be more efficiently taken up by antigen-presenting cells and cross-presented on MHC class I molecules.

In a second report along these lines by the group of Matthew Albert [45], evidence was provided that efficient

cross-presentation of viral antigens required macroautophagy in antigen donor cells. In this study, two cellular sources were used to provide antigen for cross-presentation experiments, wild-type mouse embryonic fibroblasts (MEFs) and Bax/Bak^{-/-} MEFs. While wild-type MEFs can undergo caspase-dependent apoptosis, Bax/Bak^{-/-} MEFs are unable to perform this type of cell death, but upregulate macroautophagy under the applied treatment conditions. Both cell types were infected with influenza A virus, then treated with proapoptotic agents and adoptively transferred *in vivo* for priming experiment. Mice immunized with Bax/Bak^{-/-} MEFs showed a significantly higher CD8⁺ T cell response specific to both HA_{518–526} and NP_{366–374} immunodominant CD8⁺ T cell epitopes, compared with mice immunized with wild-type MEFs. These findings again suggest that macroautophagy might package antigen efficiently for cross-presentation by DCs. How macroautophagic substrates, however, leave antigen donor cells for cross-presentation remains to be determined.

Viral evasion from macroautophagy

The important role of macroautophagy during immune responses is further underscored by its regulation during viral infections. Three *in vivo* systems have linked macroautophagy to immune control of viral infections. These included with HSV one DNA, and with Sindbis virus and vesicular stomatitis virus (VSV) two RNA virus. Neurovirulence of both HSV and Sindbis virus was attenuated when macroautophagy was elevated either by mutation of the macroautophagy inhibiting ICP34.5 HSV gene product [46], or by overexpression of Atg6/Beclin-1, driving macroautophagy, in Sindbis virus-infected mice [47]. Furthermore, Sindbis virus infection of the CNS is augmented by macroautophagy inhibition via either recombinant viruses encoding a dominant negative Atg5 protein, recombinant viruses deleting floxed Atg5 upon infection via cre-recombinase expression or wild-type virus infection of mice with conditional Atg5 deletion in neurons [48]. In addition, VSV infection induced macroautophagy in *Drosophila* flies and restricted VSV infection in this model organism [49]. These studies did not implicate antigen processing for MHC presentation via macroautophagy in the protection from virus infection. However, during HSV infection, macroautophagy in myeloid DCs was found to be required for efficient priming of protective CD4⁺ T cell responses [36]. These studies implicate that macroautophagy restricts viral replication *in vivo*.

Another indication that macroautophagy significantly limits viral replication is that successful viral pathogens have developed immune escape mechanisms to target this pathway. In this respect, two checkpoints of macroautophagy

are inhibited by viruses. These are autophagosome formation and autophagosome fusion with lysosomes. Interestingly, DNA viruses seem to primarily compromise autophagosome formation, while RNA viruses block autophagic cargo degradation in lysosomes. With respect to the inhibition of autophagosome formation, herpesviruses have proven a rich source of macroautophagy inhibiting proteins. Within the α -herpesviruses, HSV encodes the late ICP34.5, which blocks autophagosome formation [50, 51]. ICP34.5 inhibits autophagosome generation by binding to Atg6/Beclin-1 [46]. Interestingly, HSV with a mutant ICP34.5 protein, lacking the Atg6/Beclin-1 interacting domain, shows increased neurovirulence after CNS injection in mice [46]. Within the β -herpesviruses, human cytomegalovirus also inhibits macroautophagy [52]. However, the molecular mechanism of this regulation is unknown so far. Finally, the group of γ -herpesviruses contains several members that compromise autophagosome formation, namely Kaposi sarcoma associated herpesvirus (KSHV) and murine γ -herpesvirus (MHV-68). KSHV encodes a B cell lymphoma 2 protein (Bcl-2) homologue, encoded by ORF-16 of the virus, which also interacts with Atg6/Beclin-1 and inhibits autophagosome formation [53]. In addition, the same virus compromises autophagosome formation also via its viral FLICE inhibitory protein (v-FLIP) protein, encoded by ORF71/K13 [54]. KSHV v-FLIP impairs Atg3, the E2-like enzyme of Atg8/LC3 lipidation. This mechanism prevents Atg8/LC3-dependent autophagosome formation. Therefore, KSHV expresses with v-FLIP a macroautophagy inhibitor during latent, and with v-Bcl-2 one during lytic infection. The closest rodent virus to KSHV, MHV-68, also expresses a Bcl-2 homologue, M11, which interacts with Atg6/Beclin-1 to inhibit autophagosome formation [55]. Deletion of the macroautophagy blocking, but not the apoptosis compromising domain, impairs the establishment of chronic infection by this virus [56]. Collectively, these data suggest that herpesviruses inhibit autophagosome formation in order to inhibit restriction of viral infection and viral antigen presentation.

In addition to autophagosome generation as a first checkpoint of macroautophagy, degradation of these vesicles via fusion with lysosomes is targeted by RNA viruses. Already, early electron microscopy studies of poliovirus-infected cells reported the accumulation of double-membrane engulfed vesicles [57]. This stabilization of autophagosomes could be induced by the viral proteins 2BC and 3A [58], even so the molecular mechanism underlying this presumed block in autophagosome degradation remains unclear. Formation of these poliovirus stabilized vesicles is dependent on some Atg proteins because siRNA-mediated silencing of Atg8/LC3 and Atg12 inhibits their generation, and they seem to represent, therefore, a subtype of autophagosomes [58]. The stabilized membranes support viral replication and are thought to function as a scaffold for viral replication. In

addition, inhibition of autophagosome degradation might enhance virus particle exocytosis [58]. Furthermore, the stabilization of autophagosome membranes for viral replication has been proposed for other RNA viruses, like the flaviviruses hepatitis C virus (HCV) and dengue virus [59–63]. Moreover, even RNA viruses that do not use autophagosomal membranes for their replication seem to benefit from inhibition of autophagosome degradation for their release of infectious viruses from infected cells [64]. Along these lines, the HIV was found to block autophagosome maturation. Particularly, its Nef protein interacts with Atg6/Beclin-1 blocking degradation of macroautophagic cargo, including HIV particles. These, therefore, are exocytosed at increased rates. This benefit for HIV in blocking autophagosome degradation was only observed in macrophages, but not T cells. Similarly, a slight benefit of macroautophagy regulation for the replication of the segmented RNA virus influenza virus was observed only in canine kidney epithelial cells, but not in human lung epithelia or mouse embryonic fibroblasts [65, 66]. Irrespective of a benefit for viral replication, influenza infection leads to autophagosome accumulation in a broad range of cell types [66]. The virus achieves this by blocking autophagosome fusion with lysosomes via its matrix protein 2. This macroautophagy block enhances apoptotic cell death of infected cells and might regulate immunogenic viral protein release. Taken together, these studies suggest that viruses hijack macroautophagy to prevent their degradation and viral antigen presentation, or even use autophagic membranes for their own replication and exit from infected cells. Depending on the need of the virus, they block autophagosome generation or degradation with DNA viruses targeting preferentially the first, and RNA viruses the second checkpoint of macroautophagy, respectively.

Conclusions

Macroautophagy supports adaptive immune responses by facilitating antigen processing for MHC presentation to T cells. Since autophagosomes direct cytoplasmic content for lysosomal degradation and MHC class II molecules present lysosomal products to CD4⁺ T cells, most studies suggest a role for macroautophagy in antigen presentation to these helper T cells. However, some studies also describe that macroautophagy might even assist MHC class I antigen presentation to CD8⁺ T cells. While intracellular antigen presentation by MHC class II molecules seems to benefit from macroautophagic substrate delivery to MHC class II loading compartments, the mechanisms by which extracellular antigen delivery to these MHCs is supported by macroautophagy, by which autophagosome content can again escape to the cytosol for classical MHC class I antigen

presentation, and by which antigen can be efficiently packaged by macroautophagy for cross-presentation remain unclear. The evidence that viruses, however, target this pathway for their immune escape suggests that it might be important for immune control. Thus, macroautophagy should be harnessed to enhance immunotherapies and assist antigen processing during vaccinations.

Acknowledgements Our research is supported by the National Cancer Institute (R01CA108609 and R01CA101741), the Foundation for the National Institutes of Health (Grand Challenges in Global Health), and the Swiss National Science Foundation (310030_126995).

References

- Rammensee H, Bachmann J, Emmerich NP, Bachor OA, Stevanovic S (1999) SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 50(3–4):213–219
- Peaper DR, Cresswell P (2008) Regulation of MHC class I assembly and peptide binding. *Annu Rev Cell Dev Biol* 24:343–368
- Hammer GE, Kanaseki T, Shastri N (2007) The final touches make perfect the peptide-MHC class I repertoire. *Immunity* 26(4):397–406
- Bevan MJ (1976) Cross-priming for a secondary cytotoxic response to minor H antigens with H-2 congenic cells which do not cross-react in the cytotoxic assay. *J Exp Med* 143(5):1283–1288
- Amigorena S, Savina A (2010) Intracellular mechanisms of antigen cross presentation in dendritic cells. *Curr Opin Immunol* 22(1):109–117
- Saveanu L, Carroll O, Weimershaus M et al (2009) IRAP identifies an endosomal compartment required for MHC class I cross-presentation. *Science* 325(5937):213–217
- Houde M, Bertholet S, Gagnon E et al (2003) Phagosomes are competent organelles for antigen cross-presentation. *Nature* 425(6956):402–406
- Guermonez P, Saveanu L, Kleijmeer M et al (2003) ER-phagosome fusion defines an MHC class I cross-presentation compartment in dendritic cells. *Nature* 425(6956):397–402
- Goldszmid RS, Coppens I, Lev A et al (2009) Host ER-parasitophorous vacuole interaction provides a route of entry for antigen cross-presentation in *Toxoplasma gondii*-infected dendritic cells. *J Exp Med* 206(2):399–410
- Burgdorf S, Kautz A, Bohnert V, Knolle PA, Kurts C (2007) Distinct pathways of antigen uptake and intracellular routing in CD4 and CD8 T cell activation. *Science* 316(5824):612–616
- Trombetta ES, Mellman I (2005) Cell biology of antigen processing in vitro and in vivo. *Annu Rev Immunol* 23:975–1028
- Mizushima N, Klionsky DJ (2007) Protein turnover via autophagy: implications for metabolism. *Annu Rev Nutr* 27:19–40
- Yla-Anttila P, Vihinen H, Jokitalo E, Eskelinen EL (2009) 3D tomography reveals connections between the phagophore and endoplasmic reticulum. *Autophagy* 5(8):1180–1185
- Hayashi-Nishino M, Fujita N, Noda T et al (2009) A subdomain of the endoplasmic reticulum forms a cradle for autophagosome formation. *Nat Cell Biol* 11(12):1433–1437
- English L, Chemali M, Duron J et al (2009) Autophagy enhances the presentation of endogenous viral antigens on MHC class I molecules during HSV-1 infection. *Nat Immunol* 10(5):480–487
- Ohsumi Y (2001) Molecular dissection of autophagy: two ubiquitin-like systems. *Nat Rev Mol Cell Biol* 2(3):211–216
- Nakatogawa H, Ichimura Y, Ohsumi Y (2007) Atg8, a ubiquitin-like protein required for autophagosome formation, mediates membrane tethering and hemifusion. *Cell* 130(1):165–178
- Thurston TL, Ryzhakov G, Bloor S, von Muhlinen N, Randow F (2009) The TBK1 adaptor and autophagy receptor NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* 10(11):1215–1221
- Zheng YT, Shahnazari S, Brech A et al (2009) The adaptor protein p62/SQSTM1 targets invading bacteria to the autophagy pathway. *J Immunol* 183(9):5909–5916
- Ponpuak M, Davis AS, Roberts EA et al (2010) Delivery of cytosolic components by p62 endows autophagosomes with unique anti-microbial properties. *Immunity* 32:329–341
- Itakura E, Kishi C, Inoue K, Mizushima N (2008) Beclin 1 forms two distinct phosphatidylinositol 3-kinase complexes with mammalian Atg14 and UVRAG. *Mol Biol Cell* 19(12):5360–5372
- Matsunaga K, Saitoh T, Tabata K et al (2009) Two Beclin 1-binding proteins, Atg14L and Rubicon, reciprocally regulate autophagy at different stages. *Nat Cell Biol* 11(4):385–396
- Zhong Y, Wang QJ, Li X et al (2009) Distinct regulation of autophagic activity by Atg14L and Rubicon associated with Beclin 1-phosphatidylinositol-3-kinase complex. *Nat Cell Biol* 11(4):468–476
- Liang C, Feng P, Ku B et al (2006) Autophagic and tumour suppressor activity of a novel Beclin1-binding protein UVRAG. *Nat Cell Biol* 8(7):688–699
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ (2008) Autophagy fights disease through cellular self-digestion. *Nature* 451(7182):1069–1075
- Schmid D, Pypaert M, Münz C (2007) MHC class II antigen loading compartments continuously receive input from autophagosomes. *Immunity* 26:79–92
- Dengjel J, Schoor O, Fischer R et al (2005) Autophagy promotes MHC class II presentation of peptides from intracellular source proteins. *Proc Natl Acad Sci USA* 102:7922–7927
- Nedjic J, Aichinger M, Emmerich J, Mizushima N, Klein L (2008) Autophagy in thymic epithelium shapes the T-cell repertoire and is essential for tolerance. *Nature* 455(7211):396–400
- Kasai M, Tanida I, Ueno T et al (2009) Autophagic compartments gain access to the MHC class II compartments in thymic epithelium. *J Immunol* 183(11):7278–7285
- Zhou D, Li P, Lott JM et al (2005) Lamp-2a facilitates MHC class II presentation of cytoplasmic antigens. *Immunity* 22:571–581
- Lee DY, Sugden B (2008) The latent membrane protein 1 oncogene modifies B-cell physiology by regulating autophagy. *Oncogene* 27(20):2833–2842
- Paludan C, Schmid D, Landthaler M et al (2005) Endogenous MHC class II processing of a viral nuclear antigen after autophagy. *Science* 307(5709):593–596
- Leung CS, Haigh TA, Mackay LK, Rickinson AB, Taylor GS (2010) Nuclear location of an endogenously expressed antigen, EBNA1, restricts access to macroautophagy and the range of CD4 epitope display. *Proc Natl Acad Sci USA* 107(5):2165–2170
- Riedel A, Nimmerjahn F, Burdach S et al (2008) Endogenous presentation of a nuclear antigen on MHC class II by autophagy in the absence of CRM1-mediated nuclear export. *Eur J Immunol* 38(8):2090–2095
- Leib DA, Alexander DE, Cox D, Yin J, Ferguson TA (2009) Interaction of ICP34.5 with Beclin 1 modulates herpes simplex virus type 1 pathogenesis through control of CD4⁺ T-cell responses. *J Virol* 83(23):12164–12171
- Lee HK, Mattei LM, Steinberg BE et al (2010) In vivo requirement for Atg5 in antigen presentation by dendritic cells. *Immunity* 32(2):227–239
- Jagannath C, Lindsey DR, Dhandayuthapani S et al (2009) Autophagy enhances the efficacy of BCG vaccine by increasing

- peptide presentation in mouse dendritic cells. *Nat Med* 15(3):267–276
38. Dorfel D, Appel S, Grunebach F et al (2005) Processing and presentation of HLA class I and II epitopes by dendritic cells after transfection with in vitro transcribed MUC1 RNA. *Blood* 105(8):3199–3205
 39. Blander JM, Medzhitov R (2006) Toll-dependent selection of microbial antigens for presentation by dendritic cells. *Nature* 440(7085):808–812
 40. Sanjuan MA, Dillon CP, Tait SW et al (2007) Toll-like receptor signalling in macrophages links the autophagy pathway to phagocytosis. *Nature* 450(7173):1253–1257
 41. Cooney R, Baker J, Brain O et al (2010) NOD2 stimulation induces autophagy in dendritic cells influencing bacterial handling and antigen presentation. *Nat Med* 16(1):90–97
 42. Travassos LH, Carneiro LA, Ramjeet M et al (2009) Nod1 and Nod2 direct autophagy by recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol* 11(1):55–62
 43. Blanchet FP, Moris A, Nikolic DS et al (2010) Human immunodeficiency virus-1 inhibition of immunoamphisomes in dendritic cells impairs early innate and adaptive immune responses. *Immunity* 32:654–669
 44. Li Y, Wang LX, Yang G et al (2008) Efficient cross-presentation depends on autophagy in tumor cells. *Cancer Res* 68(17):6889–6895
 45. Uhl M, Kepp O, Jusforgues-Saklani H et al (2009) Autophagy within the antigen donor cell facilitates efficient antigen cross-priming of virus-specific CD8⁺ T cells. *Cell Death Differ* 16(7):991–1005
 46. Orvedahl A, Alexander D, Tallozy Z et al (2007) HSV-1 ICP34.5 confers neurovirulence by targeting the Beclin 1 autophagy protein. *Cell Host Microbe* 1:23–35
 47. Liang XH, Kleeman LK, Jiang HH et al (1998) Protection against fatal Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. *J Virol* 72(11):8586–8596
 48. Orvedahl A, MacPherson S, Sumpter R Jr et al (2010) Autophagy protects against Sindbis virus infection of the central nervous system. *Cell Host Microbe* 7(2):115–127
 49. Shelly S, Lukinova N, Bambina S, Berman A, Cherry S (2009) Autophagy is an essential component of *Drosophila* immunity against vesicular stomatitis virus. *Immunity* 30(4):588–598
 50. Tallozy Z, Jiang W (2002) Virgin HWt, et al. Regulation of starvation- and virus-induced autophagy by the eIF2alpha kinase signaling pathway. *Proc Natl Acad Sci USA* 99(1):190–195
 51. Alexander DE, Ward SL, Mizushima N, Levine B, Leib DA (2007) Analysis of the role of autophagy in replication of herpes simplex virus in cell culture. *J Virol* 81(22):12128–12134
 52. Chaumorcel M, Souquere S, Pierron G, Codogno P, Esclatine A (2008) Human cytomegalovirus controls a new autophagy-dependent cellular antiviral defense mechanism. *Autophagy* 4(1):46–53
 53. Pattingre S, Tassa A, Qu X et al (2005) Bcl-2 antiapoptotic proteins inhibit Beclin 1-dependent autophagy. *Cell* 122(6):927–939
 54. Lee JS, Li Q, Lee JY et al (2009) FLIP-mediated autophagy regulation in cell death control. *Nat Cell Biol* 11(11):1355–1362
 55. Ku B, Woo JS, Liang C et al (2008) Structural and biochemical bases for the inhibition of autophagy and apoptosis by viral BCL-2 of murine gamma-herpesvirus 68. *PLoS Pathog* 4(2):e25
 56. Xiaofei E, Hwang S, Oh S et al (2009) Viral Bcl-2-mediated evasion of autophagy aids chronic infection of gammaherpesvirus 68. *PLoS Pathog* 5(10):e1000609
 57. Dales S, Eggers HJ, Tamm I, Palade GE (1965) Electron microscopic study of the formation of poliovirus. *Virology* 26:379–389
 58. Jackson WT, Giddings TH Jr, Taylor MP et al (2005) Subversion of cellular autophagosomal machinery by RNA viruses. *PLoS Biol* 3(5):e156
 59. Sir D, Chen WL, Choi J et al (2008) Induction of incomplete autophagic response by hepatitis C virus via the unfolded protein response. *Hepatology* 48(4):1054–1061
 60. Dreux M, Gastaminza P, Wieland SF, Chisari FV (2009) The autophagy machinery is required to initiate hepatitis C virus replication. *Proc Natl Acad Sci USA* 106(33):14046–14051
 61. Lee YR, Lei HY, Liu MT et al (2008) Autophagic machinery activated by dengue virus enhances virus replication. *Virology* 374(2):240–248
 62. Khakpoor A, Panyasrivanit M, Wikan N, Smith DR (2009) A role for autophagolysosomes in dengue virus 3 production in HepG2 cells. *J Gen Virol* 90(Pt 5):1093–1103
 63. Panyasrivanit M, Khakpoor A, Wikan N, Smith DR (2009) Colocalization of constituents of the dengue virus translation and replication machinery with amphisomes. *J Gen Virol* 90(Pt 2):448–456
 64. Kyei GB, Dinkins C, Davis AS et al (2009) Autophagy pathway intersects with HIV-1 biosynthesis and regulates viral yields in macrophages. *J Cell Biol* 186(2):255–268
 65. Zhou Z, Jiang X, Liu D et al (2009) Autophagy is involved in influenza A virus replication. *Autophagy* 5(3):321–328
 66. Gannage M, Dormann D, Albrecht R et al (2009) Matrix protein 2 of influenza A virus blocks autophagosome fusion with lysosomes. *Cell Host Microbe* 6(4):367–380