



Segregation distortion of *Brassica carinata* derived black rot resistance in *Brassica oleracea*

Muhammet Tonguç¹, Elizabeth D. Earle² & Phillip D. Griffiths^{1,*}

¹Department of Horticultural Sciences, Cornell University NYSAES, Geneva, N.Y. 14456, U.S.A.; ²Department of Plant Breeding, Cornell University, 514 Bradfield Hall Ithaca, N.Y. 14853, U.S.A.; (*author for correspondence, e-mail: pdg8@cornell.edu)

Received 1 April 2003; accepted 20 August 2003

Key words: cabbage, Ethiopian mustard, RAPD, *Xanthomonas campestris*

Summary

Three segregating F₂ populations were developed by self-pollinating 3 black rot resistant F₁ plants, derived from a cross between black rot resistant parent line 11B-1-12 and the susceptible cauliflower cultivar 'Snow Ball'. Plants were wound inoculated using 4 isolates of *Xanthomonas campestris* pv. *campestris* (Xcc) race 4, and disease severity ratings of F₂ plants from the three populations were scored. A total of 860 arbitrary oligonucleotide primers were used to amplify DNA from black rot resistant and susceptible F₂ plants and bulks. Eight RAPD markers amplified fragments associated with completely disease free plants following black rot inoculation, which segregated in frequencies far lower than expected. Segregation of markers with black rot resistance indicates that a single, dominant major gene controls black rot resistance in these plants. Stability of this black rot resistance gene in populations derived from 11B-1-12 may complicate introgression into *B. oleracea* genotypes for hybrid production.

Introduction

Black rot is a bacterial disease of *Brassica oleracea* L. caused by *Xanthomonas campestris* pv. *campestris* (Xcc). Under natural conditions Xcc infects through hydathodes, moves through the midrib veins of the leaf and clogs vessels with polysaccharides producing V-shaped lesions associated with the disease (Cook et al., 1952).

Black rot resistance is particularly important in the *Brassica* vegetables (*B. oleracea*) from which several accessions and varieties have been identified as resistant (Bain, 1952; Hunter et al., 1987). However, *B. oleracea* derived resistance is incomplete and controlled by quantitative trait loci (Camargo et al., 1995; Vicente et al., 2002). For effective hybrid production simple genetic control of resistance is desirable. Resistance has been identified in related *Brassica* species (Guo et al., 1991; Westman et al., 1999; Taylor et al., 2002; Tonguç & Griffiths, 2003) including *B. carinata* accession PI 199947 (previously identified as *B. napus*

PI 199947). Black rot resistance derived from the accession has been associated with a single dominant gene, *Xca 1* (Guo et al., 1991; Taylor et al., 2002; Vicente et al., 2002).

Related *Brassica* species can be used to introgress disease resistance and other agronomic and horticultural traits into *B. oleracea* even though chromosome numbers differ. This is due to high tolerance of aneuploidy and common ancestry in diploids. Resistance identified from PI 199947 was used to transfer black rot resistance to *B. oleracea* by protoplast fusion with a rapid-cycling *B. oleracea* followed by backcrosses to *B. oleracea* (Hansen & Earle, 1995; Earle, 1997). Marker studies were previously undertaken with segregating populations derived from resistant somatic hybrids. Reciprocal backcross populations were used to study inheritance of the introgressed chromosomal fragment and to identify molecular markers linked to black rot resistance (Zhou et al., 1997). The proportion of resistant plants was significantly lower than expected for backcross generations and five linked markers

were found. It was suggested that aneuploidy was the cause of distorted segregation, and repeated backcrosses would eliminate extra chromosomal fragments and stabilize the transmission of the resistance.

Molecular markers provide a useful tool for plant breeders to identify resistance genes, and correlate their presence with disease severity symptoms following inoculation. Randomly amplified polymorphic DNA (RAPD) markers utilize random pieces of genomic DNA to create polymorphisms (Williams et al., 1990), and have been used to identify a number of disease resistance genes in different crop species such as pea (Tiwari et al., 1998) and broccoli (Giovannelli et al., 2002).

When evaluating germplasm for resistance to black rot, the race specificity of the resistance becomes very important. Six races of Xcc have been identified, and races 1 and 4 have been found to be the most common races worldwide (Vicente et al., 2001). Resistance to the most common races appeared to be present in several B-genome *Brassicaceae* including *B. carinata*. Resistance derived from PI 199947 provides complete protection against races 1 and 4 of Xcc following wound inoculation (Taylor et al., 2002; Vicente et al., 2002).

The aim of the current study was to identify RAPD markers associated with black rot resistance in *B. oleracea* breeding lines derived from *B. carinata* accession PI 199947 and evaluate their segregation in F₂ populations relative to disease symptoms.

Materials and methods

Plant material

The breeding line 11B-1-12 (Figure 1) was derived from protoplast fusion between susceptible rapid-cycling *B. oleracea* and a resistant *B. carinata* accession PI 199947, followed by a backcross of a resistant somatic hybrid to white-flowered broccoli (Hansen & Earle, 1995). Additional generations of backcrosses and selfs produced line 11B-1-12 (Ren & Earle, unpublished data). Resistant F₁ hybrids were selected following hybridization with the black rot susceptible cauliflower cultivar 'Snow Ball' and self-pollinated to produce segregating F₂ populations. Plants from 3 F₂ populations that segregated resistant plants along with cauliflower controls ('Snow Ball' and 'Delira'), cabbage controls ('Atlantis' and 'Bartolo') and broccoli controls ('Marathon' and 'Titleist') were grown




Figure 1. Generation of segregating F₂ populations and pedigree of the resistant parent.

in plastic pots (10 cm diameter) containing 'Cornell mix' (Boodley & Sheldrake, 1982) in a greenhouse. Plants were grown at 23/20 °C day/night with a 14 h photoperiod under 1000 W metal halide lamps.

DNA isolation

Genomic DNA was isolated from the first true leaf according to Doyle & Doyle (1990). Extracted DNA was quantified with a ThermoSpectronic spectrophotometer (BioMate, Pittsford, N.Y.) and final concentration of DNA was adjusted to 40 ng/μl.

Plant inoculations

Plants were inoculated twice; first at 21 days, and the second at the 56 days using four race 4 Xcc isolates, obtained from D. Reed (Reeds Seeds, Cortland, N.Y.). Isolates were grown on YDCP medium (Shelton & Hunter, 1985) for 2 days and used to wound inoculate the F₂ populations and control plants. The wound inoculation was performed by dipping 2 needles into bacteria and piercing a true leaf with infected needles

Table 1. Disease severity rating scale of Xcc 14 days after wound inoculation

Disease rating	Symptoms
1	No symptoms
2	Minimal symptom development from the point of infection
3	Development of symptoms up to 1 cm ² from the point of infection
4	Symptom development extending to leaf margins
5	Symptom development along leaf margins followed by leaf abscission

Table 2. RAPD markers linked to black rot resistance, their sequences and the sizes of the linked fragments

Primer name	Primer sequence (5'–3')	Fragment size (bp)
OPAB04	GGCACGCGTT	575
UBC 72	GAGCACGGGA	1500
UBC 322	GCCGCTACTA	400
UBC 66	GAGGGCGTGA	500
UBC 205	CGGTTTGAA	900
UBC 121	ATACAGGGAG	360
UBC 320	CCGGCATAGA	450
UBC 327	ATACGGCGTC	300

on either side of the midrib. All four isolates were used to inoculate leaves of the same plants. Plants were rated after 35 days and diseased leaves were removed. Plants were allowed to develop new leaves for 21 days prior to re-inoculation at 56 days and at a second rating at 70 days. After the each inoculation all plants were placed in a 100% humidity mist chamber for 48 h at 20 °C. Plants were rated according to Table 1, and plants with a rating of 1 were considered to be completely resistant.

Nuclear DNA content

Estimation of nuclear DNA content was conducted with a flow cytometry, EPICS Profile Analyzer (Coulter Electronics, Hialeah, Fla.). Samples and standards were prepared and analyzed as described in Arumuganathan & Earle (1991).

RAPD analysis

DNA bulks (Michelmore et al., 1991) were prepared from 8 symptomless and 10 completely susceptible plants following inoculation with black rot in a segregating F₂ population. Arbitrary 10-mer oligonucleotide primers from Operon Technologies Inc., (Alameda,

Calif.) and University of British Columbia (Vancouver, B.C., Canada) were used to amplify DNA to identify RAPD markers associated with symptomless plants (Williams et al., 1990). Each reaction consisted of 8 µl of sterile dH₂O, 4 µl of 5X PCR buffer (0.5 M Tris, pH 8.3, 10 mM MgCl₂, 10 mM Tartrazine, 14% w/v Ficoll), 2 µl of genomic DNA, 2 µl of 0.13 mM of each of the dNTPs, 2 µl of 0.1 mM primer and 2 µl of *Taq* polymerase. One drop of mineral oil was added to the top of each well. PCR reactions were performed using a RoboCycler[®] thermalcycler (Stratagene, La Jolla, Calif.). The PCR cycle parameters were 40 cycles of 1 min denaturation at 94 °C, 1.5 min annealing at 35 °C and 2 min extension at 72 °C. PCR products were separated by gel electrophoresis in 1.5% agarose gels. Electrophoresis was performed in IX TAE buffer for 70 min at 83 V (constant voltage). Gels were stained with ethidium bromide and visualized under UV light with a Gel Doc 2000 system and pictures were digitally recorded with 'Quantity One' software (Bio-Rad Lab., Hercules, Calif.).

Statistical analysis

Association between markers and disease ratings were calculated with Gamma (γ) statistics (Agresti, 1984).

$$\gamma = C-D/C+D,$$

$$\text{where } C = \sum_{i < k} \sum_{j < l} n_{ij}n_{kl}$$

$$\text{and } D = \sum_{i < k} \sum_{j > l} n_{ij}n_{kl}.$$

C and D are the sum of all of the pairwise products of the cells, i = the number of individuals with presence or absence of the markers with a marker score of j , and k and l are the ordered variables defining rows and columns of a cell.

Recombination frequencies were calculated with the formulae: $r = R/N$, where r = recombination fre-




Figure 2. Disease severity ratings for black rot inoculated plants in populations P1 (A), P2 (B) and P3 (C).

quency, R = number of recombinants, and N = total number of individuals tested. The standard error of recombination frequency was estimated as $SE_r = [r(1-r)/N]^{1/2}$ (Adams & Joly, 1980). Mendelian segregation of resistance was investigated with the chi-square goodness-of-fit test.

Results

The majority of plants from the parent line 11B-1-12 were symptomless when needle inoculated; however, some plants were susceptible. The same susceptibility was also observed in some F_1 plants derived from resistant 11B-1-12 parent plants suggesting that the resistance was heterozygous and/or not stable in the F_1 progeny. Three resistant F_1 plants were chosen and

Table 3. Association of RAPD markers with black rot resistant as measured using γ statistics

	P1		P2		P3	
	Rating 1	Rating 2	Rating 1	Rating 2	Rating 1	Rating 2
OPAB04	0.78 ***	0.88 ***	0.89 ***	0.95 ***	1.0 ***	1.0 ***
UBC 72	0.87 ***	1.0 ***	1.0 ***	1.0 ***	0.91 ***	0.86 ***
UBC 322	0.88 ***	1.0 ***	1.0 ***	1.0 ***	1.0 ***	1.0 ***
UBC 66	0.87 ***	1.0 ***	1.0 ***	1.0 ***	0.91 ***	0.86 ***
UBC 205	0.88 ***	1.0 ***	1.0 ***	1.0 ***	0.82 ***	0.74 ***
UBC 121	0.88 ***	1.0 ***	1.0 ***	1.0 ***	0.91 ***	0.86 ***
UBC 320	0.88 ***	1.0 ***	ND	ND	ND	ND
UBC 327	0.92 ***	1.0 ***	1.0 ***	0.69 **	1.0 ***	1.0 ***

** significant at $p = 0.01$, and *** significant at $p = 0.001$, respectively.

ND = No segregation of the marker.

Table 4. Recombination frequencies between RAPD markers and black rot resistance in populations P1, P2 and P3, and χ^2 -values for the deviation from the expected segregation ratio of 3:1

	P1		P2		P3	
	r \pm SE (cM)	χ^2	r \pm SE (cM)	χ^2	r \pm SE (cM)	χ^2
OPAB04	7 \pm 2	205.0***	3 \pm 2	232.3***	0.0	232.2***
UBC 72	3 \pm 2	205.0***	0.0225.3***	1 \pm 0.09	218.6***	
UBC 322	4 \pm 2	205.0***	0.0225.3***	1 \pm 0.09	225.4***	
UBC 66	3 \pm 2	93.0***	0.0225.3***	1 \pm 0.09	218.6***	
UBC 205	3 \pm 2	93.0***	0.0225.3***	2 \pm 1	211.8***	
UBC 121	3 \pm 2	93.0***	0.0225.3***	1 \pm 0.09	211.8***	
UBC 320	3 \pm 2	93.0***	ND	–	ND	–
UBC 327	3 \pm 2	205.0***	0.0225.3***	5 \pm 3	148.2***	

*** significant at $p = 0.001$.

ND = No segregation of the marker.

self-pollinated to produce 3 segregating F_2 populations (P1, P2, P3). The proportions of resistant plants in the F_2 populations significantly deviated from the 3:1 ratio, expected for a single dominant gene in an F_2 population [P1 $\chi^2 = 201.7$ ($p > 0.001$), P2 $\chi^2 = 225.3$ ($p > 0.001$) and P3 $\chi^2 = 239.0$ ($p > 0.001$)]. F_3 progeny derived from self pollinating F_2 plants also failed to meet expected segregation ratios for a single dominant homozygous or heterozygous resistance gene, suggesting multiple gene control, pleiotropy or chromosomal instability causing segregation distortion from expected frequencies. Control cultivars were completely susceptible with slower disease development in broccoli cultivars. Plants initially rated 2–3 after 14 days (Figure 2) showed slower disease development, but were considered susceptible to black rot infection if disease symptoms were allowed additional time to develop.

Flow cytometry analysis was carried out to determine genome size in 11B-1-12. Standard 'Green Comet'

has diploid nuclear DNA content of 1.3 pg. The diploid chromosome number of *B. oleracea* is 18, and DNA content for an average *B. oleracea* chromosome is 0.07 pg. Flow cytometry analysis of resistant plants revealed that those plants also had DNA content of 1.3 pg, which fell into the expected range for $2n = 18$ (data not shown).

A total of 860 Operon and UBC primers were used to screen resistant and susceptible bulks. RAPD primers produced 3 to 15 scoreable fragments with an average of 8.1 fragments per primer. 150 RAPD polymorphisms were identified between resistant and susceptible bulks and used to evaluate population P1. Twenty marker polymorphisms were identified in the P1 population. Of these polymorphisms 8 markers showed strong association with black rot resistance (Table 2). These markers were used to screen all populations, and significant associations between markers and the symptomless phenotype were calculated (Table 3). Recombination frequencies were estimated for resist-




Figure 3. RAPD polymorphism generated by UBC 327 is inherited from *B. carinata* (PI 199947) in resistant plants in F_2 populations.

ant locus and for each marker in the three populations, as well as χ^2 -values (Table 4). All linked markers significantly deviated from the expected 3:1 ratio. All primers were used to amplify donor DNA, and we found that they also amplified the corresponding fragments in PI 199947 (Figure 3). All primers gave clear scoreable fragments in all populations. UBC 320 did not produce any polymorphisms between resistant and susceptible individuals in the populations P2 and P3, and there were differences between the populations for the recombination frequencies of the linked markers. All markers segregated in P1; however, there was no recombination between the black rot resistance and six markers in P2. One marker (OPAB04) did not show any recombination in P3. The highest recombination frequency was observed for the marker OPAB04 in the population P1 (Table 4).

Discussion

The presence of dominant RAPD markers was associated with complete black rot resistance in wound inoculated seedlings. As with the disease severity ratings, the RAPD markers associated with black rot

resistance showed extreme segregation distortion. This suggests that a single dominant resistant gene for black rot resistance is present in these segregating populations, but did not segregate in expected ratios. However it was not possible to confirm monogenic segregation of resistance in F_2 populations due to a distorted segregation. Causes of segregation distortion could include: multiple gene control, pleiotropy, gametic abortion, any chromosomal instability such as aneuploidy or an unstable translocation. The segregation distortion of RAPD markers linked to the resistance in F_2 populations provides strong evidence that multiple gene control, pleiotropy and aneuploidy are not responsible for the distortion of resistance as linked RAPD markers would have segregated in 75% of the F_2 progenies, although presence of a second gene, very closely linked to the *Xca 1* in the genome of *B. carinata*, and/or minor genes are possible. Seed set was also normal indicating that gametes were stable and viable in these populations.

In a previous study Zhou et al. (1997) identified five RAPD primers linked to the black rot resistance; however, none of the reported markers were linked to black rot resistance in this study. Eight markers were linked to symptomless plants in these populations. Association of RAPD fragments and disease rating scores was highly significant (Table 3), and recombination frequencies were generally low. Different recombination frequencies were observed for linked RAPD markers in F_2 populations (Table 4). All markers came from the resistant donor parent and therefore they amplify the B-genome of *B. carinata*. These results suggested that the introgressed genetic material from *B. carinata* might not be stable in *B. oleracea* genotypes.

Resistant plants grown to flowering in the greenhouse produced normal flowers and did not appear to exhibit gametic inviability. When they were self or cross-pollinated, plants produced normal seeds and formed full siliques. No bud pollination was necessary to produce selfed seeds, which suggested that black rot resistant plants were self-compatible. Irregular meiotic behavior might be present in resistant plants as a result of an unstable chromosome translocation that reduces the transmission of resistance in selfed and backcross generations.

Brassica species contain homologous segments within and between genomes resulting from extensive duplications of chromosomal segments and gene ordering (Kianian & Quiros, 1992; Struss et al., 1996; Truco et al., 1996). The B and C *Brassica* genomes

are moderately homologous (Truco et al., 1996), and it is possible to produce interspecific hybrids between *Brassica* species to transfer various traits into cultivated germplasm. In some populations derived from interspecific crosses it is difficult to obtain stable recombinant lines or the transferred resistance can be lost during backcross generations (Plieske et al., 1998). Similar transmission problems have been encountered with other species involving interspecific hybridizations (Heijbroek et al., 1988). One way to circumvent the problem is to obtain plants in which the chromosomal segment carrying the resistance gene of interest is translocated to a recurrent parent chromosome. This process would stabilize the introgressed segment of donor DNA and consequently populations would not exhibit segregation distortion (Heijbroek et al., 1988; Struss et al., 1996).

Loss of linked markers from a previous study (Zhou et al., 1997) suggests that the size of the introgressed segment in these populations is reduced. In the present study, loss of the linked fragment from UBC 320 in the populations P2 and P3 might also indicate that a chromosomal segment has been lost among resistant plants originated from two different F₁ plants. Different recombination rates were observed between the *Xca 1* locus and linked markers in different populations. This result suggests that recombination between B and C-genomes occurs but it was not sufficient to stabilize B-genome derived resistance in a C-genome background.

It will be necessary to stabilize this black rot resistance gene in *B. oleracea* genotypes for it to be used in commercial breeding. One approach could be to observe meiotic behavior of resistant plants and select plants with most regular meiosis to produce selfed and backcross generations (Heijbroek et al., 1988; Chèvre et al., 1997). Another approach might be to order the RAPD markers relative to each other and the resistance locus and follow segregation or loss of RAPD markers in further generations. Radiation is known to cause chromosomal translocations in plants. Irradiation of seeds or floral parts has previously been used to stabilize foreign DNA in cultivated plants (Knott, 1968). It might be possible to use the same approach in *B. oleracea* plants carrying *B. carinata* chromosome segments to stabilize the transmission of the resistance locus.

Acknowledgements

We thank John Barnard for help in statistical analysis in this work, and Jian-ping Ren for the development of resistant parent used in this study. This work was supported by a scholarship for Muhammet Tonguç from the Turkish Ministry of Education and N.Y. State IPM Program.

References

- Adams, W.T. & R.J. Joly, 1980. Linkage relationships among twelve allozyme loci in loblolly pine. *J Hered* 1: 199–202.
- Agresti, A., 1984. *Analysis of Ordinary Data*. Wiley, New York, N.Y.
- Arumugunathan, K. & E.D. Earle, 1991. Estimation of nuclear DNA content of plants by flow cytometry. *Plant Mol Biol Rep* 9: 229–241.
- Bain, D.C., 1952. Reaction of *Brassica* seedlings to black rot. *Phytopathology* 42: 497–500.
- Boodley, J.W. & R. Sheldrake, Jr., 1982. Cornell peat-lite mixes for commercial plant growing. *New York Agr Exp Sta Agr Info Bul* 43.
- Camargo, L.E.A., P.H. Williams & T.C. Osborn, 1995. Mapping of quantitative loci controlling resistance of *Brassica oleracea* to *Xanthomonas campestris* pv. *campestris* in the field and greenhouse. *Phytopathology* 85: 1296–1300.
- Chèvre, A.M., P. Barret, F. Eber, P. Dupuy, H. Brun, X. Tanguy & M. Renard, 1997. Selection of stable *Brassica napus*-*B. juncea* recombinant lines resistant to black leg (*Leptosphaeria maculans*). 1. Identification of molecular markers, chromosomal and genomic origin of the introgression. *Theor Appl Genet* 95: 1104–1111.
- Cook, A.A., J.C. Walker & R.H. Larson, 1952. Studies on the disease cycle of black rot of crucifers. *Phytopathology* 42: 162–167.
- Doyle, J. & J. Doyle, 1990. Isolation of plant DNA from fresh tissue. *Focus* 12: 13–15.
- Earle, E.D., 1997. Crucifer vegetables with resistance to blackrot and *Alternaria* leaf spot. 1996 New York State Vegetable Project Reports Relating to IPM. NYS IPM Publication #121, Cornell Cooperative Extension.
- Giovannelli, J.L., M.W. Farnham, M. Wang & A.E. Strand, 2002. Development of sequence characterized amplified region markers linked to downy mildew resistance in broccoli. *J Amer Soc Hort Sci* 127: 597–601.
- Guo, H., M.H. Dickson & J.E. Hunter, 1991. *Brassica napus* sources of resistance to black rot in crucifers and inheritance of resistance. *HortScience* 26: 1545–1547.
- Hansen, L.N. & E.D. Earle, 1995. Transfer of resistance to *Xanthomonas campestris* pv. *campestris* into *Brassica oleracea* L. by protoplast fusion. *Theor Appl Genet* 91: 1293–1300.
- Heijbroek, W., A.J. Roelands, J.H. de Jong, C. van Hulst, A.H.L. Schoone & R.G. Munning, 1988. Sugar beets homozygous for resistance to beet cyst nematode (*Heterodera schachtii* Schm.), developed from monosomic additions of *Beta procumbens* to *B. vulgaris*. *Euphytica* 38: 121–131.
- Hunter, J.E., M.H. Dickson & J.W. Ludwig, 1987. Sources of resistance to black rot of cabbage expressed in seedlings and adult plants. *Plant Dis* 71: 263–266.

- Kianian, S.F. & C.F. Quiros, 1992. Generation of a *Brassica oleracea* composite map, linkage arrangements among various populations and evolutionary implications. *Theor Appl Genet* 84: 544–554.
- Knott, D.R., 1968. Translocations involving *Triticum* chromosomes and *Agropyron* chromosomes carrying rust resistance. *Can J Genet Cytol* 10: 695–696.
- Michelmore, R.W., I. Paran & R.V. Kesseli, 1991. Identification of markers linked to disease resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA* 88: 9828–9832.
- Plieske, J., D. Struss & G. Röbbelen, 1998. Inheritance of resistance derived from the B-genome of *Brassica* against *Phoma lingam* in rapeseed and the development of molecular markers. *Theor Appl Genet* 97: 929–936.
- Shelton, A.M. & J.E. Hunter, 1985. Evaluation of the potential of the flea beetle *Phyllotreta cruciferae* to transmit *Xanthomonas campestris* pv. *campestris*, causal agent of black rot of crucifers. *Can J Plant Path* 7: 308–310.
- Struss, D., C.F. Quiros, J. Plieske & G. Röbbelen, 1996. Construction of *Brassica* B genome synteny groups based on chromosomes extracted from three different sources by phenotypic, isozyme and molecular markers. *Theor Appl Genet* 93: 1026–1032.
- Taylor, J.D., J. Conway, S.J. Roberts, D. Astley & J.G. Vicente, 2002. Sources and origin of resistance to *Xanthomonas campestris* pv. *campestris* in *Brassica* genomes. *Phytopathology* 92: 105–111.
- Tiwari, K.R., G.A. Penner & T.D. Warkentin, 1998. Identification of coupling and repulsion phase RAPD markers for powdery mildew resistance gene *er-1* in pea. *Genome* 41: 440–444.
- Tonguç, M. & P.D. Griffiths, 2003. Evaluation of *Brassica carinata* accessions for resistance to black rot (*Xanthomonas campestris* pv. *campestris*). *HortScience* (in press).
- Truco, M.J., J. Hu, J. Sadowski & C.F. Quiros, 1996. Inter- and intra-genomic homology of the *Brassica* genomes: implications for their origin and evolution. *Theor Appl Genet* 93: 1225–1233.
- Vicente, J.G., J. Conway, S.J. Roberts & J.D. Taylor, 2001. Identification and origin of *Xanthomonas campestris* pv. *campestris* races and related pathovars. *Phytopathology* 91: 492–499.
- Vicente, J.G., J.D. Taylor, A.G. Sharpe, I.A.P. Parkin, D.J. Lydiate & G.J. King, 2002. Inheritance of race-specific resistance to *Xanthomonas campestris* pv. *campestris* in *Brassica* genomes. *Phytopathology* 92: 1134–1141.
- Westman, A.L., S. Kresovich & M.H. Dickson, 1999. Regional variation in *Brassica nigra* and other weedy crucifers for disease reaction to *Alternaria brassicicola* and *Xanthomonas campestris* pv. *campestris*. *Euphytica* 106: 253–259.
- Williams, J.G.K., A.R. Kubelik, K.J. Livak, J.A. Rafalski & S.V. Tingey, 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 18: 6531–6535.
- Zhou, Z., N.F. Weeden & M.H. Dickson, 1997. The expression of a resistant gene to black rot in progeny of the protoplast fusion broccoli (*B. oleracea*). *Cruciferae Newsl* 19: 109–110.