

Pediatr Cardiol (2006) 27:652–654
DOI 10.1007/s00246-005-6001-2

IMAGES IN PEDIATRIC CARDIOLOGY

Cantrell's Syndrome *Forme Fruste* in a Newborn Diagnosed by Transthoracic Echocardiography and Cardiac Magnetic Resonance Imaging

W. Knirsch · A. Dodge-Khatami · D. Bolz ·
E. Valsahgiacomo Büchel

© Springer Science+Business Media, Inc. 2005

A newborn was referred to our hospital because of prominent cardiac pulsations in the subxyphoid region associated with supraumbilical midline cutaneous lesions (Fig. 1). Echocardiography revealed mesocardia with a ventricular septal defect, a secundum atrial septal defect, and an unroofed coronary sinus. The morphological correlate for the pulsatile tumor was a left ventricular diverticulum diagnosed by transthoracic echocardiography (Fig. 2). Cardiac magnetic resonance imaging (MRI) confirmed the presence of a left ventricular diverticulum originating from the apex of the heart and ruled out the presence of a diaphragmatic hernia (Fig. 3).

Cardiac surgical repair, performed at 4 weeks of age, confirmed the preoperative findings (Fig. 4). The left ventricular diverticulum was a continuation of the apex of the left ventricle and was totally covered by pericardium. We found a midline muscular defect of the abdominal wall and a missing linea alba, as well as a

missing xyphoid, but no evidence of diaphragmatic hernia or additional midline malformations.

This represents a *forme fruste* of Cantrell's Syndrome. Cantrell's pentalogy includes a midline, supraumbilical abdominal wall defect, a defect of the lower sternum, a deficiency of the anterior diaphragm, a defect in the diaphragmatic pericardium, and congenital intracardiac defects [1]. In this case, the definition of the topographic localization of the diverticulum performed by MRI and of its relation to the diaphragm was essential in planning the surgical repair.

References

1. Cantrell JR, Haller JA, Ravitch MM (1958) A syndrome of congenital heart defects involving the abdominal wall, sternum, diaphragm, pericardium and heart. *Surg Gynecol Obstet* 107:602–614

W. Knirsch (✉) · E. V. Büchel
Division of Pediatric Cardiology, University Children's
Hospital Zurich, Steinwiesstr-75, CH 8032, Zurich,
Switzerland
E-mail: walter.knirsch@kispi.unizh.ch


A. Dodge-Khatami
Division of Cardiovascular surgery, University Children's
Hospital Zurich, Switzerland

D. Bolz
Division of Pediatric Cardiology, University Children's
Hospital Basel, Switzerland

Fig. 1 Preoperative status showing the subxyphoid-to-supraumbilical covered midline defect of the abdominal wall



Fig. 2 (Left) Two-dimensional echocardiogram of the left ventricular diverticulum (*D*, arrows) and small left ventricle (*LV*). **(Right)** Color Doppler showing blood flow (*red*) into the diverticulum (*D*)






Fig. 3 Contrast-enhanced MR angiography demonstrating the left ventricular diverticulum (*D*) with “vermiform” structure originating from the apex of the left ventricle (*LV*) and its relationship to the diaphragm (*asterisks*) (lateral projection)

Fig. 4 Intraoperative situs of the left ventricular diverticulum (*D*). *LV*, left ventricle, *RV*, right ventricle

