

Massimiliano M. Siccoli
Philipp O. Valko
Dirk M. Hermann
Claudio L. Bassetti

Central periodic breathing during sleep in 74 patients with acute ischemic stroke – Neurogenic and cardiogenic factors

Received: 6 November 2007
Received in revised form: 20 March 2008
Accepted: 15 April 2008
Published online: 13 November 2008

■ **Abstract** *Objectives* The aims of our study were 1) to better characterize central periodic breathing during sleep (CPBS) and its clinical relevance in acute stroke, 2) to better define the role of brain damage in its pathogenesis. *Methods* We included 74 consecutive patients admitted within 96 hours after stroke onset. Stroke severity at admission, stroke outcome at discharge and stroke topography were assessed. ECG and transesophageal echocardiography were performed. Nocturnal breathing was assessed with an ambulatory device the first night after admission. CPBS severity was represented as absolute time and percentage of recording time. *Results* Age was 63 ± 13 (25–82), 49 (66%) were male. Thirty (41%) patients showed CPBS during $\geq 10\%$ and 7 (9%) during $\geq 50\%$ of recording time. CPBS severity was associated with age ($p = 0.017$), stroke severity

($p = 0.008$), ECG abnormalities ($p = 0.005$) and lower left ventricular ejection fraction ($p < 0.0001$). CPBS severity was higher in patients with extensive hemispheric strokes ($n = 6$, $p < 0.0001$), and lower in patients with partial strokes involving the left insula ($n = 5$, $p < 0.0001$) and the mesencephalon ($n = 5$, $p = 0.002$). *Conclusions* CPBS is frequent in acute ischemic stroke and is associated with older age, stroke severity/extension, and lower left ventricular function. The lower occurrence of CPBS in left insular and mesencephalic stroke suggests a major role of distinct brain areas in the modulation of respiratory phenomena accompanying acute stroke.

■ **Key words** Cheyne-Stokes respiration · ischemic stroke · sleep · sleep apnea · left ventricular function · autonomic nervous system · insula

M. M. Siccoli, MD · P. O. Valko, MD ·
D. M. Hermann, MD ·
Prof. C. L. Bassetti, MD (✉)
Dept. of Neurology
University Hospital of Zurich
Frauenklinikstrasse 26
8091 Zurich, Switzerland
Tel.: +41-44/255-5503
Fax: +41-44/255-4649
E-Mail: claudio.bassetti@usz.ch

Introduction

Central periodic breathing (CPB) – or Cheyne-Stokes breathing, characterized by cyclic fluctuations in breathing drive in a gradual waxing-and-waning fashion [1] – results from a variety of conditions affecting the neuronal network involved in respiratory control.

CPB during wakefulness and/or sleep have been reported in patients with brainstem or bilateral extensive hemispheric stroke and impaired level of consciousness

[2–4]. More recent studies in stroke patients report a variable prevalence (6–38%) for CPB during sleep (CPBS) [4–8], a spontaneous recovery of this breathing pattern after stroke [6, 9, 10] and an association with large ischemic lesions and poor functional outcome [11, 12].

In both patients with and without stroke, CPBS has been linked to increased respiratory CO_2 sensitivity [13, 14] and left ventricular dysfunction [15, 16]. CPBS has also been reported in patients with unilateral lesions of variable topography without disturbed level of con-

sciousness or overt heart failure [4, 10, 17]. Although these observations clearly suggest a major role of distinct brain areas in the regulation of respiratory response to ischemic brain damage, the mechanisms leading to CPBS in association with a new-onset ischemic brain lesion remain poorly known.

Following a preliminary observation on three patients (University Hospital of Berne) [10], we performed a systematic and prospective assessment of CPBS in a new larger series of patients with the aim 1) to better characterize this breathing pattern and its clinical relevance in acute stroke and 2) to better define the role of brain damage in its pathogenesis.

Methods

■ Patients

Consecutive adult patients with first-ever acute ischemic stroke were prospectively included. The study design was approved by the local ethical committee and written informed consent was obtained from patients or relatives. Patients with very severe stroke (National Institutes of Health Stroke Scale [18] [NIHSS] > 20), intracerebral/subarachnoid hemorrhage, coma/stupor, or life-threatening medical conditions were excluded.

■ Clinical and radiological evaluation

Cardiovascular risk factors including family history, arterial hypertension (defined as BP > 140/90 mmHg measured ≥ 3 times prior to stroke), diabetes (defined as fasting glucose level > 140 mg/dl), smoking status, hypercholesterolemia (defined as cholesterol level > 250 mg/dl), obesity (defined as body mass index (BMI) ≥ 25 kg/m²) and previous history of heart failure/coronary heart disease were assessed.

Clinical stroke assessment included NIHSS at admission and modified Rankin disability Scale [19] at hospital discharge.

An ECG was performed at admission, Holter ECG and transesophageal echocardiography during the hospitalization.

Stroke topography was assessed in the acute phase according to clinical and radiological criteria by CT scan examination and/or MR imaging. A detailed topographic description of the ischemic region according to clinically identifiable subtypes [20] and vascular territories [21] was performed.

■ Nocturnal breathing

Overnight respirography was performed with a validated [9] portable device (Autoset® Embletta PDS, ResMed) during the first night after admission. The recording was considered to be successful if ≥ 60 minutes with nasal flow and at least one chest band signal were present. The analysis was performed automatically and corrected manually according to previously described standard criteria [5]. Chaotic signals automatically scored from the computer as events were carefully checked manually, and, if not clearly classifiable, were not considered as respiratory events and excluded. Apnea was defined by a cessation of oro-nasal airflow ≥ 10 seconds, hypopnea by a reduction of oro-nasal airflow ≥ 10 seconds by $\geq 50\%$, or $\geq 30\%$ when associated with an oxygen desaturation $\geq 4\%$. Apnea hypopnea index (AHI) was defined by the mean number of apneas and hypopneas per hour, apnea index (AI) by the mean number of apneas per hour between lights off and on. Sleep apnea was defined by AHI ≥ 10 /h, mild by AHI 10–30/h,

moderate-severe by AHI > 30/h. A differentiation between obstructive (OAI) and central (CAI) apnea index was undertaken according to standard criteria. The percentage of time with oxygen saturation < 90%, as well as the oxygen desaturation index (ODI) were calculated. The magnitude of the oxygen desaturation considered for ODI was $\geq 4\%$. Body position was not monitored. CPBS was defined as ≥ 3 cycles of regular crescendo-decrescendo breathing associated with reduction of $\geq 50\%$ in nasal airflow and respiratory effort lasting ≥ 10 seconds [1, 22, 23]. A cycle was defined from the end of one apnea/hypopnea to the end of the next. The periodic pattern of CPBS had to be recognized visually in both the nasal pressure and the respiratory effort signal in order to prevent over-interpretation of the data. Episodes with reduced airflow but clearly maintained respiratory effort were classified as obstructive, and not as central periodic breathing. Severity of CPBS was represented as absolute time and as percentage of total recording time. No patients received oxygen or sedative medication in the hours before or during the recording.

■ Statistics

Statistical analysis was performed with SPSS software package 12.0. Continuous data were presented as mean and standard deviation, categorical data as numbers and percentage. Student's unpaired t-tests were used for comparison of patients groups and for testing of continuous variables. Categorical data were compared with the Mann-Whitney rank-sum tests or the χ^2 tests. Correlations were calculated using the Pearson or Spearman tests as appropriate. A p-value < 0.05 was considered to be statistically significant. Multiple stepwise regression models with entry value set to 0.05 and removal value set to 0.1 in the univariate analysis were tested using stroke outcome (mRS) as dependent variable. Duration and severity of CPBS and stroke severity at admission (NIHSS) were candidate predictors.

Results

■ Clinical data

Seventy-four consecutive adult patients with first-ever acute ischemic stroke were included, three underwent systemic thrombolysis. The main characteristics of all patients are summarized in Table 1.

■ Cardiological data

ECG abnormalities (Sinus brady-/tachycardia, ST-segment/T-waves abnormalities, AV-blocks of different degree) were found in 41 (55%) patients, 10 patients (13%) had atrial fibrillation. Transesophageal echocardiography was performed in 55 (74%) patients, left ventricular ejection fraction (LVEF) was $61 \pm 9\%$ [33–80]. Eighteen (33%) of these patients had a LVEF < 60%, six (11%) of < 50%.

■ Radiological data

Stroke was localized on the right side in 40 (54%) patients and on the left side in 34 (46%) patients. Sixty-one (82%) patients had an hemispheric stroke and 13 pa-

Table 1 Main results of all patients

	All (n = 74)
Male/female (%)	49/25 (66/34)
Age (years)	63 ± 13 [25–82]
Body mass index (kg/m ²)	28 ± 4 [20–39]
BMI > 25 (%)	26 (35)
Arterial hypertension	36 (49)
Positive family history for vascular diseases	37 (50)
Diabetes	40 (54)
Smoking	31 (42)
Hypercholesterinemia	30 (40)
Coronary heart disease/heart failure	38 (51)
NIH Stroke Scale on admission	6 ± 5 [0–20]
Modified Rankin Disability Scale at discharge	2 ± 1 [0–5]
Apnea–hypopnea index (/h)	20 ± 20 [0–82]
Apnea index (/h)	10 ± 14 [0–70]
Obstructive apnea index (/h)	3 ± 7 [0–57]
Central apnea index (/h)	7 ± 12 [0–61]
Oxygen desaturation index (/h)	17 ± 20 [0–84]
Total recording time (min)	598 ± 150 [0–776]
CPBS duration (min)	102 ± 155 [0–651]
CPBS severity, % of recording time (min)	16 ± 22 [0–90]

Values are mean ± SD [range in brackets] or numbers (% in brackets)
 CPBS central periodic breathing during sleep

tients (18%) a brainstem stroke. In 55 (74%) patients the ischemic lesion was localized in the middle cerebral artery territory, in six patients (8%) in the posterior cerebral artery territory. There were no strokes in the anterior cerebral artery territory. Stroke topographies and the corresponding respiratory parameters and CPBS severity are shown in Table 3.

Table 2 Selection of relevant demographic, cardiological and clinical stroke parameters in all patients with CPBS and in two subgroups of patient with CPBS severity of different degree compared to patients without CPBS

n = 74	No CPBS (n = 21)	CPBS (n = 53)	CPBS 10–50% (n = 23)	CPBS ≥ 50% (n = 7)
Age (years)	53 ± 15 [25–82]	66 ± 10 [45–82]*	67 ± 11 [47–82]*	69 ± 10 [52–79]*
History of hypertension	14 (67)	23 (44)	9 (39)*	3 (43)
Diabetes	9 (43)	30 (57)	12 (52)	6 (86)
CAD/HF	7 (37)	29 (55)	12 (52)	5 (71)
NIH Stroke Scale day 1	6 ± 4 [0–17]	6 ± 5 [1–20]	5 ± 5 [1–20]	11 ± 8 [4–20]
mRS at discharge	2 ± 2 [0–4]	2 ± 1 [0–5]	2 ± 1 [0–4]	3 ± 2 [0–5]
LVEF	65 ± 8 [55–80]	60 ± 10 [33–77]	56 ± 9 [33–70]*	50 ± 4 [45–55]*
ECG abnormalities	7 (33)	33 (62)*	17 (74)*	5 (71)
AF on admission	2 (10)	7 (13)	4 (17)	2 (29)

Values are means ± SD [ranges in brackets] or numbers (% in brackets)

CPBS central periodic breathing during sleep; CAD/HF history of coronary heart disease or heart failure; mRS modified Rankin disability scale; LVEF left ventricular ejection fraction; ECG abnormalities electrocardiographic abnormalities at admission; CPBS central periodic breathing during sleep

* numbers in bold = statistically significant (p < 0.05) compared to patients without CPBS

■ Respirographic data

Respirography was performed 45 ± 26 hours [4–96] after stroke onset. CPBS was found in 53 (72%) patients, in 30 (41%) patients during ≥ 10% and in seven (9%) patients during ≥ 50% of recording time, by a mean total recording time of 598 ± 150 minutes [0–776]. The main respiratory results are summarized in Table 1. Sleep apnea (defined as AHI > 10/h) was found in 41 (55%) patients (AHI 33 ± 19/h [11–82], OAI 5 ± 9/h [0–57], CAI 12 ± 14/h [0–61], ODI 27 ± 21/h [5–84]). Severity of CPBS correlated with severity of sleep apnea (r = 0.755, p < 0.0001), central apneas (r = 0.798, p < 0.0001), and oxygen desaturations (r = 0.661, p < 0.0001).

Patients with CPBS were older, had a higher occurrence of ECG abnormalities on admission, and a lower LVEF compared to patients without CPBS (Table 2). Accordingly, severity of CPBS correlated with age (r = 0.280, p = 0.017), stroke severity at admission (r = 0.309, p = 0.009), occurrence of ECG abnormalities (r = 0.337, p = 0.005), and lower LVEF (r = -0.542, p < 0.0001). Conversely, there was no significant correlation between the pre-stroke cardiovascular risk factors (positive family history for cardiovascular disease, arterial hypertension, diabetes, smoking, hypercholesterinemia, obesity, history of heart failure/coronary heart disease) and CPBS severity. No significant correlation between CPBS severity and stroke outcome at discharge was found after taking into account stroke severity on admission.

Concerning stroke topography, CPBS was much more frequent/severe in total anterior circulation strokes (TACS), especially left-sided (p < 0.0001). CPBS was more frequent in patients with stroke in the frontal/pre-frontal region, deep MCA territory, and thalamic/hypothalamic region, the difference in CPBS severity compared to other stroke topographies did not reach however statistical significance. Patients with stroke in the mes-

Table 3 Nocturnal breathing parameters and stroke topography (selection of most representative localizations regarding CPBS severity)

Stroke topography	N=74	AHI (/h)	CAI (/h)	CPBS%
TACS	6 (8)	41 ± 35	17 ± 23	54 ± 39*
PACS	42 (57)	18 ± 17	7 ± 11	12 ± 17
LACS	7 (9)	27 ± 19	12 ± 10	23 ± 25
POCS	19 (26)	16 ± 19	3 ± 7	13 ± 21
Frontal/precentral region (MCA)	10 (14)	30 ± 20	17 ± 16	19 ± 19
Deep MCA	18 (24)	20 ± 16	8 ± 10	18 ± 23
Right insula	5 (7)	17 ± 12	8 ± 7	12 ± 8
Left insula	5 (7)	4 ± 5*	1 ± 1*	0 ± 0*
Thalamus/hypothalamus	3 (4)	18 ± 14	5 ± 6	27 ± 20
Pons	5 (7)	33 ± 24	4 ± 5	12 ± 10
Mesencephalon	5 (7)	7 ± 2*	1 ± 1*	4 ± 5*

Values are mean ± SD or numbers (% in brackets)

AHI Apnea–hypopnea index; CAI central apnea index; CPBS% central periodic breathing in sleep, % of total recording time

* numbers in bold = statistically significant ($p < 0.05$) compared to stroke with different topography

encephalic ($n=5$) and left insular region ($n=5$) had a significantly lower severity of sleep apnea (both topographies $p < 0.0001$), central apneas (both topographies $p < 0.0001$), and CPBS (mesencephalic $p = 0.002$, left insular region $p < 0.0001$, respectively) compared to patients with other stroke localizations, despite the absence of significant differences in stroke severity, cardiovascular risk factors and left ventricular function. A summary of sleep apnea/CPBS severity according to stroke topography is reported in Table 3.

Discussion

The aim of our study was to describe characteristics and clinical relevance of CPBS in acute ischemic stroke, and to better define the role of brain damage in its pathogenesis. Our main results are:

- CPBS is frequent in the first few days after acute ischemic stroke.
- CPBS is associated with a) older age, b) ECG abnormalities and c) lower left ventricular function.
- CPBS occurs more frequently in patients with (left-sided) severe/extensive hemispheric stroke, and less frequently in patients with stroke involving the left insular or mesencephalic region.

■ Frequency of CPBS in acute ischemic stroke

CPBS was found in 53 (72%) patients, in 30 (41%) during $\geq 10\%$ of recording time. In previous studies investigating breathing disorders during sleep in the early stroke phase (≤ 72 hours after stroke onset), performed

with different portable devices [6, 11, 12, 24] or with full polysomnography [7], CPBS was observed in 23–38% of patients [7, 11, 12, 24], and was present during $\geq 10\%$ of recording time in 12–28% of patients [6, 24]. In contrast, in the studies performed with full polysomnography during the subacute stroke phase (after a mean time of 9 to 44 days after stroke) [4, 5, 16], lower frequencies of CPBS (6–19%) were reported. Parra et al. [6] showed an improvement of CPBS after 3 months. Only in one study performed with full polysomnography [8] after a mean time of 12 days after stroke CPBS was documented during $\geq 10\%$ of recording time in 37% of patients. However, this higher frequency of CPBS might be explained by the fact that 70% of these patients underwent polysomnography early (< 10 days) following stroke and that $> 50\%$ of them had coronary heart disease or congestive heart failure.

Thus, the high frequency of CPBS in our study may be in part explained with the short latency between stroke onset and respirography. In addition, the methodology used in the recording of nocturnal breathing (highly sensitive XactTrace sensor) may have contributed to these results. The left ventricular function was similar or higher in our series compared to previous studies [4, 16] and does not explain the higher CPBS frequency found. Our patients were in part younger and had milder strokes compared to previous studies [4–7, 11, 12, 16, 24] and to a general stroke population [25, 26] despite the absence of major differences in cardiovascular risk profile or stroke topography. However, considering that CPBS has been described in association with large and severe strokes [11, 12], the younger age and the milder stroke severity are very unlikely to explain the higher CPBS frequency observed in our series.

■ CPBS and left ventricular function

We found an association of CPBS with age, ECG abnormalities and lower left ventricular function in patients with acute ischemic stroke. In our series, although only six (11%) out of 55 patients undergoing echocardiography had a LVEF $< 50\%$, we observed a significant inverse linear correlation between CPBS severity and LVEF. Our results suggest an association between CPBS severity and lower LVEF even in the absence of a cardiac dysfunction. Similarly, Nopmaneejumruslers et al. [16] found a significant association between CPBS and reduced ($< 40\%$) LVEF after stroke, but also a normal LVEF in 14 patients with CPBS, which could not be explained by the authors.

■ CPBS and stroke topography/type

In line with previous studies [4, 11, 12, 17], we found that CPBS is much more frequent and severe in patients with TACS. CPBS was also more pronounced in strokes localized in distinct brain areas (Table 3), although the difference to other topographies did not reach statistical significance. This observation, especially if extensive, confirms that brain lesions involving autonomic (insula) or volitional (prefrontal region, capsula interna, thalamus) structures participating in respiratory control may favour CPBS after stroke [10].

The most interesting, to our best knowledge unreported observation was the considerably lower frequency of central apneas and CPBS in patients with circumscribed (not extensive) stroke involving the left insula and the mesencephalon despite the absence of a significant difference in stroke severity, demographics, cardiovascular risk profile and left ventricular function compared to other topographies.

The insula is with the amygdala, the hypothalamus, the periaqueductal grey (mesencephalon), and the ventrolateral medulla part of the central autonomic network [27], which plays a central role in the integration and processing of autonomic and behavioural information essential for survival [28]. Severe cardiovascular and autonomic disturbances have been reported in human and animals in association with insular lesions including stroke, predominantly on the right side [29–33]. Otherwise, left-sided insular strokes lead rather to a disturbed cardioinhibitory/cardioregulatory function with an autonomic shift toward a pro-arrhythmic condition [29].

Hence, we suggest – in analogy – that a shift in the autonomic (sympathetic-parasympathetic) balance may be responsible for the lower frequency of central apneas and CPBS in patients with circumscribed left insular (and mesencephalic) strokes. This hypothesis is supported by recent observations showing functional changes in brain areas responsible for autonomic respiratory control (including insula) in patients with sleep-disordered breathing [34].

Interestingly, extensive left hemispheric strokes (also involving the insula) presented with severe CPBS. The considerable variability in CPBS severity observed in patients with stroke of different extension involving the left insula may be in part explained by the complex somatotopic representation of the respiratory-related area in the insular cortex [35] and by the disruption of the insular afferent/efferent autonomic regulatory pathways, leading – depending on its extension – to different changes in the respiratory patterns. However, probably due to our assessment methods, we could not identify further findings supporting this hypothesis. Obviously, considering the small sample size in the mentioned topographic subgroups, a type 1 error cannot be excluded. The suggested associations between stroke in distinct

brain areas and severity of CPBS shall be regarded as an hypothesis, and need to be confirmed in further studies including a larger number of patients.

■ CPBS and stroke severity/outcome

Previous studies showed an association between CPBS, worse functional outcome, and increased mortality after stroke [12]. In our study we did not find – possibly due to the sample size and short observation time – this prognostic association after accounting for the initial stroke severity. Our results show that patients with extensive unilateral hemispheric brain damage were more prone to develop central apneas and CPBS in the acute stroke phase. Whether CPBS may per se lead to a worse stroke evolution or (rather) represent a consequence of stroke related to its severity and extension remains however still unclear. Considering that our patients mostly presented with mild-moderate stroke, extensive hemispheric strokes (TACS) might be underrepresented in our series. This may be considered, along with the younger age of the patients compared to previous studies and to a general stroke population, as a possible limitation of the study.

The fact that no P_{CO_2} measurement during the acute phase and no follow-up respirographies have been performed should be mentioned as further limitation of our study. These examinations might provide further useful information in understanding the pathophysiology of CPBS in acute stroke.

Our observations strongly emphasize the role of a central neurogenic component – in addition to the cardiac and respiratory component – and the differential role of distinct brain areas in the pathogenesis of CPBS following acute brain damage, showing the tight link existing between brain, cardiovascular and respiratory system.

Further studies including a larger number of patients, P_{CO_2} measurements, follow-up respirographies and an assessment of autonomic function (or of its surrogate markers) are needed to confirm our observations and to better define the pathophysiological background and the clinical implications of these findings.

■ **Conflict of interest** The authors have no conflict of interest.

■ **Acknowledgements** C. Selic, MD, and S. Birkmann, MD, Department of Neurology, University Hospital of Zurich, assisted in performing patient examinations, recording and scoring of respirographies. E. Werth, PhD, and R. Poryazova, MD, Department of Neurology, University Hospital of Zurich assisted in the recording and scoring of respirographies.

References

- No authors listet (1999) Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. The Report of an American Academy of Sleep Medicine Task Force. *Sleep* 22: 667–689
- Plum F, Swanson AG (1959) Central neurogenic hyperventilation in man. *AMA Arch Neurol Psychiatry* 81: 535–549
- Lee MC, Klassen AC, Resch JA (1974) Respiratory pattern disturbances in ischemic cerebral vascular disease. *Stroke* 5:612–616
- Bassetti C, Aldrich MS, Quint D (1997) Sleep-disordered breathing in patients with acute supra- and infratentorial strokes. A prospective study of 39 patients. *Stroke* 28:1765–1772
- Bassetti C, Aldrich MS (1999) Sleep apnea in acute cerebrovascular diseases: final report on 128 patients. *Sleep* 22:217–223
- Parra O, Arboix A, Bechich S, et al. (2000) Time course of sleep-related breathing disorders in first-ever stroke or transient ischemic attack. *Am J Respir Crit Care Med* 161:375–380
- Iranzo A, Santamaria J, Berenguer J, Sanchez M, Chamorro A (2002) Prevalence and clinical importance of sleep apnea in the first night after cerebral infarction. *Neurology* 58:911–916
- Bassetti C, Aldrich MS, Chervin RD, Quint D (1996) Sleep apnea in patients with transient ischemic attack and stroke: a prospective study of 59 patients. *Neurology* 47:1167–1173
- Bassetti CL, Milanova M, Gugger M (2006) Sleep disordered breathing and acute stroke: diagnosis, risk factors, treatment, evolution and outcome. *Stroke* 37:967–972
- Hermann DM, Siccoli M, Kirov P, Gugger M, Bassetti CL (2007) Central periodic breathing during sleep in acute ischemic stroke. *Stroke* 38: 1082–1084
- Rowat AM, Wardlaw JM, Dennis MS (2007) Abnormal breathing patterns in stroke: relationship with location of acute stroke lesion and prior cerebrovascular disease. *J Neurol Neurosurg Psychiatry* 78:277–279
- Rowat AM, Dennis MS, Wardlaw JM (2006) Central periodic breathing observed on hospital admission is associated with an adverse prognosis in conscious acute stroke patients. *Cerebrovasc Dis* 21:340–347
- Solin P, Roebuck T, Johns DP, Walters EH, Naughton MT (2000) Peripheral and central ventilatory responses in central sleep apnea with and without congestive heart failure. *Am J Respir Crit Care Med* 162:2194–2200
- Javaheri S (1999) A mechanism of central sleep apnea in patients with heart failure. *N Engl J Med* 341: 949–954
- Sin DD, Fitzgerald F, Parker JD, Newton G, Floras JS, Bradley TD (1999) Risk factors for central and obstructive sleep apnea in 450 men and women with congestive heart failure. *Am J Respir Crit Care Med* 160:1101–1106
- Nopmaneejumrulsers C, Kaneko Y, Hajek V, Zivanovic V, Bradley TD (2005) Cheyne-Stokes Respiration in Stroke: Relationship to Hypocapnia and Occult Cardiac Dysfunction. *Am J Respir Crit Care Med* 171:1048–1052
- Nachtmann A, Siebler M, Rose G, Sitzer M, Steinmetz H (1995) Cheyne-Stokes respiration in ischemic stroke. *Neurology* 45:820–8821
- Brott T, Adams HP Jr, Olinger CP, et al. (1989) Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 20:864–870
- Rankin J (1957) Cerebral vascular accidents in patients over the age of 60. II. Prognosis. *Scott Med J* 2:200–215
- Bamford J, Sandercock P, Dennis M, Burn J, Warlow C (1991) Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 337:1521–1526
- Tatu L, Moulin T, Bogousslavsky J, Duvernoy H (1998) Arterial territories of the human brain: cerebral hemispheres. *Neurology* 50:1699–1708
- Meoli AL, Casey KR, Clark RW, et al. (2001) Hypopnea in sleep-disordered breathing in adults. *Sleep* 24:469–470
- Kushida CA, Littner MR, Morgenthaler T, et al. (2005) Practice parameters for the indications for polysomnography and related procedures: an update for 2005. *Sleep* 28:499–521
- Turkington PM, Bamford J, Wanklyn P, Elliott MW (2002) Prevalence and predictors of upper airway obstruction in the first 24 hours after acute stroke. *Stroke* 33:2037–2042
- Rothwell PM, Coull AJ, Giles MF, et al. (2004) Change in stroke incidence, mortality, case-fatality, severity, and risk factors in Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet* 363:1925–1933
- Schulz UG, Rothwell PM (2003) Differences in vascular risk factors between etiological subtypes of ischemic stroke: importance of population-based studies. *Stroke* 34:2050–2059
- Benarroch EE (1993) The central autonomic network: functional organization, dysfunction, and perspective. *Mayo Clin Proc* 68:988–1001
- King AB, Menon RS, Hachinski V, Cechetto DF (1999) Human forebrain activation by visceral stimuli. *J Comp Neurol* 413:572–582
- Oppenheimer SM, Kedem G, Martin WM (1996) Left-insular cortex lesions perturb cardiac autonomic tone in humans. *Clin Auton Res* 6:131–140
- Christensen H, Boysen G, Christensen AF, Johannesen HH (2005) Insular lesions, ECG abnormalities, and outcome in acute stroke. *J Neurol Neurosurg Psychiatry* 76:269–271
- Oppenheimer S (2006) Cerebrogenic cardiac arrhythmias: cortical lateralization and clinical significance. *Clin Auton Res* 16:6–11
- Laowattana S, Zeger SL, Lima JA, Goodman SN, Wittstein IS, Oppenheimer SM (2006) Left insular stroke is associated with adverse cardiac outcome. *Neurology* 66:477–483
- Ay H, Koroshetz WJ, Benner T, et al. (2006) Neuroanatomic correlates of stroke-related myocardial injury. *Neurology* 66:1325–1329
- Henderson LA, Macey KE, Macey PM, Woo MA, Yan-Go FL, Harper RM (2006) Regional brain response patterns to Cheyne-Stokes breathing. *Respir Physiol Neurobiol* 150:87–93
- Aleksandrov VG, Aleksandrova NP, Bagaev VA (2000) Identification of a respiratory related area in the rat insular cortex. *Canadian journal of physiology and pharmacology* 78:582–586