



## Improvement of cassava shoot organogenesis by the use of silver nitrate *in vitro*

Peng Zhang<sup>1</sup>, Salak Phansiri<sup>2</sup> & Johanna Puonti-Kaerlas<sup>1,\*</sup>

<sup>1</sup>Institute for Plant Sciences, ETH-Zentrum, CH-8092 Zürich, Switzerland; <sup>2</sup>Central Lab, Kasetsart University Research and Development Institute, Kasetsart University, Chatuchak, Bangkok 10900, Thailand (\*requests for offprints and present address: European Patent Office, D-80298 Munich, Germany. E-mail: puontik@yahoo.com)

Received 11 April 2000; accepted in revised form 24 April 2001

**Key words:** *Manihot esculenta* (Crantz), regeneration, selection, silver nitrate

### Abstract

To improve the regeneration efficiency of cassava (*Manihot esculenta* Crantz) *in vitro*, the effect of silver nitrate (AgNO<sub>3</sub>) on shoot organogenesis from somatic cotyledons was assessed. Adding AgNO<sub>3</sub> to the regeneration medium improved the regeneration frequency and reduced callus formation in all tested cultivars. Both the extent of the response to and the optimum concentration of AgNO<sub>3</sub> were cultivar dependent. In the model cultivar MCol22, the use of AgNO<sub>3</sub> at concentrations between 4 and 12 mg l<sup>-1</sup> increased shoot organogenesis frequency and the number of shoot primordia per explant, the maximum effect being observed on a medium containing 12 mg l<sup>-1</sup> AgNO<sub>3</sub>. At this concentration, the frequency of shoot organogenesis was enhanced from 60 to 90%, while callus formation decreased from 100 to 5%. The highest shoot organogenesis rates were obtained by supplementing the medium with 2 and 1 mg l<sup>-1</sup> AgNO<sub>3</sub> in KU50 and Hanatee, respectively, while cultivar T5 showed an optimum response at 4 mg l<sup>-1</sup>. The shoots regenerated from explants cultured on a medium containing AgNO<sub>3</sub> were more elongated than those cultured on a medium without AgNO<sub>3</sub>. The application of AgNO<sub>3</sub> did not change the dose response of shoot organogenesis for the selective agents hygromycin and mannose.

**Abbreviations:** BA – 6-benzylaminopurine; CBM – cassava basic medium; CEM – cassava elongation medium; CMM – cassava embryo maturation medium; COM – cassava shoot organogenesis medium; IBA – indole-3-butyric acid

### Introduction

Cassava (*Manihot esculenta* Crantz) is a perennial tropical root crop used as a staple by more than 500 million people world-wide (for a review see Puonti-Kaerlas, 1998). Genetic engineering of cassava has high potential as an efficient complement to traditional breeding in improving agriculturally valuable traits such as pest and virus resistance or improved root quality. However, the transformation efficacy of cassava, like many other crops, depends on plant regeneration efficiency. Direct shoot induction (shoot organogenesis) from cotyledons of somatic embryos (somatic cotyledons) of cassava is a rapid regeneration system minimising somaclonal variation (Li et al., 1998) and has already been demonstrated to

be efficient for genetic transformation using particle bombardment (Zhang et al., 2000) as well as for transformation using *Agrobacterium* (Li et al., 1996; Puonti-Kaerlas et al., 1997). The regeneration efficiency is nonetheless genotype-dependent, varying between 5 and 70% (Puonti-Kaerlas, unpublished), which may constrain the use of genetic engineering in cultivars with low regeneration capacity.

Silver nitrate has been shown to be effective in improving somatic embryogenesis and plant regeneration in a number of crop species including *Brassica* spp. (Palmer, 1992; Zhang and Ling, 1995, 1996; Eapen and George, 1997; Kuvshinov et al., 1999), maize (Vain et al., 1989a,b; Carvalho et al., 1997), muskmelon (Yadav et al., 1996), cucumber (Roustan et al., 1992; Mohiuddin et al., 1997), cowpea (Brar

et al., 1999), peanut (Pestana et al., 1999), wheat (Lashermes, 1992), rice (Lentini et al., 1995) and barley (Castillo et al., 1998). As  $\text{Ag}^+$  ions can prevent a wide variety of ethylene-induced plant responses, including growth inhibition and senescence, the effect is assumed to be mediated via the inhibition of the physiological action of ethylene (Beyer et al., 1984), a potential inhibitor of many plant regeneration systems (Vain et al., 1989a; Chraibi et al., 1991; Kong and Yeung, 1994). In this paper we present the results of the first study on the effect of  $\text{AgNO}_3$  on cassava shoot organogenesis from somatic cotyledons. Our study shows that  $\text{AgNO}_3$  can be used to improve the *in vitro* regeneration frequencies of cassava without affecting adversely the efficiency of selectable markers required for selection of transgenic plants.

## Materials and methods

### *Somatic embryogenesis*

One Colombian (MCol22) and three Thai cultivars (T5, KU50 and Hanatee) obtained from CIAT, Colombia and Kasetsart University, Thailand, respectively, were used in this study. The plant material was maintained as shoot cultures on CBM (MS salts and vitamins [Murashige and Skoog, 1962] supplemented with 2% sucrose and  $2 \mu\text{M}$   $\text{CuSO}_4$ , solidified with 0.6% plant agar (Duchefa, pH 5.8) at  $26^\circ\text{C}$  with a 16-h photoperiod (fluorescent tubes,  $200 \mu\text{M m}^{-2} \text{s}^{-1}$ ) and subcultured at 4-week intervals. Apical shoot meristems and 2–5-mm long immature leaf lobes were isolated from shoot cultures and cultured on CIM medium (CBM supplemented with  $12 \text{ mg l}^{-1}$  picloram) at  $26^\circ\text{C}$  in the dark. After 2 weeks, the developing embryos were transferred to fresh CIM medium to initiate cycling secondary embryo cultures. After two to three cycles on CIM medium, the somatic embryos were harvested and cultured onto CMM medium (CBM with  $0.1 \text{ mg l}^{-1}$  BA) at  $26^\circ\text{C}$  with a 16-h photoperiod ( $90\text{--}110 \mu\text{M m}^{-2}\text{s}^{-1}$ ) for production of maturing somatic embryos.

### *Shoot organogenesis*

Green cotyledons were collected from somatic embryos cultured on maturation medium (CMM) for 10–15 days, cut into 5-mm<sup>2</sup> pieces and transferred to COM medium (CBM with  $1 \text{ mg l}^{-1}$  BA and  $0.5 \text{ mg l}^{-1}$  IBA) for shoot organogenesis at  $26^\circ\text{C}$  in the dark. Shoot primordia developing on the explants were

transferred after 3 weeks to CEM (CBM supplemented with  $0.4 \text{ mg l}^{-1}$  BA) for shoot elongation.

### *Silver nitrate test*

COM medium supplemented with  $\text{AgNO}_3$  was used for assessing the effect of silver nitrate. An  $8 \text{ mg ml}^{-1}$  stock solution of  $\text{AgNO}_3$  (Merck, Darmstadt) was filter sterilised, stored at  $4^\circ\text{C}$  and added into the autoclaved medium. The effect of various silver nitrate concentrations (0, 1, 2, 4, 8 and  $12 \text{ mg l}^{-1}$ ) on callus formation and on shoot regeneration ability of cotyledon explants was tested. Fifty explants per cultivar were cultured per treatment. Each experiment had three replicates and was repeated 2 times. After 3 weeks, the status of shoot organogenesis and callus formation were recorded. Three different degrees of organogenesis and callus formation were used to evaluate the response of the cotyledon explants to silver nitrate. Organogenesis degree 1 (OD1) indicates less than five shoot primordia per explant, organogenesis degree 2 (OD2) between five and 10 shoot primordia per explant and organogenesis degree 3 (OD3) more than 10 shoot primordia per explant. In the evaluation of callus development, callus degree 1 (CD1) indicates callus size less than 0.25 cm in diameter, callus degree 2 (CD2) callus size between 0.25 and 0.5 cm in diameter and callus degree 3 (CD3) indicates the size of calli larger than 0.5 cm in diameter.

### *Plant regeneration*

The developing shoot primordia and regenerating shoots were detached from the explants and transferred to CEM medium for shoot elongation. In one experiment, the effect of  $\text{AgNO}_3$  on the competence of shoot primordia to produce elongating shoots was evaluated. Twenty clusters of shoot primordia per treatment were transferred to CEM for elongation. After 4 weeks, the number of elongated shoots was recorded. The shoots were transferred to CBM for rooting and further growth.

### *Effect of silver nitrate on selection agents*

In order to test whether  $\text{AgNO}_3$  influences the efficiency of hygromycin and mannose used for selection of transgenic cassava shoots, cotyledon explants were cultured on COM supplemented with  $8 \text{ mg l}^{-1}$   $\text{AgNO}_3$  and either with  $20 \text{ mg l}^{-1}$  hygromycin or  $20 \text{ g l}^{-1}$  mannose. Fifty explants of MCol22 were used in each

Table 1. The effect of silver nitrate on the frequency of shoot organogenesis (SO)

Cultivars		Concentration of AgNO <sub>3</sub> (mg l <sup>-1</sup> )					
		0	1	2	4	8	12
MCol22	SO	63.3±5.0 <sup>b</sup>	63.3±8.1 <sup>b</sup>	66.0±4.0 <sup>b</sup>	80.7±5.0 <sup>a</sup>	85.3±4.2 <sup>a</sup>	88.7±4.2 <sup>a</sup>
	OD1	42.7±2.3 <sup>a</sup>	30.7±6.1 <sup>bc</sup>	20.7±5.0 <sup>d</sup>	23.3±5.0 <sup>cd</sup>	36.7±4.2 <sup>ab</sup>	34.7±7.0 <sup>ab</sup>
	OD2	20.0±3.5 <sup>b</sup>	21.3±6.1 <sup>b</sup>	24.7±5.0 <sup>ab</sup>	32.7±3.1 <sup>a</sup>	26.7±5.8 <sup>ab</sup>	24.0±6.0 <sup>ab</sup>
	OD3	0.70±1.2 <sup>d</sup>	11.3±4.2 <sup>c</sup>	20.7±3.1 <sup>b</sup>	24.7±6.1 <sup>ab</sup>	22.0±2.0 <sup>b</sup>	30.0±7.2 <sup>a</sup>
KU50	SO	47.3±14.2 <sup>d</sup>	72.7±3.1 <sup>bc</sup>	87.3±3.1 <sup>a</sup>	80.7±4.2 <sup>ab</sup>	62.7±6.1 <sup>c</sup>	49.3±5.0 <sup>d</sup>
	OD1	28.7±4.2 <sup>a</sup>	25.0±6.1 <sup>a</sup>	24.0±4.0 <sup>a</sup>	21. ±4.2 <sup>a</sup>	26.7±5.0 <sup>a</sup>	20.7±3.1 <sup>a</sup>
	OD2	14.7±6.1 <sup>c</sup>	28.0±4.0 <sup>ab</sup>	34.0±3.5 <sup>a</sup>	32.0±3.5 <sup>a</sup>	21.3±6.1 <sup>bc</sup>	18.0±6.0 <sup>c</sup>
	OD3	4.0±4.0 <sup>c</sup>	19.3±5.0 <sup>ab</sup>	26.0±5.3 <sup>a</sup>	27.3±7.0 <sup>a</sup>	14.7±7.0 <sup>b</sup>	10.7±2.3 <sup>bc</sup>
Hanatee	SO	50.0±5.3 <sup>c</sup>	67.3±4.2 <sup>a</sup>	60.7±3.0 <sup>ab</sup>	52.0±8.0 <sup>bc</sup>	43.3±5.0 <sup>c</sup>	43.3±4.2 <sup>c</sup>
	OD1	36.0±5.3 <sup>ab</sup>	37.3±4.2 <sup>a</sup>	26.7±3.1 <sup>c</sup>	30.0±3.5 <sup>bc</sup>	25.3±4.2 <sup>cd</sup>	19.3±3.1 <sup>d</sup>
	OD2	8.0±2.0 <sup>c</sup>	17.3±1.2 <sup>a</sup>	19.3±2.3 <sup>a</sup>	10.7±3.1 <sup>bc</sup>	10.0±2.0 <sup>bc</sup>	13.3±2.3 <sup>b</sup>
	OD3	6.0±2.0 <sup>c</sup>	12.7±2.3 <sup>ab</sup>	14.7±2.3 <sup>a</sup>	11.3±3.1 <sup>ab</sup>	8.0±3.5 <sup>bc</sup>	10.7±3.1 <sup>abc</sup>
T5	SO	25.3±6.4 <sup>c</sup>	32.0±8.0 <sup>bc</sup>	32.7±4.2 <sup>bc</sup>	47.3±5.0 <sup>a</sup>	40.7±7.0 <sup>ab</sup>	38.0±3.5 <sup>ab</sup>
	OD1	19.3±2.3 <sup>b</sup>	24.7±5.0 <sup>ab</sup>	25.3±2.3 <sup>ab</sup>	28.7±3.1 <sup>a</sup>	26.0±5.3 <sup>a</sup>	23.3±3.1 <sup>ab</sup>
	OD2	6.0±5.3 <sup>b</sup>	7.3±3.1 <sup>ab</sup>	8.7±1.2 <sup>ab</sup>	12.0±2.0 <sup>a</sup>	12.0±2.1 <sup>a</sup>	12.0±4.0 <sup>a</sup>
	OD3	0 <sup>b</sup>	0 <sup>b</sup>	2.0±2.0 <sup>b</sup>	6.7±1.2 <sup>a</sup>	2.7±1.1 <sup>b</sup>	2.7±3.0 <sup>b</sup>

The numbers indicate the mean number of responding explants (SO) as percent of all explants, and the distribution of the response between different organogenesis degrees (OD1, OD2 and OD3). Letters a–d indicate significant differences ( $p=0.05$ ) within each row using an LSD test.

treatment with three replicas. After 4 weeks, the number of developing shoots was recorded. For statistical significance, the data were analysed by the LSD test at the 5% level.

## Results

Differences were observed between the four cultivars in their competence for organogenesis under standard conditions (Table 1). MCol22 had the highest organogenesis frequency (63%); while in T5, only 25% of the explants were able to produce shoot primordia. The organogenesis frequency of KU50 and Hanatee was about 50%. In most of the explants organogenesis degree one (less than five shoot primordia per explant) was dominant. At the same time, large amounts of non-morphogenic callus were produced from the cut edges of the explants (Figure 1).

### Enhancement of organogenesis frequency

After 4 weeks of culture on COM supplemented with different concentrations of AgNO<sub>3</sub>, different responses could be observed among the tested cultivars both in shoot organogenesis frequencies and the numbers of developing shoot primordia per explant

(Table 1). In MCol22, the frequency of shoot organogenesis and organogenesis degree increased with increasing AgNO<sub>3</sub> concentrations of up to 12 mg l<sup>-1</sup>. On a medium containing 12 mg l<sup>-1</sup> AgNO<sub>3</sub>, 89% of the explants produced shoot primordia, and 30% of these had more than 10 shoot primordia per explant (organogenesis degree 3); whereas in the controls, the organogenesis frequency was 63%, and only 0.7% of the explants showed organogenesis degree 3. In KU50 and Hanatee, the highest increase in organogenesis rate could be obtained by supplementing the medium with 2 and 1 mg l<sup>-1</sup> AgNO<sub>3</sub> respectively; while in T5, the best response was observed using 4 mg l<sup>-1</sup> AgNO<sub>3</sub> in the medium. Concentrations higher than 8 mg l<sup>-1</sup> had a slightly inhibitory effect on shoot organogenesis of the Thai cultivars.

### Inhibition of callus formation

On COM, 100% of the somatic cotyledon explants of MCol22, KU50 and T5 formed callus, more than 70% of which was degree 3 (callus size larger than 0.5 cm in diameter) (Table 2). In Hanatee, 89% of the explants produced callus and 73% of this was degree 3. In all cultivars, increasing concentrations of AgNO<sub>3</sub> reduced callus formation, both in terms of callus frequency and of callus degree. In MCol22, callus frequency was decreased from 100% (control) to 5%




Figure 1. The effect of silver nitrate on shoot organogenesis and callus formation in MCol22; left, without silver nitrate; right, with 4 mg l<sup>-1</sup> silver nitrate.

(12 mg l<sup>-1</sup> AgNO<sub>3</sub>) and the frequency of callus degree 3 was reduced from 98 to 0% (Figure 1). In Hanatee, callus formation was reduced to 25% by 4 mg l<sup>-1</sup> and totally inhibited by 8 mg l<sup>-1</sup> AgNO<sub>3</sub>; while in KU50 and in T5, the effect was less pronounced, but still significantly different from the controls.

#### *Improvement of shoot elongation*

Shoot primordia of MCol22 induced on COM with or without 4 mg l<sup>-1</sup> AgNO<sub>3</sub> were transferred individually onto CEM to test the effect of silver nitrate on shoot elongation. In all the tested cultivars, shoot primordia induced on COM with AgNO<sub>3</sub> produced more elongating shoots than those induced on a medium without AgNO<sub>3</sub> (Table 3).

#### *Effect of silver nitrate on the dose response to selective agents*

Hygromycin and mannose, used for production of transgenic cassava were tested for their efficacy in inhibiting shoot organogenesis of MCol22 cotyledon explants cultured on COM supplemented with AgNO<sub>3</sub>. Addition of 8 mg l<sup>-1</sup> silver nitrate did not affect the sensitivity of shoot primordia regeneration to hygromycin or mannose (Table 4).

#### *Discussion*

Many reports have demonstrated the positive effect of AgNO<sub>3</sub> on plant tissue culture, although in some plants and tissue types no positive or even an inhibitory effect has been reported (Puonti-Kaerlas, 1991;

Palmer, 1992; Jun et al., 1997; Teo et al., 1997; Kumar et al., 1998; El Meskaoui and Tremblay, 1999). In the case of cassava, it has been shown that the maturation and regeneration of secondary somatic embryos of one cultivar could be improved by the use of 16 mg l<sup>-1</sup> AgNO<sub>3</sub> and 0.25 mg l<sup>-1</sup> ABA (Zhu et al., 1998). We show here that the capacity of shoot organogenesis in a number of cassava cultivars can be improved by supplementing the medium with AgNO<sub>3</sub>. The optimum concentration of AgNO<sub>3</sub> was found to differ among the different cultivars. In MCol22, using 4–12 mg l<sup>-1</sup> AgNO<sub>3</sub> increased the frequency of shoot organogenesis from 63% (control) to over 80%. At the same time, the number of regenerated shoots per explant increased as well, with concomitant inhibition of callus growth. The optimum AgNO<sub>3</sub> concentrations for shoot regeneration in the other cultivars were lower than in MCol22; 2 mg l<sup>-1</sup> in KU50, 1 mg l<sup>-1</sup> in Hanatee and 4 mg l<sup>-1</sup> in T5. Hence, different cassava cultivars appear to have quantitatively different responses to AgNO<sub>3</sub> and the optimal concentration needs be determined separately for each cultivar. The genotype and the developmental stage of explants have been shown to affect the response of other plant species to silver nitrate as well (Palmer et al., 1992; Evans and Batty, 1994; Hyde and Phillips, 1996; Mohiuddin et al., 1997; Santos et al., 1997). Of the cassava lines tested in this study, MCol22 appears to have a broader optimum for AgNO<sub>3</sub> than the Thai cultivars. In contrast to MCol22, the most efficient concentrations for improving shoot regeneration in the Thai cultivars are lower than those required for maximal suppression of callus growth. The highest number of

Table 2. The effect of silver nitrate on the frequency of callus formation (CF) and the callus degrees (CD1, CD2 and CD3)

Cultivars	Concentration of AgNO <sub>3</sub> (mg l <sup>-1</sup> )						
	0	1	2	4	8	12	
MCol22	CF	100 <sup>a</sup>	92.0±4.0 <sup>a</sup>	24.0±8.0 <sup>b</sup>	20.7±6.1 <sup>b</sup>	16.0±4.0 <sup>b</sup>	4.7±2.3 <sup>c</sup>
	CD1	0 <sup>c</sup>	26.6±4.2 <sup>a</sup>	12.0±8.0 <sup>b</sup>	13.3±3.1 <sup>b</sup>	12.7±1.2 <sup>b</sup>	4.0±2.0 <sup>c</sup>
	CD2	2.0±3.5 <sup>cd</sup>	36.7±5.0 <sup>a</sup>	10.0±2.0 <sup>b</sup>	7.3±4.2 <sup>bc</sup>	3.3±4.2 <sup>cd</sup>	0.7±1.1 <sup>d</sup>
	CD3	98.0±3.5 <sup>a</sup>	28.7±3.1 <sup>b</sup>	2.0±2.0 <sup>c</sup>	0 <sup>c</sup>	0 <sup>c</sup>	0 <sup>c</sup>
KU50	CF	100 <sup>a</sup>	98.7±2.3 <sup>a</sup>	78.7±4.2 <sup>b</sup>	56.0±7.2 <sup>c</sup>	31.3±10.1 <sup>d</sup>	32.0±15.6 <sup>d</sup>
	CD1	4.0±2.0 <sup>c</sup>	11.3±3.0 <sup>b</sup>	16.6±3.1 <sup>b</sup>	24.0±4.0 <sup>a</sup>	15.3±3.1 <sup>b</sup>	14.7±4.2 <sup>b</sup>
	CD2	7.3±1.2 <sup>b</sup>	14.0±7.2 <sup>ab</sup>	17.3±2.3 <sup>ab</sup>	19.3±3.1 <sup>a</sup>	10.0±2.0 <sup>ab</sup>	17.3±12.9 <sup>ab</sup>
	CD3	88.7±3.1 <sup>a</sup>	73.3±6.1 <sup>b</sup>	44.7±7.0 <sup>c</sup>	12.7±7.0 <sup>d</sup>	6.0±6.0 <sup>de</sup>	0 <sup>e</sup>
Hanatee	CF	89.3±6.1 <sup>a</sup>	62.0±5.3 <sup>b</sup>	40.7±5.0 <sup>c</sup>	25.3±3.1 <sup>d</sup>	0 <sup>e</sup>	0 <sup>e</sup>
	CD1	5.3±2.3 <sup>c</sup>	34.0±3.5 <sup>a</sup>	32.0±2.0 <sup>a</sup>	20.0±2.0 <sup>b</sup>	0 <sup>d</sup>	0 <sup>d</sup>
	CD2	11.3±2.3 <sup>b</sup>	20.0±4.0 <sup>a</sup>	7.3±2.3 <sup>c</sup>	4.7±1.2 <sup>c</sup>	0 <sup>d</sup>	0 <sup>d</sup>
	CD3	72.7±10.1 <sup>a</sup>	8.0±2.0 <sup>b</sup>	1.3±1.2 <sup>bc</sup>	0.7±1.1 <sup>bc</sup>	0 <sup>c</sup>	0 <sup>c</sup>
T5	CF	100 <sup>a</sup>	100 <sup>a</sup>	100 <sup>a</sup>	91.3±3.0 <sup>b</sup>	90.7±4.2 <sup>b</sup>	89.3±5.0 <sup>b</sup>
	CD1	4.0±4.0 <sup>d</sup>	18.0±5.3 <sup>c</sup>	38.0±6.0 <sup>b</sup>	37.3±2.3 <sup>b</sup>	43.3±5.0 <sup>ab</sup>	47.3±4.2 <sup>a</sup>
	CD2	26.7±6.1 <sup>b</sup>	38.7±10.1 <sup>a</sup>	38.0±5.3 <sup>a</sup>	30.0±7.2 <sup>a</sup>	28.0±3.5 <sup>a</sup>	29.3±3.1 <sup>a</sup>
	CD3	69.3±4.6 <sup>a</sup>	42.0±8.7 <sup>b</sup>	24.0±4.0 <sup>c</sup>	24±5.3 <sup>c</sup>	19.3±4.2 <sup>cd</sup>	12.7±3.0 <sup>d</sup>

Letters a–e indicate significant differences ( $p=0.05$ ) within each row using an LSD test.

responding explants, however, was obtained in all cultivars at the same concentration as the highest number of developing shoot primordia.

Histological studies in *Brassica parachinensis* showed that AgNO<sub>3</sub> influenced the mode of plant regeneration, allowing direct shoot primordia development without an intermediate callus phase (Zhang and Ling, 1995, 1996). Also in *Albizia procera* the use of silver nitrate has been reported to promote callus-free shoot regeneration (Kumar et al., 1998). In our experiments with cassava, supplementing COM with AgNO<sub>3</sub> altered the regeneration mode from organogenesis involving a callus phase to that without an intervening callus phase (Figure 1). Callus formation is often considered to inhibit plant regeneration and, in cassava, reproducible plant regeneration from callus has only been reported for one cultivar (Mussio et al., 1998). We assume therefore that the increased capacity of shoot organogenesis is at least partially related to the inhibition of callus formation. The correlation between organogenesis frequency and callus frequency in MCol22 is  $-0.82$ , indicating that organogenesis and callus formation are indeed inversely correlated.

The elongation of shoots was more efficient from primordia induced on a regeneration medium containing AgNO<sub>3</sub> than from those cultured on a medium

without AgNO<sub>3</sub> (Table 3). Thus, the use of AgNO<sub>3</sub> during shoot primordia induction also has a positive effect on later plant development at the shoot elongation step. Silver nitrate has been shown in *Brassica rapa* to improve the frequency of shoot organogenesis from seedling explants pre-treated with AgNO<sub>3</sub> during seed germination (Burneet et al., 1994). In some cases, the use of AgNO<sub>3</sub> has reduced or inhibited rooting of the regenerated shoots (Castillo et al., 1998; Madsen et al., 1998) but in cassava no adverse effects on rooting ability could be observed.

The mode of action of AgNO<sub>3</sub> in plant tissue culture is assumed to be associated with the physiological effects of ethylene, silver ions acting as a competitive inhibitor of ethylene action rather than inhibiting ethylene synthesis per se. Ethylene production may in fact increase in plant cultures treated with silver nitrate (Pua et al., 1993, 1999; Lee et al., 1997; Zhang et al., 1998). Until now, the mechanism of ethylene action on plant tissue culture has not been elucidated. While ethylene may act as an inhibitor in some plant regeneration systems (Vain et al., 1989b; Chraibi et al., 1991; Kong and Yeung, 1994), it may on the other hand function as stimulator in others (Hatanaka et al., 1995; Nissen, 1994). Biochemical studies suggested that a transition metal is involved in binding of ethylene to receptors, since addition of CuSO<sub>4</sub> to

Table 3. The frequency of shoot elongation from four cassava cultivars on CEM from shoot primordia induced either on COM or AgCOM (COM with 4 mg l<sup>-1</sup> silver nitrate)

Cultivar	Medium used for shoot primordia induction	No. of shoot primordia clusters	No. of elongating shoots	Shoot elongation frequency (%)
MCol22	COM	20	8	40
	AgCOM	20	12	60
T5	COM	20	4	20
	AgCOM	20	5	25
KU50	COM	20	8	40
	AgCOM	20	11	55
Hanatee	COM	20	3	15
	AgCOM	20	9	45

Table 4. The influence of silver nitrate (Ag) on inhibition of shoot primordia development by hygromycin (Hm) and mannose (Man) in MCol22

Type of medium	Shoot organogenesis frequency (%)
COM	63.0±4.0 <sup>a</sup>
COM+Hm	0.7±1.2 <sup>b</sup>
AgCOM+Hm	1.3±1.2 <sup>b</sup>
COM+Man	2.5±2.3 <sup>b</sup>
AgCOM+Man	1.0±1.7 <sup>b</sup>

Letters a–b indicate significant differences ( $p=0.05$ ) using an LSD test.

membranes isolated from ETR1-expressing yeast resulted in up to a 20-fold increase in ethylene binding (Bleecker, 1997). Very interestingly and surprisingly, Ag<sup>+</sup> ions also increased receptor affinity for ethylene in the yeast system, suggesting that silver ions act by interfering with intramolecular signal transduction rather than by ligand binding (Bleecker, 1997). Other studies showed that silver ions may interact with polyamines, which have been shown to promote organogenesis and embryogenesis (Feirer et al., 1984; Meijer and Simmonds, 1988; Pua et al., 1999) since ethylene and polyamines compete for the same precursor, SAM (*S*-adenosyl methionine). Few studies have addressed the interaction between silver ions, ethylene and polyamines (Pua et al., 1999), and so far the nature of these interactions is not known.

The enhancement of *in vitro* shoot organogenesis by AgNO<sub>3</sub> should not interfere with transformation or selection efficiencies, so as not to prevent the use of AgNO<sub>3</sub> in transgenic plant production. Our results show that silver nitrate did not influence the dose re-

sponse of cassava tissues to the selective agents hygromycin and mannose. Shoot organogenesis is inhibited by 20 mg l<sup>-1</sup> hygromycin or 20 g l<sup>-1</sup> mannose both in the presence and absence of AgNO<sub>3</sub>. The addition of AgNO<sub>3</sub> in shoot regeneration medium has been shown to be essential for regeneration of transformed tissues in *Brassica napus* and *Brassica oleracea* (DeBlock et al., 1989), *Brassica campestris* (Mukhopadhyay et al., 1992; Kuvshinov et al., 1999) and *Brassica rapa* (Radke et al., 1992). The present study shows that it is also possible to improve the frequencies of shoot organogenesis in cassava by supplementing the regeneration medium with AgNO<sub>3</sub>. This should allow higher transformation frequencies and more efficient regeneration of transgenic cassava plants in the future.

### Acknowledgements

We would like to thank Prof. Dr Ingo Potrykus for his keen interest and his generous support. The skilful assistance of Mr Bertrand Hankoua is acknowledged. This project was partially supported by the Swiss Development Agency (SDC) through the Swiss Centre of International Agriculture (ZIL).

### References

- Beyer EM, Morgan JPW & Yang SF (1984) Ethylene. In: Wilkins MB (ed) *Advanced Plant Physiology*. (pp 111–126) Pitman Press, Bath, UK
- Bleecker AB (1997) The ethylene binding site of the ETR1 protein. In: Kanellis A, Chang C, Kende H & Grierson D (eds) *Biology and Biotechnology of the Plant Hormone Ethylene* (pp 63–70) Kluwer Academic Publishers, Dordrecht, The Netherlands

- Brar MS, Moore MJ, Al Khayri JM, Morelock TE & Anderson EJ (1999) Ethylene inhibitors promote *in vitro* regeneration of cowpea (*Vigna unguiculata* L.). *In Vitro Cell. Dev. Biol.* 35: 222–225
- Burnett L, Arnoldo M & Yarrow S (1994) Enhancement of shoot regeneration from cotyledon explants of *Brassica rapa* ssp. *oleifera* through pretreatment with auxin and cytokinin and use of ethylene inhibitors. *Plant Cell Tiss. Org. Cult.* 37: 253–258
- Carvalho CHS, Bohorova N, Bordallo PN, Abreu LL, Valicente FH, Bressan W & Paiva E (1997) Type II callus production and plant regeneration in tropical maize genotypes. *Plant Cell Rep.* 17: 73–76
- Castillo AM, Egana B, Sanz JM & Cistue L (1998) Somatic embryogenesis and plant regeneration from barley cultivars grown in Spain. *Plant Cell Rep.* 17: 902–906
- Chi GL, Lin WS, Lee JEE & Pua EC (1994) Role of polyamines on *de novo* shoot morphogenesis from cotyledons of *Brassica campestris* ssp. *pekinensis* (Lour) Olsson *in vitro*. *Plant Cell Rep.* 13: 323–329
- Chraïbi BKM, Latche A, Roustan JP & Fallot J (1991) Stimulation of shoot regeneration from cotyledons of *Helianthus annuus* by ethylene inhibitors, silver and cobalt. *Plant Cell Rep.* 10: 204–207
- DeBlock M, DeBrouwer D & Tennig P (1989) Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the *bar* and *neo* genes in the transgenic plants. *Plant Physiol.* 91: 694–701
- Eapen S & George L (1997) Plant regeneration from peduncle segments of oil seed *Brassica* species: influence of silver nitrate and silver thiosulfate. *Plant Cell Tiss. Org. Cult.* 51: 229–232
- El Meskaoui A & Tremblay FM (1999) Effects of sealed and vented gaseous microenvironments on the maturation of somatic embryos of black spruce with a special emphasis on ethylene. *Plant Cell Tiss. Org. Cult.* 56: 201–209
- Evans JM & Batty NP (1994) Ethylene precursors and antagonists increase embryogenesis of *Hordeum vulgare* L. anther culture. *Plant Cell Rep.* 13: 676–678
- Feirer RP, Mingnon G & Litray JD (1984) Arginine decarboxylase and polyamines required for embryogenesis in the wild carrot. *Science* 233: 1433–1435
- Hatanaka T, Sawabe E, Azuma T, Uchida N & Yasuda T (1995) The role of ethylene in somatic embryogenesis from leaf discs of *Coffea canephora*. *Plant Sci.* 107: 199–204
- Hyde CL & Phillips GC (1996) Silver nitrate promotes shoot development and plant regeneration of chile pepper (*Capsicum annum* L.) via organogenesis. *In Vitro Cell. Dev. Biol.* 32: 72–80
- Jun JH, Yae BW, Hwang JH & Shin YU (1997) Plant regeneration from leaf tissue of *Malus domestica* cv. 'Fuji' *in vitro*. *RDA J. Hort. Sci.* 39: 102–105
- Kong L & Yeung EC (1994) Effects of ethylene and ethylene inhibitors on white spruce somatic embryo maturation. *Plant Sci.* 104: 71–80
- Kumar S, Sarkar AK & Kunhikannan C (1998) Regeneration of plants from leaflet explants of tissue culture raised safed siris (*Albizia procera*). *Plant Cell Tiss. Org. Cult.* 54: 137–143
- Kuvshinov V, Koivu K, Kanerva A & Pehu E (1999) *Agrobacterium tumefaciens*-mediated transformation of greenhouse-grown *Brassica rapa* ssp. *oleifera*. *Plant Cell Rep.* 18: 773–777
- Lashermes P (1992) Improved anther culture method for obtaining direct regeneration in wheat (*Triticum aestivum* L.). *J. Genet. Breed.* 46: 99–102
- Lee T, Huang MEE & Pua EC (1997) High frequency shoot regeneration from leaf disc explants of garland chrysanthemum (*Chrysanthemum coronarium* L.) *in vitro*. *Plant Sci.* 126: 219–226
- Lentini Z, Reyes P, Martinez CP & Roca WM (1995) Androgenesis of highly recalcitrant rice genotypes with maltose and silver nitrate. *Plant Sci.* 110: 127–138
- Li H-Q, Sautter C, Potrykus I & Puonti-Kaerlas J (1996) Genetic transformation of cassava (*Manihot esculenta* Crantz). *Nat. Biotechnol.* 14: 736–740
- Li H-Q, Huang YW, Liang CY, Guo JY, Liu HX, Potrykus I & Puonti-Kaerlas J (1998) Regeneration of cassava plants via shoot organogenesis. *Plant Cell Rep.* 17: 410–414
- Madsen MH, Nauerby B, Frederiksen CG & Wyndaele R (1998) Regeneration of pea (*Pisum sativum* L.) by the thin cell layer nodal system: Influence of explant culture media on rooting and plantlet formation. *Acta Agric. Scand. Sect. B Soil Plant Sci* 48: 58–64
- Meijer EGM & Simmonds J (1988) Polyamine levels in relation to growth and somatic embryogenesis of *Medicago sativa*. *J. Exp. Bot.* 203: 787–794
- Mohiuddin AKM, Chowdhury MKU, Abdullah Zaliha C & Napis S (1997) Influence of silver nitrate (ethylene inhibitor) on cucumber *in vitro* shoot regeneration. *Plant Cell Tiss. Org. Cult.* 51: 75–78
- Mukhopadhyay A, Arumugam N, Nandakumar PBA, Pradhan AK, Gupta V & Pental D (1992) *Agrobacterium* mediated genetic transformation of oilseed *Brassica campestris*: transformation frequency is strongly influence by the mode of shoot regeneration. *Plant Cell Rep.* 11: 506–513
- Murashige T & Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* 15: 473–497
- Mussio I, Chaput MH, Serraf I, Ducreux G & Sihachakr D (1998) Adventitious shoot regeneration from leaf explants of an African clone of cassava (*Manihot esculenta* Crantz) and analysis of the conformity of regenerated plants. *Plant Cell Tiss. Org. Cult.* 53: 205–211
- Nissen P (1994) Stimulation of somatic embryogenesis in carrot by ethylene: effects of modulators of ethylene biosynthesis and action. *Physiol. Plant.* 92: 397–403
- Palmer CE (1992) Enhanced shoot regeneration from *Brassica campestris* by silver nitrate. *Plant Cell Rep.* 11: 541–545
- Pestana MC, Lacorte C, de Freitas VG, de Oliveira DE & Mansur E (1999) *In vitro* regeneration of peanut (*Arachis hypogaea* L.) through organogenesis: effect of culture temperature and silver nitrate. *In Vitro Cell. Dev. Biol.* 35: 214–216
- Pua EC & Chi GL (1993) *De novo* shoot morphogenesis and plant growth of mustard (*Brassica juncea*) *in vitro* in relation to ethylene. *Physiol. Plant.* 88: 467–474
- Pua EC, Deng X & Koh ATC (1999) Genotypic variability of *de novo* shoot morphogenesis of *Brassica oleracea* *in vitro* in response to ethylene inhibitors and putrescine. *J. Plant Physiol.* 155: 598–605
- Puonti-Kaerlas J (1991) Tissue culture and genetic transformation of pea (*Pisum sativum* L.). *Acta Universitatis Upsaliensis, Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science* (p. 340)
- Puonti-Kaerlas J (1998) Cassava biotechnology. In: Tombs MP (ed) *Biotechnology and Genetic Engineering Reviews*. Vol. 15 (pp 329–364) Intercept Ltd, Andover, Hants, UK
- Puonti-Kaerlas J, Li HQ, Sautter C & Potrykus I (1997) Production of transgenic cassava (*Manihot esculenta* Crantz) via organogenesis and *Agrobacterium*-mediated transformation. *African J. Root Tuber Crops* 2: 181–186

- Radke SE, Turner JC & Facciotti D (1992) Transformation and regeneration of *Brassica rapa* using *Agrobacterium tumefaciens*. *Plant Cell Rep.* 11: 499–505
- Roustan Jean P, Latche A & Fallois J (1992) Enhancement of shoot regeneration from cotyledons of *Cucumis melo* by silver nitrate, an inhibitor of ethylene action. *J. Plant Physiol.* 140: 485–488
- Santos KGB, Mundstock E & Bodanese Zanettini MH (1997) Genotype-specific normalization of soybean somatic embryogenesis through the use of an ethylene inhibitor. *Plant Cell Rep.* 16: 859–864
- Teo W, Lakshmanan P, Kumar P, Goh CJ & Swarup S (1997) Direct shoot formation and plant regeneration from cotyledon explants of rapid-cycling *Brassica rapa*. *In Vitro Cell. Dev. Biol.* 33: 288–292
- Vain P, Flament P & Soudain P (1989a) Role of ethylene in embryogenic callus initiation and regeneration in *Zea mays* L. *J. Plant Physiol.* 135: 537–540
- Vain P, Yean H & Flament P (1989b) Enhancement of production and regeneration of embryogenic type II callus in *Zea mays* L. by silver nitrate. *Plant Cell Tiss. Org. Cult.* 18: 143–152
- Yadav RC, Saleh Mohamed T & Grumet R (1996) High frequency shoot regeneration from leaf explants of muskmelon. *Plant Cell Tiss. Org. Cult.* 45: 207–214
- Zhang P & Ling DH (1995) Enhancement of plant regeneration rate of *Brassica parachinensis* cultured *in vitro*. *Acta Bot. Sinica* 37: 902–908
- Zhang P & Ling DH (1996) Histological studies of plant regeneration modes of *Brassica parachinensis* by affecting AgNO<sub>3</sub> and ABA. *J. Trop. Subtrop. Bot.* 4(1): 71–76
- Zhang FL, Takahata Y & Xu JB (1998) Medium and genotype factors influencing shoot regeneration from cotyledonary explants of Chinese cabbage (*Brassica campestris* L. ssp. *pekinensis*). *Plant Cell Rep.* 17: 780–786
- Zhang P, Legris G, Coulin P & Puonti-Kaerlas J (2000) Production of stably transformed cassava plants via particle bombardment. *Plant Cell Rep.* 19: 939–945
- Zhu J, Huang YW & Liang CY (1998) Improvement of plant regeneration from cyclic secondary somatic embryos in cassava (*Manihot esculenta* Crantz). *J. Trop. Subtrop. Bot.* 6(2): 144–151