

# Use of a silicone patch in the management of severe ischemic small bowel volvulus

A Dahl-Farhoumand, J Birraux, A Darani, BE Wildhaber

Geneva, Switzerland

A newborn was transferred to our unit for a suspicion of volvulus. An emergency laparotomy showed, in the context of malrotation, a 540° volvulus, with the totality of the small bowel dark purple. There was no improvement more than 30 minutes after the bowel was detorsed (Fig. 1). It was highly suspected that no viable small bowel was present, and we decided to give supportive care for 48 hours and then reassess the situation. To close the laparotomy, we used an abdominal patch consisting of a transparent silicone foil (Perthese, Laboratories Perouse Implant, Borel, France) stitched to the muscular sheath, in order to allow observation of the evolution of the small bowel and decompression of the abdominal cavity (Fig. 2). Against all odds, the

neonate improved clinically, with a progressive better coloration of the bowel visible through the silicone foil, thus the planned second look at day two was postponed. Seven days after the first operation, re-exploration was undertaken: the small bowel was completely viable (Fig. 3). A Ladd's procedure with appendectomy was performed. Recovery was complete and uneventful. A follow-up of two years have shown no complications and an excellent evolution of a well thriving boy.

## Discussion

This newborn with small bowel volvulus and with an appearance of severe ischemia of the totality of the small bowel was considered to have a very poor




Fig. 1. Appearance of the ischemic intestine after more than 30 minutes of detorsion during the first laparotomy.




Fig. 2. The small bowel visible through the patch.



Fig. 3. Appearance of the small bowel on laparotomy 7 days later.

**Author Affiliations:** Unit of Pediatric Surgery, University Hospital, Geneva, Switzerland (Dahl-Farhoumand A, Birraux J, Darani A, Wildhaber BE)

**Corresponding Author:** Agnes Dahl-Farhoumand, Department of Pediatric surgery, Geneva University Hospitals, Rue Willy Donze, 6, 1211 Geneva 14, Switzerland (Tel: 00 41 22 3824772; Fax: 00 41 22 3825484; Email: agnesingeneva@gmail.com)

doi: 10.1007/s12519-011-0270-z

©Children's Hospital, Zhejiang University School of Medicine, China and Springer-Verlag Berlin Heidelberg 2011. All rights reserved.

prognosis.<sup>[1]</sup> How could we have identified whether the bowel was still viable and would recover? There have been a few attempts to find a per-operative sensitive and specific marker for gut viability. Intravenous fluorescein and Wood's lamp illumination have been used to assess bowel viability after aortic reconstruction.<sup>[2]</sup> Some studies in animals<sup>[3]</sup> and some in human colon during aortic surgery<sup>[4]</sup> have used mucosal oxygen saturation as a measure of ischemia. Visible light spectroscopy oximeter was used to measure mucosal capillary hemoglobin oxygen saturation.<sup>[5]</sup> None of these methods is used routinely on a wide scale yet, but they seem to be promising. For the moment the surgeon uses his own experience to judge the viability of the intestine. He may decide to perform a second look approach to evaluate the need to further resect bowel when there is suspected disease progression, as in acute mesenteric ischemia.<sup>[6]</sup> Yet, few studies have related the second look approach in patients with bowel obstruction.<sup>[7]</sup> The timing of the second look operation might be difficult to determine. We advocate the use of a transparent silicone foil as a practical visual aid, through which the intestinal evolution can be observed. A further advantage of the silicone foil is the associated enlargement of the abdominal cavity and thus the prevention of a possible abdominal compartment syndrome.<sup>[8]</sup>

**Funding:** None.

**Ethical approval:** Not needed.

**Competing interest:** None declared.

**Contributors:** Dahl-Farhoumand A wrote the main body of the

article under the supervision of Birraux J, Darani A and Wildhaber BE. Wildhaber BE is the guarantor.

## References

- 1 Messineo A, MacMillan JH, Palder SB, Filler RM. Clinical factors affecting mortality in children with malrotation of the intestine. *J Pediatr Surg* 1992;27:1343-1345.
- 2 Bergman RT, Gloviczki P, Welch TJ, Naessens JM, Bower TC, Hallett JW Jr, et al. The role of intravenous fluorescein in the detection of colon ischemia during aortic reconstruction. *Ann Vasc Surg* 1992;6:74-79.
- 3 Erikoglu M, Kaynak A, Beyatli EA, Toy H. Intraoperative determination of intestinal viability: a comparison with transserosal pulse oximetry and histopathological examination. *J Surg Res* 2005;128:66-69.
- 4 Lee ES, Bass A, Arko FR, Heikkinen M, Harris EJ, Zarins CK, et al. Intraoperative colon mucosal oxygen saturation during aortic surgery. *J Surg Res* 2006;136:19-24.
- 5 Benaron DA, Parachikov IH, Cheong WF, Friedland S, Rubinsky BE, Otten DM, et al. Design of a visible-light spectroscopy clinical tissue oximeter. *J Biomed Opt* 2005;10:44005.
- 6 Yanar H, Taviloglu K, Ertekin C, Ozcinar B, Yanar F, Guloglu R, et al. Planned second-look laparoscopy in the management of acute mesenteric ischemia. *World J Gastroenterol* 2007;13:3350-3353.
- 7 Pearl JP, Rosen MJ. Second-look laparoscopy after laparoscopic relief of strangulated small bowel obstruction. *Surg Laparosc Endosc Percutan Tech* 2009;19:241-243.
- 8 Hoffman MA, Johnson CL, Moore T, Pearl RH. Management of catastrophic neonatal midgut volvulus with a silo and second-look laparotomy. *J Pediatr Surg* 1992;27:1336-1339.

*Received September 6, 2010*

*Accepted after revision October 12, 2010*