

Extremes (2006) 9:177–192
DOI 10.1007/s10687-006-0027-6

Extreme VaR scenarios in higher dimensions

Paul Embrechts · Andrea Höing

Received: 18 January 2006 / Revised: 17 September 2006 /
Accepted: 5 October 2006 / Published online: 23 February 2007
© Springer Science + Business Media, LLC 2007

Abstract For a sequence of random variables X_1, \dots, X_n , the dependence scenario yielding the worst possible Value-at-Risk at a given level α for $X_1 + \dots + X_n$ is known for $n = 2$. In this paper we investigate this problem for higher dimensions. We provide a geometric interpretation highlighting the dependence structures which imply the worst possible scenario. For a portfolio (X_1, \dots, X_n) with given uniform marginals, we give an analytical solution sustaining the main result of Rüschendorf (*Adv. Appl. Probab.* 14(3):623–632, 1982). In general, our approach allows for numerical computations.

Keywords Value-at-Risk · Dependent risks · Copulas

AMS 2000 Subject Classifications Primary—G10; Secondary—IM01, IM12, IM52

1 Introduction

Throughout this paper, we use the language of quantitative risk management (QRM) as for instance explained in McNeil et al. (2005). A random variable X is typically referred to as a (one-period) risk corresponding to the unknown value of an underlying financial or insurance instrument at the end of a given time period, viewed from today. For a portfolio (X_1, \dots, X_n) of risks with given marginal distributions, we consider the problem of finding the worst possible Value-at-Risk at the level α for $X_1 + \dots + X_n$ under all possible dependence scenarios for the random variables X_1, \dots, X_n . We denote

P. Embrechts · A. Höing (✉)
Department of Mathematics, ETH Zürich, CH-8092 Zürich, Switzerland
e-mail: ahoeing@math.ethz.ch

this value by $\text{wVaR}_\alpha(\sum_{i=1}^n X_i)$. The definition of Value-at-Risk is given in Section 2. The latter question has been widely studied in the literature, often formulated in terms of the best possible lower bound for the distribution function of the sum; see for instance Section 6.2 in McNeil et al. (2005) and references therein. In risk management this question is motivated by the fact that the worst-case dependence scenario does not occur under comonotonic dependence; see Fallacy 3 in Embrechts et al. (2002). We do not emphasize this issue further. Recent publications on this subject, which also widely discuss the role of comonotonicity, are Denuit et al. (2005), Embrechts et al. (2003), Embrechts and Puccetti (2006) and Embrechts et al. (2005), where the problem is considered for non-decreasing functionals. A more practically oriented paper highlighting the importance of such questions is Neslehova et al. (2006). While the above cited papers provide bounds on $\text{wVaR}_\alpha(\sum_{i=1}^n X_i)$ and fully explain the two-dimensional situation finding a worst dependence scenario in terms of copulas, they all fail to catch the nature of the copula solving the problem in higher dimensions. In this paper we describe this extreme dependence scenario extending some geometrical arguments introduced in Embrechts et al. (2005) for $n = 2$. This allows us to numerically answer the question at hand and, for uniform marginal distributions, to provide an analytical solution equivalent to that presented in Rüschendorf (1982). The latter is the only known analytical result for continuous marginals. Some applications of our results are given in Section 4.

2 Preliminaries and Fundamental Results

We briefly summarize the basic tools used in the literature and recall the fundamental results on the problem of bounding the Value-at-Risk. All the theorems are formulated for the sum of risks assuming no information about their interdependence. For further discussions regarding more general functionals and the assumption of partial dependence information, we refer to the papers cited in the introduction.

2.1 Value-at-Risk and Copulas

For risk management purposes we assume X_1, \dots, X_n to have distribution functions F_1, \dots, F_n with losses represented in their right tails; i.e. losses correspond to positive values of the X_i 's.

Definition 1 Let X be a random variable with distribution F_X . For $0 < \alpha < 1$ the Value-at-Risk at probability level α of X is its α -quantile, i.e. $\text{VaR}_\alpha(X) := F_X^{-1}(\alpha) := \inf\{x \in \mathbb{R}: F_X(x) \geq \alpha\}$.

In risk management applications, typical values for α are 0.95 or 0.99 in the case of market or credit risk and $\alpha = 0.999$ for operational risk.

Given the joint distribution function $F(\mathbf{x}) = \mathbf{P}(X_1 \leq x_1, \dots, X_n \leq x_n)$, $\mathbf{x} \in \mathbb{R}^n$, the problem of calculating $\text{VaR}_\alpha(\sum_{i=1}^n X_i)$ reduces to a computational issue. In what follows we assume full knowledge about the marginals but no prior information on the dependence structure. In this context, the idea of copula allows for a precise formulation of the problem separating F into one part describing the dependence structure and another part containing the information on the marginals. We refer to Nelsen (1999) for the basic results about copulas.

Definition 2 A n -dimensional copula C is a distribution function on $[0, 1]^n$ with uniform- $(0, 1)$ marginals. We denote their class by \mathcal{C}^n .

Remark 1 A copula can be equivalently defined as a function $C : [0, 1]^n \rightarrow [0, 1]$ satisfying $C(u_1, \dots, u_{i-1}, 0, u_{i+1}, \dots, u_n) = 0$, $C(1, \dots, 1, u_i, 1, \dots, 1) = u_i$ and $\sum_{i_1=1}^2 \dots \sum_{i_n=1}^2 (-1)^{i_1+\dots+i_n} C(u_{1,i_1}, \dots, u_{n,i_n}) \geq 0$ for $\mathbf{u}, \mathbf{v} \in [0, 1]^n$ with $\mathbf{u} \leq \mathbf{v}$ and $u_{j,1} = u_j, u_{j,2} = v_j, j = 1, \dots, n$.

Sklar’s Theorem yields that, for a $C \in \mathcal{C}^n$ and marginal distributions F_1, \dots, F_n , the function $F(\mathbf{x}) := C(F_1(x_1), \dots, F_n(x_n))$ is a distribution with these marginal distributions. Conversely, for any joint distribution function with given marginals, there is a copula linking them. It is unique if the marginals are continuous. Any copula C lies between the so called lower and upper Fréchet bounds $W(\mathbf{u}) := (\sum_{i=1}^n u_i - n + 1)^+$ and $M(\mathbf{u}) := \min_{1 \leq i \leq n} u_i$ implying countermonotonic (if $n = 2$), respectively comonotonic dependence for the random variables. Taking $\Pi(\mathbf{u}) := \prod_{i=1}^n u_i$ we obtain independence. Finally, we want to stress that the lower Fréchet bound is not a copula for $n \geq 3$.

2.2 Bound on wVaR and Known Optimality Results

Let F^- denote the left-continuous version of a distribution function F , i.e. $F^-(x) = \mathbf{P}(X < x) = F(x-)$. For $C \in \mathcal{C}^n$, univariate distributions F_1, \dots, F_n and $s \in \mathbb{R}$ we define

$$\sigma_{C,+}(F_1, \dots, F_n)(s) := \int_{\{\sum_{i=1}^n x_i < s\}} dC(F_1(x_1), \dots, F_n(x_n)),$$

$$\tau_{C,+}(F_1, \dots, F_n)(s) := \sup_{x_1, \dots, x_{n-1} \in \mathbb{R}} C \left(F_1(x_1), \dots, F_{n-1}(x_{n-1}), F_n^- \left(s - \sum_{i=1}^{n-1} x_i \right) \right),$$

where $\sigma_{C,+}(F_1, \dots, F_n)(s) = \mathbf{P}(X_1 + \dots + X_n < s)$ for a portfolio (X_1, \dots, X_n) with marginal distributions F_1, \dots, F_n and copula C . The following result yields distributional bounds for $\sigma_{C,+}(F_1, \dots, F_n)(s)$ and $\text{VaR}_\alpha(\sum_{i=1}^n X_i)$ when no information about the underlying dependence structure is available. A more general version can be found in Embrechts et al. (2003, Theorem 3.1) and Embrechts and Puccetti (2006, Theorem 3.1), where results are given for non-decreasing functionals in the presence of partial information.

Proposition 1 *Let (X_1, \dots, X_n) have marginal distributions F_1, \dots, F_n and copula $C \in \mathcal{C}^n$. Then for every real s and every $\alpha \in (0, 1)$ we have that*

$$\sigma_{C,+}(F_1, \dots, F_n)(s) \geq \tau_{W,+}(F_1, \dots, F_n)(s) \tag{1}$$

implying $\text{VaR}_\alpha(\sum_{i=1}^n X_i) \leq \text{wVaR}_\alpha(\sum_{i=1}^n X_i) \leq \tau_{W,+}(F_1, \dots, F_n)^{-1}(\alpha)$.

Note that for practical applications, the F_i 's are assumed to be known but C is unknown. A long history exists about the sharpness of these bounds. Makarov (1981) provided the first result for the sum of two random variables. Later, using a geometric approach, Frank et al. (1987) restated the result using the copula language. The pointwise best possible nature of the bounds in the two-dimensional case was finally proved in Williamson and Downs (1990) for non-decreasing functionals. Below we reformulate their optimality theorem for the sum. More historical references can be found in the introduction of Embrechts and Puccetti (2006).

Proposition 2 *Let (X_1, X_2) have marginal distributions F_1, F_2 and define $C_{\tilde{\alpha}} \in \mathcal{C}^2$ for $\tilde{\alpha} \in [0, 1]$ as*

$$C_{\tilde{\alpha}}(u_1, u_2) := \begin{cases} \max\{\tilde{\alpha}, W(u_1, u_2)\} & \text{if } (u_1, u_2) \in [\tilde{\alpha}, 1]^2, \\ M(u_1, u_2) & \text{otherwise.} \end{cases} \tag{2}$$

Then, choosing $\tilde{\alpha} = \alpha(s) := \tau_{W,+}(F_1, F_2)(s)$, we obtain $\sigma_{C_{\tilde{\alpha}},+}(F_1, F_2)(s) = \alpha(s)$. Hence, for any $\alpha \in (0, 1)$, $\text{wVaR}_\alpha(X_1 + X_2) = \tau_{W,+}(F_1, F_2)^{-1}(\alpha)$ is attained under $C_{\tilde{\alpha}}, \tilde{\alpha} = \alpha$.

Remark 2

- (a) Observe that, given some $C_L \in \mathcal{C}^2$, a similar result holds assuming partial information $C \geq C_L$ on the unknown copula C and substituting $W(u_1, u_2)$ by $C_L(u_1, u_2)$.
- (b) Taking $C_L(\mathbf{u}), \mathbf{u} \in [0, 1]^n, n \geq 3$ instead of $C_L(u_1, u_2)$, Eq. 2 is not a copula. In the no information case, this immediately follows from the fact that the lower Fréchet bounds is not a copula for $n \geq 3$. In the presence of partial information, we refer to the example by Geiss and Páivinen reported in Embrechts and Puccetti (2006).

Without mentioning the idea of copulas, Rüschendorf (1982) gave the same result stated by Frank et al. (1987) extending it for the sum of n uniform random variables.

Proposition 3 *The best possible lower bound on the distribution of $\sum_{i=1}^n X_i$ with (X_1, \dots, X_n) having standard uniform marginals is $\min\{(2s/n - 1)^+, 1\}$ for $s \in (0, n)$. This implies $\text{wVaR}_\alpha(\sum_{i=1}^n X_i) = n(1 + \alpha)/2$ for $\alpha \in (0, 1)$.*

Till now, this and a similar expression for binomial marginals are the only known analytical results for the multidimensional problem.

3 Worst Value-at-Risk Scenarios for the Multidimensional Problem

The above result of Rüschendorf (1982) provides sharpness of the bounds for the n -dimensional problem for uniform marginals. An analytical generalization of Eq. 2, replacing $W(u_1, u_2)$ by $W(\mathbf{u})$, $\mathbf{u} \in [0, 1]^n$, $n \geq 3$, does not lead to sharp bounds for the multidimensional case. Below we take a more geometric approach. From this point of view, the problem at hand consists in maximizing the probability of the set $G_s := \{\mathbf{x} \in \mathbb{R}^n : x_1 + \dots + x_n \geq s\}$. We transport the problem onto the unit square through $h : \mathbb{R}^n \rightarrow [0, 1]^n$, $h(\mathbf{x}) := (F_1(x_1), \dots, F_n(x_n))$ and denote

$$A_s := h(G_s) = \{\mathbf{u} \in [0, 1]^n : F_1^{-1}(u_1) + \dots + F_n^{-1}(u_n) \geq s\}. \tag{3}$$

By definition, for $s \in \mathbb{R}$, we have that $\text{wVaR}_{\alpha(s)}(X_1 + \dots + X_n) = s$, when $1 - \alpha(s) := \sup_{C \in \mathcal{C}^n} (1 - \sigma_{C,+}(F_1, \dots, F_n)(s))$. Let \mathbf{P}_C denote the probability induced by a copula C . The worst Value-at-Risk dependence scenario at level α therefore solves the equality

$$\mathbf{P}_C (A_{\text{wVaR}_{\alpha}(X_1+\dots+X_n)}) = 1 - \alpha. \tag{4}$$

3.1 Geometrical Properties of $C_{\tilde{\alpha}} \in \mathcal{C}^2$ with $\tilde{\alpha} = \tau_{W,+}(F_1, F_2)(s)$

In the two-dimensional case, applying Proposition 2, we immediately see that $C_{\tilde{\alpha}}$ satisfies Eq. 4 if $\tilde{\alpha} = \alpha$. Moreover, for a uniform portfolio (U_1, U_2) , Embrechts et al. (2005, Proposition 9) yields that this is the only copula putting measure $1 - \alpha$ on $A_{\text{wVaR}_{\alpha}(U_1+U_2)}$ with $\text{wVaR}_{\alpha}(U_1 + U_2) = 1 + \alpha$. Therefore, in this case the density of C_{α} in $A_{1+\alpha}$ is concentrated on the boundary $\underline{H}_{\alpha} = \underline{A}_{1+\alpha}$; see Fig. 1 (left).

Figure 1 highlights the geometric idea underlying the worst scenario $C_{\tilde{\alpha}}(u_1, u_2)$. The gray areas represent the sets A_s for a uniform portfolio ($s=1+\alpha=1.25$) and a Lognormal(0.4, 1) portfolio ($s=4$), respectively. The boundary of A_s can be written as $\underline{A}_s := \{(F_1(t), F_2(s - t)), t \in \mathbb{R}\}$. We denote




Fig. 1 Sets A_s and boundaries \underline{A}_s for a two-dimensional uniform portfolio for $s=1.25$ (left) and Lognormal(0.4, 1) portfolio for $s=4$ (right). Together we plot the support \underline{H}_{α} of $C_{\tilde{\alpha}}$ for $\tilde{\alpha} = \alpha = \tau_{W,+}(F_1, F_2)(s)$ and the (upper) supports for $\tilde{\alpha} = \alpha_1 < \alpha$ and $\tilde{\alpha} = \alpha_2 > \alpha$

$\underline{H}_{\tilde{\alpha}}$ the support of $C_{\tilde{\alpha}}$ restricted to $[\tilde{\alpha}, 1]^2$. In general dimensions, we refer to the support restricted to $[\tilde{\alpha}, 1]^n$ as upper support. The solution of the problem for the uniform portfolio leads then to an optimizing copula, which upper support coincides with the boundary $\underline{A}_{1+\alpha}$.

Remark 3 The choice $\tilde{\alpha} = \tau_{W,+}(F_1, F_2)(s)$ in Proposition 2 implies that $\underline{H}_{\tilde{\alpha}}$ lies in A_s and is tangent to \underline{A}_s . Since $C_{\tilde{\alpha}} \in \mathcal{C}^2$, the density on $\underline{H}_{\tilde{\alpha}}$ is uniformly distributed and proportional to its length $l(\underline{H}_{\tilde{\alpha}})$, say. Therefore C_{α} maximizes the density on A_s . In fact, a different choice of $\tilde{\alpha}$ would decrease the probability of A_s . Trivially $l(\underline{H}_{\alpha_2}) < l(\underline{H}_{\alpha})$ for $\alpha_2 > \alpha$, whereas the shape of \underline{A}_s implies

$$l(\underline{H}_{\alpha_1}) = l(\underline{H}_{\alpha}) + 2\sqrt{2}(\alpha - \alpha_1) - l(\underline{H}_{\alpha} \cap A_s^c) < l(\underline{H}_{\alpha}) \quad \text{if } \alpha_1 < \alpha.$$

3.2 Worst VaR Scenario for a n -dimensional Uniform Portfolio

In this section we consider a uniform portfolio. Similar to the previous section, the uniform case will lead to an optimizing copula for general marginals.

The solution of the worst VaR question consists of maximizing the probability of a certain region of \mathbb{R}^n . As illustrated in the previous section, we transport the problem onto the n -dimensional unit cube and investigate the shape of the support of the copulas putting maximal measure on Eq. 3. For a n -dimensional uniform portfolio and $n-1 \leq s \leq n$, the region of the space where the probability has to be maximized is $A_s = \{\mathbf{u} \in [0, 1]^n : \sum_{i=1}^n u_i \geq s\}$ with boundary $\underline{A}_s = \{\mathbf{u} \in [0, 1]^n : \sum_{i=1}^n u_i = s\}$. For $0 \leq s \leq n-1$ the problem has a trivial solution. Because of the uniformity of the marginals, the upper support of a copula maximizing the probability of A_s has to lie in $A_s \cap [\tilde{\alpha}, 1]^n$ for some appropriate $\tilde{\alpha} \geq \alpha^* := s - (n-1)$ with $\tilde{\alpha} = \alpha^*$ when $n = 2$.

Theorem 1 *Let $\tilde{\alpha} \in [\alpha^*, 1)$ and $C_{\tilde{\alpha}} : [0, 1]^n \rightarrow [0, 1]$ be a function with support in $\{\mathbf{u} \in [0, \tilde{\alpha}]^n : u_1 = \dots = u_n\} \cup \{\mathbf{u} \in [\tilde{\alpha}, 1]^n : \sum_{i=1}^n u_i \geq s\}$ for $s \in [n-1, n]$. A necessary condition for $C_{\tilde{\alpha}}$ to be a copula is that $\tilde{\alpha} = \tilde{\alpha} := 2s/n - 1$, i.e. that the support in $[\tilde{\alpha}, 1]^n$ lies in $H_{\tilde{\alpha}} := \{\mathbf{u} \in [\tilde{\alpha}, 1]^n : \sum_{i=1}^n u_i \geq n(1 + \tilde{\alpha})/2\}$.*

Proof Assume $\tilde{\alpha} \in [\alpha^*, \tilde{\alpha}]$ and $C_{\tilde{\alpha}} \in \mathcal{C}^n$ with corresponding measure $\mu_{\tilde{\alpha}}$. Let $S_i := \{\mathbf{u} \in [0, 1]^n : \alpha^* \leq u_i \leq \tilde{\alpha}\}$, $i = 1, \dots, n-1$, $S_n := \{\mathbf{u} \in [0, 1]^n : 1 - (\tilde{\alpha} - \alpha^*) \leq u_n \leq 1\}$ and set $E_i = S_i \cap A_s^n$, $i = 1, \dots, n$. By definition, $E_i \subset E_n$ for all i . Since $C_{\tilde{\alpha}}$ is a copula with upper support in $\{\mathbf{u} \in [\tilde{\alpha}, 1]^n : \sum_{i=1}^n u_i \geq s\}$, $\mu_{\tilde{\alpha}}(E_1) = \dots = \mu_{\tilde{\alpha}}(E_{n-1}) = \tilde{\alpha} - \alpha^*$ whereas $\mu_{\tilde{\alpha}}(E_n) = \tilde{\alpha} - \alpha^*$. It immediately follows that $\mu_{\tilde{\alpha}}(E_i) = 0$ for $i = 1, \dots, n-1$ and $\tilde{\alpha} = \alpha^* = \tilde{\alpha}$. \square

Remark 4

(a) Geometrically, Theorem 1 implies that if $C_{\tilde{\alpha}} \in \mathcal{C}^n$, the set

$$\underline{H}_{\tilde{\alpha}} := [\tilde{\alpha}, 1]^n \cap \underline{A}_{\frac{n}{2}(1+\tilde{\alpha})} = \left\{ \mathbf{u} \in [\tilde{\alpha}, 1]^n : \sum_{i=1}^n u_i = \frac{n}{2}(1 + \tilde{\alpha}) \right\} \quad (5)$$

is symmetric with respect to its center $((1 + \tilde{\alpha})/2, \dots, (1 + \tilde{\alpha})/2)$.

(b) Observe that the analytic generalization of (2) with $W(u_1, u_2)$ replaced by $W(\mathbf{u})$ has upper support $\{\mathbf{u} \in [\tilde{\alpha}, 1]^3 : u_1 + u_2 + u_3 = 2\}$.

Next we extend the two-dimensional result of Embrechts et al. (2005, Proposition 9) to general dimensions and we provide the existence of a copula with support as in Eq. 5.

Theorem 2 Assume $\tilde{C}, \tilde{C} \in \mathcal{C}^n$ to have supports on H_0^n and H_0^n , respectively. Let $\mu_{\tilde{C}}$ be the measure induced by \tilde{C} . Then for $H_0^{n+} := \{\mathbf{u} \in [0, 1]^n : \sum_{i=1}^n u_i > n/2\}$ and $H_0^{n-} := \{\mathbf{u} \in [0, 1]^n : \sum_{i=1}^n u_i < n/2\}$ it holds that

$$\mu_{\tilde{C}}(H_0^{n-}) = 0 \Leftrightarrow \mu_{\tilde{C}}(H_0^{n+}) = 0,$$

and the two copulas have the same support.

Proof Assume $\mu_{\tilde{C}}(H_0^{n-}) = 0$ and $\mu_{\tilde{C}}(H_0^{n+}) > 0$. Consider the independence copula Π with support $[0, 1]^n$. Since $\tilde{C}, \tilde{C} \in \mathcal{C}^n$, there exist operators $\nu, \tilde{\nu} : \mathcal{C}^n \rightarrow \mathcal{C}^n$ with $\varphi := \tilde{\nu} \circ \nu^{-1} \neq Id$ such that $\tilde{C} = \nu(\Pi)$ and $\tilde{C} = \tilde{\nu}(\Pi)$. It follows that $\mu_{\varphi(\tilde{C})}(H_0^{n-}) = 0$ and $\mu_{\varphi(\tilde{C})}(H_0^{n+}) > 0$. On the contrary, in order to preserve the uniformity of the marginals, any operator $\tilde{\varphi} : \mathcal{C}^n \rightarrow \mathcal{C}^n$ with $\mu_{\tilde{\varphi}(\tilde{C})}(H_0^{n+}) > 0$, implies $\mu_{\tilde{\varphi}(\tilde{C})}(H_0^{n-}) > 0$, which concludes the proof. \square

Theorem 3 Let $C_{\tilde{\alpha}} : [0, 1]^n \rightarrow [0, 1]$ have support $H_{\tilde{\alpha}}$ as in Eq. 5 on $[\tilde{\alpha}, 1]^n$. Then there exists a sequence of copulas $C_{N,\tilde{\alpha}} \in \mathcal{C}^n, N \in 2\mathbb{N} + 1$ such that

$$C_{\tilde{\alpha}}(\mathbf{u}) := \begin{cases} \lim_{N \rightarrow \infty} C_{N,\tilde{\alpha}}(\mathbf{u}) & \text{if } \mathbf{u} \in [\tilde{\alpha}, 1]^n, \\ M(\mathbf{u}) & \text{otherwise.} \end{cases}$$

is a copula.

Proof Without loss of generality, we consider $\tilde{\alpha} = 0$ with $H_0 = [0, 1]^n \cap \underline{A}_{n/2}$. For $N \in 2\mathbb{N} + 1$ we consider the partition $I := [0, 1] = \cup_{k=1}^N I_k$, where $I_k := [\frac{k-1}{N}, \frac{k}{N}]$. We identify the set $I_{k_1} \times \dots \times I_{k_n}$ with the point (k_1, \dots, k_n) and define its measure as follows. For any $k = 1, \dots, (N+1)/2$ and $1 \leq \bar{k} < k$ we set the functions

$$g_1(k) := \left| \{I_k \times I^{n-1}\} \cap H_0^{(N)} \right|, \quad g_2(k, \bar{k}) := \left| \{I_k \times I_{\bar{k}} \times I^{n-2}\} \cap H_0^{(N)} \right|,$$

where $H_0^{(N)} := \{(k_1, \dots, k_n) \in \{1, \dots, N\}^n : \frac{n}{2} - \frac{1}{N} < \sum_{i=1}^n k_i \leq \frac{n}{2} + \frac{1}{N}\}$. Then we define

$$f_0^{(N)}(k_1, \dots, k_n) := \begin{cases} f^* \left(\min_{1 \leq d \leq n} k_d \right) & \text{if } (k_1, \dots, k_n) \in H_0^{(N)}, \\ 0 & \text{otherwise,} \end{cases}$$

where $f^*(k) := (\frac{1}{N} - (n - 1) \sum_{1 \leq \bar{k} < k} g_2(k, \bar{k}) f^*(\bar{k})) (g_1(k) - (n - 1) \sum_{1 \leq \bar{k} < k} g_2(k, \bar{k}))^{-1}$.

The above construction defines a copula on the grid $\{k_1, \dots, k_n\}^N$. The set $I_k \times I^{n-1}$ denotes the k th slice of $[0, 1]^n$ along the first dimension. Therefore $g_1(k)$ counts the number of points on such a slice which also lie in $H_0^{(N)}$. Similarly $g_2(k, \bar{k})$ counts the number of points on $H_0^{(N)}$ which are on the k th slice along the first dimension and on the \bar{k} th slice along the second one. By the symmetry of the support, we could define these functions using any other two dimensions. Moreover, by definition, all the slices have width $1/N$. The idea is then to consider $I_0 \times I^{n-1}$ and weight each point in order to have total measure $1/N$. In doing this, by symmetry, we assign a measure to all the points lying on an equivalent slice for any of the other dimensions. We continue with $I_1 \times I^{n-1}$ assigning a weight only to the missing points. To do this we only have to take into account the missing points on slice k , i.e. $(n - 1) \sum_{1 \leq \bar{k} < k} g_2(k, \bar{k}) f^*(\bar{k})$. By the symmetry of $H_0^{(N)}$, we only evaluate slices $k = 1, \dots, (N+1)/2$. Using f^* we finally assign probability weights to the points with respect of the marginal constraints.

For any N , by construction, the function $C_{N,0} : [0, 1]^n \rightarrow [0, 1]$ defined through

$$C_{N,0}(\mathbf{u}) := \sum_{k_1=1}^{k(u_1)} \cdots \sum_{k_n=1}^{k(u_n)} f_0^{(N)}(k_1, \dots, k_n), \quad k(u) := \sup \left\{ k \geq 1 : \frac{k}{N} \leq u \right\}$$

is a copula. Setting $C_0(\mathbf{u}) := \lim_{N \rightarrow \infty} C_{N,0}(\mathbf{u})$ we then obtain that

$$\begin{aligned} C_0(u_1, \dots, u_{i-1}, 0, u_{i+1}, \dots, u_n) &= \lim_{N \rightarrow \infty} C_{N,0}(u_1, \dots, u_{i-1}, 0, u_{i+1}, \dots, u_n) \\ &\leq \lim_{N \rightarrow \infty} 1/N = 0, \\ C_0(1, \dots, 1, u_i, 1, \dots, 1) &= \lim_{N \rightarrow \infty} C_{N,0}(1, \dots, 1, u_i, 1, \dots, 1) \\ &= \lim_{N \rightarrow \infty} k(u_i)/N = u_i, \end{aligned}$$

and for $\mathbf{u}, \mathbf{v} \in [0, 1]^n$ with $\mathbf{u} \leq \mathbf{v}$ (componentwise) and $u_{j,1} = u_j, u_{j,2} = v_j, j = 1, \dots, n$

$$\begin{aligned} &\sum_{i_1=1}^2 \cdots \sum_{i_n=1}^2 (-1)^{i_1+\dots+i_n} C_0(u_{1,i_1}, \dots, u_{n,i_n}) \\ &= \lim_{N \rightarrow \infty} \sum_{i_1=1}^2 \cdots \sum_{i_n=1}^2 (-1)^{i_1+\dots+i_n} C_{N,0}(u_{1,i_1}, \dots, u_{n,i_n}) \geq 0. \end{aligned}$$

It follows that the conditions given in Remark 1 are satisfied and $C_0 \in \mathcal{C}^n$. \square




Fig. 2 Sets $A_{3(1+\tilde{\alpha})/2}$, $H_{\tilde{\alpha}}$ and $\underline{H}_{\tilde{\alpha}}$ for a three-dimensional uniform portfolio. The set $\underline{H}_{\tilde{\alpha}}$ with $\tilde{\alpha} = \alpha$ is the upper support of any copula leading to $wVaR_{\alpha}(X_1 + X_2 + X_3)$

Remark 5 Similarly as in the above proof, it is possible to construct other copulas with support $\underline{H}_{\tilde{\alpha}}$. We denote the family of the copulas sharing this support by $\mathcal{C}_{\tilde{\alpha}}^n$.

By Theorems 1, 2 and 3, any copula putting probability $1 - \tilde{\alpha} = 1 - (2s/n - 1)$ on $A_{n(1+\tilde{\alpha})/2}$ has support $\underline{H}_{\tilde{\alpha}}$ as in Eq. 5. Figure 2 illustrates these results in the three-dimensional case.

As a consequence we obtain the result of Rüschendorf (1982) given in Proposition 3. We restate it here using the language of copulas.

Corollary 1 *Let $C_{\tilde{\alpha}} \in \mathcal{C}_{\tilde{\alpha}}^n$ with $\tilde{\alpha} = \alpha$. Then*

$$\mathbf{P}_{C_{\tilde{\alpha}}} \{X_1 + \dots + X_n < s\} = \alpha \tag{6}$$

for $s = n(1 + \alpha)/2$ and the best possible lower bound on the distribution function of $X_1 + \dots + X_n$ for uniform marginals is $\min\{(2s/n - 1)^+, 1\}$ for $s \in (0, n)$.

Proof The worst dependence scenario for Value-at-Risk at level α satisfies Eq. 4. Taking $C_{\tilde{\alpha}} \in \mathcal{C}_{\tilde{\alpha}}^n$, we obtain $\mathbf{P}_{C_{\tilde{\alpha}}}(A_{n(1+\tilde{\alpha})/2}) = \mathbf{P}_{C_{\tilde{\alpha}}}(\underline{H}_{n(1+\tilde{\alpha})/2}) = 1 - \tilde{\alpha}$. Then equality (4) is satisfied for $\tilde{\alpha} = \alpha$ and $wVaR_{\alpha}(X_1 + \dots + X_n) = n(1 + \alpha)/2$ which implies Eq. 6. \square

3.3 Worst VaR Scenario for a General Portfolio

Relying on the solution for the uniform case studied in the previous section, we provide an answer for a general portfolio with marginals F_1, \dots, F_n . Although we illustrate the case of a three-dimensional portfolio, our arguments remain valid in higher dimensions. We recall that, for a portfolio (X_1, X_2) , the copula leading to the worst possible Value-at-Risk $wVaR_{\alpha}(X_1 + X_2)$ is indeed the solution of the uniform case for $\tilde{\alpha} = \alpha$. This follows from the uniformity of the density on the upper support; see Fig. 1 and Remark 3. In general, the worst value $wVaR_{\alpha}$ is not attained under a $C_{\tilde{\alpha}} \in \mathcal{C}_{\tilde{\alpha}}^n$ with $\tilde{\alpha} = \alpha$.

Example 1 Consider the portfolio (X_1, X_2, X_3) for $X_i \sim \text{Pareto}(1/\xi_i)$ with tail distribution function $F_i(x) = (1 + x)^{-1/\xi_i}$, $i = 1, 2, 3$. Assume $\xi_i = 0.7$, $i = 1, 2, 3$. In Fig. 3 (left) we illustrate the surface \underline{A}_s for $s = 21.4$ together with the upper support $\underline{H}_{\tilde{\alpha}}$ of $C_{\tilde{\alpha}}$ for $\tilde{\alpha} = 0.9$. On the right we plot \underline{A}_s for $s = 22.7$ with $\underline{H}_{\tilde{\alpha}}$ for $\tilde{\alpha} = 0.895$. Computing the probability of A_s under these two dependence structures we obtain

$$\mathbf{P}_{C_{0.9}}(X_1 + X_2 + X_3 \geq 21.4) = \mathbf{P}_{C_{0.895}}(X_1 + X_2 + X_3 \geq 22.7) = 0.1$$

and therefore the Value-at-Risk of the sum at level $\alpha = 0.9$ under $C_{0.9}$ is smaller that under $C_{0.895}$.

The message coming from Example 1 is that choosing the upper support tangent to \underline{A}_s , i.e. $\tilde{\alpha} = \alpha$, in general does not imply a worst dependence scenario. This is due to the distribution of the density on the support. Indeed the marginal constraints imply that for $n > 2$ the density is not uniformly distributed but concentrated on the border of $\underline{H}_{\tilde{\alpha}}$ and more thinly when reaching the center $(n(1 + \tilde{\alpha})/2, \dots, n(1 + \tilde{\alpha})/2)$. This can be easily seen in the three-dimensional case looking at the projection of $\underline{H}_{\tilde{\alpha}}$ on $[0, 1]^2$.

In Fig. 1 (right) we take $\tilde{\alpha} < \alpha$. This implies that the measure on the upper support is greater than $1 - \alpha$. Contrary, for $\tilde{\alpha} > \alpha$, a portion of the support does not lie in A_s . In Remark 3 we discussed the two-dimensional situation, where the density is proportional to the length of the support and every choice of $\tilde{\alpha}$ different from α leads to a better scenario for the problem at hand. In the general case, cutting some portion of the support does not necessarily imply a better scenario. In fact the increment of probability on the boundary could compensate the reduction in some other region. For $\tilde{\alpha}$ sufficiently small, we lose too much density on A_s . Trivially, $\tilde{\alpha} > \alpha$ implies a better scenario. From the solution of the uniform problem given by Theorems 1, 2 and 3 and the




Fig. 3 Surfaces \underline{A}_s for a three-dimensional Pareto(0.7) portfolio with $s = 21.4$ (left) and $s = 22.7$ (right). We plot $\underline{H}_{\tilde{\alpha}}$ for $\tilde{\alpha} = 0.9$ on the left and $\tilde{\alpha} = 0.895$ on the right

distribution of the probability on its upper support, we immediately obtain the following result.

Theorem 4 *Let (X_1, \dots, X_n) be a portfolio with marginal distributions F_1, \dots, F_n . Then $wVaR_\alpha(X_1 + \dots + X_n)$ is attained under a copula $C_{\tilde{\alpha}} \in \mathcal{C}_\alpha^n$ for some $\tilde{\alpha} \leq \alpha$ depending on the marginal distributions. Using the same notation as in Eq. 4, we have that*

$$\sup\{\mathbf{P}_{C_{\tilde{\alpha}}}(A_{wVaR_\alpha(X_1+\dots+X_n)}) : C_{\tilde{\alpha}} \in \mathcal{C}_\alpha^n, 0 < \tilde{\alpha} \leq \alpha\} = 1 - \alpha.$$

Remark 6 In contrast to the two-dimensional case, in dimensions higher than two a copula $C_{\tilde{\alpha}}$ leading to $wVaR_\alpha(X_1 + \dots + X_n)$ depends upon the choice of the marginals. In fact the region of the support where we lose probability is given by $H_{\tilde{\alpha}} \cap \{\mathbf{u} \in [0, 1]^n : F_1^{-1}(u_1) + \dots + F_n^{-1}(u_n) < s\}$ and depends on F_1, \dots, F_n .

4 Applications

In this section we apply Theorem 4 and compute the worst-possible Value-at-Risk for the sum at level α for a three-dimensional portfolio (X_1, X_2, X_3) . The random variables $X_i, i = 1, 2, 3$ are Pareto($1/\xi_i$) distributed with tails $\overline{F}_i(x) = (1 + x)^{-1/\xi_i}$. We solve the problem for $\alpha = 0.9, 0.95, 0.99$ (typically used for market or credit risk) and $\alpha = 0.995, 0.999$ (values used in operational risk) and this for various model assumptions.

Assumption I $X_i \sim \text{Pareto}(1/\xi_i)$ with $\xi_1 = \xi_2 = \xi_3 = 0.7$,

Assumption II $X_i \sim \text{Pareto}(1/\xi_i)$ with $\xi_1 = 0.7504, \xi_2 = 0.6607$ and $\xi_3 = 0.2815$,

Assumption III $X_i \sim \text{Pareto}(1/\xi_i)$ with $\xi_1 = 1.1905, \xi_2 = 1.3889$ and $\xi_3 = 1.2195$.

The main features of these assumptions are: they are all power-tailed, homogeneous as in I, or heterogeneous as in II and III. Assumption II corresponds to a finite mean situation whereas III corresponds to an infinite

Table 1 Values of $VaR_\alpha(X_1 + X_2 + X_3)$ for Assumption I under $C_\alpha, C_{\tilde{\alpha}}$ and M

$\downarrow C, \rightarrow \alpha$	0.9	0.95	0.99	0.995	0.999	0.9999
M	13.0	21.4	72.3	119.4	374.7	1, 889.9
C_α	21.4	36.7	119.5	196.0	611.1	3, 074.7
$C_{\tilde{\alpha}}$	22.7	38.6	123.8	205.2	634.3	3, 120.0
$\tilde{\alpha}$	0.895	0.948	0.989	0.9948	0.9989	0.99989

In the last row we give the values of $\tilde{\alpha}$ yielding the worst dependence structure and $wVaR_\alpha(X_1 + X_2 + X_3)$

Table 2 Values of $\text{VaR}_\alpha(X_1 + X_2 + X_3)$ for Assumption II under $C_\alpha, C_{\tilde{\alpha}}$ and M

$\downarrow C, \xrightarrow{\alpha}$	0.9	0.95	0.99	0.995	0.999	0.9999
M	9.1	16.0	53.3	87.9	278.3	1,453.4
C_α	13.6	22.7	70.5	114.1	348.1	1,749.9
$C_{\tilde{\alpha}}$	13.6	22.7	70.5	114.1	360.5	1,981.0
$\tilde{\alpha}$	0.9	0.95	0.99	0.995	0.99865	0.999865

mean model. The ξ -values chosen correspond to examples often encountered in QRM practice. For Assumptions II and III; see for instance Moscadelli (2004). Based on Theorem 4 and the upper support $\underline{H}_{\tilde{\alpha}}^n$ of $C_{\tilde{\alpha}} \in \mathcal{C}_{\tilde{\alpha}}^n$, we propose the following numerical procedure. For given $s \in \mathbb{R}$ and $\tilde{\alpha} \in (0, 1)$, analogously as in the proof of Theorem 3, for $N \in 2\mathbb{N} + 1$, we discretize the unit cube $[\tilde{\alpha}, 1]^3$ through

$$[\tilde{\alpha}, 1] = \cup_{k=1}^N I_k, \quad I_k := \left[\tilde{\alpha} + \frac{k-1}{N}, \tilde{\alpha} + \frac{k}{N} \right]$$

and we identify the set $I_{k_1} \times I_{k_2} \times I_{k_3}$ with the point $(k_1, k_2, k_3) \in \{1, \dots, N\}^3$.

Further we consider the sets $A_s^{(N)}$ and $\underline{H}_{\tilde{\alpha}}^{(N)}$ as discretized versions of A_s and $\underline{H}_{\tilde{\alpha}}$, respectively. We let $\mathbf{w} \in \mathbb{R}^{N^3}$ be a vector containing the probability weights of the points in $[0, 1]^3$. We then generate a vector $\mathbf{f} \in \mathbb{R}^{N^3}$ with entry one when the corresponding point lies on $[0, 1]^3 \setminus A_s^{(N)} \cap \underline{H}_{\tilde{\alpha}}^{(N)}$ and zero elsewhere. Similarly we create a $N^3 \times 3N$ matrix A providing the marginal restrictions. Finally we solve the optimization problem

$$\min_{\mathbf{w}} \mathbf{f}^T \mathbf{w}, \quad A\mathbf{w} = \left(\frac{1}{N}, \dots, \frac{1}{N}\right)^T, \quad \mathbf{w} \in [0, 1]^{N^3}. \tag{7}$$

It follows that $s = \mathbf{wVaR}_\alpha(X_1 + X_2 + X_3)$ at level $\alpha = \tilde{\alpha} + \mathbf{f}^T \hat{\mathbf{w}}$, where $\hat{\mathbf{w}}$ is the solution of Eq. 7. Any copula leading to \mathbf{wVaR}_α has support $\underline{H}_{\tilde{\alpha}}^{(N)}$.

We illustrate the above procedure for the Assumptions I, II and III. Together with the worst-case \mathbf{wVaR}_α , in the Tables 1, 2 and 3 we provide the values under $C_{\tilde{\alpha}}$ with $\tilde{\alpha} = \alpha$ and for the comonotonic copula M for which $\text{VaR}_\alpha(X_1 + X_2 + X_3) = \text{VaR}_\alpha(X_1) + \text{VaR}_\alpha(X_2) + \text{VaR}_\alpha(X_3)$.

Figures 4 and 5 show the densities on $A_s^{(N)} \cap \underline{H}_{\tilde{\alpha}}^{(N)}$ as functions of the parameter $\tilde{\alpha}$ for Assumptions I and II and levels $\alpha = 0.99$ and $\alpha = 0.9999$, respectively. The starting value for $\tilde{\alpha}$ is larger than α . We can observe that in both cases the densities increase linearly in $\tilde{\alpha}$ till reaching α . For the two assumptions we observe different behavior. For Assumption I, the densities

Table 3 Values of $\text{VaR}_\alpha(X_1 + X_2 + X_3)$ for Assumption III under $C_\alpha, C_{\tilde{\alpha}}$ and M

$\downarrow C, \xrightarrow{\alpha}$	0.9	0.95	0.99	0.995	0.999	0.9999
M	53.6	135.1	1,111.2	2,754.2	22,946.6	492,468.4
C_α	130.7	320.4	2,531.2	6,161.3	48,905	960,782
$C_{\tilde{\alpha}}$	144.3	351.5	2,700	6,500	52,000	980,000
$\tilde{\alpha}$	0.89	0.947	0.989	0.9943	0.99885	0.99988




Fig. 4 Densities on $A_s^{(N)} \cap H_{\tilde{\alpha}}^{(N)}$ for $s = 123.8$ ($\alpha = 0.99$) (top) and $s = 3, 120$ ($\alpha = 0.9999$) (bottom) as functions of $\tilde{\alpha}$ for Assumption I

continue to increase after α and, once a maximum is reached, they tend to zero. The $\tilde{\alpha}$ corresponding to this maxima, $\tilde{\alpha} = 0.989$ ($\alpha = 0.99$) and $\tilde{\alpha} = 0.99989$ ($\alpha = 0.9999$), give the worst dependence scenarios.

For Assumption II, the densities on $A_s^{(N)} \cap H_{\tilde{\alpha}}^{(N)}$ have a first maximum in $\tilde{\alpha} = \alpha$ and a second one for some $\tilde{\alpha} > \alpha$. In the case $\alpha = 0.9$ the worst dependence scenario is implied by the first maximum and the upper support is tangent to \underline{A}_s . For $\alpha = 0.9999$, the second maximum dominates.

In order to understand the different nature between the two assumptions, we look at the supports plotted in Fig. 6. The idea is as follows. We set the upper support tangent to A_s (with s chosen such that $\tilde{\alpha} = \alpha = 0.9$) and we shift it by taking values of $\tilde{\alpha}$ smaller than α . The set $A_s^{(N)} \cap H_{\tilde{\alpha}}^{(N)}$ is illustrated for $\tilde{\alpha} = 0.895$ and $\tilde{\alpha} = 0.885$ under Assumptions I and II. Remark that a smaller $\tilde{\alpha}$ implies a larger cut of the support and an increment of the probability on $[\tilde{\alpha}, 1]^3$. At this point we recall that the density is not homogeneous on




Fig. 5 Densities on $A_s^{(N)} \cap H_{\tilde{\alpha}}^{(N)}$ for $s = 70.5.8$ ($\alpha = 0.99$) (top) and $s = 1, 981$ ($\alpha = 0.9999$) (bottom) as functions of $\tilde{\alpha}$ for Assumption II




Fig. 6 Upper supports for Assumptions I (top) and II (bottom). In both cases we take $\alpha = 0.9$ and we consider $\tilde{\alpha} = 0.895$ (left) and $\tilde{\alpha} = 0.885$ (right)

the support and more concentrated when reaching the border. The different dynamics observed in Figs. 4 and 5 are due to the regions where the support is cut. In Assumption I (with equal marginals) the support loses probability in the center. Hence the probability on $A_s^{(N)} \cap \underline{H}_{\tilde{\alpha}}^{(N)}$ decreases only when $\tilde{\alpha}$ is small enough; see Fig. 4. On the other hand, if the tail of one distribution dominates the others, the cut arises near the border. This is the case for Assumption II for instance, where the loss of probability can not be compensated for small adjustments of $\tilde{\alpha}$. With larger movements of the parameter, the cutted region includes the central region as in Fig. 6 (bottom/right) and the probability on $A_s^{(N)} \cap \underline{H}_{\tilde{\alpha}}^{(N)}$ grows again. Besides the region where the loss of probability occurs, the shape of the set \underline{A}_s plays a role. In particular, this explains the differences arising in Assumption II. For $\alpha = 0.99$, we observe a loss of probability for any small adjustment of $\tilde{\alpha}$, which is not compensated

Table 4 Values for $wVaR_{0.9}$, $wVaR_{0.999}$, $VaR_{0.9}$ under $C_{0.9}$ and $VaR_{0.999}$ under $C_{0.999}$ with the corresponding scaling factors

ξ	$wVaR_{0.9}$	$wVaR_{0.999}$	$\frac{wVaR_{0.999}}{wVaR_{0.9}}$	$VaR_{0.9}^{(C_{0.9})}$	$VaR_{0.999}^{(C_{0.999})}$	$\frac{wVaR_{0.999}^{(C_{0.999})}}{wVaR_{0.9}^{(C_{0.9})}}$
0.7	22.7	634.3	27.9	21.4	611	28.6
0.8	31.1	1,360	43.7	29.9	1,310	43.8
0.9	43.8	2,940	67.1	41.5	2,806	67.6
1.0	60.8	6,350	104.4	57	6,006	105.4
1.1	84.3	13,800	163.7	78	12,850	164.7
1.2	116.0	30,400	262.1	106.4	27,490	258.4
1.3	160.4	65,500	408.4	144.7	58,805	406.4
1.4	221.0	145,000	656.1	196.3	125,793	640.8
1.5	304	310,000	1,019.7	266	269,087	1,011.6

by the augmentation before the second maximum. The very sharp profile of \underline{A}_ξ for $\alpha = 0.9999$ allows the initial loss to be compensated as illustrated in Fig. 5 (bottom).

As further application of our methodology, we calculate $wVaR_\alpha(X_1 + X_2 + X_3)$ for an homogeneous portfolio (X_1, X_2, X_3) . We solve the problem for $\alpha = 0.9, 0.999$ and $X_i \sim \text{Pareto}(1/\xi), i = 1, 2, 3$ for different values of ξ . The following table gives the results of our numerical computations together with the scaling factors from $wVaR_{0.9}$ to $wVaR_{0.999}$ and from $VaR_{0.9}$ under $C_{0.9}$ to $VaR_{0.999}$ under $C_{0.999}$, respectively (Table 4). We observe that the scaling curve grows exponentially as a function of the parameter ξ . It is moreover interesting to note that the scaling curve for the Value-at-Risk computed for $\tilde{\alpha} = \alpha$, i.e. with tangent upper support, does not differ significantly from the worst one.

Remark 7 The computational complexity of our numerical procedure increases exponentially with the dimension of the portfolio. Therefore, even if the values obtained are numerically not the exact worst-possible VaRs, in high dimensions the values obtained under C_α can be used as a first approximation for $wVaR_\alpha$. More work on the numerical accuracy of the above procedure is called for.

5 Conclusion

In this paper we extend the geometrical properties of the copulae leading to the worst-possible Value-at-Risk at level α for the sum of two risks. These solutions depend upon the probability level α . We solve the problem for an n -dimensional portfolio and explain how, for $n \geq 3$, any worst-case scenarios $C_{\tilde{\alpha}}$ depends upon the choice of the marginals. In particular the worst scenarios are not obtained when the upper support of $C_{\tilde{\alpha}}$ is tangent to \underline{A}_ξ . However, when the dimension of the problem becomes high, the copulae with tangent upper support turn out to be useful in order to approximate $wVaR_\alpha$. We conclude emphasizing that the results presented in this paper can be easily

restated substituting A_s by $A_s^\psi := \{\mathbf{u} \in [0, 1]^n : \psi(F_1^{-1}(u_1), \dots, F_n^{-1}(u_n)) \geq s\}$ corresponding to the Value-at-Risk optimization question for general increasing functionals ψ .

Acknowledgements The authors would like to thank an anonymous referee for various comments which led to an improvement of the paper. The corresponding author would like to thank Andrea Binda (Zurich Insurance Company), Elia Noris and Stefano Demarta (ETH Zürich) for helpful discussions.

References

- Denuit, M., Dhane, J., Goovaerts, M.J., Kaas, R.: Actuarial Theory for Dependent Risks. Wiley, Chichester (2005)
- Embrechts, P., Puccetti, G.: Bounds for functions of multivariate risks. *J. Multivar. Anal.* **97**(2), 526–547 (2006)
- Embrechts, P., McNeil, A.J., Straumann, D.: Correlation and dependence in risk management: properties and pitfalls. In: Dempster, M. (ed.) *Risk Management: Value at Risk and Beyond* (Cambridge, 1998). pp.176–223. Cambridge University Press, Cambridge (2002)
- Embrechts, P., Höing, A., Juri, A.: Using copulae to bound the Value-at-Risk for functions of dependent risks. *Finance Stoch.* **7**(2), 145–167 (2003)
- Embrechts, P., Höing, A., Puccetti, G.: Worst VaR scenarios. *Insurance Math. Econom.* **37**(1), 134–155 (2005)
- Frank, M.J., Nelsen, R.B., Schweizer, B.: Best-possible bounds for the distribution of a sum—a problem of Kolmogorov. *Probab. Theory Relat. Fields* **74**(2), 199–211 (1987)
- Makarov, G.D.: Estimates for the distribution function of a sum of two random variables when the marginal distributions are fixed. *Theory Probab. Appl.* **26**(4), 803–806 (1981)
- McNeil, A.J., Frey, R., Embrechts, P.: *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton University Press, Princeton (2005)
- Moscadelli, M.: The modelling of operational risk: experience with the analysis of the data, collected by the Basel Committee. Banca d'Italia, Temi di discussione del Servizio Studi, no. 517-July 2004
- Nelsen, R.B.: *An Introduction to Copulas*. Springer, Berlin Heidelberg New York (1999)
- Neslehova, J., Embrechts, P., Chavez-Demoulin, V.: Infinite mean models and the LDA for operational risk. *J. Oper. Risk* **1**(1), 3–25 (2006)
- Rüschendorf, L.: Random variables with maximum sums. *Adv. Appl. Probab.* **14**(3), 623–632 (1982)
- Williamson, R.C., Downs, T.: Probabilistic arithmetic. I. Numerical methods for calculating convolutions and dependency bounds. *Int. J. Approx. Reason.* **4**(2), 89–158 (1990)