

Miner Deposita (2006) 41: 259–280  
DOI 10.1007/s00126-006-0056-8

## ARTICLE

Isabelle Chambefort · Robert Moritz

## Late Cretaceous structural control and Alpine overprint of the high-sulfidation Cu–Au epithermal Chelopech deposit, Srednogie belt, Bulgaria

Received: 16 September 2005 / Accepted: 10 February 2006 / Published online: 1 April 2006  
© Springer-Verlag 2006

**Abstract** The Chelopech epithermal high-sulfidation deposit is located in the Panagyurishte ore district in Bulgaria, which is defined by a NNW alignment of Upper Cretaceous porphyry–Cu and Cu–Au epithermal deposits, and forms part of the Eastern European Banat–Srednogie belt. Detailed structural mapping and drillcore descriptions have been used to define the structural evolution of the Chelopech deposit from the Late Cretaceous to the present. The Chelopech deposit is characterized by three fault populations including ~N55, ~N110, and ~N155-trending faults, which are also recognized in the entire Panagyurishte district. Mapping and 3-D modeling show that hydrothermal alteration and orebody geometry at Chelopech are controlled by the ~N55-trending and ~N110-trending faults. Moreover, the ~N155-trending faults are parallel to the regional ore deposit alignment of the Panagyurishte ore district. It is concluded that the three fault populations are early features and Late Cretaceous in age, and that they were active during high-sulfidation ore formation at Chelopech. However, the relative fault chronology cannot be deduced anymore due to Late Cretaceous and Tertiary tectonic overprint. Structurally controlled ore formation was followed by Senonian sandstone, limestone, and flysch deposition. The entire Late Cretaceous magmatic and sedimentary rock succession underwent folding, which produced WNW-oriented folds throughout the Panagyurishte district. A subsequent tectonic stage resulted in overthrusting of older rock units along ~NE-trending reverse faults on the Upper Cretaceous magmatic and sedimentary host rocks of the high-sulfidation epithermal deposit at Chelopech. The three fault populations contemporaneous

with ore formation, i.e., the ~N55-, ~N110- and ~N155-trending faults, were reactivated as thrusts or reverse faults, dextral strike-slip faults, and transfer faults, respectively, during this event. Previous studies indicate that the present-day setting is characterized by dextral transtensional strike-slip tectonics. The ~NE-trending overthrust affecting the Chelopech deposit and the reactivation of the ore-controlling faults are compatible with dextral strike-slip tectonics, but indicate local transpression, thus revealing that the Chelopech deposit might be sited at a transpressive offset within a generally transtensional strike-slip system. The early WNW-trending folds require a roughly NNE–SSW shortening, which is incompatible with the present-day dextral strike-slip tectonic setting and the ~NE-trending thrust formed during the tectonic overprint of the Chelopech deposit. This reveals a rotation of the principal stress axes after Late Cretaceous high-sulfidation ore formation and post-ore deposition of sedimentary rocks. The nature of the sedimentary rocks interlayered and immediately covering the Upper Cretaceous magmatic rocks hosting the Chelopech deposit indicates sedimentation and associated volcanism in an extensional setting immediately before ore formation. It is concluded that the Chelopech deposit was formed when the tectonic setting changed from extensional during Late Cretaceous basin sedimentation and magmatism, to compressional producing WNW-trending folds under a roughly NNE–SSW compression, possibly in a sinistral strike-slip system. Thus, like other world-class, high-sulfidation epithermal deposits, the Chelopech deposit was formed at the end of an extensional period or during a transient period of stress relaxation, which are particularly favorable tectonic settings for the formation of high-sulfidation epithermal deposits. The exceptional preservation of the Upper Cretaceous Chelopech epithermal deposit is explained by the combined deposition of a thick Senonian sedimentary sequence on top of the Upper Cretaceous magmatic host rocks of the deposit, and the later overthrust of older rock units on top of the deposit. Our study at Chelopech supports previous studies stating that post-ore basin sedimentation and tectonic processes provide the favorable environment

*Editorial handling:* B. Lehmann

I. Chambefort (✉) · R. Moritz  
Section des Sciences de la Terre,  
University of Geneva,  
rue des Maraichers, 13,  
1205 Geneva, Switzerland  
e-mail: Isabelle.Chambefort@terre.unige.ch  
Tel.: +41-22-3796821  
Fax: +41-22-3793210

to preserve old epithermal deposits from erosion. The tectonic evolution of the Chelopech deposit is similar to that of the entire Panagyurishte ore district. This coherence of the magmatic, hydrothermal, and tectonic events from north to south suggests that the ore deposits of the entire Panagyurishte ore district were formed in a similar tectonic environment.

**Keywords** Chelopech Cu–Au high-sulfidation epithermal deposit · Panagyurishte ore district · Bulgaria · Structural control · Geodynamic evolution

---

## Introduction

Precious metal epithermal ore deposits have generally a poor preservation potential due to their shallow depth of formation, and the likelihood of their erosion increases with age. Therefore, epithermal deposits are most commonly preserved in volcanic arcs of late Cenozoic age (Cooke and Simmons 2000; Hedenquist et al. 2000; Kesler et al. 2004). However, there are a number of older epithermal deposits, some of them with major economic importance, that have been preserved from erosion, such as the Upper Cretaceous deposits at Pueblo Viejo, Dominican Republic (Muntean et al. 1990; Kesler et al. 2003), and in the Camagüey district, Cuba (Kesler et al. 2004), the Jurassic deposits of Patagonia, Argentina (Shalamuk et al. 1997), the Paleozoic epithermal deposits in northeastern Queensland and New South Wales, Australia (Wood et al. 1990; Bobis et al. 1995; Masterman et al. 2002), and in the Western Tianshan, Xinjiang Province, China (Long et al. 2005). Preservation of pre-Cenozoic epithermal deposits is typically attributed to postmineralization tectonic processes and/or burial (e.g., Masterman et al. 2002; Kesler et al. 2004), subsequent to the extensional tectonic to near-neutral stress settings, which are considered as favorable environments for the development of epithermal deposits in magmatic arcs (Tosdal and Richards 2002; Sillitoe and Hedenquist 2003; Kesler et al. 2004; Tosdal 2004).

The Upper Cretaceous Banat–Timok–Srednogie belt (BTS) belt is a major ore province in Eastern Europe, linked to subduction-related magmatism during the convergence between Africa and Eurasia (Jankovic 1997; Berza et al. 1998; Ciobanu et al. 2002; Heinrich and Neubauer 2002). This belt hosts some of the major operating European gold mines exploiting high-sulfidation epithermal deposits, including Bor in the Serbian Timok district (Jankovic 1990; Jankovic et al. 1998) and Chelopech in the Bulgarian Panagyurishte district, which is comparable in tonnage and grade with Cenozoic world-class deposits of the circum-Pacific region, such as El Indio in Chile, Lepanto in the Philippines, and Pierina in Peru (Moritz et al. 2004). The Carpathian–Balkan arc is currently a major target for mineral exploration in Europe (Danielson 2005). Therefore, understanding the geologic environments that were favorable for the preservation of epithermal deposits in the Upper Cretaceous BTS belt is a key for successful mineral exploration programs in a

geologic setting that has undergone successive orogenic deformation events.

In this contribution, we present new data obtained at the high-sulfidation epithermal Chelopech deposit, during detailed structural, surface, and underground mapping from drill-core studies and 3-D reconstructions using GEMCOM data provided by the mine staff at Chelopech. These data allow us to unravel the geologic evolution of the Chelopech deposit within the Upper Cretaceous volcano-sedimentary arc, from the early magmatic regime and the favorable ore forming events to the younger geologic events that created a favorable environment for its preservation. Finally, the structural evolution proposed for the Chelopech area will be discussed with respect to previous structural studies carried out in the Panagyurishte ore district (Popov and Popov 2000; Antonov and Jelev 2001; Ivanov et al. 2001; Kouzmanov et al. 2002; Jelev et al. 2003) and recent geochronological data obtained on the Panagyurishte ore deposits and magmatic rocks (Kamenov et al. 2004; Von Quadt et al. 2005).

---

## Geodynamic models for the Banat–Timok–Srednogie belt

The Upper Cretaceous BTS belt is a segment of the Alpine–Balkan–Carpathian orogen (Neugebauer et al. 2001; Heinrich and Neubauer 2002; Neubauer 2002). This orogen resulted from the convergence between Africa and Europe and the closure of the Tethys in the past 100 Ma (Dabovski et al. 1991; Ricou et al. 1998). Magmatism in the BTS belt has been generally related to north to northeast subduction of the African Plate below the Eurasian Plate (Dewey et al. 1973; Radulescu and Sandulescu 1973; Herz and Savu 1974; Hsü et al. 1977; Ivanov 1988; Lips 2002). Recent studies on magmatic rocks in the belt have revealed that the Late Cretaceous magmatism has a calc-alkaline composition (Dupont et al. 2002; Stoykov et al. 2002; Kamenov et al. 2003a,b, 2004), and an island or continental arc setting has been proposed. Ricou et al. (1998) advanced that the Srednogie belt was formed during back-arc opening, linked to postcollisional subduction during the Senonian after migration of the subduction zone to the southwest, where it encountered the dense oceanic lithosphere of the Vardar basin. The existence of the Vardar Ocean is still the subject of debate.

In previous contributions, Boncevic (1976) and Popov et al. (1979) interpreted the Banat–Srednogie belt as a rift-related structure. Popov (1987, 2002), and Popov et al. (2001) proposed that the Late Cretaceous rifting was related to the emplacement of a large, sheet-like, mantle diapir, associated with postcollisional collapse. According to the latter author, the large-scale geodynamic setting changed at the end of the Turonian, resulting in uplift, and during the Senonian the regional tectonic setting was characterized by a stage of subsidence in an extensional environment, with the formation of horsts and grabens controlled by transcurrent oblique faults. The Late Cretaceous sedimentation, associated with crustal exten-

sion, was characterized by turbiditic sedimentation under submarine conditions. The late-rift stage, which is common in the entire Alpine–Balkan–Carpathian–Dinaride (ABCD) region except in the Western Alps, was associated with Alpine deformation and the formation of small intramontane grabens (Neubauer 2002; Ilic et al. 2005).

By contrast, Boccaletti et al. (1974), Berza et al. (1998), Neubauer (2002), and Neubauer et al. (2003a,b) suggested postcollision slab break-off as a trigger for Late Cretaceous magmatism and that ore genesis was contemporaneous with the formation of postcollisional collapse basins.


## Geology and metallogeny of the Panagyurishte ore district

The roughly east–west trending Srednogie tectonic zone is the Bulgarian segment of the BTS belt, located between the Balkan Zone in the north and the Rhodope Massif and the Sakar–Strandja Zone in the south (Fig. 1a; Boncev 1988; Ivanov 1988). The NNW-trending Panagyurishte ore district, approximately 60 to 90 km east of Sofia, belongs to the Central Srednogie zone (Fig. 1a) and has supplied

approximately 95% of the Bulgarian Cu and Au production up to the present (Mutafchiev and Petrunov 1996, unpublished report).

The pre-Mesozoic basement rocks of the Panagyurishte district (Fig. 1b) consist of two-mica migmatites, amphibolites, and gneisses of uncertain Precambrian age, referred to as the Pirdop Group (Dabovski 1988), Srednogie-type metamorphic rocks (Cheshitev et al. 1995) or the pre-Rhodopian Supergroup (Katskov and Iliev 1993), late Precambrian to Cambrian phyllites, chlorite schists and diabases of the Berkovitsa Group (Haydoutov 2001), and Paleozoic gabbrodiorites, quartz–diorites, tonalites, and granodiorites–granites (Dabovski et al. 1972; Kamenov et al. 2002). All these units underwent ductile deformation and associated low-grade metamorphism at ~100 Ma (Velichkova et al. 2004) and are unconformably covered by Turonian conglomerate and sandstone, containing metamorphic rock fragments and coal-bearing interbeds (Aiello et al. 1977; Moev and Antonov 1978; Stoykov and Pavlishina 2003).

These early sedimentary rocks are in turn crosscut and covered by Upper Cretaceous magmatic rocks, where subvolcanic and effusive rocks are predominant in the



**Fig. 1** a Major tectonic zones of Bulgaria (after Ivanov 1988). b Simplified geology of the Panagyurishte ore district (modified after Popov and Popov 2000). c Structural map of the Panagyurishte ore district (modified after Popov and Popov 2000 and Ivanov and Dimov 2002)

north, whereas intrusive rocks become more abundant in the south, which reveals a deeper erosional level in the southernmost part of the district (Fig. 1b). Andesites predominate in the northern and central Panagyurishte district, whereas dacites are more abundant in its southern sector (Boccaletti et al. 1978; Stanisheva-Vassileva 1980). Rhyodacites and rhyolites only occur in the central and southern Panagyurishte district (Dimitrov 1983; Nedialkov and Zartova 2002). Small, subvolcanic dacite, quartz monzodiorite, and granodiorite intrusions (mostly <1 km<sup>2</sup> in size) are comagmatic with the Cretaceous volcanic rocks. Larger sized, northwest-elongated, syntectonic, Upper Cretaceous granodioritic–granitic intrusions are restricted to the southernmost Panagyurishte district along the Iskar–Yavoritsa Shear Zone (Ivanov et al. 2001; Peytcheva et al. 2001), which corresponds to the transition between the Srednogorie zone and the Rhodopes (Fig. 1b). Argillaceous limestone, calcarenite, and sandstone, with abundant volcanic rock fragments, are interbedded with the Upper Cretaceous volcanic rocks. These sedimentary rocks are also Turonian in age, based on paleontological dating by Stoykov and Pavlishina (2003), instead of early Senonian as reported previously (Aiello et al. 1977; Moev and Antonov 1978). Aiello et al. (1977) proposed that this sedimentation could have occurred in an intra-arc basin, related to the destabilization of the volcanic edifice. The interbedded volcano-sedimentary rock assemblage is transgressively overlain by Santonian–Campanian red limestone of the Mirkovo Formation, and Campanian–Maastrichtian calcarenite and mudstone flysch of the Chugovista Formation (Fig. 1c; Aiello et al. 1977; Moev and Antonov 1978; Popov 2001). The Senonian flysch units were deposited as outer-fan lobes within one or more basins, which were closely related to the eruptive centers (Aiello et al. 1977) and have probably preserved the volcanic rocks from erosion notably in the northern part of the district (Stoykov and Pavlishina 2003).

The major ore deposit types of the Panagyurishte ore district (Fig. 1b,c) include high-sulfidation epithermal Cu–Au deposits hosted by volcanic and subsidiary sedimentary rocks (Petrunov 1995; Popov and Popov 1997; Strashimirov and Popov 2000; Kouzmanov et al. 2002, 2004; Moritz et al. 2004), and porphyry–Cu deposits hosted by the apical parts of subvolcanic to hypabyssal intrusions, and locally by volcanic and crystalline basement country rocks (Bogdanov 1986; Strashimirov and Popov 2000; Strashimirov et al. 2002; Von Quadt et al. 2002; Popov et al. 2003; Tarkian et al. 2003). There is generally a close spatial association of porphyry–Cu and high-sulfidation epithermal deposits in the Panagyurishte ore district (Fig. 1b,c; Petrunov et al. 1991; Kouzmanov 2001; Tsonev et al. 2000; Kouzmanov et al. 2001; Strashimirov et al. 2002; Moritz et al. 2004), which is also recognized in other parts of the BTS belt, such as at Bor in Serbia (Jankovic 1990; Jankovic et al. 1998). The porphyry–Cu and high-sulfidation Cu–Au deposits of the Panagyurishte district are aligned along a north–northwest oriented trend, which is oblique with respect to the east–west trending Srednogorie zone in Bulgaria (Fig. 1a). There is a southward

decrease in the age of the calc-alkaline magmatism and the associated ore deposits from about 91–92 Ma in the north at Elatsite and Chelopech, through 89–90 Ma at Medet and Assarel, about 86 Ma at Elshitsa and Vlaykov Vruh, and 78 Ma at Capitan Dimitriev in the southernmost part of the Panagyurishte district (Fig. 1b; Von Quadt et al. 2005).

---

### Geodynamic setting of the Panagyurishte ore district

The Panagyurishte tectonic evolution started during the Mesozoic and has been affected by several tectonic phases during the Cenozoic. The southern part of the district is bound by Tertiary intrusive rocks of the Rhodope Massif, which are separated from the Srednogorie belt by the Maritsa Fault (Fig. 1c); whereas the Stara Planina Paleozoic granitic rocks form the northern border, along the Balkan deep fault (Fig. 1c).

According to Ivanov (1988), the sedimentation in the Stara Planina and Srednogorie Zone started during the Triassic and is marked by shallower facies (conglomerate and sandstone), which overlies folded Paleozoic basement rocks. Recently, Velichkova et al. (2004) determined that these basement rocks were metamorphosed and ductilely deformed at ca. 100 Ma, before the formation of the Srednogorie back-arc basin. During the Senonian, the Balkan was marked by the opening of the Srednogorie back-arc basin (Ricou et al. 1998), which initiated volcanism and detrital sedimentation.

---

### Late Cretaceous tectonics

Recently, Kamenov et al. (2003a,b, 2004) and Von Quadt et al. (2003, 2005) proposed that Late Cretaceous sedimentary basin formation was initiated along strike–slip faults and suggested that the Late Cretaceous magmatism and ore formation was related to slab rollback to the south during northward subduction. This scenario is supported by the fact that there is a progressive age decrease of magmatic and ore forming events, starting in the north at about 92 Ma and ending at about 78 Ma in the south of Panagyurishte ore district (Kamenov et al. 2003a,b, 2004; Von Quadt et al. 2003, 2005). Lips (2002) contends that slab rollback operated only from approximately 30 Ma to the present day, and that it was the back-arc extension that initiated Late Cretaceous magmatism. According to Lips (2002), the low density and young age of the subducted lithosphere argue against a subducted slab-detachment scenario (e.g., Neubauer 2002) during the Late Cretaceous.

Jelev et al. (2003) suggested that the Late Cretaceous volcanism in the Chelopech area may have formed in pull-apart basins, which initiated along a NE-trending fault segment. The detrital sedimentary sequence in the Chelopech area, which is associated and covered by Upper Cretaceous magmatic rocks, has been dated by Stoykov and Pavlishina (2003) as Turonian, suggesting a rapid development of the pull-apart sedimentary basins and linked volcanism.

Palynological data on the flysch of the Chugovista Formation reveal a Maastrichtian age for these marine sedimentary rocks, which overlie the Turonian volcano-sedimentary succession (Stoykov and Pavlishina 2004). The deposition of a flysch succession during the Maastrichtian suggests syn-orogenic detrital terrigenous sedimentation in basins, which is probably linked to the Rhodopian shortening and a change of the orientation of the principal stress axes from an extensional to a compressional regime.


### The high-sulfidation Cu–Au epithermal Chelopech deposit

The Chelopech deposit is located in the northernmost part of the Panagyurishte ore district, about 7–8 km southeast of the major Elatsite porphyry-Cu deposit. The Chelopech deposit is hosted by Upper Cretaceous volcanic and volcano-sedimentary units, transgressively overlying Precambrian and Paleozoic metamorphic rocks (Figs. 2a and 3a). The Upper Cretaceous rock sequence consists of detrital sedimentary rocks derived from the basement, and andesitic, dacitic to trachyandesitic subvolcanic bodies, lava flows, agglomerate flows, tuffs, and epiclastic rocks. They are




transgressively covered by sandstone, argillaceous limestone, and the terrigenous flysch sequence of the Chugovista Formation (Fig. 2a). The orebodies are hosted by (1) a subvolcanic body with an andesitic texture and composition, associated with phreatomagmatic breccia, and (2) sedimentary rocks with oolitic, biodetrital and sandstone layers interbedded with (3) volcanic tephra-tuff containing accretionary lapilli and pumices (Jacquat 2003; Moritz et al. 2003; Chambefort 2005).

Laterally outward from the orebodies, there are four alteration assemblages at Chelopech, with (1) an innermost silicic zone with massive silica, sparsely developed vuggy silica, disseminated pyrite and aluminum–phosphate–sulfate (APS) minerals; (2) a quartz–kaolinite–dickite zone with pyrite, APS minerals, and anatase; (3) a widespread quartz–sericite alteration zone; and (4) an external propylitic zone. Below the present mining level (about 400 m below surface), samples from 2-km deep drillholes reveal that the alteration evolves into a diaspore–pyrophyllite–alunite–zünite–rutile–APS mineral assemblage (Petrunov 1989, 1995; Georgieva et al. 2002).

The mineralization is characterized by three successive ore stages (Petrunov 1994, 1995; Jacquat 2003). The first Fe–S stage consists of disseminated pyrite, which can become locally very abundant and results in a total



**Fig. 2** Surface maps of the Chelopech area. **a** Geological map of the Chelopech area. **b** Schematic structural map of the Chelopech deposit. Arrows show the location of the cross-sections in Fig. 3



QUATERNARY

□ Conglomerate

UPPER CRETACEOUS UNITS

□ Chugovista Formation, Flysch

■ Mirkovo Formation, Reddish limestone

■ Chelopech Formation, 2 mica sandstone

□ Resedimented and syn-eruptive breccia

■ Syn-eruptive volcanogenic breccia

□ Turonian sandstone

□ Andesite

□ Phreatomagmatic breccia

□ Volcanic tuff and sedimentary rocks

□ Dacite of the Petrovden zone

□ Silicified zone (mine data, personal communication)

□ Mixed injection and secondary polymictic hydrothermal breccia

□ Secondary mineralized breccia

□ Massive sulfide orebodies

□ Base-metal-gold vein, Vozdol

PALEOZOIC UNITS

□ Mica gneiss, Pyrdop Formation

□ Cambrian metamorphic rocks

□ F1, F2, F3 fault systems

◀ **Fig. 3** Cross-sections perpendicular to  $F_1$  and located **a** on Chelopech mine and **b** on Charlodere occurrence, **c** perpendicular to  $F_3$  and  $F_2$  and containing the exploitation blocks, 151, 150, 17, 18. The numbers 1 and 2 of the Chelopech thrust movements correspond to a compressional stage followed by the extensional stage

replacement of the host rocks to form massive sulfide orebodies, particularly in more permeable rock units such as clastic and calcareous sedimentary rocks and volcanic tuffs. This early stage is followed and partly brecciated by an intermediate Cu–As–S stage, predominantly as veins, which is the Au-bearing event and constitutes the economic ore mined at Chelopech. The mineral assemblage includes enargite, luzonite, covellite, goldfieldite, chalcopyrite, tennantite, bornite, and native gold, which constitutes the main Au carrier (Bonev et al. 2002). The late uneconomic Pb–Zn–S stage consists of galena, sphalerite, pyrite, chalcopyrite, and barite veins. The Chelopech Mine produces approximately 700,000 t/year of ore and contains 25 Mt measured and indicated reserves at 4 g/t Au and 1.5% Cu (Dundee Precious Metals Inc data; <http://www.dundeeprecious.com/>).

The Chelopech deposit is accompanied by two mineralized occurrences. The Charlodere occurrence is located about 1 km to the northeast of the Chelopech mine (Fig. 2a) and is considered as an exhumed part of the Chelopech deposit by Popov et al. (2000a). The authors considered that the Charlodere orebodies are lenslike to shear-zone types and probably controlled by radial faults. This occurrence is hosted by a strongly altered breccia of a volcanic origin and a massive andesitic body. The rocks are affected by a propylitic alteration, with the occurrence of hydrothermal biotite and chlorite, grading into quartz–sericitic alteration and an advanced argillic alteration, including alunite (Lerouge et al. 2003). The second occurrence is the polymetallic Vozdol prospect, about 1 km to the north–northeast of the Chelopech deposit (Fig. 2a), which consists of base-metal sulfides, quartz, carbonate, barite, and fluorite veins, surrounded by an alteration zone comprising carbonate, adularia, and sericite. This occurrence was considered as a low-sulfidation system by Mutafchiev and Petrunov (1996, unpublished report) and Popov et al. (2000a), and would be reclassified as an intermediate-sulfidation occurrence according to the new terminology of Hedenquist et al. (2000). The spatial association of the polymetallic Vozdol occurrence and the Chelopech deposit is analogous to other base-metal veins at the periphery of high-sulfidation systems (Sillitoe 1999; Hedenquist et al. 2000, 2001).

## The Chelopech structural setting

### Description of the fault system

The investigations on the Chelopech tectonic system discussed in this contribution are limited to the north by the Petrovden fault and to the south by the Chelopech thrust (Fig. 2). Figure 1c presents a structural map for the

Panagyurishte ore district based on data from Popov and Popov (2000), whereas Figs. 2b and 4 show in detail the new interpretative surface and underground structural maps of the Chelopech deposit. The Chelopech deposit is overlain by folded Upper Cretaceous sedimentary rocks of the Chelopech syncline, including the Chugovista, Mirkovo, and Chelopech Formations (Fig. 3a). The axial plane of the syncline has a N110–120° trend in the western part and has been affected by Tertiary deformation in the eastern part, described as Pyrenean deformation by Antonov and Jelev (2001). This late deformation induced a change in the plane orientation from N110 to N70°, associated with the Vozdol syncline affecting the sandstone of the Chelopech Formation (Fig. 2b).

Three principal fault orientations have been recognized in the Panagyurishte ore district, which are also present in the Chelopech area (Popov and Popov 2000; Ivanov and Dimov 2002; Jelev et al. 2003; this study):

- (1) ~NE-trending faults, termed  $F_1$  in this study, which include the Chelopech thrust (Fig. 2b)
- (2) ~E–W-trending faults, termed  $F_2$  in this study, essentially subvertical and which are parallel to the regional E–W trend of the Srednogorie belt
- (3) ~N155 faults, termed  $F_3$  in this study, parallel to the regional NNW-alignment of the ore deposits of the Panagyurishte ore district

These three fault orientations are identical to the major fault orientations described by Popov and Kovachev (1996) in the Chelopech area. They are also recognized in the entire Panagyurishte district (Fig. 1c).

Figure 5 is a geological and structural map of the Chelopech deposit by Popov and Kovachev (1996), and shows that these authors have also described volcanotectonic radial and concentric faults in the central part of the volcanic edifice of Chelopech, in addition to the regional NNW-uniform-trending faults. The activation of these faults and the associated intense fracturing and brecciation have been attributed by these authors to the formation of a caldera and the intrusion of several subvolcanic bodies along with the development of mineralization. The authors described post-ore volcanic rocks, termed as Vozdol Member, which are interpreted to intersect hydrothermally altered rocks and the orebodies in the northwestern part of the deposit (Fig. 5). However, these concentric faults and post-ore volcanic rocks have been neither recognized during our field investigations nor by the previous study of Jelev et al. (2003).

Our investigations at Chelopech show that:

$F_1$  faults (highlighted in red in all figures) have a ~N20 to N55 orientation, and dip approximately 40 to 50° to the south (Figs. 2b and 4). They are thrusts or reverse faults (Fig. 6a,c,d) and are generally associated with meter-scale to several tens of meter-sized folds with ~NE-trending axes (Fig. 6b). Shear sense indicators are recognized along the fault planes with P and Y shears and deflection of the foliation (Fig. 6a,c,d), which are characteristic of compression in a brittle regime (Passchier and Trouw 1998, p. 128). The folds are developed in the Upper Cretaceous




Fig. 4 Interpretative structural map of the Chelopech mine, level 405

sedimentary cover and are thus post-volcanic structures, with the N10 to N55-trending fold axes (Fig. 7, surface data stereograph). This fault system is also characterized in some places by sinistral strike-slip movement along the major Chelopech Thrust, as shown by the thrust slices at the Charlodere occurrence and in the southern part of the Chugovista Valley (Fig. 2b).

$F_2$  faults (highlighted in blue in all figures) have a  $\sim$ N90 to N110 trend with a subvertical to  $60^\circ$  southward dip. They are subparallel to the E–W orientation of the Srednogorie belt (Figs. 1c, 2b, and 4) and are conjugate dextral and sinistral strike-slip faults. The sense of movement along the  $F_2$  faults is not possible to determine from surface exposures; however, in underground exposures, they display essentially a dextral strike-slip movement (Fig. 4). On the basis of their orientations, this fault family can be considered as identical to the WNW-oriented thrusts attributed by Antonov and Moev (1978) and Popov and Popov (2000) to the so-called Laramian phase, which is Maastrichtian–Tertiary in age. However, no evidence of thrust characteristics has been found in this study.

$F_3$  faults, highlighted in green in all figures, are characterized by a  $\sim$ N135 to N170 trend with a vertical to  $60^\circ$  dip to both the east and west (Figs. 2b and 4). They form a conjugate system, with the dextral  $F_3$  faults having a

NE-directed dip and the conjugate sinistral  $F_3$  faults having a SW-directed dip.  $F_3$  faults are subparallel to the regional ore deposit alignment in the Panagyurishte ore district (Fig. 1c). Popov and Popov (2000), Popov et al. (2001), and others described similar fault orientations, which were defined as Laramian and Illyrian by these authors and which correspond to Maastrichtian–Tertiary, N135–N170-trending strike-slip movements.

Figure 7 displays equal area stereographs for the three principal fault generations identified during surface and underground mapping.  $F_1$  is the most abundant fault system and, together with the Chelopech Thrust, determines the present-day geometry of the southern flank of the Chelopech syncline (Figs. 2b and 3a). Fault orientations measured in the mine galleries exhibit a strong dispersion. Therefore, for the sake of clarity, underground faults are plotted on separate stereographs for each of the mining blocks (Fig. 7). The three fault types were observed in every mining block that was mapped.  $F_3$  faults are predominant in block 17 on level 395 (Figs. 4 and 7), the orebody of this block is completely overprinted by this fault system.  $F_1$  and  $F_2$  fault types are also present, notably on level 405, where faults exhibit a wide variety of orientations. Figure 8a shows the structural relationships of the orebody in block 17 on level 405 with the different fault




Fig. 5 Underground geology of the Chelopech deposit according to Popov and Kovachev (1996)

populations. The orebody is essentially delimited by faults.  $F_1$  and  $F_2$  faults determine the western and southern borders of the block and are displaced by  $F_3$  faults.

The mining blocks 17 and 18 are less faulted than the blocks in the southwestern part of the mine such as blocks 103, 150, and 151 (Figs. 4 and 8). Faults in block 151 on levels 400 and 405 are predominantly of the  $F_1$  type. Figure 8b displays a detailed structural map from block 151 on level 405. The southern limit of the orebody is controlled by an extensive  $F_1$  fault zone. The orebody is essentially developed in altered volcanic tuff and sedimentary rocks. Similar to blocks 17 and 18, the orebody of block 151 is overprinted by  $F_3$  faults. The southern limits of each orebody are generally more intensively faulted than the northern ones (Fig. 8b, Arizanov, personal communication, 2001).


#### Relative fault chronology


The chronology of the different fault generations and the geometric relationships among them are not obvious in the field. Different cross-sections have been produced to determine the geometric relationships (Fig. 3). They are based on surface and underground mapping, drillcore descriptions, 3-D GEMCOM models and preexisting

cross-sections of the Chelopech deposit (Mutafchiev and Petrunov 1996, unpublished report; Popov and Kovachev 1996). These cross-sections were oriented perpendicular to a given fault set. For example, the cross-sections of Fig. 3a, b are perpendicular to the orientation of the  $F_1$  fault system, whereas the cross-section of Fig. 3c is perpendicular to the orientation of the  $F_2$  and  $F_3$  fault systems. One has to keep in mind that the Panagyurishte district has been subject to various deformation stages (Popov and Popov 2000; Antonov and Jelev 2001). Therefore, the present-day kinematics revealed by the faults is only representative of the latest stage(s) of the tectonic evolution of the Panagyurishte district, particularly the dextral strike-slip tectonics described by Ivanov et al. (2001) and late-stage fault relaxation.

It appears that  $F_1$  faults are characterized by a late-stage reverse sense of movement associated with maximum vertical displacements of several hundred meters and which also overprint the Late Cretaceous post-mineralization sedimentary rocks covering the ore deposit, including the Mirkovo and Chugovista Formations (Fig. 3a). The silicified zone, which defines the innermost part of the alteration underground, has been displaced along the  $F_1$  faults. The Chelopech deposit has probably been tilted during this Tertiary deformation stage, which was defined as the Pyrenean Stage by Popov and Kovachev (1996).

**Fig. 6** Outcrop-scale structures associated with the  $F_1$  fault system. **a** Shear sense indicators in the flysch of the Chugovista Formation. **b** Fold in flysch of the Chugovista Formation, Chugovista valley. **c, d** Thrust structures in the mine galleries





**Fig. 7** Fault plane projections on equal area stereonets, lower hemisphere, for surface faults and faults in each of the exploitation blocks (*numbers* indicate the exploitation block and the level, respectively). Faults in mine galleries are displayed as pole projections for the sake of clarity



Detailed mapping of blocks 17 and 151 on level 405 (Fig. 8) shows that movement along the  $F_1$  and  $F_3$  faults displaced different parts of the orebodies with respect to each other and, in some places, crosscut the  $F_2$  faults (Fig. 8a). Our mapping shows that the long axes of the orebodies, the breccia bodies, and the silicified zones are subparallel to the orientation of the  $F_1$  and  $F_2$  faults (Fig. 4). Therefore, we conclude that the  $F_1$  and  $F_2$  fault systems controlled the emplacement of the orebodies during the Late Cretaceous, with the  $F_1$  faults being reactivated as thrusts or reverse faults during the Alpine orogeny, after magmatism and ore formation. The  $F_2$  faults exhibit the same orientation as the E–W-elongation of the Srednogorie belt, which is considered as a subduction-related volcanic belt (see above). Therefore it is coherent to consider this fault system as directly linked to Upper Cretaceous magmatism and thus mineralization. Furthermore, the  $F_2$  fault system has also been reactivated and displaced during Tertiary strike-slip movements along the  $F_1$  faults (Figs. 2b and 4).

Figure 3b shows a cross-section through the Charlodere occurrence (Fig. 2), where the orebodies and the silicified zones have a NE–SW elongation with a northern dip, which contrasts with the southern dip of the Chelopech orebodies (Fig. 3a). The Charlodere orebodies occur along  $F_1$  faults and have been affected by the reactivation of the  $F_1$  faults in a thrust movement after their formation, like the Chelopech orebodies. On the basis of drillcore description, the Charlodere silicified and mineralized zones are affected by NE–SW oriented faults. This cross-section shows the structural control of the mineralization and overprinting by the reactivation of the same  $F_1$  faults during the post-mineralization tectonic event. The present-day surface location of the Charlodere occurrence (Fig. 3b) can be explained if we consider that the Charlodere mineralization is a slice of the Chelopech deposit, which has been uplifted and exposed on the surface as a consequence of dextral strike-slip thrust movements as a positive flower structure (Woodcock and Fischer 1986) along the “Chelopech Thrust” (Figs. 2b and 3a,b).

Figure 3c displays a cross-section perpendicular to the  $F_3$  faults, along the mining blocks 151, 150, 17, and 18, respectively, from SW to NE. It clearly shows that the  $F_3$  faults and also the  $F_2$  faults overprint the Chelopech deposit with a normal sense of movement and delimit the present-day geometry of the deposit. The silicified zone was displaced by  $F_3$  faults. However, the cross-section on Fig. 3c still clearly reveals a subhorizontal geometry of the silicified zone, which is akin to typical, subhorizontal “mushroom shapes” of alteration zones described in numerous epithermal high-sulfidation deposits (e.g., Sillitoe 1997, 1999; Corbett and Leach 1998; Hedenquist et al. 2000), and likely reveals a lithological control of the alteration. Thus, although the deposit appears to be crosscut by the  $F_3$  faults, the displacement along them was relatively small, as shown by SW–NE sections.

The  $F_1$  faults were reactivated during deformation events postdating magmatism and mineralization. The  $F_3$  faults are interpreted as transfer respectively tear faults, which accommodated thrusting during reactivation of the  $F_1$  faults (Fig. 9). The  $F_3$  faults are characteristic of transtensional strike-slip faults, associated with a normal movement. Although one may attribute a late timing to the  $F_3$  faults based on present-day crosscutting relationships among faults, the regional north–northwest ore deposit alignment of the Panagyurishte district is parallel to the orientation of the  $F_3$  faults. Therefore, they are linked to a regional ore deposit control that was certainly active at the start of ore formation in the northern Panagyurishte ore district, i.e., when the Chelopech deposit was formed.

In conclusion, the relative fault chronology among  $F_1$ ,  $F_2$ , and  $F_3$  cannot be deduced anymore due to the Late Cretaceous–Tertiary overprint as a consequence of shortening across the Rhodopes and the Srednogorie zone (Ivanov 1988; Ricou et al. 1998), the Tertiary compressional tectonics at Chelopech, described as late Pyrenean structures by Popov and Popov (2000) and Antonov and Jevlev (2001), and the late dextral strike-slip tectonics (Ivanov et al. 2001). However, the geological relationship



- Hydrothermal breccia
- Altered dacite
- Phreatomagmatic breccia
- Ore body (cut off Cu 3%)
- Mineralized breccia with massive sulfide ore
- Altered sandstone
- Volcanic tuff and associated sedimentary rocks

◀ **Fig. 8** a Detailed map of block 17 on level 405. b Detailed map of block 151 on level 405


with orebody and alteration zone geometries indicates that all three fault families were active during ore formation.

The fault history presented in this study has some implications for exploration. Indeed, because of the Tertiary compressional tectonics, undiscovered mineralized zones of the Chelopech deposit could have been overthrust by presently mined orebodies. Furthermore, Tertiary transtensional dextral displacements along NNW-oriented  $F_3$  faults, i.e., parallel to the Panagyurishte alignment, may have also affected the mineralized system. Thus, undiscovered orebodies may be located at deeper structural levels northwesterly with respect to the present mine, such as beneath block 19W (Fig. 4) and in the southeastern parts of the Chelopech syncline (Fig. 3a).

### Geodynamic evolution of the Chelopech deposit and regional overview: a discussion


#### Tectonic evolution at Chelopech

Based on the stratigraphic and the geometric field relationships presented above, a summary of the tectonic evolution of the Chelopech area can be given as follows, from oldest to youngest (Fig. 10):



**Fig. 9** Interpretative structural map of the western part of the Chugovista Valley, showing the relationship between  $F_1$  and  $F_3$  faults. The numbers 1 and 2 of the Chelopech thrust movements correspond to a compressional stage followed by the extensional stage

- (1) The geometry of the alteration and mineralized zones and the shape of the hydrothermal breccia bodies show that the Chelopech hydrothermal system was controlled by the  $\sim N55$ -trending  $F_1$  and the  $\sim N110$ -trending  $F_2$  faults. In addition, the  $\sim N155$ -trending  $F_3$  faults are parallel to the regional NNW-trending ore distribution in the Panagyurishte district. Therefore, we conclude that these faults are early features, Late Cretaceous in age, which were present before deposition of the Upper Cretaceous sandstone, limestone and flysch of the Chelopech, Mirkovo and Chugovista Formations recognized in the Chelopech syncline (Fig. 3), and before the Tertiary tectonic overprint. Our study clearly documents the early nature of these faults and the control they have played during high-sulfidation ore formation.
- (2) The entire Upper Cretaceous rock succession was folded after deposition of the Upper Cretaceous sandstone, limestone, and flysch (turbiditic sandstone) of the Chelopech, Mirkovo, and Chugovista Formations, thus producing the WNW-oriented Chelopech syncline (Figs. 2b and 10b) after the Maastrichtian. These folds, attributed to the Laramian tectonic phase by Antonov and Moev (1978), Ivanov (1988), Popov and Popov (2000), Popov (2001), and Jelev et al. (2003), are recognized in the entire Panagyurishte district; thus, they are regional features.
- (3) The WNW-oriented folds were followed by thrusting along the  $\sim NE$ -trending Chelopech Thrust (Fig. 2b) and accompanied by folds with  $\sim NE$ -oriented axial planes in the Maastrichtian sedimentary cover rocks, such as the Vozdol syncline (Figs. 2a and 10c). The early  $F_1$  and  $F_2$  faults, which have controlled high-sulfidation ore formation, were reactivated as reverse faults or thrusts and dextral faults, respectively, during this compressional overprint, which also tilted the Chelopech deposit to the northwest. The  $F_3$  faults were reactivated as transfer faults during Tertiary thrusting along the  $F_1$  faults. This event, described as the Pyrenean and Illyrian phase by Popov and Kovachev (1996), Popov and Popov (2000), and Jelev et al. (2003), has only a limited development in the Panagyurishte district and besides the Chelopech area; they have only been recognized within the Mechit and Raina Knyaginya synclines, south of the town of Strelcha (Fig. 1b,c; Popov and Popov 2000).
- (4) During the Neogene kinematics, the Chelopech area was characterized by an extensional regime (Ivanov et al. 2001, Velichkova et al. 2004). Some of the  $F_1$  faults were reactivated as normal faults, as documented by the change in kinematics along the Chelopech Thrust to a normal fault along the southern margin of the Chelopech syncline (Figs. 2b and 10d). This extensional regime is also recognized in the entire Panagyurishte ore district with the formation of horsts and grabens, the formation of Cenozoic sedimentary basins and uplift of Paleozoic basement (Fig. 1b,c; Ivanov et al. 2001). The  $\sim N100$ -oriented dextral strike-slip faults, such as the Maritsa fault (Fig. 1c; Ivanov et al. 2001), are parallel to the  $F_2$



◀ **Fig. 10** Schematic tectonic evolution of the Chelopech deposit. **a** Late Cretaceous sinistral transtensional duplex system and setting of the magmatism and the mineralization. **b** Late-Maastrichtian to Tertiary Alpine deformation phase, formation of WNW-trending thrusts ( $F_2$ ) and associated N110°-trending folds (formation of the Chelopech syncline). **c** Tertiary Alpine Eocene phase with formation of a dextral transpressional duplex, thrust slices ( $F_1$ ) and associated folds post-dating ore formation, the folds have a NE-oriented axial plane. **d** Late Alpine-neotectonic dextral transtensional duplex, extensional system, reactivation of the Chelopech thrust fault as a normal fault

faults and are contemporaneous with movement along the  $F_3$  strike-slip faults.

Our investigation, documenting the structural control of the high-sulfidation orebodies and their tectonic overprint, together with previous structural investigations in the Panagyurishte district allow us to address the variation of the stress states during the geological evolution of this district. According to Ivanov et al. (2001), the present-day extensional regime in the Panagyurishte district (stage 4 above) is related to dextral transtensional strike-slip tectonics along the ~WNW-oriented Iskar–Yavoritsa Shear Zone (Fig. 1b). The formation of the ~NE-oriented Chelopech Thrust (Fig. 2b) and the ~NE-oriented Vozdol Syncline (Fig. 2b), recognized during stage 3, would be compatible with such a dextral strike-slip tectonic setting and a roughly N–S to NW–SE oriented compression ( $\sigma_1$ ) based on classical structural models (Woodcock and Fischer 1986; Sylvester 1988), although they record transpression rather than transtension. The orientation of the WNW-oriented folds of stage 2, such as the Chelopech syncline and affecting the Late Cretaceous magmatic and sedimentary rocks in the Panagyurishte district, are incompatible with dextral strike-slip tectonics along the ~WNW-oriented Iskar–Yavoritsa Shear Zone, according to classical structural models (Woodcock and Fischer 1986; Sylvester 1988). These folds require a roughly NNE-oriented compression ( $\sigma_1$ ), which could be compatible with a regional sinistral strike-slip tectonic setting (Woodcock and Fischer 1986; Sylvester 1988) within the Srednogorie belt. Therefore, the geometric relationships reveal that there has been a rotation of the orientation of the principal stress axes in the Panagyurishte district, from a roughly NNE-oriented compression during folding of the Late Cretaceous rocks covering and hosting the Chelopech deposit, toward a roughly N–S to NW–SE compression, as recorded by the younger NE-trending Chelopech Thrust and the present-day dextral transtensional strike-slip tectonics described by Ivanov et al. (2001). This interpretation agrees with the studies of Antonov and Jelev (2001) about post-mineralization tectonics at Chelopech. There is no clear structural evidence to constrain the tectonic setting during Late Cretaceous volcanism and sedimentation of the host rocks of the Chelopech deposit, as well as during ore formation. The nature of the sedimentary host rocks described by Aiello et al. (1977) and Moev and Antonov (1978) indicate sedimentation and, by association volcanism in an extensional basin setting before high-sulfida-

tion, ore formation. It can only be speculated whether the  $F_1$  and  $F_2$  faults, which are the principal faults controlling the mineralization at Chelopech, were already active during Late Cretaceous basin formation and magmatism. If this were the case, then their orientations would be compatible with sinistral strike–slip tectonics (compare the  $F_1$ ,  $F_2$ , and  $F_3$  fault orientations at Chelopech with Figs. 6e and 8a in Woodcock and Fischer 1986), which would have been transtensional during basin sedimentation and switched to transpressional during the WNW-trending folding of the Late Cretaceous sedimentary and volcanic rocks after (or during?) high-sulfidation ore formation.

#### Relationship between ore formation events and the tectonic evolution of the Panagyurishte ore district

To discuss and understand the geodynamic evolution of the Chelopech deposit, it is necessary to consider the deposit on a regional scale, with respect to both the E–W oriented Srednogorie belt and the NNE–SSW ore deposit alignment of the Panagyurishte ore district.

The Panagyurishte ore deposits are hosted by Late Cretaceous magmatic rocks, which throughout the Panagyurishte district, are progressively covered by the Campanian reddish limestone of the Mirkovo Formation (Popov and Popov 2000; Popov 2001; Popov et al. 2003; Stoykov and Pavlishina 2003; Bogdanov et al. 2004). The limestone of the Mirkovo Formation is unaltered and contains at the bottom, a volcano-sedimentary conglomeratic layer with altered and mineralized fragments (Popov and Popov 2000; Popov 2001). This observation indicates that the ore deposits of the Panagyurishte ore district were formed before the sedimentation of the Upper Cretaceous Mirkovo Formation (Popov and Popov 2000) and, therefore, before Late Cretaceous and Tertiary tectonic deformation.

In the entire Panagyurishte ore district, the Late-Cretaceous volcano-sedimentary and overlying limestone and flysch units have been affected by WNW-oriented folds at the end of the Maastrichtian (Laramian stage of Karagjuleva et al. 1974; Antonov and Moev 1978; Popov and Kovachev 1996; Popov and Popov 2000; and Popov 2001). Numerous faults accompany the Maastrichtian–Tertiary folds, the most abundant being WNW-trending thrusts (Fig. 1c), such as the Panagyurishte and the Elshitsa faults, and associated diagonal NNW-oriented faults partly interpreted by Popov and Popov (2000), and Popov (2001) as reactivated older faults. Locally, the WNW-oriented folds were overprinted by Tertiary northeast-oriented folds and faults at Vozdol and Chelopech, respectively (Fig. 2b), and Mechit and Raina Knyaginya synclines, south of the town of Strelcha (Fig. 1b,c; Popov and Popov 2000; Pyrenean deformation of Karagjuleva et al. 1974; Antonov and Moev 1978; Ivanov 1988; Popov and Kovachev 1996; Antonov and Jevlev 2001; Popov and Popov 2000; Popov 2001, 2002). At the present-day, the Panagyurishte district is composed of different horst and graben structures delimited by regional transform faults with a ~N110–120 trend

resulting from a transtensional, dextral strike–slip tectonics (Fig. 1c; Ivanov et al. 2001, personal communication).


The sequence of early ore formation, followed by Late Cretaceous deposition of sedimentary rocks (i.e., the Mirkovo Formation), Maastrichtian–Tertiary folding and thrusting, and late-stage transtensional dextral strike–slip deformation events, is identical throughout the Panagyurishte ore district and is comparable to the evolution we have described for the Chelopech area. Therefore, our literature review suggests a rotation of the principal stress axes after the ore formation events and before the present-day dextral strike–slip tectonic setting for the entire Panagyurishte district, because the WNW-oriented folds were formed after the emplacement of the ore deposits and because the orientation of these folds is incompatible with the orientation of the principal stress axes required for dextral strike–slip tectonics based on classical tectonic models (Woodcock and Fischer 1986; Sylvester 1988).

There is also a coherent orientation of the orebodies among the high-sulfidation deposits across the Panagyurishte district, including the Radka, Krassen, and Elshitsa deposits of the southern part of the district (Fig. 1b). The Radka orebodies follow a ~N120 orientation with a northern dip, i.e., parallel to the  $F_2$  fault system (Fig. 11a; from Popov and Popov 1997; Tsonev et al. 2000; Kouzmanov et al. 2002; Popov 2002; Bogdanov et al. 2004). The Krassen deposit is limited between two subparallel WNW-trending faults, and the lenticular orebodies, which are hosted by pipe-like tectonic breccia zone, have the same orientation with a dip of about 50° to NE (Bogdanov et al. 2004). The Elshitsa deposit is also hosted by a N110–115°-oriented dacitic subvolcanic body, which is parallel to the  $F_2$  fault system and the Elshitsa orebodies have the same orientation (Fig. 11b; from Chipchakova and Stefanov 1974; Kouzmanov 2001). According to Popov (2001), the Elshitsa volcano–tectonic system is characterized by an intense fracturing along N120–N130 and N60–N80-trending faults, and magmatic bodies follow the same trend. Thus, it appears that the Panagyurishte high-sulfidation epithermal deposits and associated magmatic rocks are locally controlled by the same  $F_1$  and  $F_2$  fault systems as in Chelopech and are aligned regionally along the NNW-oriented  $F_3$  fault pattern (Fig. 12). Therefore, we conclude that the ore deposits throughout the Panagyurishte district were formed under approximately similar stress conditions, i.e., under similar orientations of the principal stress axes. The coherent sequence of magmatic, hydrothermal and tectonic events, and orebody orientations from North to South, suggests that the ore deposits of the entire Panagyurishte ore district were formed in a similar tectonic environment as the one we have deduced for the Chelopech area.

---

#### Late Cretaceous tectonic setting linked with syn-arc magmatism and associated mineralization

Figure 13 is modified from Tosdal and Richards (2001) and displays a schematic diagram illustrating the relationship




**Fig. 11** a Cross-section of the Radka deposit (after Popov and Popov 1997; Tsonev et al. 2000; Kouzmanov et al. 2002). b. Cross-section of the Elshitsa deposit (after Chipchakova and Stefanov 1974; Popov et al. 2000b)

among convergent margin tectonics, upper plate structure, and magmatism during the evolution of the Panagyurishte ore district. According to Tosdal and Richards (2001, 2002), the development of porphyry–Cu and high-sulfidation epithermal deposits is favored during a relaxation of regional stress conditions. The major arc-parallel structures will be closed when the subduction setting is characterized by arc-normal compression, while during extensional or transtensional stress periods, these structures provide enhanced permeability, therefore favoring magma emplacement in the crust. The intersections of different fault generations in a transtensional regime will favor the development of strike-slip duplexes (Woodcock and

Fischer 1986) and can result in the formation of pull-apart basins.

By analogy to the model of Tosdal and Richards (2001, 2002), the WNW–E-oriented  $F_2$  faults of the Panagyurishte ore district are interpreted as major arc parallel faults during oblique subduction, and the NE-oriented  $F_1$  faults are considered as strike-slip faults delimiting the major arc parallel faults (Figs. 12 and 13), allowing the formation of pull-apart basins and eventually resulting in the emplacement of magmatic rocks and ore deposits at Elatsite and Chelopech.

Progressive southward migration of magmatism along a deep-seated structure, parallel to the orientation of the



**Fig. 12** Principal tectonic orientations of the Panagyurishte ore district in the Srednogorie belt. U–Pb ages for Elatsite and Capitan Dimitriev are from Von Quadt et al. (2003, 2005)


NNW-oriented  $F_3$  faults (Fig. 13b), is explained as a consequence of slab rollback (Kamenov et al. 2003a, 2004; Von Quadt et al. 2003), or a slab retreat (Handler et al. 2004; Von Quadt et al. 2005). Two additional pull-apart basins bound by strike–slip faults were formed within a tectonic setting characterized by strike–slip duplex south of the Chelopech–Elatsite stripe, the first one being the Assarel–Panagyurishte stripe (Fig. 13b), and the second one the southernmost Elshitsa–Radka stripe (Fig. 13c). Sedimentation of the volcanogenic sandstone of the Chelopech Formation is probably linked to the destruction of the volcanic edifice at Chelopech (Stoykov and Pavlishina 2003) during on-going extension as a result of the south-directed slab rollback or retreat.

## Conclusions

Our investigation shows that the emplacement of the orebodies and the alteration zones of the Chelopech high-sulfidation deposit have been controlled by ~N55 and ~N110-trending faults at a time when the tectonic setting changed from extensional, during Late Cretaceous basin sedimentation and magmatism, to compressional with WNW-oriented folding of the Upper Cretaceous host and cover rocks of the ore deposit, indicating a roughly NNE–

SSW compression. Thus, the Chelopech deposit is structurally controlled like other world-class high-sulfidation epithermal deposits (e.g., Mitchell and Leach 1991; Corbett and Leach 1998; Hedenquist et al. 1998), and was formed at the end of an extensional period or during a transient period of stress relaxation, which are particularly favorable tectonic settings for the formation of high-sulfidation epithermal deposits according to Tosdal and Richards (2002), Sillitoe and Hedenquist (2003), and Kesler et al. (2004). Although evidence is weak, the fault and fold orientations are compatible with an W to WNW-trending sinistral strike–slip system, subparallel to the regional Srednogorie orientation, switching from transtensional during basin sedimentation to late transpressional after deposition of Late Cretaceous sedimentary rocks and after ore formation. Similar to JeleV et al. (2003), we found no evidence to support the existence of concentric faults and a caldera scenario during Late Cretaceous magmatism as proposed in early studies by Popov and Kovachev (1996).

The Late Cretaceous magmatic, sedimentary, and ore-forming events were followed by a rotation of the orientation of the principal stress axes, whereby the Panagyurishte district was affected by a W to WNW-trending dextral transensional strike–slip system, which controlled the formation of Cenozoic sedimentary basins



**Fig. 13** Relationship between convergent margin tectonics, upper plate structure, and magmatism during the Panagyurishte ore district evolution. Localized zones of extension may be optimized during transtensional strain along arc-related structures (*MASH* melting, assimilation, storage and homogenization; modified from Tosdal and

Richards 2001). **a** Development of the Chelopech–Elatsite strike-slip system, **b** migration of the strike system to the South and development of the Panagyurishte–Assarel stripe, and **c** development of the southern Elshitsa–Radka strike-slip system

(Ivanov et al. 2001). The Chelopech ore deposit is unique among all the high-sulfidation deposits of the Panagyurishte district, because it is the only one where older rock units have been overthrust on the host rocks of the orebodies (Moritz et al. 2004; Chambefort 2005). The NE orientation of this late overthrust is compatible with the regional dextral strike-slip system recognized by Ivanov et al. (2001), but records local transpression, which possibly reveals that the Chelopech deposit is located at a transpressive offset within a generally transtensional strike-slip system.

Magmatism and hydrothermal ore formation at Chelopech developed very rapidly, as indicated by the Turonian age of both the sedimentary rocks interlayered with and immediately covering the Late Cretaceous volcanic rocks (Stoykov and Pavlishina 2003, 2004). The subsequent deposition of a thick Senonian sandstone, limestone, and flysch sequence, followed by overthrust of older rock units along the Chelopech Thrust, provided the favorable environment for preserving the Late Cretaceous epithermal deposit. Our study at Chelopech supports previous studies stating that basin sedimentation and post-ore tectonic processes are necessary to preserve old epithermal deposits from erosion (e.g., Masterman et al. 2002; Kesler et al. 2004).

The local structural evolution deduced from the Chelopech deposit is similar to the geologic evolution of the Panagyurishte ore district. Ore deposits of the central and southern parts of the Panagyurishte ore district are essentially controlled by WNW-trending faults, parallel to the orientation of the  $F_2$  fault system. The Panagyurishte magmatism was developed from north to south in three different stripes, probably in a sinistral transpressional strike-slip system, as a result of slab rollback or retreat linked to oblique convergence, during a period of approximately 14 Ma.

**Acknowledgements** This study was supported by the Swiss Science Foundation through the research grants 21-59041.99 and 200020-101853, in conjunction with the SCOPES Joint Research Project 7BUPJ62276.00/1. The authors would like to thank the staff of the Geology Department and particularly chief geologist A. Arizanov from the Chelopech Mine, Bulgaria, for arranging access to the mine and sharing geological information. M. Sartori is gratefully acknowledged for discussion and suggestions. R. Petrunov, S. Georgieva, S. Stoykov, and S. Jacquat are acknowledged for discussion and field work. We would like to thank R. Tosdal for his discussion, and F. Neubauer is thanked for his critical review, which allowed us to improve the manuscript. The authors are grateful to B. Lehmann for his editorial comments. This is a contribution to the ABCD-GEODE research program supported by the European Science Foundation.

## References

- Aiello E, Bartolini C, Boccaletti M, Gocev P, Karagjuleva J, Kostadinov V, Manetti P (1977) Sedimentary features of the Srednogorie zone (Bulgaria): an Upper Cretaceous intra-arc basin. *Sediment Geol* 19:39–68
- Antonov M, Moev M (1978) Structure of the Eastern part of the Sturzel-Chelopech strip. *Ann Univ Min Geol* 23:31–50 (in Bulgarian)
- Antonov M, Jelev V (2001) Transpression and oblique fold and thrust structures in Chelopech ore field (Bulgaria). *Ann Univ Min Geol* 43–44(1):45–49
- Berza T, Constantinescu E, Vlad, SN (1998) Upper Cretaceous magmatic series and associated mineralisation in the Carpathian-Balkan Orogen. *Resour Geol* 48:291–306
- Bobis RE, Jaireth S, Morrison GW (1995) The anatomy of a Carboniferous epithermal ore shoot at Pajingo, Queensland: setting, zoning, alteration and fluid conditions. *Econ Geol* 90:1776–1798
- Boccaletti M, Manetti P, Peccerillo A (1974) The Balkanids as an instance of back-arc thrust belt: possible relation with the Hellenids. *Geol Soc Amer Bull* 85:1077–1084
- Boccaletti M, Manetti P, Peccerillo A, Stanisheva-Vassileva G (1978) Late Cretaceous high-potassium volcanism in Eastern Srednogorie, Bulgaria. *Geol Soc Amer Bull* 89:439–447
- Bogdanov B (1986) Copper ore deposits in Bulgaria. *Schriftenreihe der Erdwissenschaftlichen Kommissionen by Osterreichische Akademie der Wissenschaften* 8:103–112
- Bogdanov K, Tsonev D, Popov K (2004) Mineral assemblages and genesis of the Cu–Au epithermal deposits in the southern part of the Panagyurishte ore district, Bulgaria. *Bull Geol Soc Of Greece, Proceedings of the 10th International congress, Thessaloniki, April 2004*
- Boncev ES (1976) Lineament-geosynclinal zones—zones of impermanent riftogenesis. *Geol Balk*
- Boncev ES (1988) Notes sur la tectonique alpine des Balkans. *Bull Soc Geol Fr* 4:241–249 (in French)
- Bonev IK, Kerestdjian T, Anatassova R, Andrew CJ (2002) Morphogenesis and composition of native gold in the Chelopech volcanic-hosted Au–Cu epithermal deposit, Srednogorie zone, Bulgaria. *Miner Depos* 37:614–629
- Chambefort I (2005) The Cu–Au Chelopech deposit Panagyurishte district, Bulgaria: volcanic setting, hydrothermal evolution and tectonic overprint of a Late Cretaceous high-sulfidation epithermal deposit. PhD thesis, University of Geneva
- Cheshitev G, Milanova V, Sapounov I, Choumachenko P (1995) Explanatory note to the geological map of Bulgaria in scale 1:100 000, Teteven map sheet, Avers, Sofia (in Bulgarian)
- Chipchakova S, Stefanov D (1974) Genetic types of argillites at the height of Golyamo Petelovo and in Elshitsa-West copper-pyrite deposit in the Panagyurishte ore district. In: Alexiev E, Mincheva-Stefanova J, Radonova T (eds) *Mineral genesis*. *Geol Inst Bulg Acad Sci* 437–454 (in Bulgarian)
- Ciobanu CL, Cook NG, Stein H (2002) Regional setting and Re–Os age of ores at Ocna de Fier Dognecea (Romania) in the context of the banatitic magmatic and metallogenic belt. *Miner Depos* 37:541–567
- Cooke D, Simmons S (2000) Characteristics and genesis of epithermal gold deposits. In: Hagemann SG, Brown PE (eds) *Gold in 2000*. *Rev Econ Geol* 13:221–244
- Corbett GJ, Leach TM (1998) Southwest Pacific rim gold–copper systems: structure, alteration, and mineralization. *Econ Geol Special Publication* 6:237
- Dabovski C (1988) Precambrian in the Srednogorie zone (Bulgaria). In: Cogne J, Kozhoukharov D, Krautner HG (eds) *Precambrian in younger fold belts*. *Essex* 841–847
- Dabovski C, Zagorchev I, Rouseva M, Chounev D (1972) Paleozoic granitoids in the Sushtinska Sredna Gora. *Annual UGP* 16:57–92 (in Bulgarian)
- Dabovski C, Harkovska A, Kamenov B, Mavrudchiev B, Stanisheva-Vassileva G, Yanev Y (1991) A geodynamic model of the Alpine magmatism in Bulgaria. *Geol Balk* 21:3–15
- Danielson V (2005) Carpathian arc key to Europe's mining renaissance. *Northern Miner* 91
- Dewey J, Pitman W, Bonnin J (1973) Plate tectonics and evolution of the Alpine system. *Geol Soc Amer Bull* 84:3137–3180

- Dimitrov C (1983) Senonian initial volcanic rocks south of Panagyurishte and Strelca. *Rev Bulg Geol Soc* 44:95–128 (in Bulgarian)
- Dupont A, Vander Auwera J, Pin C, Marincea S, Berza T (2002) Trace element and isotope (Sr, Nd) geochemistry of porphyry and skarn mineralizing Late Cretaceous intrusions from Banat, western South Carpathians, Romania. *Miner Depos* 37:568–586
- Georgieva S, Velinova N, Petrunov R, Moritz R, Chambefort I (2002) Aluminium Phosphate–Sulfate minerals in the Chelopech Cu–Au deposit: Spatial development, chemistry and genetic significance. *Geohim Mineral Petrol* 39:39–51
- Handler R, Neubauer F, Velichkova SH, Ivanov Z (2004)  $^{40}\text{Ar}/^{39}\text{Ar}$  age constraints on the timing of magmatism and post-magmatic cooling in the Panagyurishte region, Bulgaria. *Schweiz Mineral Petrogr Mitt* 84:119–132
- Haydoutov I (2001) The Balkan island-arc association in west Bulgaria. *Geol Balc* 31(1/2):109–110
- Hedenquist JW, Arribas A Jr, Reynolds TJ (1998) Evolution of an intrusion-centered hydrothermal system: Far Southeast-Lepanto porphyry and epithermal Cu–Au deposits, Philippines. *Econ Geol* 93:373–404
- Hedenquist JW, Arribas AR, Gonzalez-Urien E (2000) Exploration for epithermal gold deposits. In: Hagemann SG, Brown PE (eds) *Gold in 2000*. *Rev Econ Geol* 13:245–277
- Hedenquist JW, Claveria RJR, Villafuerte GP (2001) Types of sulfide-rich epithermal deposits and their affiliation to porphyry systems: Lepanto–Victoria–Far Southeast deposits, Philippines, as examples. *ProEXPLOR Congreso*, Lima, Perú, 29 April 2001 (released on CD)
- Heinrich CA, Neubauer F (2002) Cu–Au–Pb–Zn–Ag metallogeny of the Alpine–Balkan–Carpathian–Dinaride geodynamic province. *Miner Depos* 37:533–540
- Herz N, Savu H (1974) Plate tectonic history of Romania. *Geol Soc Amer Bull* 85:1429–1440
- Hsü KJ, Nachev IK, Vuchev VT (1977) Geologic evolution of Bulgaria in light of plate tectonics. *Tectonophysics* 40:245–256
- Ilic A, Neubauer F, Handler R (2005) Late Paleozoic–Mesozoic tectonics of the Dinarides revisited: implications from  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of detrital white micas. *Geology* 33:233–236
- Ivanov Z (1988) Aperçu général sur l'évolution géologique et structurale du massif des Rhodope dans le cadre des Balkanides. *Bull Soc Geol Fr* IV(8):227–240
- Ivanov Z, Dimov D (2002) Structure and tectonic evolution of the Panagyurishte Upper Cretaceous strip. *GEODE Workshop on the Srednogorie zone*, abstr vol, Sofia, 8
- Ivanov Z, Henry B, Dimov D, Georgiev N, Jordanova D, Jordonova N (2001) New model for Upper Cretaceous magma emplacement in the southwestern parts of Central Srednogorie–Petrostructural and AMS data. *Rom J Miner Deposits*, special volume, ABCD-GEODE 2001 workshop, Geodynamics and ore deposit evolution of the Alpin–Balkan–Carpathian–Dinaride Province, Vata Bai, Romania, 8–12 June 2001, 79:60
- Jacquat S (2003) Etude Paragénétique et Géochimique du Gisement Epithermal d'Or et de Cuivre de type "High-Sulfidation" de Chelopech, Bulgarie. *Diplôme d'Ingénieur Géologue*, University of Geneva (in French)
- Jankovic S (1990) Types of copper deposits related to volcanic environment in the Bor district, Yugoslavia. *Geol Rundsch* 79:467–478
- Jankovic S (1997) The Carpatho–Balkanides and adjacent area: a sector of the Tethyan Eurasian metallogenic belt. *Miner Depos* 32:426–433
- Jankovic S, Herrington RJ, Kozelj D (1998) The Bor and Madjanpek copper–gold deposits in the context of the Bor metallogenic zone (Serbia, Yugoslavia). In: Porter TM (ed) *Porphyry and hydrothermal copper and gold deposits; a global perspective*. Conference proceedings Australian Mineral Foundation, Glenside, South Australia, pp 169–178
- Jelev V, Antonov M, Arizanov A, Arnaudova R (2003) On the genetic model of Chelopech volcanic structure (Bulgaria). 50 years, *Ann Univ Min Geol, Geol Geophysics*, Sofia 46:47–51
- Kamenov BK, Von Quadt A, Peytcheva I (2002) New insight into petrology, geochemistry and dating of the Vejen pluton, Bulgaria. *Geohim Mineral Petrol* 39:3–25
- Kamenov BK, Moritz R, Nedialkov R, Peytcheva I, Von Quadt A, Stoykov S, Yanev Y, Zartova A (2003a) Petrology of the Late Cretaceous ore–magmatic centers from the Central Srednogorie, Bulgaria: magma evolution and sources. Final GEODE-ABCD (2003) workshop, Geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride Province, Seggauberg, Austria, abstr vol, 30
- Kamenov BK, Von Quadt A, Peytcheva I (2003b) New petrological, geochemical and isotopic data bearing on the genesis of Capitan–Dimitriev pluton, Central Srednogorie, Bulgaria. Final GEODE-ABCD (2003) workshop, Geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride Province, Seggauberg, Austria, abstr vol, 31
- Kamenov B, Yanev Y, Nedialkov R, Moritz R, Peytcheva I, Von Quadt A, Stoykov S, Zartova A (2004) An across-arc petrological transect through the Central Srednogorie Late-Cretaceous magmatic centers in Bulgaria. *Bulg Geol Soc Annual Scientific Conference "Geology 2004"*, 16–17 December, Proceedings, 35–37
- Karagjuleva J, Kostadinov V, Tzankov TZ, Goëv P (1974) Structure of the Panagyurishte strip east of the Topolnica river. *Geol Inst Bull, Geotectonics*, Sofia 23:231–301 (in Bulgarian with English abstract)
- Katskov N, Iliev K (1993) Panagyurishte map sheet. In: Explanatory note to the geological Map of Bulgaria in scale 1:100000, p 53 (in Bulgarian)
- Kesler SE, Russell N, McCurdy K (2003) Trace-element content of the Pueblo Viejo precious-metal deposits and their relation to other high-sulfidation epithermal systems. *Miner Depos* 38:668–682
- Kesler SE, Hall CM, Russell N, Pinero E, Sanchez CR, Perez RM, Moreira J (2004) Age of the Camaguey Gold–Silver district, Cuba: tectonic evolution and preservation of epithermal mineralization in volcanic arcs. *Econ Geol* 99:869–886
- Kouzmanov K (2001) Genèse de la concentration en métaux de base et précieux de Radka et Elshitsa (zone de Sredna Gora, Bulgarie): une approche par l'étude minéralogique, isotopique et des inclusions fluides. Ph.D. thesis, University of Orléans, France, p 437
- Kouzmanov K, Moritz R, Chiaradia M, Fontignie D, Ramboz C (2001) Sr and Pb isotope study of Au–Cu epithermal and porphyry-Cu deposits from the Southern part of the Panagyurishte district. In: Piestrzynski A et al (eds) *Mineral deposits at the beginning of the 21st century*, Proceedings 6th biennial SGA meeting, Krakow, Poland, pp 539–542
- Kouzmanov K, Bailly L, Ramboz C, Rouer O, Beny J-M (2002) Morphology, origin and infrared microthermometry of fluid inclusions in pyrite from the Radka epithermal copper deposit, Srednogorie zone, Bulgaria. *Miner Depos* 37:599–613
- Kouzmanov K, Ramboz C, Bailly L, Bogdanov K (2004) Genesis of high-sulfidation vinciennite-bearing Cu–As–Sn (Au) assemblage from the Radka copper epithermal deposit, Bulgaria: evidence from mineralogy and infrared microthermometry of enargite. *Can Miner* 42:1501–1521
- Lerouge C, Bailly L, Fléhoc C, Petrunov R, Kunov A, Georgieva S, Velinov I, Hikov A (2003) Preliminary results of a mineralogical and stable isotope study of alunite in Bulgaria—Constraints on its origin. In: Eliopoulos DG et al (eds) *Mineral exploration and sustainable development*, Proceedings 7th biennial SGA meeting, Athens, Greece, 24–28 August 2003, Millpress Rotterdam, pp 495–498
- Lips ALW (2002) Correlating magmatic–hydrothermal ore deposit formation over time with geodynamic processes in SE Europe. In: Blundell DJ, Neubauer F, Von Quadt A (eds) *The timing and location of major ore deposits in an evolving orogen*. *Geol Soc London Spec Publ* 204:69–79

- Long X, Hayward N, Begg G, Minlu F, Fangzheng W, Pirajno F (2005) The Jinxi–Yelmand high-sulfidation epithermal deposit, Western Tianshan, Xinjiang Province, P.R. China. *Ore Geol Rev* 26:17–37
- Masterman GJ, White NC, Wilson CJL, Pape D (2002) High-sulfidation gold deposits in ancient volcanic terranes: insights from the Mid-Paleozoic Peak Hill deposit, NSW. *Soc Econ Geol Newsletter* 51:10–16
- Mitchell AHG, Leach TM (1991) Epithermal gold in the Philippines; island arc metallogenesis, geothermal systems and geology. Acad Press, London, pp 457
- Moev M, Antonov M (1978) Stratigraphy of the Upper Cretaceous in the eastern part of the Sturguel–Chelopech strip. *Ann Univ Min Geol* 23:7–30 (in Bulgarian)
- Moritz R, Jacquat S, Chambefort I, Fontignie D, Petrunov R, Georgieva S, Von Quadt A (2003) Controls of ore formation at the high-sulphidation Au–Cu Chelopech deposit, Bulgaria: evidence from infrared fluid inclusion microthermometry of enargite and isotope systematics of barite. In: Eliopoulos DG et al (eds) Mineral exploration and sustainable development, Proceedings 7th biennial SGA meeting, Athens, Greece, 24–28 August 2003, Millpress Rotterdam, 1209–1212
- Moritz R, Kouzmanov K, Petrunov R (2004) Late Cretaceous Cu–Au epithermal deposits of the Panagyurishte district, Srednogie zone, Bulgaria. *Schweiz Mineral Petrogr Mitt* 84:79–99
- Muntean JL, Kesler SE, Russell N, Polanco J (1990) Evolution of the Monte Negro acid sulfate Au–Ag deposit, Pueblo Viejo, Dominican Republic: important factors in grade control. *Econ Geol* 85:1738–1758
- Nedialkov R, Zartova A (2002) Magmatism of the Assarel area and its ore generating capability. In: Moritz R, Von Quadt A (eds) GEODE workshop on the Srednogie zone, Sofia, Bulgaria, April 2002, Abstr vol 13
- Neubauer F (2002) Contrasting Late Cretaceous with Neogene ore provinces in the Alpine–Balkan–Carpathian–Dinaride collision belt. In: Blundell DJ, Neubauer F, Von Quadt A (eds) The timing and location of major ore deposits in an evolving orogen, *Geol Soc London Spec Publ* 204:81–102
- Neubauer F, GEODE-ABCD Geodynamics Working Group (2003) Late Cretaceous and Tertiary geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride orogen. Final GEODE-ABCD (2003a) workshop, Geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride Province, Seggau, Austria, abstr vol 40
- Neubauer F, Heinrich C, GEODE-ABCD Geodynamic Working Group (2003b) Late Cretaceous and Tertiary geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride orogen. In: Eliopoulos DG et al (eds) Mineral exploration and sustainable development, Proceedings 7th biennial SGA meeting, Athens, Greece, 24–28 August 2003, Millpress Rotterdam, pp 1133–1136
- Neugebauer J, Greiner B, Appel E (2001) Kinematics of the Alpine–West Carpathian orogen and palaeogeographic implications. *J Geol Soc (Lond)* 158:97–110
- Passchier CW, Trouw RAJ (1998) *Microtectonics*. Springer, Berlin Heidelberg New York, p 289
- Petrunov R (1989) Hypogene sulphate–sulphide zoning in a copper–pyrite deposit. *Comptes rendus Bulg Acad Sciences* 42:71–74
- Petrunov R (1994) Mineral paragenesis and physicochemical conditions of ore-forming in the Chelopech deposit. PhD Thesis, Geology Institute, Sofia, p 178 (in Bulgarian)
- Petrunov R (1995) Ore mineral paragenesis and zoning in the deposit of Chelopech. *Geohim Mineral Petrol* 30:89–98 (in Bulgarian)
- Petrunov R, Dragov P, Neykov H (1991) Polyelemental (with As, Sn, V, Bi, Ag, Te, Ge, Se etc.) mineralization in Assarel porphyry–copper deposit. *Rev Bulg Geol Soc* 52:1–7 (in Bulgarian)
- Peytcheva I, Von Quadt A, Kamenov B, Ivanov Z, Georgiev N (2001) New isotope data for Upper Cretaceous magma emplacement in the southern and south-western parts of Central Srednogie. In: ABCD-GEODE 2001, Workshop, Vata Bai, Romania, Abstr vol, 82–83
- Peytcheva I, Von Quadt A, Kouzmanov K, Bogdanov K (2003) Timing of magmatism and mineralization in Elshitsa and Vlaykov Vruh Cu (Au) deposits of Central Srednogie, Bulgaria: constraints from U–Pb zircon and rutile geochronology and Hf–zircon and Sr whole-rock tracing. Final GEODE-ABCD 2003 workshop, Geodynamics and ore deposit evolution of the Alpine–Balkan–Carpathian–Dinaride Province, Seggau, Austria, abstr vol, 46
- Popov PN (1987) Tectonics of the Banat–Srednogie rift. *Tectonophysics* 143:209–216
- Popov K (2001) Geology of the southern part of the Panagyurishte ore region. *Ann Univ Min Geol* 43–44:51–63
- Popov P (2002) Alpine geotectonic evolution and metallogeny of the Eastern part of the Balkan peninsula. *Ann Univ Min Geol* 45:33–38
- Popov P, Kovachev V (1996) Geology, composition and genesis of the ore mineralization in the central part of the Elatsite–Chelopech ore field. Proceeding of the annual Meeting Sofia, UNESCO-ICGP Project 356, Plate tectonic aspects of the Alpine Metallogeny in the Carpatho–Balkan region, 1:159–170
- Popov P, Popov K (1997) Metallogeny of the Panagyurishte ore region. In: Romic K, Konzulovic R (eds) Symposium “Ore deposits Exploration” Proceedings Belgrade 327–338
- Popov P, Popov K (2000) General geologic and metallogenic features of the Panagyurishte ore region. In: Strashimirov S, Popov P (eds) Geology and Metallogeny of the Panagyurishte ore region (Srednogie zone, Bulgaria), ABCD-GEODE 2000 workshop, guide book to excursions, Sofia, 1–7
- Popov PN, Bairaktarov I, Marinov T (1979) Magmatism and structure of the eastern part of the Burgas ore region. I: characterization of the magmatic formations. *Geol Balk* 9:109–124
- Popov P, Petrunov R, Strashimirov S, Kanazirski M (2000a) Elatsite – Chelopech ore field. In: Strashimirov S, Popov P (eds) Geology and Metallogeny of the Panagyurishte ore region (Srednogie zone, Bulgaria), ABCD-GEODE 2000 workshop, guide book to excursions, Sofia, 8–18
- Popov P, Tsonev D, Kanazirski M (2000b) Elshitsa ore field. In: Strashimirov S, Popov P (eds) Geology and metallogeny of the panagyurishte ore region (Srednogie zone, Bulgaria), ABCD-GEODE 2000 workshop, guide book to excursions, Sofia, pp 40–50
- Popov P, Radichev R, Dimovski S (2001) Geology and evolution of the Elatsite–Chelopech porphyry–copper–massive sulfide ore field. *Ann Univ Min Geol* 43–44:31–44 (in Bulgarian)
- Popov P, Strashimirov S, Popov K, Petrunov R, Kanazirski M, Tzonev D (2003) Main features in geology and metallogeny of the Panagyurishte ore region. 50 Years *Ann Univ Min Geol* 46:119–125
- Radulescu D, Sandulescu M (1973) The plate tectonics concept and geological structure of the Carpathians. *Tectonophysics* 16:155–161
- Ricou L-E, Burg J-P, Godfriaux I, Ivanov Z (1998) Rhodope and Vardar: the metamorphic and the olistostromic paired belts related to the Cretaceous subduction under Europe. *Geodynamica Acta* 11(6):285–309
- Shalamuk, IB, Zbia M, Genini A, Fernandez RR (1997) Jurassic epithermal Au–Ag deposits of Patagonia, Argentina. *Ore Geol Rev* 12:173–186
- Sillitoe RH (1997) Characteristics and controls of the largest porphyry copper–gold and epithermal gold deposits in the circum-Pacific region. *Aust J Earth Sci* 44:373–388
- Sillitoe RH (1999) Styles of high-sulfidation gold, silver and copper mineralisation in porphyry and epithermal environments. In: Weber G (ed) *Aust Inst Mineral Metallo, PacRim’99*, Bali, Indonesia, 10–13 October, Proceedings, pp 29–44

- Sillitoe RH, Hedenquist JW (2003) Linkages between volcanotectonic settings, ore-fluid composition, and epithermal precious metal deposits. In: Simmons SF, Graham I (eds) *Volcanic, geothermal, and ore-forming fluids: rulers and witnesses of processes within the earth*. Soc Econ Geol Spec Publ 10:315–343
- Stanisheva-Vassileva G (1980) The Upper Cretaceous magmatism in Srednogorie zone, Bulgaria: a classification attempt and some implications. *Geol Balc* 10:15–36
- Stoykov S, Pavlishina P (2003) New data for Turonian age of the sedimentary and volcanic succession in the southeastern part of Etropole Stara Planina Mountain, Bulgaria. *Comptes Rendus Bulg Acad Sciences* 56:55–60
- Stoykov S, Pavlishina P (2004) Stratigraphy and palynological assessments of the Upper Cretaceous sedimentary and volcanic formations in the region of Chelopech, Central Srednogorie zone, Bulgaria. *Bulg Geol Soc Annual Scientific Conference "Geology 2004"*, 16–17 December, Proceedings, pp 77–79
- Stoykov S, Yanev Y, Moritz R, Katona I (2002) Geological structure and petrology of the late cretaceous Chelopech volcano, Srednogorie magmatic zone. *Geohim Mineral Petrol* 39:27–38
- Strashimirov S, Popov P (2000) Geology and metallogeny of the Panagyurishte ore region (Srednogorie zone, Bulgaria). ABCD-GEODE 2000 workshop, guide book to excursions, Sofia, pp 50
- Strashimirov S, Petrunov R, Kanazirski M (2002) Porphyry-copper mineralisation in the central Srednogorie zone, Bulgaria. *Miner Depos* 37:587–598
- Sylvester AG (1988) Strike-slip faults. *Geol Soc Amer Bull* 100:1666–1703
- Tarkian M, Hünken U, Tokmakchieva M, Bogdanov K (2003) Precious-metal distribution and fluid inclusion petrography of the Elatsite porphyry copper deposit, Bulgaria. *Miner Depos* 38:261–281
- Tosdal RM (2004) Tectonics of porphyry copper and epithermal deposits as constrained by vein geometry. In: Cooke DR, Deyell C, Pongratz J (eds) *24 ct Au Workshop, CODES Special Publication 5*, University of Tasmania, Australia, pp 41–44
- Tosdal RM, Richards JP (2001) Magmatic and structural controls on the development of porphyry Cu ± Mo ± Au deposits. In: Richards JP, Tosdal RM (eds) *Structural controls on ore genesis*. *Rev Econ Geol* 14:157–181
- Tosdal RM, Richards JP (2002) Tectonic setting: a critical link in the formation of porphyry-Cu and epithermal deposits. In: Vearncombe S (ed) *Applied structural geology for mineral exploration and mining*. Abstr vol, 23–25 Sept. 2002, Kalgoorlie, Western Australia. *Bul Aust Instit of Geoscientists* 36:209–211
- Tsonev D, Popov K, Kanazirski M, Strashimirov S (2000) Radka ore field. In: Starshimirov S, Popov P (eds) *Geology of the Panagyurishte ore region (Srednogorie zone Bulgaria)*, ABCD-GEODE 2000 workshop, Guidebook to excursions, Sofia, pp 32–39
- Velichkova SH, Handler R, Neubauer F, Ivanov Z (2004) Variscan to Alpine tectonothermal evolution of the Central Srednogorie unit, Bulgaria: constraints from  $^{40}\text{Ar}/^{39}\text{Ar}$  analysis. *Schweiz Mineral Petrogr Mitt* 84:133–151
- Von Quadt A, Peytcheva I, Kamenov B, Fanger L, Heinrich C, Frank M (2002) The Elatsite porphyry copper deposit in the Panagyurishte ore district, Srednogorie zone, Bulgaria: U–Pb zircon geochronology and isotope-geochemical investigations of magmatism and ore genesis. In: Blundell DJ, Neubauer F, Von Quadt A (eds) *The timing and location of major ore deposits in an evolving orogen*, *Geol Soc London Spec Publ* 204:119–135
- Von Quadt A, Peytcheva I, Cvetkovic V (2003) Geochronology, geochemistry and isotope tracing of the Cretaceous magmatism of East Serbia and Panagyurishte district (Bulgaria) as part of the Apuseni–Timok–Srednogorie metallogenic belt in Eastern Europe. In: Elioupolos DG et al (eds) *Mineral exploration and sustainable development*, Proceedings 7th biennial SGA meeting, Athens, Greece, 24–28 August 2003, Millpress Rotterdam, pp 407–410
- Von Quadt A, Moritz R, Peytcheva I, Heinrich C (2005) Geochronology and geodynamics of Upper Cretaceous magmatism and Cu–Au mineralization in the Panagyurishte region of the Apuseni–Banat–Timok–Srednogorie belt (Bulgaria). *Ore Geol Rev* 27:95–126
- Wood DG, Porter RG, White NC (1990) Geological features of some Paleozoic epithermal gold occurrences in northeastern Queensland. *Aust J Geoch Expl* 36:413–433
- Woodcock NH, Fischer M (1986) Strike-slip duplexes. *J Struct Geol* 8:725–735