

Primary Hemiarthroplasty for Proximal Humeral Fractures in the Elderly: Long-Term Functional Outcome and Social Implications

Michael Dietrich¹, Christoph Meier², Daniela Zeller¹, Patrick Grueninger¹, Roger Berbig³, Andreas Platz¹

Abstract

Background: Primary shoulder hemiarthroplasty is an established treatment modality for complex fractures of the proximal humerus. Long-term functional outcome is often disappointing. However, little is known about social implications particularly in the elderly.

Methods: A single-institution case series of consecutive geriatric patients (age > 70 years) treated with shoulder hemiarthroplasty for complex fractures of the proximal humerus between 1994 and 1997 was analysed. Postoperative morbidity, long-term function, radiological outcome and social implications were evaluated.

Results: Seventy-seven patients fulfilled the study criteria. Median age at the time of operation was 80 years (range 70–93 years). Systemic and local postoperative complications were observed in 8% including 2 patients (3%) with revision surgery. Postoperative mortality was 1%. Forty-eight patients (62%) were available for follow-up (median 49 months, range 25–80 months), 22 (29%) died from causes unrelated to hemiarthroplasty before follow-up and 7 patients (9%) did not attend follow-up examination. Median Constant-Murley score was 41 points (range 17–77 points). Long-term results concerning pain were satisfying. The Oxford shoulder score ranged from 14 to 40 (median 30). Forty-one patients (85%) still lived in their original environment and managed their daily life independently despite poor shoulder function. Four patients (8%) lived in a retirement home and 3 (6%) in a nursery home. Eighty percent of our patients

were still able to use public transportation, do the daily shopping and wash their whole body by themselves.

Conclusion: Most patients managed their daily life independently despite poor shoulder function.

Key Words

Hemiarthroplasty · Elderly · Shoulder · Social implications · Outcome

Eur J Trauma Emerg Surg 2007;33:512–9

DOI 10.1007/s00068-007-6134-5

Introduction

The complex fracture of the proximal humerus as described by Neer is a severely disabling injury [1]. It is a typical injury of the elderly which becomes more common as the population ages and remains physiologically active. The overall incidence of these fractures is 70/100,000 people a year and raises up to 405/100,000 yearly including only the female older than 70 years of age [2].

The severity of this injury is due to the associated vascular compromise. Disruption of the major blood supply leaves the proximal humerus susceptible to avascular necrosis [3]. In addition to vascular compromise, these fractures are often associated with significant comminution, which makes stable open reduction and internal fixation extremely difficult, particularly in old patients with osteoporotic bone.

¹Department of Surgery, Stadtspital Triemli, Zurich, Switzerland,

²Division of Trauma Surgery, University Hospital Zurich, Zurich, Switzerland,

³SportClinic Zurich, Zurich, Switzerland.

Received: June 30, 2006; revision accepted: October 13, 2006;

Published Online: May 19, 2007

Thus, the management of these fractures is still a challenging problem [4] and hemiarthroplasty is a widely applied technique for these fracture patterns. The reported functional results following primary hemiarthroplasty for fractures of the proximal humerus vary, whereas satisfactory pain relief can usually be achieved [1, 5–8].

Several groups reported patients with relatively pain free shoulders and good functional results after acute shoulder hemiarthroplasty for three- and four-part fractures of the proximal humerus [1, 5, 8]. However, none of these studies dealt with patients having an average age of over 80 years. Other reports showed disappointing functional results, especially when the procedure was performed in elderly patients [9, 10]. Furthermore, little is known about social implications of impaired shoulder function particularly in the elderly. It was our interest to study postoperative morbidity, long-term glenohumeral function and the social implication of primary hemiarthroplasty for displaced fractures of the proximal humerus in geriatric patients.

Patients and Methods

Inclusion Criteria

The trauma registry was reviewed for fractures of the proximal humerus treated with primary hemiarthroplasty between January 1994 and December 1997. Only patients at an age of 70 years and older were included in our study. Younger patients, preceding osteosynthesis for primary treatment and pathological fractures were all excluded from further evaluation.

Operative Technique and Rehabilitation

The operative technique was standardized and followed the guidelines by the manufacturer of the prosthesis model used exclusively in this study (Neer II design: Howmedica® Shoulder prosthesis, Shannon Industrial Estate, Co. Clare, Ireland). All patients were operated in beach-chair position under general anesthesia. A deltopectoral approach was performed in each case. The head fragment was removed. The tuberosities were identified, mobilised and stay sutures were placed at the bone–tendon junction. The canal was prepared by hand and drill holes placed in the shaft on either side of the biceps groove. Non-resorbable sutures (3-Ethicon®-Mersilene, Polyester, Johnson & Johnson®, UK) were pulled through the holes before cementing. A trial component was inserted to determine proper size of head and shaft and the correct height of the prosthesis. The humeral component was

cemented in all cases in 25° of retro-version. The tuberosities were then reconstructed using heavy, non-absorbable 3-Mersilene sutures. The goal was to attach the tuberosity fragments to the humeral shaft, to each other, and to the fin of the prosthesis. Cancellous bone graft, taken from the removed humeral head, was placed between the tuberosities and the diaphysis to facilitate bone union.

Rehabilitation was started at the first postoperative day with exercises for wrist and elbow. Pendulum exercises of the shoulder were initiated 1 week after the intervention and followed the principles of early passive motion emphasized by Neer [1]. Free range of motion with elevation of more than 90° and strengthening was started 6 weeks after surgery.

Outcome Measures and Follow-Up

Clinical radiological records were reviewed for associated injuries, intra- and postoperative complications and duration of stay in hospital. Co-morbidities of these elderly patients were classified analogue the Charlson Index [11]. The initial preoperative radiographs were reviewed by one of the authors (Christoph Meier) and classified by the Neer and AO systems [12, 13].

All patients eligible for this study were contacted and called up for follow-up. At follow-up, all patients were examined by a single investigator (Michael Dietrich). Both investigators (Christoph Meier and Michael Dietrich) did not participate in any of the operations.

Assessment of the long-term result included the Constant-Murley shoulder score [14] of the operated and contra lateral side, the Oxford shoulder score [15], and radiographs of the shoulder.

Pain and shoulder function were measured using the well-established Constant-Murley shoulder score. The Oxford shoulder score was completed by the patients unaided at the time of follow-up. This questionnaire contains 12 items, each of which has five categories of response. Each item is scored from 1 to 5, from least to most difficulty or severity, and combined to produce a single score with a range from 12 (least difficulties) to 60 (most difficulties). It has been shown to be consistent and reproducible regarding the patients' perception of shoulder problems [15, 16]. The test specifically asks for important functions of daily activities (e.g., ability to use knife and fork, to hang up clothes, to use public transportation, etc.).

Radiographic outcome was assessed on a true anteroposterior view of the shoulder in neutral rotation

and a lateral Y-view. Radiographs were evaluated with specific attention to bone union and adequacy of tuberosity reduction and stem fixation. Loosening of the prosthesis was defined as radio lucent lines greater than 2 mm or progressive lines at the prosthetic bone–cement interface [17]. Proximal migration of the prosthesis was evaluated by measurement of the distance between the summit of the prosthetic head and the line of sclerosis of the acromion on the anteroposterior radiograph in neutral rotation [18]. A correlation between a rotator cuff tear and narrowing of the subacromial space to less than 7 mm, indicating proximal migration of the humerus, is well established in the literature [19–21].

Statistical Analysis

Results were analysed using the statistical program SPSS® version 11.0 (SPSS, Chicago, IL, USA). Data are presented as median and range.

For continuous data compared between shoulders, statistical analysis was performed using Wilcoxon signed ranks test. Independent groups were compared using Mann–Whitney test. To analyse any correlation between continuous data, Spearman's rank correlation was performed. Spearman rank correlation coefficient was put in brackets. The level of significance was set at $p < 0.05$.

Results

A total of 77 patients (10 male, 67 female) qualified to enter the study. Median age at the time of surgery was 80 years (range, 70–93 years). Sixty-one patients (79%) had at least one pre-existing co-morbidity, 8 (10%) patients had three or more (Table 1).

All patients sustained a low energy trauma. Most patients fell over obstacles at home ($n = 34$, 44%) or by crossing a street ($n = 25$, 32%). Five patients (6%) fell due to a cerebral absence, 6 (8%) fell under the influence of alcohol and 3 (4%) times icy ground was

claimed to be the reason for the incident. Four patients (5%) broke their arm when getting attacked by a criminal.

The dominant shoulder was affected in 62%. Associated fractures were diagnosed in 15 cases (20%) including 4 contra-lateral upper limb fractures (2 radius, 1 humerus, 1 olecranon), 8 lower limb fractures (5 proximal femur fractures, 2 pubic ramus fractures, 1 tibial fracture), 1 fracture of a facial bone as well as 2 patients with fractured ribs.

In 65 patients (84%) preoperative radiographs were available and suitable for fracture classification. According to the Neer classification, 11 patients (17%) were treated for a two-part, 26 (40%) for a three-part and 28 (43%) for a four-part fracture. Among these fractures were 8 (12%) fracture-dislocations. Using the AO-classification to categorize these fractures, 7 (11%) A, 30 (46%) B and 28 (43%) C-fractures were observed.

All operations were performed by one of three councillors. They all show a record of more than 10 years of operative experience in shoulder surgery.

Median duration of hospital stay was 17 days (range, 6–98 days). No intraoperative complications occurred. Postoperative morbidity was 8% ($n = 6$), including 3 (4%) systemic complications (1 cardiac stroke, 1 apoplexy, 1 cardio-pulmonary failure). Local complications ($n = 3$, 4%) included 2 patients (3%) who had to undergo revision surgery (1 postoperative hematoma, 1 displaced prosthesis) and 1 patient (1%) with postoperative thrombosis of the subclavian vein. One patient suffering from a known aortic valve stenosis (1%) died 7 days after trauma (6 days following shoulder- and hip arthroplasty) as a result of cardio-pulmonary failure.

Follow-Up

Median follow-up was 49 months (range, 25–80 months). Twenty-two patients (29%) of the original cohort had already died from causes unrelated to hemiarthroplasty before follow-up. Five patients (6%) were unable to participate at the examination due to health problems unrelated to previous shoulder surgery and 2 patients (3%) could not be located. Thus, physical examination and questioning was performed in a total of 48 patients (87% of all patients alive, 62% of the initial cohort). At follow-up, median age was 83 years (range, 72–97 years).

Constant-Murley Shoulder Score

Overall median Constant-Murley shoulder score score was 41 points (range, 17–77 points) on the fractured side and 77 points (range, 33–100 points) on the con-

Table 1. Pre-existing relevant co-morbidities (analogue Charlson index [11]).

Co-morbidity	n	%
Cardiovascular	53	68
COPD	16	21
Diabetes	21	27
Nephropathy	3	4
Carcinoma	2	3
Cerebral	9	12




Figure 1. Oxford shoulder score. Detailed results of two different functions regarding an activity positioned in the plane in front of the body and an activity which requires elevation and abduction.

tra-lateral side ($p < 0.01$, correlation coefficient 0.47). Median anterior elevation of the affected shoulder was 70° (range, 25° – 115°) and lateral abduction was measured 70° as well (range, 25° – 120°). Only 5 patients (10%) were able to lift their arm more than 90° in anterior elevation or lateral abduction. Power was also markedly reduced. Median isometric power of 6.6 pounds (range, 0–18 pounds) compared with 12 pounds (range, 2–25 pounds) on the contra-lateral side was observed ($p < 0.01$, correlation coefficient 0.72). Median pain score was 10 points, no patient complained about severe shoulder pain. Sixty-five percent of the patients had no or mild pain (Table 2). Three patients (6%) reported to require analgesia (NSAR only) on a daily basis, 6 (13%) took them occasionally and 39 patients (81%) did not require any analgesia at all. Adapted for age and sex, median overall Constant-Murley shoulder score was 62% (range, 24–100%). In median, patients reached 52% of the unaffected shoulder function regarding Constant-Murley shoulder score. Our results clearly showed that

Table 2. Severity of pain (Constant-Murley shoulder score).

Pain	n	%
Severe (0 points)	0	0
Moderate (5 points)	17	35
Mild (10 points)	19	40
None (15 points)	12	5




Figure 2. Example of follow-up radiographs (a: anteroposterior view, b: lateral Y-view) demonstrating marked narrowing of subacromial space and absorbed tuberosities.

patients with the best functional results had the least pain and vice versa ($p < 0.01$).

No significant correlation between fracture classification and outcome regarding function or pain was seen.

Oxford Shoulder Score

Median Oxford shoulder score was 30 (range, 14–40, Figure 1). Despite reduced range of motion of the affected shoulder, there was no limitation in function of the hand. All daily duties positioned in the plane in front of their body could be managed properly (e.g., working with fork and knife). In contrast, the most common limitations in case of fair or poor functional results were hair combing or hanging up clothes (Table 3).

Social Implications

Forty-one patients (85%) were still able to live by themselves in their original environment, some of them supported by occasional aid to pursue their daily activities. Four patients (8%) lived in a retirement home, 3 (6%) in a nursery home. Eighty percent of the patients were able to use public transportation, do the daily shopping, eat with utensil as well as they were able to wash their whole body by themselves.

Radiographic Results

Forty patients (52%) of those 48 who were physically examined were available also for radiological follow-up. Eight patients had to be visited outside of the

Table 3. Oxford shoulder score (12-item shoulder questionnaire).

Scoring category (compromise)	Normal to moderate 1–3 points		Severe or impossible 4–5 points	
	n	%	n	%
Worst pain from shoulder	44	92	4	8
Doing household, shopping alone	40	83	8	17
Trouble with dressing	47	98	1	2
Carrying a tray of food	41	85	7	15
Using a knife and fork	46	96	2	4
Combing hair	24	50	24	50
Trouble with transport	44	92	4	8
Usual level of shoulder pain	43	90	5	10
Hanging clothes in wardrobe	25	52	23	48
Work interference due to pain	44	92	4	8
Washing under both arms	48	100	0	0
Pain in bed at night	40	83	8	17

hospital and therefore did not get any X-ray. In 7 individuals (18%) ectopic bone formation was seen. No signs for implant loosening were evident on follow-up radiographs. Union of both the tuberosities and parts of them in correct position was observed in 21 patients (52%). Nineteen patients (48%) presented with at least one absorbed, ununited or totally malpositioned tuberosity (Figure 2). In 7 cases (18%), no tuberosities could be identified on the radiographs indicating complete resorption.

Comparing the group with parts of both tuberosities identified in a correct position ($n = 21$, 52%) with the group with at least one tuberosity completely resorbed ($n = 19$, 48%), a significant statistical difference was found regarding overall median Constant score [46 points (range, 28–77 points) vs. 31 points (range, 17–58 points), $p < 0.01$]. They also claimed less pain [median Constant-Murley shoulder score for pain, 10 points (range, 5–15 points) vs. 8 points (range, 5–15 points), $p < 0.01$]. Better isometric power was as well observed in those patients [median Constant-Murley shoulder score for power, 8 pounds (range, 2–18 pounds) vs. 5 pounds (range, 0–10 pounds) $p < 0.01$]. The sole radiological absence of the greater tuberosity was clearly associated with impaired shoulder function [Constant-Murley score 45 (range, 21–77) vs. 30.5 (range, 17–55), $p = 0.01$].

Proximal migration of the prosthesis as demonstrated by an acromiohumeral distance less than 7 mm on the anteroposterior radiograph was present in 22 patients (55%). These patients showed less isometric power [median Constant score for power, 5.5 pounds (range, 2–14 pounds) vs. 7 pounds (range,

0–18 pounds), $p = 0.03$]. They had a smaller range of motion [median Constant-Murley shoulder score for function, 12 points (range, 4–24 points) vs. 8 points (range, 2–28 points), $p = 0.03$], but there was no difference in pain perception in those patients [median Constant-Murley shoulder score for pain, 10 points (range, 5–15 points) vs. 10 points (range, 5–15 points)].

All 7 patients (18%) with both tuberosities absorbed on the radiographs showed a narrowing of the subacromial space to less than 5 mm. In radiographies that showed an absorbed greater tuberosity ($n = 12$, 30%) subacromial space was significantly narrower [median, 3.5 mm (range, 0–8 mm) vs. median, 7.5 mm (range, 2–18 mm), $p = 0.02$].

Discussion

In geriatric patients requirements for successful surgical treatment of proximal humeral fractures include “one-time surgery”, low postoperative morbidity even in high-risk patients, acceptable function and good pain relief. In many individuals, pain can be a greater problem than the restriction in range of motion, especially in elderly patients with relatively reduced demands for physical performance [9]. Primary shoulder hemiarthroplasty fulfilled all these requirements in our series. It was safe and associated with low postoperative morbidity in respect to age and co-morbidities of our geriatric patients. By replacing the humeral head, common problems of head-preserving therapies such as ischemic necrosis of the humeral head, implant loosening or secondary fracture displacement are avoided [22]. Furthermore, secondary implantation, as salvage procedure following failed osteosynthesis or non-operative treatment may lead to increased postoperative morbidity compared with primary hemiarthroplasty [5, 23, 24].

The late finding of superior migration of the prosthesis and its adverse effect on shoulder function has been described previously. Accurate tuberosity reconstruction was found to have a great impact on functional outcome [17, 18, 25, 26]. Boileau et al. [18] reported final tuberosity malposition in 50% of their patients, resulting in superior migration of the prosthesis, stiffness, weakness and persistent pain. Among other factors, age and sex (women over 75 years of age) significantly correlated with failure of tuberosity osteosynthesis leading to poor outcomes. In the present investigation, mainly patients incorporating these “risk factors” for poor outcome (median age 80 years, 87% female) were studied. Analysis of our results showed that cranial migration of the prosthesis was

Table 4. Comparison of the current study with other investigations regarding functional outcome of hemiarthroplasty for fractures of the proximal humerus. (FU: Follow-up; CS: Constant-Murley shoulder score; NA: not available).

Author	Year	Prosthesis	n	Age	FU (mts)	CS	Elevation (°)
Tanner and Cofield [5]	1982	Neer-Type	49	69	38	NA	NA
Hawkins and Switlyk [28]	1991	Neer-II, 3M [®] , St Paul	29	64	48	NA	72
Moeckel et al. [6]	1991	Modular component	22	70	36	NA	119
Wretenberg and Eklund [9]	1997	Neer-3M [®] , Rotherham	18	82	42	NA	55
Hoellen et al. [22]	1997	Global shoulder-Prot [®]	30	74	12	48	NA
Dimakopoulos et al. [29]	1997	Biomodular, 3M [®]	38	56	37	NA	130
Ballmer and Hertel [30]	1998	NA	34	70	42	NA	101
Zyto et al. [27]	1998	Neer-II, 3M [®]	27	71	39	46	70
Boss and Hintermann [31]	1999	Neer-II modular	20	77	32	52	99
Prakash et al. [32]	2002	Neer-II, 3M [®] , DePuy [®]	22	69	33	NA	93
Boileau et al. [18]	2002	Aequalis [®]	66	66	27	56	101
Becker et al. [33]	2002	DePuy [®]	27	67	48	45	89
Schittko et al. [34]	2003	OrTra [®] -Prothese	43	NA	NA	54	54
Robinson et al. [26]	2003	3M [®] , Osteonics [®]	138	68	76	64	NA
Kralinger et al. [25]	2004	DePuy [®] and others	167	70	29	55	NA
Schmal et al. [35]	2004	EPOCA [®] C.O.S.	20	70	14	52	NA
Current study	2006	Neer-II, Howmedica [®]	77	83	49	41	70

associated with poor function and less isometric power. However, we did not find an association of prosthesis migration with pain. In contrast, tuberosity complications, such as resorption, malreduction or migration, significantly correlated with worse range of motion, reduced power and more pain. Furthermore, no correlation between fracture classification and outcome regarding function or pain was observed in our series. This is in accordance with findings of other investigations [6, 27] (Table 4).

Functional results after shoulder hemiarthroplasty for fractures of the proximal humerus have generally been disappointing with a marked impairment of the active range of motion in most investigations found in the literature (Table 3). However, these studies are difficult to compare as they differ in patient demographics and outcome measures. As a matter of fact, a statistical correlation between outcome and age was recently demonstrated [6, 18, 26]. With a median age of 80 years at surgery and 83 years at follow-up, the present investigation contains one of the oldest patient cohorts with one of the longest follow-up time periods.

The well-established Constant-Murley shoulder score contains both subjective and objective elements [14]. Objective measurements of the active range of motion and power are allocated 65 points together, out of a total of 100. Subjective assessment of pain and activities of daily living are allocated 35 points. In the elderly, the range of motion and power may be of inferior relevance compared with the perception of

pain and the ability to manage the daily life independently. Whereas most reports found in the literature investigated shoulder function after hemiarthroplasty [22, 23, 25–27, 29, 30, 33], social implications following this procedure are poorly studied. A search of the literature revealed only one study, which addressed these issues in a similar patient group and follow-up compared with our investigation [9]. However, its small number of patients limited this study and social implications of the impaired shoulder function were mentioned but not addressed in detail. In the herein presented investigation the clinical evaluation was focused on the patients' abilities to master their daily life. The Oxford shoulder score proved to be a very suitable tool for assessing the patient's perception of shoulder problems and their impact on daily routine [15, 16]. It contains only 12 simple questions regarding pain and common activities, each coming with 5 possible answers to choose from. In an evaluation of 155 different musculoskeletal outcomes measures and instruments, the Oxford shoulder score was validated against the Constant-Murley score, the SF-36 and the Health Assessment Questionnaire [36]. Its overall quality as an outcome instrument (methodological evaluation and clinical utility) scored 9 points out of 10. In our investigation, the Oxford shoulder score proved to be easy to complete unaided even by elderly patients.

Our results demonstrate, that even with marked impairment of shoulder function, most geriatric patients could preserve fundamental activities of daily

living, which enabled them to pursue a self-contained life in their original environment. Furthermore, hemiarthroplasty provided satisfactory pain relief with the majority of the patients having none to moderate shoulder pain and only 17% suffering from shoulder pain in bed at night at long-term follow-up.

With modern techniques and implants, such as locked compression plates or nail systems which provide angular stability, osteosynthesis instead of hemiarthroplasty, has gained increasing popularity, even for pluri-fragmental fractures in osteoporotic bone. Whether this recent development changes the indication for shoulder hemiarthroplasty in geriatric patients has not been conclusively answered in the literature, yet. Further studies, comparing not only functional results, but long-term pain relief, social implications and costs are required to evaluate the roles and indications for shoulder hemiarthroplasty and head-preserving techniques in modern fracture treatment.

Acknowledgment

The authors would like to thank Professor B. Seifert, Department of Biostatistics, ISPM, University of Zurich, for providing assistance with all the statistical issues of this manuscript.

References

- Neer CS. Displaced proximal humeral fractures. Part II. Treatment of three-part, four-part displacement. *J Bone Joint Surg (Am)* 1970;52-A:1090-103.
- Lind T, Kroner K, Jensen J. The epidemiology of fractures of the proximal humerus. *Arch Orthop Trauma Surg* 1989;108:285-7.
- Neumann K, Muhr G, Breiffuss H. Primary humerus head replacement in dislocated proximal humeral fracture. Indications, technique, results. *Orthopade* 1992;21:140-7.
- Mills HJ, Horne G. Fractures of the proximal humerus in adults. *J Trauma* 1985;25:801-5.
- Tanner MW, Cofield RH. Prosthetic arthroplasty for fractures and fracture-dislocations of the proximal humerus. *Clin Orthop* 1983;179:116-28.
- Moeckel BH, Sculco TP, Alexiades MM, Dossick PH, Inglis AE, Ranawat CS. Modular hemiarthroplasty for fractures of the proximal part of the humerus. *J Bone Joint Surg (Am)* 1992;74-A:884-9.
- Goldman RT, Koval KJ, Cuomo F, Gallagher MA, Zuckerman JD. Functional outcome after humeral head replacement for acute three- and four-part proximal humeral fractures. *J Shoulder Elbow Surg* 1995;4:81-6.
- Compito CA, Self EB, Bigliani LU. Arthroplasty and acute shoulder trauma. Reasons for success and failure. *Clin Orthop Relat Res* 1994;307:27-36.
- Wretenberg P, Eklund A. Acute hemiarthroplasty after proximal humerus fracture in old patients. A retrospective evaluation of 18 patients followed for 2-7 years. *Acta Orthop Scand* 1997;68:121-3.
- Movin T, Sjoden GO, Ahrengart L. Poor function after shoulder replacement in fracture patients. A retrospective evaluation of 29 patients followed for 2-12 years. *Acta Orthop Scand* 1998;69:392-6.
- Charlson ME, Pompei P, Ales KL, Mac Kenzie CR. A new method of classification prognostic comorbidity in longitudinal studies: development and validation. *J Chron Dis* 1987;40:373-83.
- Neer CS. Displaced proximal humeral fractures: part I classification and evaluation. *J Bone Joint Surg (Am)* 1970;52:1077-89.
- Jakob RP, Kristiansen T, Mayo K, Ganz R, Muller ME. Classification and Aspects of Fractures of the Proximal Humeri. In: Bateman JE, Welsh RP (eds). *Surgery of the Shoulder*. St Louis, Decker 1984, pp 330-43.
- Constant CR, Murley AH. A clinical method of functional assessment of the shoulder. *Clin Orthop Relat Res* 1987;214:160-4.
- Dawson J, Fitzpatrick R, Carr A. Questionnaire on the perceptions of patients about shoulder surgery. *J Bone Joint Surg (Br)* 1996;78-B:593-600.
- Dawson J, Hill G, Fitzpatrick R, Carr A. The benefits of using patient-based methods of assessment. *J Bone Joint Surg Surg (Br)* 2001;83-B:877-82.
- Mighell MA, Kolm GP, Collinge CA, Frankle MA. Outcomes of hemiarthroplasty for fractures of the proximal humerus. *J Shoulder Elbow Surg* 2003;12:569-77.
- Boileau P, Krishnan SG, Tinsi L, Walch G, Coste JS, Mole D. Tuberosity malposition and migration: reasons for poor outcomes after hemiarthroplasty for displaced fractures of the proximal humerus. *J Shoulder Elbow Surg* 2002;11:401-12.
- Cotton RE, Rideout DF. Tears of the humeral rotator cuff. A radiological and pathological necropsy survey. *J Bone Joint Surg* 1964;46:314-28.
- Golding FC. The shoulder - the forgotten joint. *Br J Radiol* 1962;35:149-58.
- Azzoni R, Cabitza P. Sonographic versus radiographic measurement of the subacromial space width. *Chir Organi Mov* 2004;89:143-50.
- Hoellen IP, Bauer G, Holbein O. Prosthetic humeral head replacement in dislocated humerus multi-fragment fracture in the elderly - an alternative to minimal osteosynthesis? *Zentralb Chir* 1997;122:994-1001.
- Boileau P, Trojani C, Walch G, Krishnan SG, Romeo A, Sinnerton R. Shoulder arthroplasty for the treatment of the sequelae of fractures of the proximal humerus. *J Shoulder Elbow Surg* 2001;10:299-308.
- Antuna SA, Sperling JW, Sanchez-Sotelo J, Cofield RH. Shoulder hemiarthroplasty for proximal humeral malunions: long-term results. *J Shoulder Elbow Surg* 2002;11:122-9.
- Kralinger F, Schwaiger R, Wambacher M, Farrell E, Menth-Chiari E, Hubner C, Resch H. Outcome after primary hemiarthroplasty for fracture of the head of the humerus. A retrospective multicentre study of 167 patients. *Bone Joint Surg* 2004;86-B:217-9.
- Robinson CM, Page RS, Hill RMF, Sanders LD, Court-Brown CM, Wakefield AE. Primary hemiarthroplasty for treatment of proximal humeral fractures. *J Bone Joint Surg* 2003;85-A:1215-23.
- Zyto K, Wallace WA, Frostick SP, Preston BJ. Outcome after hemiarthroplasty for three- and four-part fractures of the proximal humerus. *J Shoulder Elbow Surg* 1998;7:85-9.
- Hawkins RJ, Switlyk P. Acute prosthetic replacement for severe fractures of the proximal humerus. *Clin Orthop Relat Res* 1991;289:156-60.

29. Dimakopoulos P, Potamitis N, Lambiris E. Hemiarthroplasty in the treatment of comminuted intraarticular fractures of the proximal humerus. *Clin Orthop Relat Res* 1997;341:7–11.
30. Ballmer FT, Hertel R. Indications and results of shoulder prosthetics in complex proximal humerus fractures. *Ther Umsch* 1998;55:197–202.
31. Boss AP, Hintermann B. Primary endoprosthesis in comminuted humeral head fractures in patients over 60 years of age. *Int Orthop* 1999;23:172–4.
32. Prakash U, McGurty DW, Dent JA. Hemiarthroplasty for severe fractures of the proximal humerus. *J Shoulder Elbow Surg* 2002;11:428–30.
33. Becker R, Pap G, Machner A, Neumann WH. Strength and motion after hemiarthroplasty in displaced four-fragment fracture of the proximal humerus: 27 patients followed for 1–6 years. *Acta Orthop Scand* 2002;73:44–9.
34. Schittko A, Braun W, Ruter A. Experiences with the OrTra-prosthesis in primary prosthetic replacement of fractures of the humeral head – indication, technique and results. *Zentralbl Chir* 2003;128:12–6.
35. Schmal H, Klemt C, Sudkamp NP. Evaluation of shoulder arthroplasty in treatment of four-fragment fractures of the proximal humerus. *Unfallchirurg* 2004;107:575–82.
36. Suk M, Hanson BP, Norvell DC, Helfet DL. Shoulder, Oxford Shoulder Score. In: Suk M, Hanson BP, Norvell DC, Helfet DL (eds). *AO handbook. Musculoskeletal outcomes measures and instruments*. Davos, AO Publishing 2005, pp 76–9.

Address for Correspondence

Michael Dietrich, MD
Department of Surgery
Stadtspital Triemli
Birmensdorferstrasse 497
8063 Zurich
Switzerland
Phone (+41/44) 466-1079, Fax -2601
e-mail: michael.dietrich@triemli.stzh.ch