Eur. Phys. J. A **28**, s02, 1 (2006)
DOI: 10.1140/epja/i2006-10-116-7

EPJ A direct
electronic only

Erratum

Beta decay of ^{57}Zn

A. Jokinen[1,2,a], A. Nieminen[1], J. Äystö[1,2], R. Borcea[3], E. Caurier[4], P. Dendooven[1], M. Gierlik[5], M. Górska[3], H. Grawe[3], M. Hellström[6], M. Karny[5], Z. Janas[5], R. Kirchner[3], M. La Commara[3], G. Martinez-Pinedo[7], P. Mayet[3], H. Penttilä[1], A. Plochocki[5], M. Rejmund[3], E. Roeckl[3], M. Sawicka[5], C. Schlegel[3], K. Schmidt[3], and R. Schwengner[8]

[1] University of Jyväskylä, Department of Physics, FIN-40351 Jyväskylä, Finland
[2] CERN, CH-1211 Geneva 23, Switzerland
[3] Gesellschaft fr Schwerionenforschung mbH, D-64291, Darmstadt, Germany
[4] Institut de Recherches Subatomiques (IN2P3-CNRS-Université Louis Pasteur), F-67037 Strasbourg Cedex 2, France
[5] Warsaw University, Institute of Experimental Physics, PL-00681 Warsaw, Poland
[6] Lund University, Division of Cosmic and Atomic and Subatomic Physics, S-22100 Lund, Sweden
[7] Department of Physics and Astronomy, University of Basel, CH-4056 Basel, Switzerland
[8] Institut für Kern- und Hadronenphysik, Forschungszentrum Rossendorf, D-01314 Dresden, Germany

Original article: EPJdirect A3, 1–11 (2002) DOI 10.1007/s1010502a0003

Received: 7 August 2006 /
Published online: 11 August 2006 –

We publish herewith the correct table 3 which was wrongly published in the original paper as a repetition of table 2.

Table 3. Adopted states in ^{57}Cu and experimental feeding of the states, extracted $\log ft$ and $B_{\rm GT}$ values of the ^{57}Zn decay in comparison with level-energy and $B_{\rm GT}$ predictions obtained from shell-model calculations. $B_{\rm GT}$ values predicted for higher ^{57}Cu energies are omitted.

E^*	I^π	Intensity	$\log ft$	$B_{\rm GT}$	E^* (th.)	$B_{\rm GT}$ (th.)
g.s.	$3/2-$					
1028(4)	$5/2-$	$< 7^{(*)}$	$> 5^{(*)}$	$< 0.06^{(*)}$	552	0.0091
1106(4)	$1/2-$					
2395(9)	$5/2-$	1.2(6)	5.5(3)	0.017^{+14}_{-8}	4126	0.0004
2525(17)	$7/2-$	10(2)	4.60(16)	0.15^{+7}_{-5}	3624	0.1537
3236(22)	$7/2-$	19(4)	4.19(15)	0.39^{+16}_{-11}	4320	0.1614
3510(25)	$(9/2+)$					
3786(25)	$(5/2,7/2,9/2)-$	7(2)	4.50(17)	0.19^{+9}_{-6}		
4208(27)	$(5/2,7/2,9/2)-$	3(1)	4.8(2)	0.10^{+6}_{-4}		
4378(27)	$(5/2,7/2,9/2)-$	1.8(8)	5.0(2)	0.06^{+5}_{-3}		
4563(20)	$(5/2,7/2,9/2)-$	5.5(15)	4.44(18)	0.22^{+11}_{-7}		
5168(32)	$(5/2,7/2,9/2)-$	2.1(8)	4.7(2)	0.12^{+8}_{-5}		
5314(28)	$7/2-$, IAS	52(6)	3.28(14)	$0.13^{(**)}$	5724	0.13

$(^*)$ See discussion in Chapter 3.; $(^{**})$ Shell-model value.

[a] *Present address*: CERN/PS-division, CH-1211 Geneva 23, Switzerland; e-mail: `ari.jokinen@cern.ch`