

Tissue distribution, antigen specificity and effector functions of $\gamma\delta$ T cells in human diseases

Gennaro De Libero

Experimental Immunology, University Hospital, Basel, Switzerland

Introduction

Since the discovery of T cells bearing the $\gamma\delta$ TCR [1], a large number of studies have described the molecular characteristics of this receptor and examined the possible functions of these cells in a wide range of vertebrate species. However, although we know the genetics of γ and δ genes in great detail, the function of $\gamma\delta$ T cells in vivo is only in part defined. In man, $\gamma\delta$ T cells constitute ~5% of circulating CD3⁺ cells and are most often CD4⁻ or CD4⁺ [2, 3]. Although $\gamma\delta$ T cells represent a small population when compared to $\alpha\beta$ T cells, they show peculiar functional characteristics that make them unique and important. Rather than discussing studies on the $\gamma\delta$ T cells in general, this chapter will focus on the physiology of human $\gamma\delta$ T lymphocytes and their possible role in diseases.

Gene organization of γ and δ loci

In the human genome the TCR γ locus spans 160 kb of DNA, and maps to chromosome 7. The γ locus comprises two constant gene segments and five joining elements, J1, JP, and JP1 located upstream of C γ 1 and JP2 J2, which are upstream of C γ 2. Fourteen variable γ (V γ) genes have been identified, including eight pseudogenes and six active genes (reviewed in [4]). These genes fall into four subgroups designated V γ I–V γ IV. Five functional genes belong to the V γ I family (V γ 2,3,4,5,8), whereas family II consists of only one gene, designated V γ 9 (V γ 2 in another nomenclature). Structural differences exist between C γ 1 and C γ 2 genes: C γ 1 is made of three exons, with exon 2 encoding the cysteine residue forming a disulfide bridge with the δ chain. C γ 2 has an allelic polymorphism with either two or three copies of an exon homologous to C γ 1 exon 2 without codons for this cysteine residue. Thus, $\gamma\delta$ receptors using C γ 2 are not disulfide linked to the δ chain. On the cell surface at least four forms of γ

Correspondence to: Gennaro De Libero, Experimental Immunology, University Hospital, Hebelstrasse 20, 4031 Basel, Switzerland,
e-mail: Gennaro.DeLibero@unibas.ch

chain can be found: the 40-kDa $\gamma 1$ chain, the differently glycosylated $\gamma 2$ chains (of 40 or 44 kDa) encoded by the $C\gamma 2$ gene with the duplication of the exon 2, and the 55-kDa form encoded by the $C\gamma 2$ gene with the triplication of exon 2.

The human δ locus is within the TCR α locus on chromosome 14. One $C\delta$ gene segment is located in front of the $J\delta$ segment cluster and is preceded by four different $J\delta$ segments. Three diversity (D) elements are also identified in front of the $J\delta$ cluster [5–7]. The number of $V\delta$ chains found expressed on the cell surface is limited to six [8]. A molecular map of the relative localization of $V\delta$ genes indicates $V\delta 2$ as the most 3' $V\delta$ gene, followed in a 3'–5' direction by $V\delta$ genes 8, 7, 5, 1, 6 and 4, while $V\delta 3$ is located at 3' of $C\delta$ [8].

The different TCR $\gamma\delta$ which can be potentially expressed approximates to the large number of 10^{18} . The δ genes show an unprecedented junctional diversity with up to three $D\delta$ segments used in tandem, together with imprecise joining and the extensive incorporation of N nucleotides [9]. Despite the small number of $V\delta$ genes, these characteristics generate TCR δ chains with an extremely high variability in the CDR3 region, thus leading to a potential TCR $\gamma\delta$ repertoire which is at least three orders of magnitude higher than the TCR $\alpha\beta$ repertoire. This property may have important implications in the antigen recognition by $\gamma\delta$ T cells.

Another striking feature of the TCR $\gamma\delta$ is the preferential association of different $V\delta$ and $V\gamma$ chains. $V\delta 2$ mostly associates with the $V\gamma 9$ chain, while $V\delta 1$ and $V\delta 3$ chains mostly associate with $V\gamma$ chains other than $V\gamma 9$. This preferential association seems to be generated by some kind of antigen selection, since mixed combinations are found in $\gamma\delta$ T cell clones isolated from thymus [10].

Maturation and selection in the thymus

Ordered and coordinated rearrangement of $V\gamma$ and $V\delta$ loci in the human fetal thymus has been reported. The initial rearrangements join $V\delta 2$ to $D\delta 3$, and downstream $V\gamma$ genes ($V\gamma 8$ and $V\gamma 9$) to upstream $J\gamma$ gene segments. These rearrangements are characterized by minimal junctional diversity. At later developmental stages there is a switch such that $V\delta 1$ is joined to upstream $D\delta$ gene segments and $V\gamma$ genes are joined to downstream $J\gamma$ gene segments. These rearrangements are characterized by extensive junctional diversity [10, 11].

How $\gamma\delta$ T cells differentiate in human thymus is not as clear as in the mouse. Mouse thymic $\gamma\delta$ T cells are positively selected [12] and autoreactive $\gamma\delta$ T cells are anergized [13] or deleted [14]. Positive selection is driven by cortical epithelial cells [15] and for some TCR $\gamma\delta$ by MHC class I-like TL molecules [16]. The p56 lck protein tyrosine kinase, the CD45 protein tyrosine phosphatase [17] and the adapter SLP-76 [18] are necessary for intrathymic $\gamma\delta$ T cell differentiation. MAP kinase activation is required for positive selection of $\alpha\beta$, but not $\gamma\delta$ T cells [19]. IL-7 [20], IL-7 receptor [21] as well as IL-2 receptor beta chain [22] are required for maturation of $\gamma\delta$ thymocytes. CD30 is required for negative selection of $\gamma\delta$ T cells [23].

Maturation and selection in other organs

Mouse $\gamma\delta$ T cells may also mature extrathymically. Intestinal maturation is inferred by the absence of Thy 1 antigen on mouse $\gamma\delta$ T cells [24], and by the presence of im-

mature T cells in the gut of normal [25] and athymic mice (reviewed in [26]). Likewise in the human gut a minor population of cells has a phenotype of T cell progenitors. These precursor cells are CD7⁺, CD2⁻, CD3⁻, CD4⁻, CD5⁻, and CD8⁻ [27], and give rise to phenotypically mature $\alpha\beta$ and $\gamma\delta$ T cells.

TCR $\gamma\delta$ transgenic or deficient mice have been instrumental in further understanding extrathymic maturation of these cells. CD3 ζ chain-deficient animals display highly reduced numbers of thymocytes, but normal numbers of intraepithelial lymphocytes (IELs), which however constitute a new population. These cells express normal levels of TCR-CD3 complexes associated with Fc ϵ RI γ homodimers. In contrast to CD3 ζ -containing IELs, these cells do not proliferate when triggered with anti-TCR mAbs or mitogens, suggesting that they mature in the intestine independently from TCR engagement [28]. In the intestine, precursor cells are Thy-1⁺, CD3⁻, CD44⁺, CD25⁺, CD45R⁻, CD24⁺. They mature into the $\gamma\delta$ T cell lineage after expression of CD3 ϵ chain and IL-2 receptor [29] and in the presence of locally produced IL-7 [30]. Expression of CD45 is also required for normal numbers of IEL $\gamma\delta$ T cells. Indeed, in CD45 null mice apparently normal intestinal T cell maturation is followed by increased intestinal apoptosis and reduced numbers of IEL $\gamma\delta$ T cells [31]. After maturation in the gut, $\gamma\delta$ T cells become CD3 bright but are still functionally immature. They acquire a potent cytolytic activity only upon interaction with antigens and continuous stimulation [32]. Most likely antigen recognition induces an oligoclonal expansion of IELs, as suggested by the clonality of TCR sequences of intraepithelial, but not intra lamina propria T cells [33]. Extrathymic selection of $\gamma\delta$ T cells has also been reported in the lung of mice [34]. These $\gamma\delta$ T cells are positively selected by strain-specific polymorphic ligands that are encoded outside of the classical H-2 region. Selection can take place in the absence of thymus and requires IL-7 [35]. Maturation of $\gamma\delta$ T cells may also occur in decidual tissue [36].

Distribution of $\gamma\delta$ T cells in human tissues

A detailed analysis of the distribution of different V γ and V δ chains in various human tissues has been partially performed. Within the lymphoid organs, $\gamma\delta$ T cells appear to be evenly distributed [2]. At variance with mouse, $\gamma\delta$ T cells are rare in human skin, and form a minority of IELs in the gut. Nevertheless, human $\gamma\delta$ T cells appear preferentially associated with epithelial cells, as seen in the mouse. In the tonsil they are mainly located in the interfollicular area, often around the small vessels and in the intraepithelial and sub epithelial zones. In these areas about 30% of cells express the TCR $\gamma\delta$ [37], while very few $\gamma\delta$ T cells are found in liver, kidney, salivary glands [38], and the bronchial tree [39]. In the thymus, which is the organ where most lymphocytes develop, $\gamma\delta$ T cells are a very minor population, constituting less than 1% of total thymocytes.

The small number of available V γ and V δ segments are not used randomly and the TCR $\gamma\delta$ repertoire is markedly restricted. The intestinal $\gamma\delta$ T cells mostly use the V δ 1 and V δ 3 chains paired with members of V γ 1 gene family [40]. Nearly all the remaining $\gamma\delta$ T cells in peripheral blood use V δ 1. Only a fraction of total thymocytes (less than 0.05%) express the V γ 9/V δ 2 heterodimer. On the contrary, V γ 9/V δ 2 T cells are very frequent in peripheral blood, in the tonsil and in the spleen, where they represent 5–10% of total lymphocytes and 50–80% of $\gamma\delta$ T cells [10, 41]. V γ 9/V δ 2 cells are rare also in the placenta and in the peripheral blood of neonates [42]. Dur-

ing the first years of life these cells expand and become the predominant circulating $\gamma\delta$ T cell population. This finding has suggested the possibility that environmental antigens stimulate and increase the number of these cells during the first years of life [42].

Antigen specificities of human $\gamma\delta$ T cells

Despite intensive efforts, the antigens stimulating human $\gamma\delta$ T cells remain in part unknown. Several types of antigen specificities have been reported. Here the most common ones, which are likely the most important, will be discussed first.

Recognition of non-peptidic phosphorylated antigens

Peripheral $\gamma\delta$ T cells react with ligands present in many different microbes. Lysates from *Mycobacterium tuberculosis*, *M. bovis*, *M. leprae*, *Listeria monocytogenes*, *Staphylococcus aureus*, group A streptococci, *Escherichia coli*, *Salmonella typhi*, *Yersinia enterocolitica*, *Francisella tularensis* and *Plasmodium falciparum* induce proliferation of human $\gamma\delta$ T cells (reviewed in [43]). In all these cases proliferating cells express the V γ 9/V δ 2 heterodimer.

The mycobacterial ligands have been extensively studied. They are protease-resistant non-peptidic compounds with a low molecular weight (<3 kDa) [44]. Constant et al. [45] isolated four different active compounds from *M. tuberculosis* cell extracts. One of these substances contains triphosphorylated thymidine linked to an unknown compound. Tanaka et al. [46] found that monoalkylphosphates may induce activation of V γ 9/V δ 2 cells and not of other $\gamma\delta$ T cell clones bearing different V γ and V δ chains. In a third publication, it was reported that *M. tuberculosis* lysates contain active compounds with phosphate and an unidentified carbohydrate [47]. In later studies a ligand was purified from culture supernatants of *M. scrofulaceum* and *M. fortuitum* and characterized as a phosphorylated molecule without nucleotide residues [48]. This compound was identified as isopentenylpyrophosphate (IPP), which is an intermediate of prenyl and cholesterol biosynthesis and is present in both prokaryotic and eukaryotic cells [48]. At the same time we reported that IPP, together with five other naturally occurring non-peptidic metabolites, stimulate the same population of $\gamma\delta$ T cells [49]. In addition, we isolated and characterized from *M. tuberculosis* cell extracts a non-peptidic molecule with a molecular mass of 262 daltons, containing a pyrophosphate but no nitrogen or cheton residues [43]. This molecule has an activity which is ~1000 times higher than IPP and could represent an intermediate in IPP synthesis in bacteria. By analysis of all active compounds reported so far it appears that the length and structure of the alkyl chain are important for immunogenicity. For example, while methylphosphate is active, dimethylphosphate and trimethylphosphate are not stimulatory [46]. We found that substitutions of IPP with a cheton in position 2 or with a hydroxyl in position 3 destroy immunogenicity. Moreover, the number and position of the phosphate groups also play an important role: while 2,3-diphosphoglyceric acid is active, neither 2-diphosphoglyceric acid nor 3-diphosphoglyceric acid are immunogenic (our unpublished data).

Recently, two new antigen reactivities have been reported for V γ 9/V δ 2 cells. Bukowski et al. [50] have shown that this population specifically recognizes several

Table 1. Ligands stimulating V γ 9/V δ 2 cells

Compound	Natural source	Structure identification	P groups	m.w.
TubAg1	<i>M. tuberculosis</i>	No	1	–
TubAg2	<i>M. tuberculosis</i>	No	1	–
TubAg3	<i>M. tuberculosis</i>	Partial	3	–
TubAg4	<i>M. tuberculosis</i>	Partial	3	–
MEP	No	Yes	1	126
Methyl-P	No	Yes	1	112
n-Propyl-P	No	Yes	1	140
iso-propyl-P	No	Yes	1	140
sec-Butyl-P	No	Yes	1	154
β -Hydroxyethyl-P	No	Yes	1	142
Phosphoglycolic acid	Yes	Yes	1	156
P-non peptidic	<i>M. smegmatis</i>	No	?	–
P-non peptidic	<i>M. fortuitum</i>	No	?	–
P-non peptidic	<i>M. tuberculosis</i>	No	?	–
IPP	<i>M. smegmatis</i>	Yes	2	216
IPP-CH ₂ OH	<i>M. fortuitum</i>	Partial	2	276
DMAPP	Yes	Yes	2	216
Farnesyl-PP	Yes	Yes	2	382
Geranyl-PP	Yes	Yes	2	314
Geranyl-geranyl-PP	Yes	Yes	2	450
Monoethyl-2'-dTTP	?	Yes	3	510
Monoethyl-2',3'-dTTP	?	Yes	3	494
Monoethyl-2'-dUTP	?	Yes	3	512
PP-non peptidic	<i>M. tuberculosis</i>	Partial	2	262
DPG	Yes	Yes	2	266
G-3-P	Yes	Yes	1	172
Xylose-1-P	Yes	Yes	1	200
Ribose-1-P	Yes	Yes	1	200
Ethylamine	Yes	Yes	0	45
n-Propylamine	Yes	Yes	0	59
n-Butylamine	Yes	Yes	0	73
iso-Propylamine	Yes	Yes	0	59
iso-Butylamine	Yes	Yes	0	73
sec-Butylamine	Yes	Yes	0	73
iso-Amylamine	Yes	Yes	0	87

P, Phosphate; MEP, monoethylphosphate; P-non peptidic, phosphorylated non peptide; IPP, isopentenylpyrophosphate; DMAPP, dymethylallylpyrophosphate; PP, pyrophosphate; 2'-dTTP, 2'-deoxythymidine triphosphate; 2',3'-dTTP, 2',3'-dideoxythymidine triphosphate; 2'-dUTP, 2'-deoxyuridine triphosphate; DPG, diphosphoglyceric acid; G-3-P, glycerol-3-phosphate

alkylamines released in millimolar amounts by different bacteria. Recognition of these compounds is mediated by the V γ 9/V δ 2 TCR and is affected by the type of the alkyl chain. Thus, alkylamines represent a new class of $\gamma\delta$ ligands widely present in nature and capable of recruiting a large number of $\gamma\delta$ T cells. A second reactivity is directed against bisphosphonates (pamidronate and alendronate), drugs usually used to contrast bone resorption. One of the most frequent side effects of this therapy is an acute-phase reaction, whose intensity appears correlated with expansion of the V γ 9/V δ 2 T cell population [51]. The mechanism by which these compounds induce proliferation of $\gamma\delta$ T cells is not clear: they could directly mimic IPP and other pre-




Fig. 1. Four possible models of interaction between $\gamma\delta$ T cell receptor, stimulatory ligand and putative antigen-presenting molecule (APM) are shown (APC antigen-presenting cell)

nylphosphates, due to a structural homology, or instead induce accumulation of IPP and other metabolic intermediates since they block the mevalonate pathway [52]. Table 1 reports the active ligands described so far.

Analysis of several $\gamma\delta$ clones pointed out that only cells expressing both V γ 9 and V δ 2 chains react to non-peptidic ligands [46]. In addition, the junctional regions of the V γ 9 or V δ 2 chains are important to confer reactivity to non-peptidic ligands [53], implying a reactivity different from that to protein superantigens and MHC-peptide complexes. However, as T cell clones with different junctional sequences cross-react with the same collection of ligands [49, 54], it has to be argued that many different $\gamma\delta$ TCR recognize a structural motif common to all these compounds. An additional important observation is that a consistent fraction of V γ 9/V δ 2 thymocytes [55] or V γ 9/V δ 2 clones isolated from postnatal thymus [10] also react to this class of li-

gands, suggesting the possibility that this receptor has intrinsic structural characteristics which confer reactivity to non-peptidic ligands and their putative antigen-presenting molecules (APM). Whether dedicated molecules exist which present non-peptidic ligands to V γ 9/V δ 2 TCR is still an open issue. Filler cells and membrane-membrane interactions are definitely required to optimally stimulate $\gamma\delta$ T cells [54, 56, 57] and only cells of human origin have this capacity (our unpublished results). These filler cells might provide necessary co-stimulation or, alternatively, they could associate non-peptidic ligands with species-specific APM, thus forming new complexes recognized by the V γ 9/V δ 2 TCR. Another possible way of interaction is initial binding of ligands to APC and induction of conformational changes of surface APM, which then make cognate interaction with the TCR. A speculative possibility is that non-peptidic ligands first bind to conserved regions of the V γ 9/V δ 2 TCR, which acquires a new conformation and the capacity to interact with APC. These four possible models of interaction are illustrated in Figure 1.

Recognition of cell surface molecules

Antigen reactivities of human $\gamma\delta$ T cells against allo-MHC molecules [58–60], CD1c [61, 62], and CD48 [63] have been described. However, only a few $\gamma\delta$ clones react to these molecules, suggesting that they are rare specificities. Also in the mouse a few alloreactive $\gamma\delta$ clones were isolated and characterized. The reactivities of these cells are quite different from those of $\alpha\beta$ T cells. One clone reacted to an unknown peptide presented by the TL 27b molecule [64], a second to the MHC class I-like TL 10b molecule [65], and a third to the MHC class II I-Ek surface glycoprotein [66], independently of associated peptides and antigen processing [67]. Site-directed mutagenesis of MHC molecules showed that the topology of $\gamma\delta$ TCR interaction with the MHC is distinct from that of $\alpha\beta$ T cells [67]. An intriguing finding is that mouse [68] and human [69] $\gamma\delta$ T cells accumulate in the decidua during normal pregnancy. Mouse $\gamma\delta$ T cells recognize non-MHC-encoded molecules present on both mouse and human trophoblast cell lines, but not other tumor cells [70]. The recognized surface structures have not been identified.

Recognition of peptides and carbohydrates

Rare human $\gamma\delta$ T cell clones have been isolated which recognize peptide-MHC complexes [59, 71–73]. In mice there are also examples of these infrequent reactivities. Polyclonal $\gamma\delta$ T cells specific for ovalbumin were induced after antigen inhalation [74]. It was not analyzed whether they were activated by the intact protein or by small peptides. A series of mouse hybridomas were reported to be stimulated by peptides derived from bacterial heat shock proteins (HSPs) [75]. Recognition of these peptides required presentation by unknown non-MHC molecules, and was affected either by amino acid substitutions in the peptide or by the polymorphism of V γ chain.

Mouse $\gamma\delta$ T cells may also recognize di- and trisaccharides coupled to class I-binding peptides [76]. The crystal structure analysis of these glycosylated peptide-MHC complexes has shown that the carbohydrates are located in the central region of the putative TCR binding site without altering the overall MHC structure [77].

These findings support a model of antigen recognition by these $\gamma\delta$ TCR characterized by cognate interaction with the sugar residues and not with MHC molecules.

Recognition of heat shock proteins

A few studies have reported the possible recognition of HSPs by human $\gamma\delta$ T cells [78–80]. In two of these studies a rabbit antiserum or an mAb specific for an unknown human 58-kDa HSP partially inhibited reactivity of freshly isolated, but not of cloned, $\gamma\delta$ T cells to a tumor line stained by these antibodies. The HSP stimulating $\gamma\delta$ cells was not identified and solid data supporting cognate interaction of the $\gamma\delta$ TCR with members of this protein family have not been published. Taken together, all these findings show that antigen recognition by $\gamma\delta$ T cells is different from that of $\alpha\beta$ T cells. Additional evidence supporting this conclusion comes from the different length of CDR3 loops which are often critical for antigen binding in Ig and appear to provide the principal peptide binding residues in $\alpha\beta$ TCRs. Comparison of the CDR3 regions of Ig H and L chains with TCR α , β , γ and δ chains, showed that Ig H and TCR δ CDR3 are the most variable in size and are significantly longer than Ig L and TCR γ chains, respectively. In contrast, TCR α and β chains pairing is constrained by nearly identical average CDR3 lengths [81]. These TCR structural differences have been related to $\gamma\delta$ TCR recognition of molecules other than peptide-MHC complexes and, in general, more to a type of antigen recognition similar to that of Ig [82].

Effector functions of $\gamma\delta$ T cells

Like $\alpha\beta$ T cells, $\gamma\delta$ T cells can be classified in different functional subsets. $\gamma\delta$ T cell clones are potent cytotoxic cells which lyse different tumor cell lines [1] and secrete different lymphokines irrespective of the expressed V chains [3]. $\gamma\delta$ T cells isolated from leprosy patients release soluble factors that positively influence development of granulomas [83]. In contrast, an inhibitory role in granuloma formation during mycobacterial infection has been attributed to $\gamma\delta$ T cells in mice lacking the δ gene [84] or treated with anti-TCR $\gamma\delta$ mAbs [85]. Another important function is protection during infections. By releasing proinflammatory lymphokines, $\gamma\delta$ T cells might facilitate activation of macrophages in the early phases of infection. This has been described in mice infected with *Candida albicans*, in which $\gamma\delta$ T cells facilitate nitric oxide production by macrophages and, thus, enhance resistance to mucosal candidiasis [86]. In addition, by killing the infected target cells, $\gamma\delta$ T cells might contribute to microbial burden, as shown with cytotoxic CD8⁺ T cells in mycobacterial and listerial infections in mice [87–89] or with human CD8⁺ TCR $\alpha\beta$ ⁺ T cells [90, 91].

$\gamma\delta$ T cells may also regulate the effector phases of the immune response. After inhalation of soluble antigens, $\gamma\delta$ T cells potently inhibit induction of IgE secretion in the mouse, possibly by blocking maturation of Th2 cells [74]. Thus, $\gamma\delta$ T cells might have a prominent role in protection against primary allergic sensitization to environmental antigens.

$\gamma\delta$ T cells may also influence the functional type of T helper cells which develop during immune responses. In mice infected with *Listeria monocytogenes*, $\gamma\delta$ T cells produce Th1-type cytokines, while they produce Th2-type cytokines in *Nippostron-*

gylus brasiliensis-infected animals [92]. In both infections $\gamma\delta$ T cells are among the first type of lymphocytes activated. As the effector phase of the immune response is influenced by the cytokine milieu in which the initial antigen priming occurs, $\gamma\delta$ T cells may have the important regulatory role of determining the later response [92]. Furthermore, a possible role during viral infection has been proposed based on the findings that V γ 9/V δ 2 T cells recognize and kill target cells infected with different types of viruses [93].

An additional important function attributed to human $\gamma\delta$ T cells is help to B cells. Indeed, it has been shown that, analogously to $\alpha\beta$ T cells, human $\gamma\delta$ T may facilitate B cell maturation, isotype switching and IgG production [94].

Role of $\gamma\delta$ T cells in animal disease models

Studies conducted in mice deficient in $\gamma\delta$ T cells by homologous recombination or treated with anti-TCR $\gamma\delta$ mAbs have provided convincing evidence of the importance of $\gamma\delta$ T cells in different disease models. $\gamma\delta$ T cells provide protection during lethal encephalitis with HSV-1 [95], in malaria after immunization with inactivated sporozoites [96], and in the early phases of listeriosis [84, 97]. In the absence of $\gamma\delta$ T cells, mice are more susceptible to high *M. tuberculosis* inocula [98] and have more severe tissue inflammatory injury after low inocula of the same bacteria [99]. In infection with *Nocardia asteroides*, $\gamma\delta$ cells induce local recruitment of neutrophils responsible for microbial clearance [100]. In autoimmune disease models, $\gamma\delta$ T cells appear to have an important regulatory role on $\alpha\beta$ T cells and other effector cells. Elimination of $\gamma\delta$ T cells worsens clinical manifestations in adjuvant arthritis [101] and lupus nephritis [102], leads to the aggravation and disease recurrence in experimental allergic encephalomyelitis [103], and mediates prevention of diabetes following mucosal insulin aerosolization in NOD mice [104].

Evidence for participation of $\gamma\delta$ T cells in human diseases

Infections

Expansion of human $\gamma\delta$ T cells has been described in bacterial, viral and parasite infections. The number of $\gamma\delta$ T cells has been found increased in infections of bacterial origin such as brucellosis [105], salmonellosis [106], tuberculosis [107] and tularemia [108]. Increase of $\gamma\delta$ T cells in hospital workers who are in close contact with tuberculosis patients has been also reported [109]. Increased numbers of circulating $\gamma\delta$ T cells have been found in parasite infections such as malaria [106, 110, 111], ehrlichiosis [112], leishmaniasis [113, 114], toxoplasmosis [115], and trypanosomiasis [116]. In most of these infections the expanded cells display a V γ 9/V δ 2 TCR.

The number of $\gamma\delta$ T cells is also altered in some viral infections. In patients with AIDS a large number of oligoclonal V δ 1⁺ cells are found in the peripheral blood [117] and of V δ 2⁺ cells in the lung [118]. These $\gamma\delta$ populations might contribute to the immune defense against opportunistic microorganisms frequently present in HIV-infected patients. Expansion of $\gamma\delta$ T cells has also been found after cytomegalovirus (CMV) infection in kidney transplant patients [119]. The expanded cells express oligoclonal V δ 1 and V δ 3 chains and proliferate when challenged with CMV-infected

fibroblasts. These findings strongly suggest that an antigen-driven proliferation occurs during this viral infection.

The exact role of the expanded $\gamma\delta$ populations in infections is not clear. $\gamma\delta$ T cells might recognize microbial ligands and contribute to protection by reducing microbial load or by killing cells infected with intracellular pathogens. $\gamma\delta$ T cells might also exert the important function of signaling the microbial presence in the early phases of infection, when a small load of microbial cells is present in tissues and other lymphocyte populations have not yet been recruited.

Tumors

One of the first identified functions of human $\gamma\delta$ T cells was the “nonspecific” cytotoxicity toward tumor targets [1]. Moreover, some human $\gamma\delta$ T cells kill Burkitt’s lymphomas, thymic lymphomas and erythroleukemia cells [54, 120, 121]. Other studies identified $\gamma\delta$ T cell clones reacting with unknown surface molecules expressed by activated B cells or EBV-transformed cells [122]. In all these cases no antigens interacting with the $\gamma\delta$ TCR have been identified. Only a single $\gamma\delta$ T cell clone that recognizes a peptide derived from the Ig expressed by a myeloma from the same patient has been described [123]. In some CNS neoplasms, including primary malignancies, metastatic cancers, and meningiomas, oligoclonal expansion of $\gamma\delta$ populations was detected [124]. $\gamma\delta$ T cells were found infiltrating cutaneous melanocytic tumors and capable of killing solid tumors [125]. $\gamma\delta$ T cells may also accumulate around epithelial [126], renal [127] and lung [128] carcinomas.

A recent finding is that many tumors of epithelial origin express MICA [129]. Since this MHC-related molecule stimulates $V\delta 1^+$ T cells [130], recognition of this surface antigen may allow $\gamma\delta$ T cell response against tumor cells.

Autoimmune and inflammatory diseases

A possible involvement of $\gamma\delta$ T cells in the pathogenesis of autoimmune diseases has been claimed. However, clear evidence of direct participation of $\gamma\delta$ T cells in autoimmune reactions is not available. Most of the studies have analyzed the relative number of $\gamma\delta$ T cells in peripheral blood or in other tissues, their phenotype and TCR repertoire. In rheumatoid arthritis the number of $\gamma\delta$ T cells is increased in the affected joints [131], mainly in those with active synovitis [132]. In multiple sclerosis $\gamma\delta$ T cells accumulate around brain lesions during the exacerbation phases [133] and in the cerebrospinal fluid where they display polyclonal $V\delta 1$ chains [134, 135]. Expansion of $\gamma\delta$ T cells has been reported in cases of autoimmune thrombocytopenia [136] and autoimmune neutropenia [137]. In this latter study it was reported that five affected patients showed monoclonal expansion of $\gamma\delta$ T cells, thus suggesting a direct correlation with the development of neutropenia. Interestingly, clonal expansion of $V\delta 1^+$ T cells was observed in an HTLV-I carrier patient with chronic neutropenia [138]. Taken together, these findings suggest that in some circumstances neutrophils might express surface molecules recognized by unique subsets of $\gamma\delta$ TCR.

The number of circulating $\gamma\delta$ T cells is increased in several inflammatory diseases. Coeliac disease (CD), an immune-mediated disorder arising from an hyper-responsiveness to gluten with histological alterations in the small intestine, is charac-

terized by increased numbers of intraepithelial and lamina propria $\gamma\delta$ T cells showing a phenotype of activated and memory cells [139, 140]. However, increased numbers of $\gamma\delta$ T cells are found also in patients with normal jejunal morphology or with latent disease [141]. Furthermore, the number of $\gamma\delta$ T cells does not directly correlate with the presence of gluten in the diet [142]. According to all these observations, it is likely that $\gamma\delta$ T cells do not react against gluten components. Interestingly, most $\gamma\delta$ T cells in duodenum of CD patients express polyclonal TCR with predominant use of V δ 1, V δ 3 and V δ 8 chain [40], and the rare J δ 3 segment [143]. It is tempting to speculate that recognition of polymorphic molecules expressed by normal epithelial cells drives expansion of $\gamma\delta$ T cells expressing these particular V δ and J δ gene segments. Candidate $\gamma\delta$ T cell ligands are the MICA and MICB molecules, which activate V δ 1+ T cells [130]. In CD, $\gamma\delta$ T cells might have an anti-inflammatory role facilitating the repair of damaged tissue. The local protective role might be exerted by release of epithelial cell growth factors such as keratinocyte growth factor [144].

In inflammatory bowel disease (IBD), the number of $\gamma\delta$ T cells has also been found to be increased in blood [145] and intestine [146]. V δ 1-D δ 3-J δ 1-bearing cells were found to be expanded in patients either with severe disease or in those with recently diagnosed or less severe forms of IBD.

Other inflammatory diseases showing increased numbers of $\gamma\delta$ T cells are Still's disease [147], hypertrophic obstructive adenoids [148], Lyme arthritis [149], dermatitis herpetiformis [150], primary Sjogren's syndrome and untreated patients with systemic lupus erythematosus [151]. In chronic cutaneous lupus erythematosus V γ 9/V δ 2 T cells were observed in close vicinity to the damaged basal keratinocyte layer, suggesting their participation in the inflammatory reaction [152]. In lupus nephritis, $\gamma\delta$ T cells help production of anti-DNA immunoglobulins [153]. In patients undergoing surgical interventions, there is infiltration of $\gamma\delta$ T cells immediately after blood circulation is reestablished in ligated ischemic arteries, suggesting specific and immediate recruitment at the site of injury [154]. $\gamma\delta$ T cells are also present in the transition zone between normal intima and fatty streaks in atherosclerotic plaques, suggesting a role in initiating the inflammatory process in atherosclerosis [155].

All these studies suggest that $\gamma\delta$ T cells have multiple roles in autoimmune and inflammatory diseases. It is likely that in some cases $\gamma\delta$ T cells participate in the pathogenesis of autoimmune diseases as they may recognize self antigens. However, it is conceivable that in other instances they may also limit inflammatory reactions and facilitate tissue repair.

Regulation of $\gamma\delta$ T cells

The property of recognizing a large variety of non-peptidic ligands present both in eukaryotic and prokaryotic cells makes V γ 9/V δ 2 cells a readily activated cell population. Their overactivation may sometimes be responsible for dangerous acute inflammatory reactions, as reported in malaria infection [110] and following therapy with bisphosphonates [51]. Therefore, this type of antigen reactivity necessitates a tight control, which is likely provided by different mechanisms, leading to a fine balance between activation and inhibition of $\gamma\delta$ T cells.

One regulatory mechanism is provided by weak agonist non-peptidic ligands which induce a state of transient energy in $\gamma\delta$ T cells after repeated stimulation. Natural compounds such as 2,3-diphosphoglyceric acid, which is present in huge

amounts (5 mM) in erythrocytes, induce unresponsiveness in $\gamma\delta$ T cells to the most active ligands [156]. This state of anergy lasts for a few days and is associated with partial tyrosine phosphorylation of the CD3-TCR complex. All the V γ 9/V δ 2 T cells are affected by this inhibitory TCR engagement [156]. Thus, this mechanism may efficiently and simultaneously shut off the response of the whole V γ 9/V δ 2 T cell population, preventing its massive activation. Such a regulation may occur in patients constantly exposed to *P. falciparum*, e.g., individuals living in malaria-endemic areas who do not expand V γ 9/V δ 2 cells in vivo [157] and do not suffer from the pathognomonic acute inflammatory reaction concomitant to the blood stage of the disease.

A second regulatory mechanism is represented by expression of inhibitory receptors (IR) shared with NK cells. These receptors recognize MHC class I molecules and inhibit the response of $\gamma\delta$ T cells more effectively when low rather than high antigen doses are present [158]. Engagement of IR facilitates recruitment of SHP-1 phosphatase to TCR-CD3 complex and affects phosphorylation of Lck and ZAP-70 kinase, but not of CD3 ζ chain upon TCR triggering [158]. These events may cause abortion of proximal TCR-mediated signaling. The role of IR is to set a higher TCR $\gamma\delta$ activation threshold and therefore to focus the response of $\gamma\delta$ T cells against APC loaded with high amounts of antigen.

In some cases, activation of $\gamma\delta$ T cells requires facilitatory mechanisms. Optimization of $\gamma\delta$ T cell response is important in tissues where a small number of $\gamma\delta$ T cells is present and their effector functions have to be maximized to be effective. An enhancing mechanism is provided by expression of the CD66a molecule. This surface glycoprotein belongs to the CD66/CEA family and makes homotypic interactions with different CD66/CEA members. Engagement of CD66a with its physiological ligands enhances the amounts of released lymphokines, while it does not induce a shift of the dose-response curve (our unpublished data). Thus, it increases the potency of $\gamma\delta$ response. As CD66 molecules are frequently expressed by epithelial cells, this interaction facilitates release of large amounts of lymphokines in epithelial tissues when small numbers of responding $\gamma\delta$ T cells are present.

The function of $\gamma\delta$ T cells in immune response

The enormous body of information derived from the reviewed studies allows us to discuss the role of $\gamma\delta$ T cells in immune response on the basis of solid experimental data. A series of hypotheses can be made based on non-peptidic antigen specificity, on effector functions and also on cell number and distribution in human diseases.

The first function attributed to $\gamma\delta$ T lymphocytes was that of sentinel cells according to their tissue distribution [159]. As $\gamma\delta$ T cells in the mouse have the tendency to accumulate in the epithelia, it was suggested that this population may participate in the early host response against invading pathogens. A sentinel function has been attributed also to human V γ 9/V δ 2 lymphocytes according to their unique antigen recognition characteristics, rather than tissue distribution [43]. V γ 9/V δ 2 cells may fulfill the function of readily recruited and alerted sentinel cells, capable of immediately signaling the presence of danger. Their activation would have important consequences such as prompt release of pro-inflammatory cytokines and chemokines capable of facilitating the onset of local inflammation.

A second important function of $\gamma\delta$ T cells may be to fill the gap between innate and acquired immunity, providing the response during the time when antigen-specif-

ic $\alpha\beta$ T cells have not yet been recruited and expanded [43, 50]. The capacity of a series of ligands to activate a large number of $\gamma\delta$ T cells, all sharing the same V γ /V δ pair, in a cross-reactive manner is unique for a lymphocyte population. This type of T cell activation is different from activation induced by peptides or bacterial and viral superantigens and, to some extent, it resembles ligand recognition by receptors expressed on the surface of innate immunity cells (e.g., CD14, CR1 and mannose receptor). In other words, V γ 9/V δ 2 receptors recognize patterns of molecules and do not make fine discriminations among them. In this respect, this subset of human $\gamma\delta$ T cells exploits an antigen-recognition strategy typical of innate immunity cells, while at the same time retaining the functions of other lymphocytes.

A third important function is to drive maturation of antigen-specific T cells into different effector subsets. As shown during the early phases of infection in mice, the lymphokine secretion pattern of $\gamma\delta$ T lymphocytes controls the subsequent functional maturation of $\alpha\beta$ T cells [92]. Although there is no clear evidence that similar mechanisms also occur in human infections, it is likely that soluble factors released by human $\gamma\delta$ T cells influence local lymphokine milieu. Therefore, it is conceivable that human $\gamma\delta$ T cells also participate in the Th1 versus Th2 functional maturation of $\alpha\beta$ T cells.

A fourth function of human $\gamma\delta$ T cells is recognition of damaged or transformed epithelial cells expressing MICA and MICB proteins, whose gene transcription is regulated by promoter heat shock elements similar to those of HSP genes. As MICA and MICB are recognized by V δ 1-bearing cells, this subset may patrol the presence of recently altered cells in epithelial tissues.

Conclusions

In conclusion, the large number of studies on human $\gamma\delta$ T cells have shown that these lymphocytes share several characteristics in common with $\alpha\beta$ T cells, and also embody many unique properties. Some investigations have perhaps suffered a constant analogy with the $\alpha\beta$ T cell population, which has precluded new and original experimental approaches. Nevertheless, this gigantic amount of work has provided solid clues for defining the role of human $\gamma\delta$ T cells in diseases. In addition, we have also learnt more about the extreme plasticity of the immune system and its polymorphic capacity to adapt and recognize foreign molecules.

Acknowledgement. This work was supported by grant no. 31-045518.95 from Swiss National Fund, and by grants from the Velux Foundation, the Krebsliga beider Basel and Schweizerische Krebsliga. I thank Dr. L. Mori for reading the manuscript.

References

1. Brenner MB, McLean J, Dialynas DP, Strominger JL, Smith JA, Owen FL, Seidman JG, Ip S, Rosen F, Krangel MS (1986) Identification of a putative second T-cell receptor. *Nature* 322:145
2. Groh V, Porcellì S, Fabbi M, Lanier LL, Picker LJ, Anderson T, Warnke RA, Bhan AK, Strominger JL, Brenner MB (1989) Human lymphocytes bearing T cell receptor $\gamma\delta$ are phenotypically diverse and evenly distributed throughout the lymphoid system. *J Exp Med* 169:1277
3. Morita CT, Verma S, Aparicio P, Martinez C, Spius H, Brenner MB (1991) Functionally distinct subsets of human $\gamma\delta$ T cells. *Eur J Immunol* 21:2999

4. Lefranc MP, Rabbitts TH (1990) Genetic organization of the human T-cell receptor γ and δ loci. *Res Immunol* 141:565
5. Chien YH, Iwashima M, Kaplan KB, Elliott JF, Davis MM (1987) A new T-cell receptor gene located within the α locus and expressed early in T-cell differentiation. *Nature* 327:677
6. Hata S, Brenner MB, Krangel MS (1987) Identification of putative human T cell receptor δ complementary DNA clones. *Science* 238:678
7. Takihara Y, Reimann J, Michalopoulos E, Ciccone E, Moretta L, Mak TW (1989) Diversity and structure of human T cell receptor δ chain genes in peripheral blood $\gamma\delta$ -bearing T lymphocytes. *J Exp Med* 169:393
8. Migone N, Padovan S, Zappador C, Giachino C, Bottaro M, Matullo G, Carbonara C, Libero GD, Casorati G (1995) Restriction of the T-cell receptor V δ gene repertoire is due to preferential rearrangement and is independent of antigen selection. *Immunogenetics* 42:323
9. Hata S, Satyanarayana K, Devlin P, Band H, McLean J, Strominger JL, Brenner MB, Krangel MS (1988) Extensive junctional diversity of rearranged human T cell receptor δ genes. *Science* 240:1541
10. Casorati G, De Libero G, Lanzavecchia A, Migone N (1989) Molecular analysis of human $\gamma\delta^+$ clones from thymus and peripheral blood. *J Exp Med* 170:1521
11. Krangel MS, Yssel H, Brocklehurst C, Spits H (1990) A distinct wave of human T cell receptor $\gamma\delta$ lymphocytes in the early fetal thymus: evidence for controlled gene rearrangement and cytokine production. *J Exp Med* 172:847
12. Lafaille JJ, Haas W, Coutinho A, Tonegawa S (1990) Positive selection of $\gamma\delta$ T cells. *Immunol Today* 11:75
13. Bonneville M, Ishida I, Itohara S, Verbeek S, Berns A, Kanagawa O, Haas W, Tonegawa S (1990) Self-tolerance to transgenic $\gamma\delta$ T cells by intrathymic inactivation. *Nature* 344:163
14. Dent AL, Matis LA, Hooshmand F, Widacki SM, Bluestone JA, Hedrick SM (1990) Self-reactive $\gamma\delta$ T cells are eliminated in the thymus. *Nature* 343:714
15. De Koning J, Di Molfetto L, Reilly C, Wei Q, Havran WL, Lo D (1997) Thymic cortical epithelium is sufficient for the development of mature T cells in *relB*-deficient mice. *J Immunol* 158:2558
16. Tsujimura K, Takahashi T, Morita A, Hasegawa NH, Iwase S, Obata Y (1996) Positive selection of $\gamma\delta$ CTL by TL antigen expressed in the thymus. *J Exp Med* 184:2175
17. Kawai K, Kishihara K, Molina TJ, Wallace VA, Mak TW, Ohashi PS (1995) Impaired development of V γ 3 dendritic epidermal T cells in *p56lck* protein tyrosine kinase-deficient and CD45 protein tyrosine phosphatase-deficient mice. *J Exp Med* 181:345
18. Pivniouk V, Tsitsikov E, Swinton P, Rathbun G, Alt FW, Geha RS (1998) Impaired viability and profound block in thymocyte development in mice lacking the adaptor protein. SLP-76. *Cell* 94:229
19. Albcrola IJ, Hogquist KA, Swan KA, Bevan MJ, Perlmutter RM (1996) Positive and negative selection invoke distinct signaling pathways. *J Exp Med* 184:9
20. Moore TA, von Freeden, Jeffrey U, Murray R, Zlotnik A (1996) Inhibition of $\gamma\delta$ T cell development and early thymocyte maturation in *IL-7*^{-/-} mice. *J Immunol* 157:2366
21. Maki K, Sunaga S, Ikuta K (1996) The V-J recombination of T cell receptor- γ genes is blocked in interleukin-7 receptor-deficient mice. *J Exp Med* 184:2423
22. Kawai K, Suzuki H, Tomiyama K, Minagawa M, Mak TW, Ohashi PS (1998) Requirement of the IL-2 receptor β chain for the development of V γ 3 dendritic epidermal T cells. *J Invest Dermatol* 110:961
23. Amakawa R, Hakem A, Kundig TM, Matsuyama T, Simard JJ, Timms E, Wakeham A, Mittrucker HW, Griesser H, Takimoto H, Schmits R, Shahinian A, Ohashi P, Penninger JM, Mak TW (1996) Impaired negative selection of T cells in Hodgkin's disease antigen CD30-deficient mice. *Cell* 84:55124
24. Guy-Grand D, Cerf-Bensussan N, Malissen B, Malassis-Seris M, Briottet C, Vassalli P (1991) Two gut intraepithelial CD8⁺ lymphocyte populations with different T cell receptors: a role for the gut epithelium in T cell differentiation. *J Exp Med* 173:471
25. Bandeira A, Mota ST, Itohara S, Degermann S, Heusser C, Tonegawa S, Coutinho A (1990) Localization of $\gamma\delta$ T cells to the intestinal epithelium is independent of normal microbial colonization. *J Exp Med* 172:239
26. Haas W, Pereira P, Tonegawa S (1993) $\gamma\delta$ cells. *Annu Rev Immunol* 11:637
27. Jarry A, Cerf-Bensussan N, Brousse N, Selz F, Guy GD (1990) Subsets of CD3⁺ (T cell receptor $\alpha\beta$ or $\gamma\delta$) and CD3⁻ lymphocytes isolated from normal human gut epithelium display phenotypical features different from their counterparts in peripheral blood. *Eur J Immunol* 20:1097

28. Liu CP, Ueda R, She J, Sancho J, Wang B, Weddell G, Loring J, Kurahara C, Dudley EC, Hayday A, Terhorst C, Huang M (1993) Abnormal T cell development in CD3- ζ -/- mutant mice and identification of a novel T cell population in the intestine. *EMBO J* 12:4863
29. Page ST, Bogatzki LY, Hamerman JA, Sweenie CH, Hogarth PJ, Malissen M, Perlmutter RM, Pullen AM (1998) Intestinal intraepithelial lymphocytes include precursors committed to the T cell receptor $\alpha\beta$ lineage. *Proc Natl Acad Sci USA* 95:9459
30. Laky K, Lefrancois L, Freedden-Jeffry U von, Murray R, Puddington L (1998) The role of IL-7 in thymic and extrathymic development of TCR $\gamma\delta$ cells. *J Immunol* 161:707
31. Yada S, Kishihara K, Kong YY, Nomoto K (1998) Differential requirements of CD45 protein tyrosine phosphatase for cytolytic activities and intrathymic and extrathymic development of intestinal intraepithelial lymphocytes. *J Immunol* 161:2208
32. Lefrancois L, Goodman T (1989) In vivo modulation of cytolytic activity and Thy-1 expression in TCR- $\gamma\delta$ intraepithelial lymphocytes. *Science* 243:1716
33. Van Kerckhove C, Russel GJ, Deusch K, Reich K, Bahn AK, DerSimonian H, Brenner MB (1992) Oligoclonality of human intestinal intraepithelial T cells. *J Exp Med* 175:57
34. Sim GK, Augustin A (1991) Extrathymic positive selection of $\gamma\delta$ T cells V γ 4J γ 1 rearrangements with "GxYS" junctions. *J Immunol* 146:2439
35. Hayes SM, Sirr A, Jacob S, Sim GK, Augustin A (1996) Role of IL-7 in the shaping of the pulmonary $\gamma\delta$ T cell repertoire. *J Immunol* 156:2723
36. Hayakawa S, Shiraishi H, Saitoh S, Satoh K (1996) Decidua as a site of extrathymic V γ 1 T-cell differentiation. *Am J Reprod Immunol* 35:233
37. Graeme CF, Bhan AK, Harris NL (1993) Immunohistochemical characterization of intraepithelial and subepithelial mononuclear cells of the upper airways. *Am J Pathol* 143:1416
38. Falini B, Flenghi L, Pileri S, Pelicci P, Fagioli M, Martelli MF, Moretta L, Ciccone E (1989) Distribution of T cells bearing different forms of the T cell receptor $\gamma\delta$ in normal and pathological human tissues. *J Immunol* 143:2480
39. Richmond I, Pritchard GE, Ashcroft T, Corris PA, Walters EH (1993) Distribution of $\gamma\delta$ T cells in the bronchial tree of smokers and non-smokers. *J Clin Pathol* 46:926
40. De Libero G, Rocci MP, Casorati G, Giachino C, Oderda G, Tavassoli K, Migone N (1993) T cell receptor heterogeneity in $\gamma\delta$ T cell clones from intestinal biopsies of patients with celiac disease. *Eur J Immunol* 23:499
41. Triebel F, Faure F, Mami-Chouaib F, Jitsukawa S, Griscelli A, Genevec C, Roman RS, Hercend T (1988) A novel human V δ gene expressed predominantly in the T γ A fraction of $\gamma\delta$ ⁺ peripheral lymphocytes. *Eur J Immunol* 18:2021
42. Parker CM, Groh V, Band H, Porcelli SA, Morita C, Fabbi M, Glass D, Strominger JL, Brenner MB (1990) Evidence for extrathymic changes in the T cell receptor $\gamma\delta$ repertoire. *J Exp Med* 171:1597
43. De Libero G (1997) Sentinel function of broadly reactive human $\gamma\delta$ T cells. *Immunol Today* 18:22
44. Pfeffer K, Schoel B, Gulle H, Kaufmann SH, Wagner H (1990) Primary responses of human T cells to mycobacteria: a frequent set of $\gamma\delta$ T cells are stimulated by protease-resistant ligands. *Eur J Immunol* 20:1175
45. Constant P, Davodeau F, Peyrat MA, Poquet Y, Puzo G, Bonneville M, Fournie JJ (1994) Stimulation of human $\gamma\delta$ T cells by nonpeptidic mycobacterial ligands. *Science* 264:267
46. Tanaka Y, Sano S, Nieves E, De Libero G, Rosa D, Modlin RL, Brenner MB, Bloom BR, Morita CT (1994) Nonpeptide ligands for human $\gamma\delta$ T cells. *Proc Natl Acad Sci USA* 91:8175
47. Schoel B, Sprenger S, Kaufmann SHE (1994) Phosphate is essential for stimulation of V γ 9V δ 2 T lymphocytes by mycobacterial low molecular weight ligand. *Eur J Immunol* 24:1886
48. Tanaka Y, Morita CT, Tanaka Y, Nieves E, Brenner MB, Bloom BR (1995) Natural and synthetic nonpeptide antigens recognized by human $\gamma\delta$ T cells. *Nature* 375:155
49. Bürk MR, Mori L, De Libero G (1995) Human V γ 9-V δ 2 cells are stimulated in a cross-reactive fashion by a variety of phosphorylated metabolites. *Eur J Immunol* 25:2052
50. Bukowski JF, Morita CT, Brenner MB (1999) Human $\gamma\delta$ T cells recognize alkylamines derived from microbes, edible plants, and tea: implications for innate immunity. *Immunity* 11:57
51. Kunzmann V, Bauer E, Wilhelm M (1999) $\gamma\delta$ T-cell stimulation by pamidronate. *N Engl J Med* 340:737
52. Fisher JE, Rogers MJ, Halasy JM, Luckman SP, Hughes DE, Masarachia PJ, Wesolowski G, Russell RG, Rodan GA, Reszka AA (1999) Alendronate mechanism of action: geranylgeraniol, an intermediate in the mevalonate pathway, prevents inhibition of osteoclast formation, bone resorption, and kinase activation in vitro. *Proc Natl Acad Sci USA* 96:133

53. Bukowski JF, Morita CT, Band H, Brenner MB (1998) Crucial role of TCR γ chain junctional region in prenyl pyrophosphate antigen recognition by $\gamma\delta$ T cells. *J Immunol* 161:286
54. De Libero G, Casorati G, Giachino C, Carbonara C, Migone N, Matzinger P, Lanzavecchia A (1991) Selection by two powerful antigens may account for the presence of the major population of human peripheral $\gamma\delta$ T cells. *J Exp Med* 173:1311
55. Kabelitz D, Bender A, Prospero T, Wesselborg S, Janssen O, Pechhold K (1991) The primary response of human $\gamma\delta^+$ T cells to *Mycobacterium tuberculosis* is restricted to V γ 9-bearing cells. *J Exp Med* 173:1331
56. Morita CT, Beckman EM, Bukowski JF, Tanaka Y, Band H, Bloom BR, Golan DE, Brenner MB (1995) Direct presentation of nonpeptide prenyl pyrophosphate antigens to human $\gamma\delta$ T cells. *Immunity* 3:495
57. Lang F, Peyrat MA, Constant P, Davodeau F, David AJ, Poquet Y, Vie H, Fournie JJ, Bonneville M (1995) Early activation of human V γ 9 V δ 2 T cell broad cytotoxicity and TNF production by nonpeptidic mycobacterial ligands. *J Immunol* 154:5986
58. Ciccone E, Viale O, Pende D, Malnati M, Battista FG, Barocci S, Moretta A, Moretta L (1989) Specificity of human T lymphocytes expressing a $\gamma\delta$ T cell antigen receptor. Recognition of a polymorphic determinant of HLA class I molecules by a $\gamma\delta$ clone. *Eur J Immunol* 19:1267
59. Spits H, Paliard X, Engelhard VH, de Vries J (1990) Cytotoxic activity and lymphokine production of T cell receptor (TCR)- $\alpha\beta^+$ and TCR- $\gamma\delta^+$ cytotoxic T lymphocyte (CTL) clones recognizing HLA-A2 and HLA-A2 mutants. Recognition of TCR- $\gamma\delta^+$ CTL clones is affected by mutations at positions 152 and 156. *J Immunol* 144:4156
60. Rivas A, Koide J, Cleary ML, Engleman EG (1989) Evidence for involvement of the γ , δ T cell antigen receptor in cytotoxicity mediated by human alloantigen-specific T cell clones. *J Immunol* 142:1840
61. Porcelli S, Brenner MB, Greenstein JL, Balk SP, Terhorst C, Bleicher PA (1989) Recognition of cluster of differentiation 1 antigens by human CD4-CD8- cytolytic T lymphocytes. *Nature* 341:447
62. Faure F, Jitsukawa S, Miossec C, Hercend T (1990) CD1c as a target recognition structure for human T lymphocytes: analysis with peripheral blood $\gamma\delta$ cells. *Eur J Immunol* 20:703
63. Del Porto P, Mami Chouaib F, Bruneau JM, Jitsukawa S, Dumas J, Harnois M, Hercend T (1991) TCT 1, a target molecule for $\gamma\delta$ T cells, is encoded by an immunoglobulin superfamily gene (Blast-1) located in the CD1 region of human chromosome 1. *J Exp Med* 173:1339
64. Van Kaer L, Wu M, Ichikawa Y, Ito K, Bonneville M, Ostrand-Rosenberg S, Murphy DB, Tonegawa S (1991) Recognition of MHC TL gene products by $\gamma\delta$ T cells. *Immunol Rev* 120:89
65. Bluestone JA, Cron RQ, Cotterman M, Houlden BA, Matis LA (1988) Structure and specificity of T cell receptor $\gamma\delta$ on major histocompatibility complex antigen-specific CD3 $^+$, CD4 $^-$, CD8 $^-$ T lymphocytes. *J Exp Med* 168:1899
66. Matis LA, Fry AM, Cron RQ, Cotterman MM, Dick RF, Bluestone JA (1989) Structure and specificity of a class II MHC alloreactive $\gamma\delta$ T cell receptor heterodimer. *Science* 245:746
67. Schild H, Mavaddat N, Litzenberger C, Ehrlich EW, Davis MM, Bluestone JA, Matis L, Draper RK, Chien YH (1994) The nature of major histocompatibility complex recognition by $\gamma\delta$ T cells. *Cell* 76:29
68. Heyborne KD, Cranfill RL, Carding SR, Born WK, O'Brien RL (1992) Characterization of $\gamma\delta$ T lymphocytes at the maternal-fetal interface. *J Immunol* 149:2872
69. Mincheva-Nilsson L, Baranov V, Yeung MM, Hammarstrom S, Hammarstrom ML (1994) Immunomorphologic studies of human decidua-associated lymphoid cells in normal early pregnancy. *J Immunol* 152:2020
70. Heyborne K, Fu YX, Nelson A, Farr A, O'Brien R, Born W (1994) Recognition of trophoblasts by $\gamma\delta$ T cells. *J Immunol* 153:2918
71. Kozbor D, Trinchieri G, Monos DS, Isobe M, Russo G, Haney JA, Zmijewski C, Croce CM (1989) Human TCR- $\gamma^+\delta^+$, CD8 $^+$ T lymphocytes recognize tetanus toxoid in an MHC-restricted fashion. *J Exp Med* 169:1847
72. Holoshitz J, Vila LM, Keroack BJ, McKinley DR, Bayne NK (1992) Dual antigenic recognition by cloned human $\gamma\delta$ T cells. *J Clin Invest* 89:308
73. Del Porto P, D'Amato M, Fiorillo MT, Tuosto L, Piccolella E, Sorrentino R (1994) Identification of a novel HLA-B27 subtype by restriction analysis of a cytotoxic $\gamma\delta$ T cell clone. *J Immunol* 153:3093
74. McMenamin C, Pimm C, McKersey M, Holt PG (1994) Regulation of IgE responses to inhaled antigen in mice by antigen-specific $\gamma\delta$ T cells. *Science* 265:1869

75. Born W, Hall L, Daltas A, Boymel J, Shinnick T, Young D, Brennan P, O'Brien R (1990) Recognition of a peptide antigen by heat shock-reactive $\gamma\delta$ T lymphocytes. *Science* 249:67
76. Abdel-Motal UM, Berg L, Rosen A, Bengtsson M, Thorpe CJ, Kihlberg J, Dahmen J, Magnusson G, Karlsson KA, Jondal M (1996) Immunization with glycosylated Kb-binding peptides generates carbohydrate-specific, unrestricted cytotoxic T cells. *Eur J Immunol* 26:544
77. Speir JA, Abdel-Motal UM, Jondal M, Wilson IA (1999) Crystal structure of an MHC class I presented glycopeptide that generates carbohydrate-specific CTL. *Immunity* 10:51
78. Haregewoin A, Soman G, Hom RC, Finberg RW (1989) Human $\gamma\delta^+$ T cells respond to mycobacterial heat-shock protein. *Nature* 340:309
79. Fisch P, Malkovsky M, Kovats S, Sturm E, Braakman E, Klein BS, Voss SD, Morrissey LW, DeMars R, Welch WJ, Boihus RLH, Sondel PM (1990) Recognition by human $V\gamma 9/V\delta 2$ T cells of a GroEL homolog on Daudi Burkitt's lymphoma cells. *Science* 250:1269
80. Kaur I, Voss SD, Gupta RS, Schell K, Fisch P, Sondel PM (1993) Human peripheral $\gamma\delta$ T cells recognize hsp60 molecules on Daudi Burkitt's lymphoma cells. *J Immunol* 150:2046
81. Rock EP, Sibbald PR, Davis MM, Chien YH (1994) CDR3 length in antigen-specific immune receptors. *J Exp Med* 179:323
82. Davis MM, Chien Y (1995) Issues concerning the nature of antigen recognition by $\alpha\beta$ and $\gamma\delta$ T-cell receptors. *Immunol Today* 16:316
83. Modlin RL, Pirmez C, Hofman FM, Torigian V, Uyemura K, Rea TH, Bloom BR, Brenner MB (1989) Lymphocytes bearing antigen-specific $\gamma\delta$ T-cell receptors accumulate in human infectious disease lesions. *Nature* 339:544
84. Mombaerts P, Arnoldi J, Russ F, Tonegawa S, Kaufmann SH (1993) Different roles of $\alpha\beta$ and $\gamma\delta$ T cells in immunity against an intracellular bacterial pathogen. *Nature* 365:53
85. Fu Y-X, Roark CE, Kelly K, Drevets D, Campbell P, O'Brien R, Born W (1994) Immune protection and control of inflammatory tissue necrosis by $\gamma\delta$ T cells. *J Immunol* 153:3101
86. Jones-Carson J, Vazquez-Torres A, van der Heyde HC, Warner T, Wagner RD, Balish E (1995) $\gamma\delta$ T cell induced nitric oxide production enhances resistance to mucosal candidiasis. *Nat Med* 1:552
87. De Libero G, Flesch I, Kaufmann SH (1988) Mycobacteria-reactive $Lyt-2^+$ T cell lines. *Eur J Immunol* 18:59
88. Flynn JL, Goldstein M, Triebold KJ, Koller B, Bloom BR (1992) MHC class I restricted T cells are required for resistance to *M tuberculosis* infection. *Proc Natl Acad Sci USA* 89:12013
89. Kägi D, Lederman B, Bürki K, Hengartner H, Zinkernagel R M (1994) $CD8^+$ T cell-mediated protection against an intracellular bacterium by perforin-dependent cytotoxicity. *Eur J Immunol* 24:3068
90. Stenger S, Mazzaccaro RJ, Uyemura K, Cho S, Barnes PF, Rosat JP, Sette A, Brenner MB, Porcelli SA, Bloom BR, Modlin RL (1997) Differential effects of cytolytic T cell subsets on intracellular infection. *Science* 276:1684
91. Stenger S, Hanson DA, Teitelbaum R, Dewan P, Niazi KR, Froelich CJ, Ganz T, Thoma-Uszynski S, Melian A, Boydan C, Porcelli SA, Bloom BR, Krensky AM, Modlin RL (1998) An antimicrobial activity of cytolytic T cells mediated by granulysin. *Science* 282:121
92. Ferrick DA, Schrenzel MD, Mulvania T, Hsieh B, Ferlin WG, Lepper H (1995) Differential production of interferon- γ and interleukin-4 in response to Th1- and Th2-stimulating pathogens by $\gamma\delta$ T cells in vivo. *Nature* 373:255
93. Bukowski JF, Morita CT, Brenner MB (1994) Recognition and destruction of virus-infected cells by human $\gamma\delta$ CTL. *J Immunol* 153:5133
94. Horner AA, Jabara H, Ramesh N, Geha RS (1995) $\gamma\delta$ T lymphocytes express CD40 ligand and induce isotype switching in B lymphocytes. *J Exp Med* 181:1239
95. Shammass R, Kodukula P, Tang Q, Hendricks RL, Bluestone JA (1997) T cell $\gamma\delta$ cells protect mice from herpes simplex virus type 1-induced lethal encephalitis. *J Exp Med* 185:1969
96. Tsuji M, Mombaerts P, Lefrancois L, Nussenzweig RS, Zavala F, Tonegawa S (1994) $\gamma\delta$ T cells contribute to immunity against the liver stages of malaria in $\alpha\beta$ T-cell-deficient mice. *Proc Natl Acad Sci USA* 91:345
97. Hiromatsu K, Yoshikai Y, Matsuzaki G, Ohga S, Muramori K, Matsumoto K, Bluestone JA, Nomoto K (1992) A protective role of $\gamma\delta$ T cells in primary infection with *Listeria monocytogenes* in mice. *J Exp Med* 175:49
98. Ladel CH, Hess J, Daugelat S, Mombaerts P, Tonegawa S, Kaufmann SH (1995) Contribution of $\alpha\beta$ and $\gamma\delta$ T lymphocytes to immunity against *Mycobacterium bovis* bacillus Calmette Guerin: studies with T cell receptor-deficient mutant mice. *Eur J Immunol* 25:838

99. D'Souza CD, Cooper AM, Frank AA, Mazzaccaro RJ, Bloom BR, Orme IM (1997) An anti-inflammatory role for $\gamma\delta$ T lymphocytes in acquired immunity to *Mycobacterium tuberculosis*. *J Immunol* 158:1217
100. King DP, Hyde DM, Jackson KA, Novosad DM, Ellis TN, Putney L, Stovall MY, Van Winkle LS, Beaman BL, Ferrick DA (1999) Cutting edge: protective response to pulmonary injury requires $\gamma\delta$ T lymphocytes. *J Immunol* 162:5033
101. Pelegri C, Kuhnlein P, Buchner E, Schmidt CB, Franch A, Castell M, Hunig T, Emmrich F, Kinne RW (1996) Depletion of $\gamma\delta$ T cells does not prevent or ameliorate, but rather aggravates, rat adjuvant arthritis. *Arthritis Rheum* 39:204
102. Peng SL, Madaio MP, Hayday AC, Craft J (1996) Propagation and regulation of systemic autoimmunity by $\gamma\delta$ T cells. *J Immunol* 157:5689
103. Kobayashi Y, Kawai K, Ito K, Honda H, Sobue G, Yoshikai Y (1997) Aggravation of murine experimental allergic encephalomyelitis by administration of T-cell receptor $\gamma\delta$ -specific antibody. *J Neuroimmunol* 73:169
104. Harrison LC, Dempsey-Collier M, Kramer DR, Takahashi K (1996) Aerosol insulin induces regulatory CD8 $\gamma\delta$ T cells that prevent murine insulin-dependent diabetes. *J Exp Med* 184:2167
105. Bertotto A, Gerli R, Spinozzi F, Muscat C, Scalise F, Castellucci G, Sposito M, Candio F, Vaccaro R (1993) Lymphocytes bearing the $\gamma\delta$ T cell receptor in acute *Brucella melitensis* infection. *Eur J Immunol* 23:1177
106. Hara T, Mizuno Y, Takaki K, Takada H, Akeda H, Aoki T, Nagata M, Ueda K, Matsuzaki G, Yoshikai Y, Nomoto K (1992) Predominant activation and expansion of V γ 9-bearing $\gamma\delta$ T cells in vivo as well as in vitro in Salmonella infection. *J Clin Invest* 90:204
107. Balbi B, Valle MT, Oddera S, Giunti D, Manca F, Rossi GA, Allegra L (1993) T-lymphocytes with $\gamma\delta^+ V\delta 2^+$ antigen receptors are present in increased proportions in a fraction of patients with tuberculosis or with sarcoidosis. *Am Rev Respir Dis* 148:1685
108. Sumida T, Maeda T, Takahashi H, Yoshida S, Yonaka F, Sakamoto A, Tomioka H, Koike T, Yoshida S (1992) Predominant expansion of V γ 9/V δ 2 cells in a tularemia patient. *Infect Immun* 60:2554
109. Ueta C, Tsuyuguchi I, Kawasumi H, Takashima T, Toba H, Kishimoto S (1994) Increase of $\gamma\delta$ T cells in hospital workers who are in close contact with tuberculosis patients. *Infect Immun* 62:5434
110. Ho M, Webster HK, Tongtawe P, Pattanapanyasat K, Weidanz WP (1990) Increased $\gamma\delta$ T cells in acute *Plasmodium falciparum* malaria. *Immunol Lett* 25:139
111. Nakazawa S, Brown AE, Maeno Y, Smith CD, Aikawa M (1994) Malaria-induced increase of splenic $\gamma\delta$ T cells in humans, monkeys, and mice. *Exp Parasitol* 79:391
112. Caldwell CW, Everett ED, McDonald G, Yesus YW, Roland WE (1995) Lymphocytosis of $\gamma\delta$ T cells in human ehrlichiosis. *Am J Clin Pathol* 103:761
113. Raziuddin S, Telmasani AW, El-Awad E, Al-Amari O, Al-Yanadi M (1992) $\gamma\delta$ T cells and the immune response in visceral leishmaniasis. *Eur J Immunol* 22:1143
114. Alalbac M, Daga A, Harms G, Morris J, Yu RC, Zwingerberger K, Chu AC (1993) Molecular analysis of the $\gamma\delta$ T-cell receptor repertoire in normal human skin and in Oriental cutaneous leishmaniasis. *Exp Dermatol* 2:106
115. Scalise F, Gerli R, Castellucci G, Spinozzi F, Fabietti G M, Crupi S, Sensi L, Britta R, Vaccaro R, Bertotto A (1992) Lymphocytes bearing the $\gamma\delta$ T-cell receptor in acute toxoplasmosis. *Immunology* 76:668
116. Flynn JN, Sileghem M (1994) Involvement of $\gamma\delta$ T cells in immunity to trypanosomiasis. *Immunology* 83:86
117. Hiuz T, Wesch D, Friese K, Reckziegel A, Arden B, Kabelitz D (1994) T cell receptor $\gamma\delta$ repertoire in HIV-1-infected individuals. *Eur J Immunol* 24:3044
118. Agostini C, Zambello R, Trentin L, Cerutti A, Bulian P, Crivellaro C, Cipriani A, Semenzato G (1994) $\gamma\delta$ T cell receptor subsets in the lung of patients with HIV-1 infection. *Cell Immunol* 153:194
119. Dechanet J, Merville P, Berge F, Bone-Mane G, Taupin JL, Michel P, Joly P, Bonneville M, Potaux L, Moreau JF (1999) Major expansion of $\gamma\delta$ T lymphocytes following cytomegalovirus infection in kidney allograft recipients. *J Infect Dis* 179:1
120. Fisch P, Malkovsky M, Braakman E, Sturm E, Bolhuis R L, Prieve A, Sosman J A, Lam VA, Sondel PM (1990) $\gamma\delta$ T cell clones and natural killer cell clones mediate distinct patterns of non-major histocompatibility complex-restricted cytotoxicity. *J Exp Med* 171:1567
121. Di Fabrizio L, Kimura Y, Ware R, Rogozinski L, Chess L (1991) Specific triggering of $\gamma\delta$ receptors by K562 cells activates the $\gamma\delta$ TCR and may regulate natural killer-like function. *J Immunol* 146:2495

122. Orsini DL, van Gils M, Kooy YM, Struyk L, Klein G, van den Elsen P, Koning F (1994) Functional and molecular characterization of B cell-responsive V δ 1⁺ $\gamma\delta$ T cells. *Eur J Immunol* 24: 3199
123. Kim HT, Nelson EL, Clayberger C, Sanjanwala M, Sklar J, Krensky AM (1995) $\gamma\delta$ T cell recognition of tumor Ig peptide. *J Immunol* 154:1614
124. Merlo A, Filgueira L, Zuber M, Juretic A, Harder F, Gratzl O, De Libero G, Heberer M, Spagnoli GC (1993) T-cell receptor V-gene usage in neoplasms of the central nervous system. A comparative analysis in cultured tumor infiltrating and peripheral blood T cells. *J Neurosurg* 78:630
125. Flageul B, Bachelez H, Boumsell L, Degos L, Bensussan A (1990) Infiltrating lymphocytes in benign and malignant naevomelanocytic lesions. *Nouv Rev Fr Hematol* 32:9
126. Maeurer MJ, Martin D, Walter W, Liu K, Zitvogel L, Halusczyk K, Rabinowich H, Duquesnoy R, Storkus W, Lotze MT (1996) Human intestinal V δ 1⁺ lymphocytes recognize tumor cells of epithelial origin. *J Exp Med* 183:1681
127. Choudhary A, Davodeau F, Moreau A, Peyrat MA, Bonneville M, Jotereau F (1995) Selective lysis of autologous tumor cells by recurrent $\gamma\delta$ tumor-infiltrating lymphocytes from renal carcinoma. *J Immunol* 154:3932
128. Zocchi MR, Ferrarini M, Migone N, Casorati G (1994) T-cell receptor V δ gene usage by tumour reactive $\gamma\delta$ T lymphocytes infiltrating human lung cancer. *Immunology* 81:234
129. Groh V, Rhinehart R, Secrist H, Bauer S, Grabstein KH, Spies T (1999) Broad tumor-associated expression and recognition by tumor-derived $\gamma\delta$ T cells of MICA and MICB. *Proc Natl Acad Sci U S A* 96:6879
130. Groh V, Steinle A, Bauer S, Spies T (1998) Recognition of stress-induced MHC molecules by intestinal epithelial $\gamma\delta$ T cells. *Science* 279:1737
131. Brennan FM, Londei M, Jackson AM, Hercend T, Brenner MB, Maini RN, Feldmann M (1988) T cells expressing $\gamma\delta$ chain receptors in rheumatoid arthritis. *J Autoimmun* 1:319
132. Jacobs MR, Haynes BF (1992) Increase in TCR $\gamma\delta$ T lymphocytes in synovia from rheumatoid arthritis patients with active synovitis. *J Clin Immunol* 12:130
133. Wucherpfennig KW, Newcombe J, Li H, Keddy C, Cuzner ML, Hafler DA (1992) $\gamma\delta$ T-cell receptor repertoire in acute multiple sclerosis lesions. *Proc Natl Acad Sci U S A* 89:4588
134. Shimonkevitz R, Colburn C, Burnham JA, Murray RS, Kotzin BL (1993) Clonal expansions of activated $\gamma\delta$ T cells in recent-onset multiple sclerosis. *Proc Natl Acad Sci USA* 90:923
135. Nick S, Pileri P, Tongiani S, Uematsu Y, Kappos L, De Libero G (1995) T cell receptor $\gamma\delta$ repertoire is skewed in cerebrospinal fluid of multiple sclerosis patients: molecular and functional analyses of antigen-reactive $\gamma\delta$ clones. *Eur J Immunol* 25:355
136. Ashihara E, Shimazaki C, Hirata T, Okawa K, Oku N, Goto H, Inaba T, Fujita N, Nakagawa M (1993) Autoimmune thrombocytopenia following peripheral blood stem cell autografting. *Bone Marrow Transplant* 12:297
137. Scott CS, Richards SJ, Sivakumaran M, Steed AJ, Norfolk DR, Milligan DW, Short M (1994) Persistent clonal expansions of CD3⁺ TCR $\gamma\delta$ ⁺ and CD3⁺ TCR $\alpha\beta$ ⁺ CD4⁺CD8⁻ lymphocytes associated with neutropenia. *Leuk Lymphoma* 14:429
138. Horiuchi T, Hirokawa M, Satoh K, Kitabayashi A, Muira AB (1999) Clonal expansion of $\gamma\delta$ T lymphocytes in an HTLV-I carrier, associated with chronic neutropenia and rheumatoid arthritis. *Ann Hematol* 78:101
139. Halstensen TS, Scott H, Brandtzaeg P (1989) Intraepithelial T cells of the TCR $\gamma\delta$ ⁺ CD8⁻ and V δ 1/J δ 1⁺ phenotypes are increased in coeliac disease. *Scand J Immunol* 30:665
140. Spencer J, Isaacson PG, Diss TC, MacDonald TT (1989) Expression of disulfide-linked and non-disulfide-linked forms of the T cell receptor $\gamma\delta$ heterodimer in human intestinal intraepithelial lymphocytes. *Eur J Immunol* 19:1335
141. Maki M, Holm K, Collin P, Savilahti E (1991) Increase in $\gamma\delta$ T cell receptor bearing lymphocytes in normal small bowel mucosa in latent coeliac disease. *Gut* 32:1412
142. Kutlu T, Brousse N, Rambaud C, Le DF, Schmitz J, Cerf Bensussan N (1993) Numbers of T cell receptor (TCR) $\alpha\beta$ ⁺ but not of TCR $\gamma\delta$ ⁺ intraepithelial lymphocytes correlate with the grade of villous atrophy in coeliac patients on a long term normal diet. *Gut* 34:208
143. Landau SB, Probert CS, Stevens CA, Balk SP, Blumberg RS (1996) Over-utilization of the J δ 3 gene-segment in Crohn's disease. *J Clin Lab Immunol* 48:33
144. Boismenu R, Havran WL (1994) Modulation of epithelial cell growth by intraepithelial $\gamma\delta$ T cells. *Science* 266:1253

145. Soderstrom K, Bucht A, Halapi E, Gronberg A, Magnusson I, Kiessling R (1996) Increased frequency of abnormal $\gamma\delta$ T cells in blood of patients with inflammatory bowel diseases. *J Immunol* 156:2331
146. McVay LD, Li B, Biancaniello R, Creighton MA, Bachwich D, Lichtenstein G, Rombeau JL, Carding SR (1997) Changes in human mucosal $\gamma\delta$ T cell repertoire and function associated with the disease process in inflammatory bowel disease. *Mol Med* 3:183
147. Hoshino T, Ohta A, Nakao M, Ota T, Inokuchi T, Matsueda S, Gouhara R, Yamada A, Itoh K, Oizumi K (1996) TCR $\gamma\delta^+$ T cells in peripheral blood of patients with adult Still's disease. *J Rheumatol* 23:124
148. Olofsson K, Hellstrom S, Hammarstrom M L (1996) Abundance of intraepithelial $\gamma\delta$ T cells in hypertrophic obstructive but not in chronically infected adenoids. *Clin Exp Immunol* 106:396
149. Vincent MS, Roessner K, Lynch D, Wilson D, Cooper SM, Tschopp J, Sigal LH, Budd RC (1996) Apoptosis of Fashigh CD4⁺ synovial T cells by borrelia-reactive Fas-ligand(high) $\gamma\delta$ T cells in Lyme arthritis. *J Exp Med* 184:2109
150. Vecchi M, Crosti L, Berti E, Agape D, Cerri A, De Franchis R (1992) Increased jejunal intraepithelial lymphocytes bearing $\gamma\delta$ T-cell receptor in dermatitis herpetiformis. *Gastroenterology* 102:1499
151. Gerli R, Agea E, Bertotto A, Tognellini R, Flenghi L, Spinozzi F, Velardi A, Grignani F (1991) Analysis of T cells bearing different isotypic forms of the $\gamma\delta$ T cell receptor in patients with systemic autoimmune diseases. *J Rheumatol* 18:1504
152. Volc-Platzer B, Anegg B, Milota S, Pickl W, Fischer G (1993) Accumulation of $\gamma\delta$ T cells in chronic cutaneous lupus erythematosus. *J Invest Dermatol* 100:84S
153. Rajagopalan S, Zordan T, Tsokos GC, Datta SK (1990) Pathogenic anti-DNA autoantibody-inducing T helper cell lines from patients with active lupus nephritis: isolation of CD4⁺8⁻ T helper cell lines that express the $\gamma\delta$ T-cell antigen receptor. *Proc Natl Acad Sci USA* 87:7020
154. Heng MK, Heng MC (1994) Heat-shock protein 65 and activated $\gamma\delta$ T cells in injured arteries. *Lancet* 344:921
155. Kleindienst R, Xu Q, Willeit J, Waldenberger FR, Weimann S, Wick G (1993) Immunology of atherosclerosis. Demonstration of heat shock protein 60 expression and T lymphocytes bearing $\alpha\beta$ or $\gamma\delta$ receptor in human atherosclerotic lesions. *Am J Pathol* 142:1927
156. Burk MR, Carena I, Donda A, Mariani F, Mori L, De Libero G (1997) Functional inactivation in the whole population of human V δ 9/V δ 2 T lymphocytes induced by a nonpeptidic antagonist. *J Exp Med* 185:91
157. Goodier M, Krause JM, Sanni A, Massougboji A, Sadeler BC, Mitchell GH, Modolell M, Eichmann K, Langhorne J (1993) $\gamma\delta$ T cells in the peripheral blood of individuals from an area of holoendemic *Plasmodium falciparum* transmission. *Trans R Soc Trop Med Hyg* 87:692
158. Carena I, Shamshiev A, Donda A, Colonna M, Libero GD (1997) Major histocompatibility complex class I molecules modulate activation threshold and early signaling of T cell antigen receptor- $\gamma\delta$ stimulated by nonpeptidic ligands. *J Exp Med* 186:1769
159. Janeway CA, Jr, Jones B, Hayday A (1988) Specificity and function of T cells bearing $\gamma\delta$ receptors. *Immunol Today* 9:73