




ELSEVIER

Contents lists available at ScienceDirect

European Journal of Pharmacology

journal homepage: www.elsevier.com/locate/ejphar

Full length article

Cardamonin (2',4'-dihydroxy-6'-methoxychalcone) isolated from *Boesenbergia rotunda* (L.) Mansf. inhibits CFA-induced rheumatoid arthritis in rats



Fui-Ling Voon^a, Mohd Roslan Sulaiman^{a,*}, Muhammad Nadeem Akhtar^c, Mohamad Fauzi Idris^a, Ahmad Akira^a, Enoch Kumar Perimal^a, Daud Ahmad Israf^a, Lee Ming-Tatt^{b,*}

^a Department of Biomedical Science, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

^b Department of Pharmaceutical Biology, Faculty of Pharmaceutical Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia

^c Faculty of Industrial Sciences & Technology, University Malaysia Pahang, Gambang, Pahang, Malaysia

ARTICLE INFO

ABSTRACT

Boesenbergia rotunda (L.) Mansf.
Antinociceptive
Anti-inflammatory

cytokines such as tumour necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), and interleukin-6 (IL-6) in vitro. Thus, the possible therapeutic effect of cardamonin in the rheumatoid arthritis (RA) joints is postulated. This study was performed to investigate the anti-arthritis properties of cardamonin in rat model of induced RA, particularly on the inflammatory and pain response of RA. Rheumatoid arthritis paw inflammation was induced by intraplantar (i.pl.) injection of complete Freund's adjuvant (CFA) in *Sprague Dawley* rats. Using four doses of cardamonin (0.625, 1.25, 2.5, and 5.0 mg/kg), anti-arthritis activity was evaluated through the paw edema, mechanical allodynia and thermal hyperalgesia responses. Enzyme-linked immunosorbent assay (ELISA) was carried out to evaluate the plasma level of TNF- α , IL-1 β , and IL-6. Histological slides were prepared from the harvested rat paws to observe the arthritic changes in the joints. Behavioral, biochemical, and histological studies showed that cardamonin demonstrated significant inhibition on RA-induced inflammatory and pain responses as well as progression of joint destruction in rats. ELISA results showed that there was significant inhibition in TNF- α , IL-1 β , and IL-6 levels in plasma of the cardamonin-treated RA rats. Overall, cardamonin possesses potential anti-arthritis properties in CFA-induced RA rat model.

