

PAPER • OPEN ACCESS

Sine-Gordon soliton as a model for Hawking radiation of moving black holes and quantum soliton evaporation

To cite this article: Leone Di Mauro Villari *et al* 2018 *J. Phys. Commun.* **2** 055016

View the [article online](#) for updates and enhancements.

Related content

- [Quantum circuit model for non-inertial objects: a uniformly accelerated mirror](#)
Daiqin Su, C T Marco Ho, Robert B Mann et al.
- [Parity horizons in shape dynamics](#)
Gabriel Herczeg
- [Magnetized Kerr/CFT correspondence](#)
Haryanto M Siahhan



PAPER

Sine-Gordon soliton as a model for Hawking radiation of moving black holes and quantum soliton evaporation

OPEN ACCESS

RECEIVED
9 April 2018REVISED
2 May 2018ACCEPTED FOR PUBLICATION
9 May 2018PUBLISHED
17 May 2018Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Leone Di Mauro Villari^{1,2}, Giulia Marcucci^{2,3} , Maria Chiara Braidotti^{2,4} and Claudio Conti^{2,3} ¹ Institute of Photonics and Quantum Sciences, Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom² Institute for Complex Systems, National Research Council, (ISC-CNR), Via dei Taurini 19, 00185, Rome Italy³ University of Rome La Sapienza, Department of Physics, Piazzale Aldo Moro 5, 00185 Rome, Italy⁴ University of L'Aquila, Department of Physical and Chemical Sciences, Via Veneto 10, I-67010 L'Aquila, ItalyE-mail: giulia.marcucci@uniroma1.it**Keywords:** Hawking Radiation, Sine-Gordon Geometrization, quantum soliton evaporation, black hole emission**Abstract**

The intriguing connection between black holes' evaporation and physics of solitons is opening novel roads to finding observable phenomena. It is known from the inverse scattering transform that velocity is a fundamental parameter in solitons theory. Taking this into account, the study of Hawking radiation by a moving soliton gets a growing relevance. However, a theoretical context for the description of this phenomenon is still lacking. Here, we adopt a soliton geometrization technique to study the quantum emission of a moving soliton in a one-dimensional model. Representing a black hole by the one soliton solution of the Sine-Gordon equation, we consider Hawking emission spectra of a quantized massless scalar field on the soliton-induced metric. We study the relation between the soliton velocity and the black hole temperature. Our results address a new scenario in the detection of new physics in the quantum gravity panorama.

1. Introduction

During the last ten years, analogue gravity systems have attracted major interest in the scientific community [1]. These models aim at providing valuable scenarios to test inaccessible features of quantum gravity, as the Hawking radiation emission by black holes (BHs) [2]. Furthermore, the recent observation of gravitational waves (GWs) emitted by colliding BHs [3, 4] shaded new light and opened unexplored roads towards the search for quantum effects in gravity [5], as the Hawking's BH evaporation [6]. Indeed, quantum BH emission might be observed by the concomitant monitoring of the BH collisions by gravitational and electromagnetic antennas. However, the collision process changes the original Hawking's framework.

Originally, Hawking considered quantum fields in a stationary BH background, the Schwarzschild metric, and discovered that BHs emit thermal radiation and evaporate. His paper appeared exactly one year after a trailblazing article by Ablowitz, Kaup, Newell and Segur (AKNS), that cast new light on nonlinear waves by establishing the general method to solve classes of nonlinear field equations [7, 8]. Surprisingly, AKNS classes generate a metric and define an event horizon (EH). Indeed, it is known in the field of the nonlinear waves that integrable systems, which can be solved exactly by the inverse scattering transform (IST), describe a Riemannian surface with constant negative curvature [9, 10].

Recently, Hawking radiation analogues from solitons were considered in a huge variety of physical contexts, including light [11–15], ultracold gases [16–19], water and sound waves [20, 21]. Here, we study the geometrization of soliton equation by considering a canonical field quantization in the classical background of the Sine-Gordon (SG) soliton metric. Indeed, the 1 + 1 dimensional Sine-Gordon (SG) equation

$$\phi_{tt} - \phi_{xx} + m^2 \sin(\phi) = 0 \quad (1)$$

is a nonlinear model that exhibits a Riemannian surface with constant negative curvature.

In this frame, the SG equation can be considered the AKNS counterpart of a two dimensional gravitational theory. Two dimensional theories of gravity are useful models to understand the quantum properties of higher-dimensional gravity. These theories capture essential features of higher-dimensional counterparts, and in particular have black hole solutions and Hawking radiation [22–25]. The link between the 1 + 1 dimensional gravity and the SG model introduces further simplifications since the quantum properties of this equation have been largely studied [26, 27]. As we shall recall in the next section, the integrability condition of SG equation determines a metric, with a coordinate singularity, which defines an EH. In particular 1 + 1 dimensional BHs can be realized as solitons of the SG equation [28] and it has been shown with a one loop perturbative computation that this BH emits thermal radiation [29, 30].

In this paper, we show that SG soliton emits thermal particles with a specific Hawking temperature, finding the way the temperature changes with the velocity of the SG-BH. Afterward, we perform two different kinds of quantization, one for a massless scalar field and another for the soliton itself, and obtain their Hawking emission spectra. In both cases, we discover that an observer on the soliton tail detects a thermal radiation with a temperature directly proportional to the soliton speed. Furthermore, we analyze the temperature detected by an observer at rest by adding a Doppler effect.

Our paper is organized as follows: in section 2 we review the geometrization of the SG model; we show the connection between a soliton solution of an AKNS system and a metric on a two dimensional surface. In section 3 we study the BH metric induced by the SG equation and introduce suitable coordinate systems for the field quantization. In section 4 we quantize massless scalar fields on the soliton background. In section 5 we quantize the SG soliton following the Faddeev semiclassical quantization [26], and show that the Sine-gordon BH evaporates. Conclusions are drawn in section 6. A short appendix furnishes a minimal mathematical background to forms and curvature.

2. Sine-Gordon geometrization

We start reviewing the way integrable nonlinear equations generates surfaces with constant negative curvature [9]. By considering the SG equation defined in equation (1), we perform the coordinate transformation

$$\chi = \frac{m}{2}(x + t), \quad \theta = \frac{m}{2}(x - t) \quad (2)$$

and get

$$\phi_{\chi\theta} = \sin \phi. \quad (3)$$

As originally stated by Ablowitz, Kaup, Newel and Segur [8], for equation (3) the following system defines the scattering problem

$$\begin{cases} \mathbf{V}_\chi = \hat{L}\mathbf{V} \\ \mathbf{V}_\theta = \hat{M}\mathbf{V} \end{cases}, \quad (4)$$

where \hat{L} and \hat{M} are 2×2 matrices, defining the Lax pair for equation (3). \mathbf{V} is a vector. This system corresponds to the integrable Pfaffian system [31] (see appendix for an introduction to forms and surfaces)

$$d\mathbf{V} = \hat{\Omega}\mathbf{V}, \quad \mathbf{V} = \begin{pmatrix} V_1 \\ V_2 \end{pmatrix}, \quad (5)$$

where $\hat{\Omega}$ is a traceless matrix

$$\hat{\Omega} = \hat{L} d\chi + \hat{M} d\theta = \begin{pmatrix} \omega_1 & \omega_2 \\ \omega_3 & -\omega_1 \end{pmatrix}, \quad (6)$$

with the matrix elements ω_i given by [9]

$$\begin{aligned} \omega_1 &= -\frac{1}{2}\lambda d\chi - \frac{1}{2\lambda} \cos(\phi) d\theta, \\ \omega_2 &= -\frac{1}{2}\phi_\chi d\chi - \frac{1}{2\lambda} \sin(\phi) d\theta, \\ \omega_3 &= \frac{1}{2}\phi_\chi d\chi - \frac{1}{2\lambda} \sin(\phi) d\theta, \end{aligned} \quad (7)$$

where λ is the spectral parameter of the SG scattering problem [8]. Following [10], the arclength of the induced Riemannian surface is written in terms of the matrix elements ω_i as follows [9, 31]

$$\begin{aligned} ds^2 &= (\omega_2 + \omega_3)^2 + (2\omega_1)^2 \\ &= \lambda^2 d\chi^2 + 2 \cos(\phi) d\chi d\theta + \frac{1}{\lambda^2} d\theta^2. \end{aligned} \tag{8}$$

Equation (8) defines the constant negative curvature metric induced by the ISM associated to the SG equation (3). By changing the coordinates set as in the following, we write the first fundamental form ds^2 as [9]

$$ds^2 = \sin^2\left(\frac{\phi}{2}\right) d\tau^2 + \cos^2\left(\frac{\phi}{2}\right) d\xi^2, \tag{9}$$

which results to be associated with a SG equation of the form

$$\phi_{\xi\xi} - \phi_{\tau\tau} = \sin(\phi), \tag{10}$$

where

$$\begin{cases} \xi = \lambda\chi + \lambda^{-1}\theta \\ \tau = \lambda\chi - \lambda^{-1}\theta \end{cases} \tag{11}$$

Thus the metric tensor is

$$\hat{g} = \begin{pmatrix} g_{\tau\tau} & g_{\xi\tau} \\ g_{\tau\xi} & g_{\xi\xi} \end{pmatrix} = \begin{pmatrix} \sin^2\frac{\phi}{2} & 0 \\ 0 & \cos^2\frac{\phi}{2} \end{pmatrix}. \tag{12}$$

However, ds^2 in equation (9) is not Lorentz invariant and it does not lead to a Schwarzschild-like metric. Following [28], in order to obtain a Minkowski-like metric, we perform a Wick rotation $\tau \rightarrow i\tau$ and obtain the elliptic SG (ESG) equation:

$$\phi_{\xi\xi} + \phi_{\tau\tau} = \sin(\phi), \tag{13}$$

whose corresponding metric is

$$ds^2 = -\sin^2\left(\frac{\phi}{2}\right) d\tau^2 + \cos^2\left(\frac{\phi}{2}\right) d\xi^2. \tag{14}$$

3. The Sine-Gordon soliton black hole

We show that the one-soliton solution of the ESG equation determines a BH metric.

The well known forward-propagating one-soliton solution of the equation (10) is

$$\phi(\xi, \tau) = 4 \arctan \{ \exp [\gamma(\xi - \beta_s\tau)] \}, \tag{15}$$

with $\gamma = (1 - \beta_s^2)^{-1/2}$ and $0 < \beta_s < 1$ the soliton velocity [32]. The backward-propagating one-soliton solution gives the same treatise with $-1 < \beta_s < 0$, by substituting β_s in $-\beta_s$ in what follows. For this reason, we can choose solution (15) without loss of generality. Equation (15) is also solution of equation (13) with

$$\gamma = (1 + \beta_s^2)^{-1/2}. \tag{16}$$

We adopt equation (16) hereafter. Substituting equation (15) in equation (14), we have

$$ds^2 = ds_{sol}^2 = -\operatorname{sech}^2(\rho) d\tau^2 + \tanh^2(\rho) d\xi^2, \tag{17}$$

with $\rho = \gamma(\xi - \beta_s\tau)$. Following [28], we adopt various coordinate transformations: first from (τ, ξ) to (\mathcal{T}, ρ) , with ρ as defined above and

$$\mathcal{T} = \tau - \frac{1}{\beta_s} \{ \tanh^{-1}[\gamma^{-1} \tanh(\rho)] - \gamma^{-1}\rho \}. \tag{18}$$

Next, we transform (\mathcal{T}, ρ) to (\mathcal{T}, r) by

$$r = \frac{1}{\gamma} \operatorname{sech}(\rho). \tag{19}$$

The result of the transformation is the line element

$$ds^2 = (\beta_s^2 - r^2) d\mathcal{T}^2 - (\beta_s^2 - r^2)^{-1} dr^2. \tag{20}$$

Equation (20) is the metric of a 1 + 1 dimensional BH with EH at $r_g := \beta_s$. Figures 1 and 2 show the EH positions $\rho_g = \operatorname{arcsech}(\gamma r_g)$ on the soliton profiles and energy densities \mathcal{E} , respectively for different velocities β_s . The energy density, at fixed t , is defined as follows [26]:




Figure 1. The Sine-Gordon soliton at fixed time $\tau = 1$, varying the velocity β_s . The positions of the EHs $\rho_g = m\gamma(\xi_g - \beta_s \tau)$ are in dashed lines.




Figure 2. The soliton energy density at various velocities β_s , at fixed time $\tau = 1$. The positions of the EHs $\rho_g = m\gamma(\xi_g - \beta_s \tau)$ are in dashed lines.

$$\mathcal{E} = \frac{1}{2}(\partial_\xi \phi_s)^2 + [1 - \cos(\phi_s)].$$

It is now convenient to introduce two new sets of coordinates: the modified *Regge-Wheeler coordinate*, that we call the *slug coordinate* in analogy with the *tortoise coordinate*, as usually reported [2, 5], and the *Kruskal-Szekeres coordinates*.

We get the slug coordinate $r^*(r)$ according to

$$dr^* = (\beta_s^2 - r^2)^{-1} dr, \tag{21}$$

so that

$$r^*(r) = \frac{1}{\beta_s} \tanh^{-1} \left(\frac{r}{\beta_s} \right) = \frac{1}{2\beta_s} \ln \left(\frac{\beta_s + r}{\beta_s - r} \right). \tag{22}$$

Equation (20) then becomes

$$ds^2 = [\beta_s^2 - r^2(r^*)][dT^2 - (dr^*)^2]. \tag{23}$$

The slug coordinate is singular at $r = \beta_s$ and it is defined on the exterior of the BH when $\rho \rightarrow \pm\infty$ and $r \rightarrow 0$. In fact, as r approaches β_s , r^* goes to $+\infty$, while far away from the BH $r^* \rightarrow 0$ as $r \rightarrow 0$.

Introducing the slug lightcone coordinates

$$\tilde{u} = T - r^*, \quad \tilde{v} = T + r^*, \tag{24}$$

we write equation (20) as

$$ds^2 = [\beta_s^2 - r^2(\tilde{u}, \tilde{v})] d\tilde{u}d\tilde{v}. \tag{25}$$

The slug lightcone coordinates are singular and they span only the exterior of the black hole. To describe the entire spacetime, we need another coordinate system. In order to be consistent with literature, we refer to them as the *Kruskal-Szekeres (KS) coordinates*. From equations (22) and (24) it follows that

$$\beta_s^2 - r^2 = (\beta_s + r)^2 \exp[\beta_s(\tilde{u} - \tilde{v})]. \tag{26}$$

The BH metric thus becomes

$$ds^2 = [\beta_s + r(\tilde{u}, \tilde{v})]^2 e^{\beta_s(\tilde{u} - \tilde{v})} d\tilde{u}d\tilde{v}. \tag{27}$$

In the KS lightcone coordinates, defined as

$$u = \frac{e^{\beta_s \tilde{u}}}{\beta_s}, \quad v = -\frac{e^{-\beta_s \tilde{v}}}{\beta_s}, \quad (28)$$

Equation (27) takes the form

$$ds^2 = [\beta_s + r(\tilde{u}, \tilde{v})]^2 d\tilde{u}d\tilde{v}, \quad (29)$$

and it is regular at $r = \beta_s$. The singularity occurring in the ESG-soliton metric is, as the Schwarzschild one, a coordinate singularity, which can be removed by a coordinate transformation. The KS coordinates, indeed, span the entire spacetime.

4. Massless scalar field quantization

We consider a field quantization on the classical soliton background metric. We first analyze a massless scalar field with the action

$$S[\phi] = \frac{1}{2} \int g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \sqrt{-g} d^2 \underline{x}, \quad (30)$$

where $g^{\mu\nu}$ represents the inverse of a general metric tensor $g_{\mu\nu}$, g is the determinant of $g_{\mu\nu}$ and $\underline{x} = (x^0, x^1)$. The action in equation (30) is conformally invariant, and in terms of lightcone slug coordinates and lightcone KS coordinates (29) it reads

$$\begin{aligned} S[\phi] &= \int \partial_{\tilde{u}} \phi \partial_{\tilde{v}} \phi d\tilde{u}d\tilde{v}, \\ S[\phi] &= \int \partial_u \phi \partial_v \phi dudv. \end{aligned} \quad (31)$$

We write the solution of the scalar field equation in terms of the lightcone slug coordinates

$$\phi = \tilde{A}(\tilde{u}) + \tilde{B}(\tilde{v}), \quad (32)$$

and in the lightcone KS coordinate as

$$\phi = A(u) + B(v), \quad (33)$$

where A , \tilde{A} and B , \tilde{B} are arbitrary smooth functions. In correspondance of the tail of the soliton, i.e., far away from the EH, the mode expansion of the field is

$$\hat{\phi} = \int_0^\infty \frac{d\Omega}{2\sqrt{\pi\Omega}} [e^{-i\Omega\tilde{u}} \hat{b}_\Omega^- + e^{+i\Omega\tilde{v}} \hat{b}_\Omega^+] + \text{left moving}. \quad (34)$$

In equation (34) the *left moving* part is given by the terms weighted by $e^{\pm i\Omega\tilde{v}}$ in the mode expansion. The vacuum state $|0_B\rangle$, defined by $\hat{b}_\Omega^- |0_B\rangle = 0$, is the *Boulware vacuum* (BV) and does not contain particles for an observer located far from the EH. However, as the slug coordinate is singular at horizon, the BV is also singular at the EH.

To obtain a vacuum state defined over the entire spacetime, we expand the field operator in terms of the KS lightcone coordinates

$$\hat{\phi} = \int_0^\infty \frac{d\omega}{2\sqrt{\pi\omega}} [e^{-i\omega u} \hat{a}_\omega^- + e^{i\omega v} \hat{a}_\omega^+] + \text{left moving}. \quad (35)$$

The creation and annihilation operators \hat{a}_ω^\pm determine the *Kruskal vacuum* (KV) state $\hat{a}_\omega^- |0_K\rangle = 0$. The KV is regular on the horizon and corresponds to true physical vacuum in the presence of the BH.

For a remote observer the KV contains particles. To determine their number density, we follow the original calculations of Hawking and Unruh with the only difference in the definition of the KS coordinates (see chapters 8 and 9 of [5] for details).

We find that the remote observer moving with the soliton tail sees particles with the thermal spectrum

$$\langle \hat{N}_\Omega \rangle = \langle 0_K | \hat{b}_\Omega^+ \hat{b}_\Omega^- | 0_K \rangle = \left[\exp\left(\frac{2\pi\Omega}{\beta_s}\right) - 1 \right]^{-1} \delta(0). \quad (36)$$

If we consider a finite volume quantization we can put $V = \delta(0)$ [5] and we obtain the number density

$$n_\Omega = \left[\exp\left(\frac{2\pi\Omega}{\beta_s}\right) - 1 \right]^{-1}, \quad (37)$$




Figure 3. Spectral radiance for massless fields varying β_s .

corresponding to the temperature

$$T_H = \frac{\beta_s}{2\pi}. \tag{38}$$

In figure 3, we show the radiance $B(\Omega) = \Omega^3 n_\Omega$. We observe that for a static soliton ($\beta_s = 0$) we get $T_H = 0$. This result may appear in contradiction with the Hawking original work, where he considered the emission from a static BH. However the result in equation (38) is coherent with the structure of the metric induced by the SG equation, where the singularity occurs for $r = r_g = \beta_s$ and no emission can be observable for $r_g = 0$. This dependence of the Hawking radiations on the translation velocity is peculiar of soliton dynamics [33] and it is related to the structure of the spectral parameter in the IST [7, 8].

4.1. Hawking temperature in the laboratory frame

Unlike the Schwarzschild BH, the ESG soliton is not static, but translates with velocity β_s . The frequency Ω seen by an observer at rest with respect to the soliton contains a Doppler shift. Letting Ω_s be the frequency emitted by the soliton in (36), the frequency measured by an observed moving with velocity $-\beta_s$ with respect to the soliton, and located at an angle θ_s with respect to the soliton direction is

$$\Omega_o = \frac{1 - \beta_s \cos \theta_s}{\sqrt{1 - \beta_s^2}} \Omega_s. \tag{39}$$

In the collinear case $\theta_s = 0$, and we have

$$\frac{\Omega_o}{\Omega_s} = \sqrt{\frac{1 - \beta_s}{1 + \beta_s}}. \tag{40}$$

The corresponding Hawking temperature is (for small β_s)

$$T_H = \frac{\beta_s}{2\pi} \sqrt{\frac{1 - \beta_s}{1 + \beta_s}} \simeq \frac{\beta_s}{2\pi} (1 - \beta_s). \tag{41}$$

This calculation also applies to a massive bosonic field, as the number density spectrum depends only on the statistics [34]. In the case of a fermionic field the theory is similar, but the number density spectrum follows the Fermi–Dirac statistics [2].

5. Soliton quantization

Previously we studied the BH evaporation following the works of Hawking and Unruh in [2, 35]. Now, we analyze a quantum perturbation of the BH metric given by the classical soliton solution of the ESG equation, and we obtain a BH evaporation without the interaction with a massless scalar field. We start from

$$\phi \simeq \phi_s + \phi_1, \tag{42}$$

where ϕ_s is the classical solution in equation (15) and ϕ_1 represents a weak field perturbation. We consider the conformally invariant action

$$S[\phi] = \int \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + \cos(\phi) \right] \sqrt{-g} \, d^2x, \tag{43}$$

which leads to a field equation

$$g^{\mu\nu} \partial_\mu \partial_\nu \phi + \sin(\phi) = 0. \tag{44}$$

The solutions of equations (10), (13) differ for a Wick rotation. In other words, one passes from the SG soliton to the ESG one by the transformation

$$\tau \rightarrow i\tau, \quad \beta_s \rightarrow -i\beta_s. \quad (45)$$

We perform the inverse Wick rotation, i.e., $\tau \rightarrow -i\tau$, $\beta_s \rightarrow i\beta_s$, passing from the ESG to the SG, and substitute equation (42) into equation (44), hence we obtain

$$g^{\mu\nu}\partial_\mu\partial_\nu\phi_1 + \cos(\phi_s)\phi_1 = 0, \quad (46)$$

where we neglect terms $O(\phi_1^2)$. This equation expresses the interaction between a massive particle and the gravitational field, because the weak quantum field ϕ_1 obeys a *generalized Klein–Gordon* equation with squared mass $\cos(\phi_s)$ depending on the soliton, and thus on the metric. Recalling equation (15), we have

$$g^{\mu\nu}\partial_\mu\partial_\nu\phi_1 + \cos\{4\arctan[\exp(\rho)]\}\phi_1 = 0. \quad (47)$$

For an observer located on the tail of the soliton ($\rho \rightarrow \infty$), the field equation reduces to

$$g^{\mu\nu}\partial_\mu\partial_\nu\phi_1 + \phi_1 = 0, \quad (48)$$

while for an observer on the horizon ($\rho \rightarrow \rho_g$), we have

$$g^{\mu\nu}\partial_\mu\partial_\nu\phi_1 + F(\rho)\phi_1 = 0, \quad (49)$$

with $F(\rho)$ given by

$$F(\rho)|_{\rho \sim \rho_g} \simeq 1 + \frac{5}{2}\gamma^2\beta_s^2 - 5\gamma^2\beta_s^2\sqrt{1 - \gamma^2v^2}(\rho - \rho_g). \quad (50)$$

Equation (50) truncated at the order zero in $\rho - \rho_g$, i.e., exactly on the horizon, leads to

$$F \simeq 1 + \frac{5}{2}\gamma^2\beta_s^2. \quad (51)$$

Due to the inverse Wick rotation, even if the action is conformally invariant, the quantization is not straightforward. We need to adapt both the slug and the KS lightcone coordinates in equations (24), (28) to the rotated system. We obtain

$$\begin{aligned} r^*(r) &= \int_0^r \frac{dr'}{\beta_s^2 + r'^2} = \frac{i}{2\beta_s} \ln\left(\frac{i\beta_s + r}{i\beta_s - r}\right), \\ \tilde{u} &= \mathcal{T} - ir^*, \quad \tilde{v} = \mathcal{T} + ir^*, \\ u &= -\frac{e^{-\beta_s\tilde{u}}}{\beta_s}, \quad v = \frac{e^{\beta_s\tilde{v}}}{\beta_s}. \end{aligned} \quad (52)$$

Since the action (43) is conformally invariant, we thus write the field equation as follows

$$\begin{aligned} \partial_{\tilde{u}}\partial_{\tilde{v}}\phi_1 + \phi_1 &= 0 \quad \text{slug lightcone,} \\ \partial_u\partial_v\phi_1 + F\phi_1 &= 0 \quad \text{K - S lightcone,} \end{aligned} \quad (53)$$

Equations (53) have exponential solution

$$\begin{aligned} \phi_1 &\propto e^{i(K-\Omega_K)\tilde{u} - i(K+\Omega_K)\tilde{v}}, \\ \phi_1 &\propto e^{i(k-\omega_k)u - i(k+\omega_k)v}, \end{aligned} \quad (54)$$

with the following dispersion relations,

$$\begin{aligned} \Omega_K &= \sqrt{K^2 + 1}, \\ \omega_k &= \sqrt{k^2 + F^2}. \end{aligned} \quad (55)$$

From now on, we omit the K and k indices. We write the quantum fields as follows

$$\begin{aligned} \hat{\phi}_0 &= \frac{1}{2\pi} \int_0^\infty \frac{d\Omega}{\sqrt{\Omega}} [\hat{b}_\Omega^- e^{i(K-\Omega)\tilde{u} - i(K+\Omega)\tilde{v}} + \hat{b}_\Omega^+ e^{-i(K-\Omega)\tilde{u} + i(K-\Omega)\tilde{v}}] \\ &= \frac{1}{2\pi} \int_0^\infty \frac{d\omega}{\sqrt{\omega}} [\hat{a}_\omega^- e^{i(k-\omega)u - i(k+\omega)v} + \hat{a}_\omega^+ e^{-i(k-\omega)u + i(k+\omega)v}], \end{aligned} \quad (56)$$

where, as in the non interacting case, the annihilation operators \hat{b}_Ω^- and \hat{a}_ω^- define the Boulware vacuum $|0_B\rangle$ and the Kruskal vacuum $|0_K\rangle$, respectively. The operators \hat{a}_ω^\pm and \hat{b}_Ω^\pm are related by the Bogolyubov transformations

$$\hat{b}_\Omega^- = \int_0^\infty d\omega (\alpha_{\Omega\omega}\hat{a}_\omega^- - \beta_{\Omega\omega}\hat{a}_\omega^+). \quad (57)$$




Figure 4. Spectral radiance for massive fields varying the soliton velocity.

By substituting this in equation (56), we find

$$\frac{1}{\sqrt{\omega}} \int_{-\infty}^{\infty} d\tilde{u}d\tilde{v} e^{i[\Omega(\tilde{u}+\tilde{v})-K(\tilde{u}-\tilde{v})]} e^{-i[\omega(u+v)+k(u-v)]} = \int_0^{\infty} \frac{d\Omega'}{\sqrt{\Omega'}} \alpha_{\Omega'\omega} [2\pi\delta(\Omega - \Omega')]^2, \tag{58}$$

hence we obtain

$$\alpha_{\omega\Omega} = \frac{1}{2\pi V} \sqrt{\frac{\Omega}{\omega}} \int d\tilde{u}d\tilde{v} e^{iu(k-\omega)-iv(k+\omega)} e^{-i\tilde{u}(K-\Omega)+iv(k+\Omega)}. \tag{59}$$

Seemingly for $\beta_{\Omega\omega}$, we have

$$\beta_{\omega\Omega} = -\frac{1}{2\pi V} \sqrt{\frac{\Omega}{\omega}} \int d\tilde{u}d\tilde{v} e^{-i\tilde{u}(k-\omega)+iv(k+\omega)} e^{-i\tilde{u}(K-\Omega)+iv(k+\Omega)}. \tag{60}$$

Using now the KS coordinate (52), after lengthy but straightforward calculations, we find

$$\alpha_{\Omega\omega} = \frac{1}{2\pi V} \frac{\Omega}{\omega} e^{\pi\Omega/\beta_s} e^{iF(\Omega,K,\omega,k)} \Gamma\left[i\frac{\Omega+K}{\beta_s}\right] \Gamma\left[i\frac{\Omega-k}{\beta_s}\right],$$

$$\beta_{\Omega\omega} = \frac{1}{2\pi V} \frac{\Omega}{\omega} e^{-\pi\Omega/\beta_s} e^{iG(\Omega,K,\omega,k)} \Gamma\left[i\frac{\Omega+K}{\beta_s}\right] \Gamma\left[i\frac{\Omega-k}{\beta_s}\right]. \tag{61}$$

It follows that $\alpha_{\Omega\omega}$ and $\beta_{\Omega\omega}$ obey the useful relation

$$|\alpha_{\Omega\omega}|^2 = e^{4\pi\Omega/\beta_s} |\beta_{\Omega\omega}|^2. \tag{62}$$

Therefore we can compute the expectation value of the b -particle number operator $\hat{N}_{\Omega} = \hat{b}_{\Omega}^+ \hat{b}_{\Omega}^-$ in the Kruskal vacuum [5], and obtain the number density

$$n_{\Omega} = \left[\exp\left(\frac{2\Omega}{T_H}\right) - 1 \right]^{-1}. \tag{63}$$

This corresponds to an emitted radiation with twice the frequency with respect to the simple massless case, of which spectral radiance $B(\Omega) = \Omega^3 n_{\Omega}$ is reported in figure 4. We observe that the Hawking temperature is equal to equation (38) for the massless scalar field. This is expected since the surface gravity is the same. For a moving observer with respect to the soliton the Hawking temperature, for small β_s , reads

$$T_H = \frac{\beta_s}{2\pi} \sqrt{\frac{1-\beta_s}{1+\beta_s}} \simeq \frac{\beta_s}{2\pi} (1 - \beta_s). \tag{64}$$

Equation (64) provides the Hawking temperature of soliton evaporation in this toy model.

6. Conclusions

We adopted the geometrization of the ESG model and reported on the connection between the one-soliton solution of the 1 + 1-dimensional elliptic Sine-Gordon equation and a metric with a Schwarzschild-like coordinate singularity. We determined the BH metric and, by suitable coordinate systems, we eliminated the singularity and obtained a regular metric on the EH. We quantized a massless scalar field and found the thermal radiation detected by an observer far away on the BH exterior. We obtained that the temperature is proportional to the soliton velocity. We analyzed the temperature detected by an observer in the laboratory frame, by a Doppler effect. We studied also the quantum soliton evaporation, and found the corresponding spectrum.

Our analysis allows to predict the Hawking radiation for a moving 1 + 1 dimensional BH and shows that the velocity affects the temperature and the corresponding emitted thermal spectrum. In a BH collisional process

one can hence expect a frequency shift of the emitted photon concomitant with the variation of spiraling velocity of the BHs. The resulting chirp of the emitted photons may have a clear and detectable signature in the electromagnetic spectrum. Analogues of these processes may be eventually simulated in the long-range interactions between optical solitons pairs recently observed over astronomical distances [36], or similar optical experiments [37, 38].

Our results may be extended to any metric induced by AKNS systems, hence to many different physical models to conceive experimentally realizable analogues for studying Hawking evaporation of moving black holes.

Acknowledgments

We are pleased to acknowledge Prof. Fabio Biancalana for invaluable discussions and for a critical reading of the manuscript. We also acknowledge I M Deen for technical support with the computational resources. CC and GM acknowledge support from the Templeton Foundation (grant number 58277), the H2020 QuantERA project QUOMPLEX (project ID 731473) and PRIN project NEMO (ref. 2015KEZNYM).

Appendix. Minimal introduction to forms, Pfaff problems and curvature

A 1-form $\Omega = X dx + T dt$ is a combination of the differentials dx and dt , which have to be retained as elements of a basis. X and T can be matrices with the same size, or also operators. A 2-form is a combination of the symbols ('exterior products') $dx \wedge dt$ and $dt \wedge dx = -dx \wedge dt$. One can obtain a 2-form from a 1-form by the differential operator d :

$$d\Omega = \frac{\partial X}{\partial t} dt \wedge dx + \frac{\partial T}{\partial x} dx \wedge dt = \left(-\frac{\partial X}{\partial t} + \frac{\partial T}{\partial x} \right) dx \wedge dt, \quad (65)$$

which can be kept in mind by letting $dx \wedge dx = dt \wedge dt = 0$, so that terms like $\partial_x X$ and $\partial_t T$ do not appear in $d\Omega$.

One can also obtain a 2-form by the exterior product $\Omega \wedge \Omega$ again by $dx \wedge dx = dt \wedge dt = 0$

$$\Omega \wedge \Omega = XT dx \wedge dt + TX dt \wedge dx = [X, T] dx \wedge dt \quad (66)$$

with $[X, T]$ the commutator.

By using forms, the AKNS integrability condition

$$\frac{\partial X}{\partial t} - \frac{\partial T}{\partial x} + [X, T] = 0, \quad (67)$$

reads as

$$d\Omega - \Omega \wedge \Omega = 0. \quad (68)$$

For some authors, using forms has the advantage of a more compact notation as the explicit coordinates x and t do not appear in (68). Equation (68) is referred to a Pfaffian integrability condition, or Pfaff problem.

Forms are directly connected to the curvature of surfaces. If one considers a surface, and a local point vector \mathbf{P} on the surface, let \mathbf{e}_1 and \mathbf{e}_2 the orthogonal tangent vectors. For infinitesimal motion on the surface $d\mathbf{P}$

$$d\mathbf{P} = \sigma^1 \mathbf{e}_1 + \sigma^2 \mathbf{e}_2, \quad (69)$$

where σ^1 and σ^2 contain the differentials of the adopted coordinates and are hence 1-form. $\sigma^1 \wedge \sigma^2$ is the elemental area on the surface. When one moves of an amount $d\mathbf{P}$, $\mathbf{e}_{1,2}$ changes of amounts $d\mathbf{e}_{1,2}$. One considers a surface such that $d\mathbf{e}_1 = \omega \mathbf{e}_2$ and $d\mathbf{e}_2 = -\omega \mathbf{e}_1$ where ω depends on the shape of the surface, contains the differentials of the coordinate systems, and is a 1-form named the *connection one form*. One finds the following equation

$$d\omega = -K \sigma^1 \wedge \sigma^2 \quad (70)$$

where K is the Gaussian curvature. ω, σ^1 and σ^2 are one forms that fix all the properties of the surface. In the particular case $K = -1$, one has from (70)

$$d\omega = \sigma^1 \wedge \sigma^2. \quad (71)$$

By using (71) and considering the matrix 1-form [10]

$$\Omega = \begin{pmatrix} -\frac{1}{2}\sigma^2 & \frac{1}{2}(\omega + \sigma^1) \\ \frac{1}{2}(-\omega + \sigma^1) & \frac{1}{2}\sigma^2 \end{pmatrix}, \quad (72)$$

one finds the Pfaff system in equation (68). In other words, considering the integrability condition (68), and retaining the element of Ω as the forms of a two-dimensional surface, equation (68) implies that the surface has a constant negative curvature $K = -1$. Hence integrability produces pseudospherical surfaces, i.e., surfaces of constant negative curvature.

ORCID iDs

Giulia Marcucci  <https://orcid.org/0000-0001-7166-3643>

Claudio Conti  <https://orcid.org/0000-0003-2583-3415>

References

- [1] Barceló C, Liberati S and Visser M 2011 *Living Rev. Relativ.* **14** 3
- [2] Hawking S W 1974 *Nature* **248** 30
- [3] Abbott B P et al 2016 *Phys. Rev. Lett.* **116** 241103
- [4] Abbott B P et al 2016 *Phys. Rev. Lett.* **118** 221101
- [5] Mukhanov V F and Winitzky S 2007 *Introduction to Quantum Effects in Gravity* (Cambridge: Cambridge University Press)
- [6] Sakalli I and Ovgun A 2016 *Gen. Relativ. Gravit.* **48** 1
Sakalli I and Ovgun A 2015 *EPL* **110** 10008
Sakalli I and Ovgun A 2015 *Eur. Phys. J. Plus* **130** 110
Sakalli I and Ovgun A 2016 *Eur. Phys. J. Plus* **131** 184
- [7] Ablowitz M, Kaup D, Newell A and Segur H 1973 *Phys. Rev. Lett.* **31** 125
- [8] Ablowitz M, Kaup D, Newell A and Segur H 1973 *Phys. Rev. Lett.* **30** 1262
- [9] Bullough R and Caudrey P 1980 *Solitons* (Berlin Heidelberg: Springer-Verlag)
- [10] Sasaki R 1979 *Phys. Lett. A* **71** 390
- [11] Leonhardt U and Philbin T G 2008 arXiv:0803.0669
- [12] Steinhauer J 2014 *Nat. Phys.* **10** 864
- [13] Bermudez D and Leonhardt U 2016 *Phys. Rev. A* **93** 053820
- [14] Tettamanti M, Cacciatori S L, Parola A and Carusotto I 2106 *EPL* **114** 60011
- [15] Di Mauro Villari L, Wright E M, Biancalana F and Conti C 2016 arXiv:1608.04905
- [16] Garay L J, Anglin J R, Cirac J I and Zoller P 2000 *Phys. Rev. Lett.* **85** 4643
- [17] Garay L J 2002 *Int. J. Theor. Phys.* **41** 2073
- [18] Becker C, Stellmer S, Soltan-Panahi P, Dorscher S, Baumert M, Richter E, Kronjager J, Bongs K and Sengstock K 2008 *Nat. Phys.* **4** 496
- [19] Volkoff T J and Fischer U R 2016 *Phys. Rev. D* **94** 024051
- [20] Unruh W G 1981 *Phys. Rev. Lett.* **46** 1351
- [21] Gerace D and Carusotto I 2012 *Phys. Rev. B* **86** 144505
- [22] Giddings S and Strominger A 1993 *Phys. Rev. D* **47** 2454
- [23] Mandal G, Sengupta A M and Wadia S R 1991 *Mod. Phys. Lett. A* **06** 1685
- [24] Witten E 1991 *Phys. Rev. D* **44** 314
- [25] Callan C G, Giddings S B, Harvey J A and Strominger A 1992 *Phys. Rev. D* **45** R1005
- [26] Faddeev L D and Korepin V 1978 *Phys. Rep.* **42** 1
- [27] Zamolodchikov A B and Zamolodchikov A B 1979 *Ann. of Phys.* **120** 253
- [28] Gegenberg J and Kunstatter G 1997 *Phys. Lett. B* **43** 274
- [29] Vaz C and Witten L 1995 *Cl. and Quant. Grav.* **12** 2607
- [30] Kim S and Won T K 1995 *Phys. Lett. B* **361** 38
- [31] Flanders H 1963 *Differential forms* (New York: Dover Publications, INC.)
- [32] Ablowitz M, Kaup D, Newell A and Segur H 1973 *Studies in Applied Mathematics* **53** 246
- [33] Martina L, Pashaev O K and Soliani G 1998 *Phys. Rev. D* **58** 084025
- [34] Crispino L C B, Higuchi A and Matsas G E A 2008 *Rev. Mod. Phys.* **80** 787
- [35] Unruh W G 1976 *Phys. Rev. D* **14** 870
- [36] Jang J K, Erkkanto M, Murdoch S G and Coen S 2013 *Nat. Photon* **7** 657
- [37] Bekenstein R, Schley R, Mutzafi M, Rotschild C and Segev M 2015 *Nat. Phys.* **11** 872
- [38] Roger T, Maitland C, Wilson K, Westerberg N, Vocke D, Wright E and Faccio D 2016 *Nat. Commun.* **7** 13492