

# ΦΥΤΟΝ

REVISTA INTERNACIONAL DE BOTÁNICA EXPERIMENTAL  
INTERNATIONAL JOURNAL OF EXPERIMENTAL BOTANYFUNDACION ROMULO RAGGIO  
Gaspar Campos 861, 1638 Vicente López (BA), Argentina  
[www.revistaphyton.fund-romuloraggio.org.ar](http://www.revistaphyton.fund-romuloraggio.org.ar)

## *In vitro* antifungal/fungistatic activity of manganese phosphite against soybean soil-borne pathogens

Actividad fungicida/fungistática *in vitro* del fosfito de manganeso contra hongos patógenos habitantes del suelo con soja

Carmona MA<sup>1</sup>, E Simonetti<sup>2,3</sup>, ME Ravotti<sup>1</sup>, MM Scandiani<sup>4</sup>, AG Luque<sup>4</sup>, NA Formento<sup>5</sup>, FJ Sautua<sup>1</sup>

**Abstract.** Soybean root and stem rots caused by soil-borne pathogens are diseases commonly found in soybean fields, and one of the most important causes of crop losses. In the present study, the mycelial sensitivity of *Fusarium virguliforme*, *F. tucumaniae*, *Sclerotinia sclerotiorum* and *Macrophomina phaseolina* was evaluated on potato dextrose agar media (25 mL) supplemented with different concentrations of manganese phosphite (MnPhi) diluted in water (0, 25, 37.5, 50, 100, 200, 300, 400, 500, 800 and 1000 µg/mL). Mycelial growth sensitivity was analyzed using logarithmic linear regression analysis. The MnPhi concentration needed to inhibit 50% of the mycelial growth (IC<sub>50</sub>) ranged from 105 µg/mL (*Fusarium* spp.) to 409 µg/mL (*M. phaseolina*). Sclerotia were completely inhibited at 500 µg/mL. The results of our study represent the first report on the direct *in vitro* fungicidal/fungistatic action of MnPhi against fungi that are causal agents of soil-borne diseases.

**Keywords:** *Glycine max*; *Fusarium virguliforme*; *Fusarium tucumaniae*; *Macrophomina phaseolina*; *Sclerotinia sclerotiorum*; Inhibitory concentration.

**Resumen.** Las pudriciones de raíz y tallo (PRYT) en el cultivo de soja causadas por patógenos habitantes del suelo son enfermedades comúnmente encontradas en campos de soja, y son una de las causas más importantes de pérdidas económicas. La sensibilidad micelial de *Fusarium virguliforme*, *Fusarium tucumaniae*, *Sclerotinia sclerotiorum* y *Macrophomina phaseolina* fue evaluada en medio de cultivo agar papa glucosa (25 mL) suplementado con diferentes concentraciones (µg/mL) de fosfito de manganeso (PhiMn) diluido en agua (0; 25; 37,5; 50; 100; 200; 300; 400; 500; 800 y 1000). La sensibilidad del crecimiento micelial fue analizada usando análisis de regresión lineal logarítmico. Se calculó la concentración de PhiMn necesaria para inhibir el 50% del crecimiento micelial (CI<sub>50</sub>). Los valores de CI<sub>50</sub> fueron desde 105 µg/mL (*Fusarium* spp.) hasta 409 µg/mL (*M. phaseolina*). La formación de esclerocios fue completamente inhibida a 500 µg/mL. Los resultados del presente estudio representan el primer reporte de la acción fungicida/fungistática *in vitro* del Phi contra los hongos que causan las PRYT en el cultivo de soja.

**Palabras clave:** *Glycine max*; *Fusarium virguliforme*; *Fusarium tucumaniae*; *Macrophomina phaseolina*; *Sclerotinia sclerotiorum*; Concentración inhibitoria.

<sup>1</sup>Fitopatología, Facultad de Agronomía, Universidad de Buenos Aires, Av. San Martín 4453, C1417DSE, Buenos Aires, Argentina.

<sup>2</sup>Microbiología Agrícola, FAUBA.

<sup>3</sup>Instituto de Investigaciones en Biociencias Agrícolas y Ambientales (CONICET/UBA), FAUBA.

<sup>4</sup>Centro de Referencia de Micología (CEREMIC), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, 2000 Rosario, Argentina.

<sup>5</sup>INTA EEA Paraná, Paraná, Entre Ríos, Argentina.

Address correspondence to: Ing. Agr. M Sc Dr. Marcelo A. Carmona, e-mail: [carmonam@agro.uba.ar](mailto:carmonam@agro.uba.ar)

Received 18.IV.2015. Accepted 17.XII.2015.

## INTRODUCTION

Soybean is the main crop in Argentina, with a planted area of 19.78 million hectares in the 2013/2014 growing season. Root and stem rots caused by soil-borne pathogens are diseases commonly found in soybean fields and one of the most important causes of crop losses (Gupta et al., 2012; Leandro et al., 2012; Walters et al., 2013). In addition to killing plants, root and stem rots can slow down or stop plant growth, causing premature defoliation and decreasing the number and weight of seeds. *Rhizoctonia solani* Kühn, *Fusarium tucumaniae* Aoki, O'Donnell, Homma & Lattanzi, *F. virguliforme* O'Donnell & T. Aoki, *Macrophomina phaseolina* (Tassi) Goidanish, *Pythium* spp., *Phytophthora sojae* (Kaufman & Gerdman) and *Sclerotinia sclerotiorum* (Lib.) De Bary, are among the most important soil-borne pathogens causing root and stem rots, and yield-reducing diseases (Hartman et al., 1999; Scandiani et al., 2010; Ploper et al., 2011; Carmona et al., 2015; Grijalba & Gally, 2015). All these soybean soil-borne pathogens attack many species of cultivated plants, and develop different survival structures such as chlamydospores, microsclerotia, sclerotia and oospores. This is why the common management strategies, such as genetic resistance, seed treatment with fungicides and crop rotation, generally fail to provide adequate control of soil-borne diseases (Dorrance et al., 2009; Gupta et al., 2012; Leandro et al., 2012).

Phosphites ( $\text{PO}_3^{-3}$ ; Phi) are phloem-mobile oxyanions of phosphoric acid ( $\text{H}_3\text{PO}_3$ ) able to elicit systemic acquired resistance in some host species such as apple, potato, and *Arabidopsis* (Percival et al., 2009; Eshraghi et al., 2011 Machinandiarena et al., 2012), and to exhibit direct toxicity against certain pathogens such as *Venturia inaequalis*, *Phytophthora cinnamomi* and *Phytophthora infestans* (Smillie et al., 1989; Guest & Grant, 1991; Wilkinson et al., 2001; Thao & Yamakawa, 2009). Thus, Phi could be an alternative to manage soybean diseases caused by soil-borne pathogens.

The objective of the present study was to evaluate the *in vitro* mycelial sensitivity of root and stem rot pathogens in soybean to manganese phosphite (MnPhi), at different concentrations.

## MATERIALS AND METHODS

The mycelial sensitivity of *F. virguliforme* (11-390-5), *F. tucumaniae* (11-389-3 and 248-11), *S. sclerotiorum* (LARP 56) [all from the Culture Collection of Centro de Referencia de Micología (CEREMIC), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Santa Fe, Argentina] and *M. phaseolina* (INTA Estación Experimental Agropecuaria Salta Culture Collection) were tested in 9-cm plastic Petri dishes. These dishes contained 25 mL of potato dextrose agar media (PDA, Merck) supplemented with different concentrations of MnPhi (ULTRA PLUS Mn -SPRAY-

TEC Fertilizantes Ltda. Brazil), 40% w/w  $\text{P}_2\text{O}_5$  (520 mg  $\text{P}_2\text{O}_5/\text{mL}$ ) diluted in water (0, 25, 37.5, 50, 100, 200, 300, 400, 500, 800 and 1000  $\mu\text{g}/\text{mL}$ ). Manganese phosphite was filtered through 0.22  $\mu\text{m}$  MILLEX®GP filter units (Millipore) and then added to the autoclaved medium. Mycelial discs of each pathogen isolate, 6 mm in diameter, taken from 7-day-old PDA colonies, were placed in the center of each Petri dish supplemented with the different MnPhi concentrations. The plates were sealed with PVC plastic film and incubated in a growth chamber at  $22 \pm 2$  °C, 12 h photoperiod, provided by three fluorescent lamps (Osram daylight 40 W), positioned 50 cm above the plates.

Colony diameters were measured with a precision ruler in two perpendicular directions, when the mycelial growth in the control (0 MnPhi  $\mu\text{g}/\text{mL}$ ) reached the dish edge three days after the inoculation with *S. sclerotiorum*, and six days after the inoculation with *M. phaseolina*. For slow-growing pathogens, such as *Fusarium* species, the diameter of the fungal colony was measured 26 days post-inoculation, even though the mycelium did not reach the dish edge.

The influence of pH on growth inhibition due to MnPhi addition was evaluated for each treatment. The pH of the PDA either supplemented or not with each concentration of MnPhi was measured with a pH meter before the addition of mycelial discs of each pathogen isolate. An additional control treatment was carried out by seeding mycelial discs of all pathogens in PDA acidified up to pH 3 with tartaric acid 10% v/v.

A completely randomized experimental design was used for the *in vitro* tests with three repetitions within each assay. Experimental units were represented by a Petri dish. The assays were repeated three times. The mycelial growth sensitivity bioassays were analyzed using logarithmic linear regression analysis. The MnPhi concentrations needed to inhibit 50% of the mycelial growth ( $\text{IC}_{50}$ ) were calculated from the generated equations. Data from the pH tests were analyzed using ANOVA, and differences between means were tested using Fisher's LSD test. Statistical analyses were carried out using INFostat software (professional version 1.1).

## RESULTS

The results from the *in vitro* bioassay showed that while the concentration of MnPhi increased, the mycelial growth of all pathogens decreased (Fig. 1). The coefficients of determination for the  $\text{IC}_{50}$  equations ranged from 0.79 to 0.98 (Table 1). The highest  $\text{IC}_{50}$  was determined for *M. phaseolina* followed by *S. sclerotiorum*. The *Fusarium* spp. isolates were the most sensitive to MnPhi, with an  $\text{IC}_{50}$  ranging from 105 to 114  $\mu\text{g}/\text{mL}$  (Fig. 1).

In the Petri dish, formation of sclerotia of *S. sclerotiorum* was completely inhibited over 500  $\mu\text{g}/\text{mL}$  and mycelial growth was null at 1000  $\mu\text{g}/\text{mL}$  (Fig. 1). A similar result was




Fig. 1. Effect of manganese phosphite (MnPhi) on *in vitro* mycelial growth of *Sclerotinia sclerotiorum* (LARP 56), *Macrophomina phaseolina* (Salta province), *Fusarium virguliforme* (11-390-5), *Fusarium tucumaniae* (CCC 248-11) and *Fusarium tucumaniae* (11-389-3) using different concentrations.

Fig. 1. Efecto *in vitro* del fosfito de manganeso (PhiMn) sobre el crecimiento micelial de *Sclerotinia sclerotiorum* (LARP 56), *Macrophomina phaseolina* (Provincia de Salta), *Fusarium virguliforme* (11-390-5), *Fusarium tucumaniae* (CCC 248-11) y *Fusarium tucumaniae* (11-389-3), usando diferentes concentraciones.

Table 1. Regression equations, coefficients of determination ( $R^2$ ) and 50% inhibitory concentrations of mycelium growth ( $IC_{50}$ ) of *Macrophomina phaseolina*, *Sclerotinia sclerotiorum*, *Fusarium virguliforme* and *Fusarium tucumaniae*.

Tabla 1. Ecuaciones de regresión, coeficientes de determinación ( $R^2$ ) y concentraciones inhibitoria del 50% del crecimiento micelial ( $CI_{50}$ ) de *Macrophomina phaseolina*, *Sclerotinia sclerotiorum*, *Fusarium virguliforme* y *Fusarium tucumaniae*.

Pathogens	Linear regression equations	n	$R^2$	P values	$IC_{50}$ ( $\mu\text{g/mL}$ )
<i>Macrophomina phaseolina</i> (Salta province)	$y = -70.20 \text{ Ln}(x) + 459.22$	12	0.80	0.0001	409
<i>Sclerotinia sclerotiorum</i> (LARP 56)	$y = -42.08 \text{ Ln}(x) + 280.47$	18	0.79	<0.0001	326
<i>Fusarium virguliforme</i> (11-390-5)	$y = -19.89 \text{ Ln}(x) + 122.19$	15	0.98	<0.0001	114
<i>Fusarium tucumaniae</i> (248-11)	$y = -15.26 \text{ Ln}(x) + 95.51$	15	0.96	<0.0001	111
<i>Fusarium tucumaniae</i> (11-389-3)	$y = -21.57 \text{ Ln}(x) + 130.58$	15	0.94	<0.0001	105

observed for *M. phaseolina*, since a concentration of 800  $\mu\text{g/mL}$  completely inhibited the mycelial growth and a concentration of 500  $\mu\text{g/mL}$  inhibited microsclerotia formation (Fig. 1). For the three isolates of *Fusarium* spp., the mycelial growth was very slow, and even in the control it failed to cover the entire plate at 26 days post-seeding (Fig. 1). Manganese

phosphite concentrations of 500  $\mu\text{g/mL}$  or above completely inhibited the mycelial growth of the three isolates of *Fusarium* spp. (Fig. 1).

The pH ranged from 5.5 (0  $\mu\text{g/mL}$  MnPhi) to 3 (1000  $\mu\text{g/mL}$  MnPhi). The mycelia of *M. phaseolina* and *S. sclerotiorum* growing on media acidified with tartaric acid showed no dif-

ferences ( $P < 0.0001$ ) with the control. By contrast, *Fusarium* spp. were partially affected (up to 30%) ( $P < 0.0001$ ) by acidification, showing total inhibition of mycelial growth at pH 4.

All pathogens were completely inhibited by MnPhi (Fig. 1). The inhibitory effect of MnPhi was more evident on *Fusarium* spp., with the lowest  $IC_{50}$  (105-114  $\mu\text{g/mL}$ ) compared to the other pathogens (326-409  $\mu\text{g/mL}$ ) (Fig. 1, Table 1).

## DISCUSSION

The results obtained are consistent with those reported by Lobato et al. (2010) and Araújo et al. (2010) in relation to *in vitro* inhibition of true fungi such as *Fusarium solani*, *Rhizoctonia solani*, *Streptomyces scabies* and *Colletotrichum gloeosporioides* by Phi. Lobato et al. (2010) found higher  $IC_{50}$  for *F. solani*, being 1280  $\mu\text{g/mL}$  for CaPhi, >3560  $\mu\text{g/mL}$  for KPhi and 680  $\mu\text{g/mL}$  for CuPhi. In the case of oomycetes such as *P. infestans* and *P. plurivora*,  $IC_{50}$  were lower. Dalio et al. (2014) found an  $IC_{50}$  of 34  $\mu\text{g/mL}$  for mycelial growth inhibition and of 2.9  $\mu\text{g/mL}$  for zoospore inhibition of *P. plurivora* by Phi.

The media supplemented with 25 and 1000  $\mu\text{g/mL}$  of MnPhi had a pH of 5 and 3, respectively. At pH values between 3 and 5, *M. phaseolina* and *S. sclerotiorum* were not significantly inhibited (Csöndes et al., 2011). Therefore, the inhibition of mycelial growth observed in the present study is mainly attributed to the effect of MnPhi and not to the acidification of the medium. In contrast, the effect of acidification by MnPhi could partially contribute to the mycelial growth inhibition of *Fusarium* species causing sudden death syndrome (Sanogo & Yang, 2001). According to Lobato et al. (2010), the activity of Phi cannot be attributed to a single reason, and probably would result from a combination of several factors such as the concentration of phosphite anion, nature of cation, acidification of the medium and type of pathogen. In this way and according to Lobato et al. (2010), when acidification is not important, the concentration of phosphite anion plays the main role in inhibiting the growth of pathogens. Most fungi tolerate a wide pH range, but the optimum range of growth is between 5.0 and 6.5. Thus, microorganisms that are in media at pHs above or below the optimum range will have impaired development (Araújo et al., 2010). However, the type of pathogen or fungal species would also be an important influential factor. In this regard, Lobato et al. (2010) determined that the pH could partially contribute to mycelial growth inhibition in other fungal species (up to 30%), including those of the *Fusarium solani* species complex.

There is still no conclusive explanation regarding the biochemical mode of action of Phi against pathogens (Dalio et al., 2014). It might be possible that *in vivo* interactions of Phi act in a direct way (Smilie et al., 1989), reducing fungal growth or generating hyphal disruption (King et al., 2010). We found no studies on the effects of Phi on the formation of sclerotia produced by pathogens.

The results of the present study represent the first report of direct *in vitro* fungicidal/fungistatic action of Phi against soybean fungi, which are causal agents of soil-borne diseases. Further *in vivo* research is needed to assess the impact of Phi applied on the soil or plant to analyze whether the control achieved in this work *in vitro* is also attainable in the plant. In this regard, Simonetti et al. (2015) tested MnPhi alone or in combination with PGPR bacteria to control *M. phaseolina* in soybean under greenhouse conditions; it was the first work to report effective control of *M. phaseolina* (up to 37%) using MnPhi as seed treatment.

In the same way, some preliminary results related to the effect of MnPhi seed treatment on field development of soybean sudden death were reported (Carmona et al., 2013).

## ACKNOWLEDGEMENTS

We thank Guadalupe Mercado Cárdenas for the provision of the *Macrophomina phaseolina* isolate. This work was funded by the grant UBACyT 20020130100604BA Universidad de Buenos Aires, Argentina, and partially by SPRAYTEC Fertilizantes Ltda., Argentina.

## REFERENCES

- Araújo, L., R.M. Valdebenito-Sanhueza & M.J. Stadnik (2010). Avaliação de formulações de fosfito de potássio sobre *Colletotrichum gloeosporioides* *in vitro* e no controle pós-infeccional da mancha foliar de *Glomerella* em macieira. *Tropical Plant Pathology* 35: 054-059.
- Carmona, M., N. Formento, A. Luque, L. Lenzi, M.F. Sautua, E. Simonetti & M. Scandiani (2013). Effect of Mn phosphite seed treatment on development of soybean sudden death syndrome caused by *Fusarium tucumaniae*. World Soybean Research Conference 2013 - Durban South Africa, abstract 227. February 17-22, 2013.
- Carmona, M., M. Gally, P. Grijalba & F. Sautua (2015). Evolución de las enfermedades de la soja en Argentina: pasado, y presente. Aportes de la Fauba al manejo integrado. *Agronomía & Ambiente, Rev. Facultad de Agronomía UBA* 35: 37-52.
- Csöndes, I., A. Cseh, J. Taller & P. Poczai (2011). The effects of temperature and pH on the growth of 45 Hungarian *Macrophomina phaseolina* isolates from sunflower fields in Hungary. *Molecular Biology Reports* 39: 3259-3269.
- Dalio, R.J.D., F. Fleischmann, M. Humez & W. Osswald (2014). Phosphite protects *Fagus sylvatica* seedlings towards *Phytophthora plurivora* via local toxicity, priming and facilitation of pathogen recognition. *Public Library of Science One* 9: e87860. doi:10.1371/journal.pone.0087860
- Dorrance, A.E., A.E. Robertson, S. Cianzo, L.J. Giesler, C.R. Grau, M.A. Draper, A.U. Tenuta & T.R. Anderson (2009). Integrated management strategies for *Phytophthora sojae* combining host resistance and seed treatments. *Plant Disease* 93: 875-882.
- Eshraghi, L., J. Anderson, N. Aryamanesh, B. Shearer, J. McComb, G.E. Hardy & P. O'Brien (2011). Phosphite primed defence responses and enhanced expression of defence genes in *Arabidopsis thaliana* infected with *Phytophthora cinnamomi*. *Plant Pathology* 60: 1086-1095.

- Grijalba, P. & M. Gally (2015). Virulence of *Phytophthora sojae* in the Pampeana Subregion of Argentina from 1998 to 2004. *Journal of Phytopathology* 163: 723-730.
- Guest, D. & B.R. Grant (1991). The complex action of phosphonates as antifungal agents. *Biological Reviews* 66: 159-187.
- Gupta, G.K., S.K. Sharma & R. Ramteke (2012). Biology, Epidemiology and Management of the Pathogenic Fungus *Macrophomina phaseolina* (Tassi) Goid with Special Reference to Charcoal Rot of Soybean [*Glycine max* (L.) Merrill]. *Journal of Phytopathology* 160: 167-180.
- Hartman, G.L., J.B. Sinclair & J.C. Rupe (1999). Compendium of Soybean Diseases, fourth ed. APS Press, The American Phytopathological Society, St. Paul, MN, USA, p. 100.
- King, M., W. Reeve, M.B. Van der Hoek, N. Williams & J. McComb (2010). Defining the phosphite-regulated transcriptome of the plant pathogen *Phytophthora cinnamomi*. *Molecular Genetics and Genomics* 284: 425-43.
- Leandro, L.F., N. Tatalovic & A. Luckew (2012). Soybean Sudden Death Syndrome—advances in knowledge and disease management. *CAB Reviews* 7:053. doi: 10.1079/PAVSNR20127053.
- Lobato, M.C., F.P. Olivieri, G.R. Daleo, A.B. Andreu (2010). Antimicrobial activity of phosphites against different potato pathogens. *Journal of Plant Diseases and Protection* 117: 102-109.
- Machinandiarena, M.F., M.C. Lobato, M.L. Feldman, G.R. Daleo, A.B. Andreu (2012). Potassium phosphite primes defense responses in potato against *Phytophthora infestans*. *Journal of Plant Physiology* 169: 1417-1424.
- Percival, G.C., K. Noviss & I. Haynes (2009). Field evaluation of systemic inducing resistance chemicals at different growth stages for the control of apple (*Venturia inaequalis*) and pear (*Venturia pirina*) scab. *Crop Protection* 28: 629-633.
- Ploper, L.D., R. Muñoz & M. Sillon (2011). Las enfermedades de la soja en Argentina. In: Muñoz R, Sillon M (Eds.) Las enfermedades de la soja y su importancia en los países del Mercosur. 1ª ed. Buenos Aires, Argentina, Editorial Hemisferio Sur SA. pp. 251-273.
- Sanogo, S. & X.B. Yang (2001). Relation of sand content, pH, and potassium and phosphorus nutrition to the development of sudden death syndrome in soybean. *Canadian Journal of Plant Pathology* 23: 174-180.
- Scandiani, M.M., T. Aoki, A.G. Luque, M.A. Carmona & K. O'Donnell (2010). First report of sexual reproduction by the soybean sudden death syndrome pathogen *Fusarium tucumaniae* in nature. *Plant Disease* 94: 1411-1416.
- Simonetti, E., N.P. Viso, M. Montecchia, C. Zilli, K. Balestrasse & M. Carmona (2015). Evaluation of native bacteria and manganese phosphite for alternative control of charcoal root rot of soybean. *Microbiological Research* 180: 40-48.
- Smillie, R., B. Grant & D. Guest (1989). The mode of action of phosphite: Evidence for both direct and indirect modes of action on three *Phytophthora* spp. in plants. *Phytopathology* 79: 921-926.
- Thao, H.T.B. & T. Yamakawa (2009). Phosphite (phosphorous acid): Fungicide, fertilizer or bio-stimulator? *Soil Science and Plant Nutrition* 55: 228-234.
- Wilkinson, C.J., B.L. Shearer, T.J. St. J. Jackson & G.E. Hardy (2001). Variation in sensitivity of Western Australian isolates of *Phytophthora cinnamomi* to phosphite *in vitro*. *Plant Pathology* 50: 83-89.