

A COHOMOLOGICAL PROOF OF PETERSON-KAC'S THEOREM ON CONJUGACY OF CARTAN SUBALGEBRAS FOR AFFINE KAC-MOODY LIE ALGEBRAS

V. CHERNOUSOV, V. EGOROV, P. GILLE, AND A. PIANZOLA

ABSTRACT. This paper deals with the problem of conjugacy of Cartan subalgebras for affine Kac-Moody Lie algebras. Unlike the methods used by Peterson and Kac, our approach is entirely cohomological and geometric. It is deeply rooted on the theory of reductive group schemes developed by Demazure and Grothendieck, and on the work of J. Tits on buildings.

Keywords: Affine Kac-Moody Lie algebra, Conjugacy, Reductive group scheme, Torsor, Laurent polynomials, Non-abelian cohomology.

MSC 2000 17B67, 11E72, 14L30, 14E20.

1. INTRODUCTION

Chevalley's theorem on the conjugacy of split Cartan subalgebras is one of the cornerstones of the theory of simple finite dimensional Lie algebras over a field of characteristic 0. Indeed, this theorem affords the most elegant proof that the root system is an invariant of the Lie algebra.

The analogous result for symmetrizable Kac-Moody Lie algebras is the celebrated theorem of Peterson and Kac [PK] (see also [Kmr] and [MP] for detailed proofs). Beyond the finite dimensional case, by far the most important Kac-Moody Lie algebras are the affine ones. These algebras sit at the “border” of finite dimensional Lie theory and they can in fact be viewed as “finite dimensional” (not over the base field but over a Laurent polynomial ring) in the sense of [SGA3]. This approach begs the question as to whether an SGA-inspired proof of conjugacy exists in the affine case. This paper, which builds in [CGP] and [GP], shows that the answer is yes. More precisely, in [P1] (the untwisted case) and [CGP] (general case) conjugacy is established for loop algebras by purely Galois cohomological methods. The step that is missing is extending this result to the “full” Kac-Moody Lie algebra. The central extension presents of course no difficulties, but the introduction of the derivation does. The present paper addresses this issue thus yielding a new cohomological proof of the conjugacy theorem of Peterson and Kac in the case of affine Kac-Moody Lie algebras.

V. Chernousov was partially supported by the Canada Research Chairs Program and an NSERC research grant.

A. Pianzola wishes to thank NSERC and CONICET for their continuous support.

Even though the conjugacy result is known (due to [PK], as mentioned above), it is relevant to point out that the cohomological methods that we are putting forward do have their advantages. Most significantly:

(a) The group under which conjugacy is achieved in our work has a very transparent structure (given in terms of Laurent polynomial points of simply connected group schemes). This is in contrast to the Kac-Moody groups used in [PK] which, in the twisted affine case, are quite difficult to “see”.

(b) The methods used in [PK] are not suitable for attacking the problem of conjugacy in “higher nullity”, for example for Extended Affine Lie Algebras (with their centres and derivations). This is definitely not the case for the ideas we present in this paper. Indeed, we believe that the methods exhibited herein will eventually lead to a proof of conjugacy for a class of Lie algebras that includes the Extended Affine Lie Algebras, EALAs for short (the conjugacy “downstairs”, namely at the level of the centreless core of the EALA, is still given by [CGP]. The difficulty, just as in the present paper, lies in how to extend the conjugacy from downstairs to “upstairs”; this is to the full EALA where the derivations play an essential role).

2. AFFINE KAC-MOODY LIE ALGEBRAS

Split case. Let \mathfrak{g} be a split simple finite dimensional Lie algebra over an algebraically closed field k of characteristic 0 and let $\mathbf{Aut}(\mathfrak{g})$ be its automorphism group. If $x, y \in \mathfrak{g}$ we denote their product in \mathfrak{g} by $[x, y]$. We also let $R = k[t^{\pm 1}]$, and $L(\mathfrak{g}) = \mathfrak{g} \otimes_k R$. We still denote the Lie product in $L(\mathfrak{g})$ by $[x, y]$ where $x, y \in L(\mathfrak{g})$.

The main object under consideration in this paper is the affine (split or twisted) Kac-Moody Lie algebra \widehat{L} corresponding to \mathfrak{g} . Any split affine Kac-Moody Lie algebra is of the form (see [Kac])

$$\widehat{L} = \mathfrak{g} \otimes_k R \oplus k c \oplus k d.$$

The element c is central and d is a degree derivation for a natural grading of $L(\mathfrak{g})$: if $x \in \mathfrak{g}$ and $p \in \mathbb{Z}$ then

$$[d, x \otimes t^p]_{\widehat{L}} = p x \otimes t^p.$$

If $l_1 = x \otimes t^p$, $l_2 = y \otimes t^q \in L(\mathfrak{g})$ are viewed as elements in \widehat{L} their Lie product is given by

$$[x \otimes t^p, y \otimes t^q]_{\widehat{L}} = [x, y] \otimes t^{p+q} + p \langle x, y \rangle \delta_{0, p+q} \cdot c$$

where $\langle x, y \rangle$ is the Killing form on \mathfrak{g} and $\delta_{0, p+q}$ is Kronecker’s delta.

Twisted case. Let m be a positive integer. Let $S = k[t^{\pm \frac{1}{m}}]$ be the ring of Laurent polynomials in the variable $s = t^{\frac{1}{m}}$ with coefficients in k . Let

$$L(\mathfrak{g})_S = L(\mathfrak{g}) \otimes_R S$$

be the Lie algebra obtained from the R -Lie algebra $L(\mathfrak{g})$ by the base change $R \rightarrow S$. Similarly we define Lie algebras

$$\tilde{L}(\mathfrak{g})_S = L(\mathfrak{g})_S \oplus kc \quad \text{and} \quad \widehat{L}(\mathfrak{g})_S = L(\mathfrak{g})_S \oplus kc \oplus kd.^1$$

Fix a primitive root of unity $\zeta \in k$ of degree m . The R -automorphism $\zeta^\times : S \rightarrow S$ given by $s \mapsto \zeta s$ generates the Galois group $\Gamma = \text{Gal}(S/R)$ which we may identify with the abstract group $\mathbb{Z}/m\mathbb{Z}$ by means of ζ^\times . Note that Γ acts naturally on $\mathbf{Aut}(\mathfrak{g})(S) = \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S)$ and on $L(\mathfrak{g})_S = L(\mathfrak{g}) \otimes_R S$ through the second factor.

Next, let σ be an automorphism of \mathfrak{g} of order m . This gives rise to an S -automorphism of $L(\mathfrak{g})_S$ via $x \otimes s \mapsto \sigma(x) \otimes s$ for $x \in \mathfrak{g}$, $s \in S$. It then easily follows that the assignment

$$\bar{i} \mapsto z_{\bar{i}} = \sigma^{-1} \in \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S)$$

gives rise to a cocycle $z = (z_{\bar{i}}) \in Z^1(\Gamma, \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S))$. This cocycle, in turn, gives rise to a twisted action of Γ on $L(\mathfrak{g})_S$. Applying Galois descent formalism we then obtain the Γ -invariant subalgebra

$$L(\mathfrak{g}, \sigma) := (L(\mathfrak{g})_S)^\Gamma = (L(\mathfrak{g}) \otimes_R S)^\Gamma.$$

This is a ‘‘simple Lie algebra over R ’’ in the sense of [SGA3], which is a twisted form of the ‘‘split simple’’ R -Lie algebra $L(\mathfrak{g}) = \mathfrak{g} \otimes_k R$. Indeed S/R is an étale extension and from properties of Galois descent we have

$$L(\mathfrak{g}, \sigma) \otimes_R S \simeq L(\mathfrak{g})_S = (\mathfrak{g} \otimes_k R) \otimes_R S.$$

Note that $L(\mathfrak{g}, id) = L(\mathfrak{g})$.

For $\bar{i} \in \mathbb{Z}/m\mathbb{Z}$, consider the eigenspace

$$\mathfrak{g}_{\bar{i}} = \{x \in \mathfrak{g} : \sigma(x) = \zeta^i x\}.$$

Simple computations show that

$$L(\mathfrak{g}, \sigma) = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_{\bar{i}} \otimes k[t^{\pm 1}]s^i.$$

Let

$$\tilde{L}(\mathfrak{g}, \sigma) := L(\mathfrak{g}, \sigma) \oplus kc \quad \text{and} \quad \widehat{L}(\mathfrak{g}, \sigma) := L(\mathfrak{g}, \sigma) \oplus kc \oplus kd.$$

We give $\widehat{L}(\mathfrak{g}, \sigma)$ a Lie algebra structure such that c is central element, d is the degree derivation, i.e. if $x \in \mathfrak{g}_{\bar{i}}$ and $p \in \mathbb{Z}$ then

$$(2.0.1) \quad [d, x \otimes t^{\frac{p}{m}}] := px \otimes t^{\frac{p}{m}}$$

and if $y \otimes t^{\frac{q}{m}} \in L(\mathfrak{g}, \sigma)$ we get

$$[x \otimes t^{\frac{p}{m}}, y \otimes t^{\frac{q}{m}}]_{\widehat{L}(\mathfrak{g}, \sigma)} = [x, y] \otimes t^{\frac{p+q}{m}} + p \langle x, y \rangle \delta_{0, p+q} \cdot c,$$

where, as before, $\langle x, y \rangle$ is the Killing form on \mathfrak{g} and $\delta_{0, p+q}$ is Kronecker’s delta.

¹Unlike $L(\mathfrak{g})_S$, these object exist over k but not over S .

2.1. **Remark.** Note that the Lie algebra structure on $\widehat{L}(\mathfrak{g}, \sigma)$ is induced by that of on $\widehat{L}(\mathfrak{g})_S$ if we view $\widehat{L}(\mathfrak{g}, \sigma)$ as a subset of $\widehat{L}(\mathfrak{g})_S$.

2.2. **Remark.** Let $\widehat{\sigma}$ be an automorphism of $\widehat{L}(\mathfrak{g})_S$ such that $\widehat{\sigma}|_{L(\mathfrak{g})_S} = \sigma$, $\widehat{\sigma}(c) = c$, $\widehat{\sigma}(d) = d$. Then $\widehat{L}(\mathfrak{g}, \sigma) = (\widehat{L}(\mathfrak{g})_S)^{\widehat{\sigma}}$.

Realization Theorem. (a) *The Lie algebra $\widehat{L}(\mathfrak{g}, \sigma)$ is an affine Kac-Moody Lie algebra, and every affine Kac-Moody Lie algebra is isomorphic to some $\widehat{L}(\mathfrak{g}, \sigma)$.*

(b) $\widehat{L}(\mathfrak{g}, \sigma) \simeq \widehat{L}(\mathfrak{g}, \sigma')$ where σ' is a diagram automorphism with respect to some Cartan subalgebra of \mathfrak{g} .

Proof. See [Kac, Theorems 7.4, 8.3 and 8.5]. \square

Let $\phi \in \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$. Since $\widetilde{L}(\mathfrak{g})_S$ is the derived subalgebra of $\widehat{L}(\mathfrak{g})_S$ the restriction $\phi|_{\widetilde{L}(\mathfrak{g})_S}$ induces a k -Lie automorphism of $\widetilde{L}(\mathfrak{g})_S$. Furthermore, passing to the quotient $\widetilde{L}(\mathfrak{g})_S/kc \simeq L(\mathfrak{g})_S$ the automorphism $\phi|_{\widetilde{L}(\mathfrak{g})_S}$ induces an automorphism of $L(\mathfrak{g})_S$. This yields a well-defined morphism

$$\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S) \rightarrow \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S).$$

Similar considerations apply to $\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g}, \sigma))$. The aim of the next few sections is to show that these two morphisms are surjective.

3. S -AUTOMORPHISMS OF $L(\mathfrak{g})_S$

In this section we construct a “simple” system of generators of the automorphism group

$$\mathbf{Aut}(\mathfrak{g})(S) = \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S)$$

which can be easily extended to k -automorphisms of $\widehat{L}(\mathfrak{g})_S$. We produce our list of generators based on a well-known fact that the group in question is generated by S -points of the corresponding split simple adjoint algebraic group and automorphisms of the corresponding Dynkin diagram.

More precisely, let \mathbf{G} be the split simple simply connected group over k corresponding to \mathfrak{g} and let $\overline{\mathbf{G}}$ be the corresponding adjoint group. Choose a maximal split k -torus $\mathbf{T} \subset \mathbf{G}$ and denote its image in $\overline{\mathbf{G}}$ by $\overline{\mathbf{T}}$. The Lie algebra of \mathbf{T} is a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$. We fix a Borel subgroup $\mathbf{T} \subset \mathbf{B} \subset \mathbf{G}$.

Let $\Sigma = \Sigma(\mathbf{G}, \mathbf{T})$ be the root system of \mathbf{G} relative to \mathbf{T} . The Borel subgroup \mathbf{B} determines an ordering of Σ , hence the system of simple roots $\Pi = \{\alpha_1, \dots, \alpha_n\}$. Fix a Chevalley basis [St67]

$$\{H_{\alpha_1}, \dots, H_{\alpha_n}, X_{\alpha}, \alpha \in \Sigma\}$$

of \mathfrak{g} corresponding to the pair (\mathbf{T}, \mathbf{B}) . This basis is unique up to signs and automorphisms of \mathfrak{g} which preserve \mathbf{B} and \mathbf{T} (see [St67, §1, Remark 1]).

Since S is a Euclidean ring, by Steinberg [St62] the group $\mathbf{G}(S)$ is generated by the so-called root subgroups $U_\alpha = \langle x_\alpha(u) \mid u \in S \rangle$, where $\alpha \in \Sigma$ and

$$(3.0.1) \quad x_\alpha(u) = \exp(uX_\alpha) = \sum_{n=0}^{\infty} u^n X_\alpha^n / n!$$

We recall also that by [St67, §10, Cor. (b) after Theorem 29], every automorphism σ of the Dynkin diagram $\text{Dyn}(\mathbf{G})$ of \mathbf{G} can be extended to an automorphism of \mathbf{G} (and hence of $\overline{\mathbf{G}}$) and \mathfrak{g} , still denoted by σ , which takes

$$x_\alpha(u) \longrightarrow x_{\sigma(\alpha)}(\varepsilon_\alpha u) \quad \text{and} \quad X_\alpha \longrightarrow \varepsilon_\alpha X_{\sigma(\alpha)}.$$

Here $\varepsilon_\alpha = \pm 1$ and if $\alpha \in \Pi$ then $\varepsilon_\alpha = 1$. Thus we have a natural embedding

$$\text{Aut}(\text{Dyn}(\mathbf{G})) \hookrightarrow \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S).$$

The group $\overline{\mathbf{G}}(S)$ acts by S -automorphisms on $L(\mathfrak{g})_S$ through the adjoint representations $ad : \overline{\mathbf{G}} \rightarrow \mathbf{GL}(L(\mathfrak{g})_S)$ and hence we also have a canonical embedding

$$\overline{\mathbf{G}}(S) \hookrightarrow \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S).$$

As we said before, it is well-known (see [P2] for example) that

$$\text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S) = \overline{\mathbf{G}}(S) \rtimes \text{Aut}(\text{Dyn}(\mathbf{G})).$$

For later use we need one more fact.

3.1. Proposition. *Let $f : \mathbf{G} \rightarrow \overline{\mathbf{G}}$ be the canonical morphism. The group $\overline{\mathbf{G}}(S)$ is generated by the root subgroups $f(U_\alpha)$, $\alpha \in \Sigma$, and $\overline{\mathbf{T}}(S)$.*

Proof. Let $\mathbf{Z} \subset \mathbf{G}$ be the center of \mathbf{G} . The exact sequence

$$1 \longrightarrow \mathbf{Z} \longrightarrow \mathbf{G} \longrightarrow \overline{\mathbf{G}} \longrightarrow 1$$

gives rise to an exact sequence in Galois cohomology

$$f(\mathbf{G}(S)) \hookrightarrow \overline{\mathbf{G}}(S) \longrightarrow \text{Ker}[H^1(S, \mathbf{Z}) \rightarrow H^1(S, \mathbf{G})] \longrightarrow 1.$$

Since $H^1(S, \mathbf{Z}) \rightarrow H^1(S, \mathbf{G})$ factors through

$$H^1(S, \mathbf{Z}) \longrightarrow H^1(S, \mathbf{T}) \longrightarrow H^1(S, \mathbf{G})$$

and since $H^1(S, \mathbf{T}) = 1$ (because $\text{Pic } S = 1$) we obtain

$$(3.1.1) \quad f(\mathbf{G}(S)) \hookrightarrow \overline{\mathbf{G}}(S) \longrightarrow H^1(S, \mathbf{Z}) \longrightarrow 1.$$

Similar considerations applied to

$$1 \longrightarrow \mathbf{Z} \longrightarrow \mathbf{T} \longrightarrow \overline{\mathbf{T}} \longrightarrow 1$$

show that

$$(3.1.2) \quad f(\mathbf{T}(S)) \hookrightarrow \overline{\mathbf{T}}(S) \longrightarrow H^1(S, \mathbf{Z}) \longrightarrow 1.$$

The result now follows from (3.1.1) and (3.1.2). \square

3.2. Corollary. *One has*

$$\text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S) = \langle \text{Aut}(\text{Dyn}(\mathbf{G})), f(U_\alpha), \alpha \in \Sigma, \overline{\mathbf{T}}(S) \rangle.$$

4. k -AUTOMORPHISMS OF $L(\mathfrak{g})_S$

We keep the above notation. Recall that for any algebra \mathfrak{A} over a field k (our main interest is in Lie algebras, but \mathfrak{A} can be arbitrary) the centroid of \mathfrak{A} is

$$\text{Ctrd}(\mathfrak{A}) = \{\chi \in \text{End}_k(\mathfrak{A}) \mid \chi(a \cdot b) = a \cdot \chi(b) = \chi(a) \cdot b \text{ for all } a, b \in \mathfrak{A}\}.$$

It is easy to check that if $\chi_1, \chi_2 \in \text{Ctrd}(\mathfrak{A})$ then both linear operators $\chi_1 \circ \chi_2$ and $\chi_1 + \chi_2$ are contained in $\text{Ctrd}(\mathfrak{A})$ as well. Thus, $\text{Ctrd}(\mathfrak{A})$ is a unital associative subalgebra of $\text{End}_k(\mathfrak{A})$. It is also well-known that the centroid is commutative whenever \mathfrak{A} is perfect (recall that a Lie algebra \mathfrak{A} is called perfect if it coincides with its derived subalgebra).

Example. Consider the k -Lie algebra $\mathfrak{A} = L(\mathfrak{g})_S$. For any $s \in S$ the linear k -operator $\chi_s : L(\mathfrak{g})_S \rightarrow L(\mathfrak{g})_S$ given by $x \rightarrow xs$ satisfy

$$\chi_s([x, y]) = [x, \chi_s(y)] = [\chi_s(x), y],$$

hence $\chi_s \in \text{Ctrd}(L(\mathfrak{g})_S)$. Conversely, it is known (see [ABP, Lemma 4.2]) that every element in $\text{Ctrd}(L(\mathfrak{g})_S)$ is of the form χ_s . Thus,

$$\text{Ctrd}(L(\mathfrak{g})_S) = \{\chi_s \mid s \in S\} \simeq S.$$

4.1. **Proposition.**

$$\begin{aligned} \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S) &\simeq \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S) \rtimes \text{Aut}_k(\text{Ctrd}(L(\mathfrak{g})_S)) \\ &\simeq \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S) \rtimes \text{Aut}_k(S). \end{aligned}$$

Proof. This isomorphism is explicitly described in [BN, Corollary 2.28]. The exact nature of the S -linear automorphisms for our particular S is given in [P2, Proposition 1]). \square \square

4.2. **Corollary.** *One has*

$$\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S) = \langle \text{Aut}_k(S), \text{Aut}(\text{Dyn}(\mathbf{G})), U_\alpha, \alpha \in \Sigma, \overline{\mathbf{T}}(S) \rangle.$$

Proof. This follows from Corollary 3.2 and Proposition 4.1. \square

5. AUTOMORPHISMS OF $\tilde{L}(\mathfrak{g})_S$

We remind the reader that the centre of $\tilde{L}(\mathfrak{g})_S$ is the k -span of c and that $\tilde{L}(\mathfrak{g})_S = L(\mathfrak{g})_S \oplus kc$. Since any automorphism ϕ of $\tilde{L}(\mathfrak{g})_S$ takes the centre into itself we have a natural (projection) mapping

$$\mu : \tilde{L}(\mathfrak{g})_S \rightarrow \tilde{L}(\mathfrak{g})_S / kc \simeq L(\mathfrak{g})_S$$

which induces the mapping

$$\lambda : \text{Aut}_{k\text{-Lie}}(\tilde{L}(\mathfrak{g})_S) \rightarrow \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S)$$

given by $\phi \rightarrow \phi'$ where $\phi'(x) = \mu(\phi(x))$ for all $x \in L(\mathfrak{g})_S$. In the last formula we view x as an element of $\tilde{L}(\mathfrak{g})_S$ through the embedding $L(\mathfrak{g})_S \hookrightarrow \tilde{L}(\mathfrak{g})_S$.

5.1. Remark. It is straightforward to check that ϕ' is indeed an automorphism of $L(\mathfrak{g})_S$.

5.2. Proposition. *The mapping λ is an isomorphism.*

Proof. See [P2, Proposition 4]. \square

In what follows if $\phi \in \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S)$ we denote its (unique) lifting to $\text{Aut}_{k\text{-Lie}}(\tilde{L}(\mathfrak{g})_S)$ by $\tilde{\phi}$.

5.3. Remark. For later use we need an explicit formula for lifts of automorphisms of $L(\mathfrak{g})_S$ induced by some “special” points in $\overline{\mathbf{T}}(S)$ (those which are not in the image of $\mathbf{T}(S) \rightarrow \overline{\mathbf{T}}(S)$). More precisely, choose the decomposition $\overline{\mathbf{T}} \simeq \mathbf{G}_{m,S} \times \cdots \times \mathbf{G}_{m,S}$ such that the canonical embedding $\mathbf{G}_{m,S} \rightarrow \overline{\mathbf{T}}$ into the i -th factor is the cocharacter of $\overline{\mathbf{T}}$ dual to α_i . As usual, we have the decomposition $\overline{\mathbf{T}}(S) \simeq \overline{\mathbf{T}}(k) \times \text{Hom}(\mathbf{G}_m, \overline{\mathbf{T}})$. The second factor in the last decomposition is the cocharacter lattice of $\overline{\mathbf{T}}$ and its elements correspond (under the adjoint action) to the subgroup in $\text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S)$ isomorphic to $\text{Hom}(Q, \mathbb{Z})$ where Q is the corresponding root lattice: if $\phi \in \text{Hom}(Q, \mathbb{Z})$ it induces an S -automorphism of $L(\mathfrak{g})_S$ (still denoted by ϕ) given by

$$X_\alpha \rightarrow X_\alpha \otimes s^{\phi(\alpha)}, \quad H_{\alpha_i} \rightarrow H_{\alpha_i}.$$

It is straightforward to check the mapping $\tilde{\phi} : \tilde{L}(\mathfrak{g})_S \rightarrow \tilde{L}(\mathfrak{g})_S$ given by

$$H_\alpha \rightarrow H_\alpha + \phi(\alpha) \langle X_\alpha, X_{-\alpha} \rangle \cdot c, \quad H_\alpha \otimes s^p \rightarrow H_\alpha \otimes s^p$$

if $p \neq 0$ and

$$X_\alpha \otimes s^p \rightarrow X_\alpha \otimes s^{p+\phi(\alpha)}$$

is an automorphism of $\tilde{L}(\mathfrak{g})_S$, hence it is the (unique) lift of ϕ .

6. AUTOMORPHISMS OF SPLIT AFFINE KAC-MOODY LIE ALGEBRAS

Since $\tilde{L}(\mathfrak{g})_S = [\hat{L}(\mathfrak{g})_S, \hat{L}(\mathfrak{g})_S]$ we have a natural (restriction) mapping

$$\tau : \text{Aut}_{k\text{-Lie}}(\hat{L}(\mathfrak{g})_S) \rightarrow \text{Aut}_{k\text{-Lie}}(\tilde{L}(\mathfrak{g})_S).$$

6.1. Proposition. *The mapping τ is surjective.*

Proof. By Proposition 5.2 and Corollary 4.2 the group $\text{Aut}_{k\text{-Lie}}(\tilde{L}(\mathfrak{g})_S)$ has the distinguished system of generators $\{\tilde{\phi}\}$ where

$$\phi \in \text{Aut}(\text{Dyn}(\mathbf{G})), \overline{\mathbf{T}}(S), \text{Aut}_k(S), f(U_\alpha).$$

We want to construct a mapping $\hat{\phi} : \hat{L}(\mathfrak{g})_S \rightarrow \hat{L}(\mathfrak{g})_S$ which preserves the identity

$$[d, x \otimes t^{\frac{p}{m}}]_{\hat{L}} = p x \otimes t^{\frac{p}{m}}$$

for all $x \in \mathfrak{g}$ and whose restriction to $\tilde{L}(\mathfrak{g})_S$ coincides with $\tilde{\phi}$. These two properties would imply that $\hat{\phi}$ is an automorphism of $\hat{L}(\mathfrak{g})_S$ lifting $\tilde{\phi}$.

If $\phi \in U_\alpha$ is unipotent we define $\hat{\phi}$, as usual, through the exponential map. If $\phi \in \text{Aut}(\text{Dyn}(G))$ we put $\hat{\phi}(d) = d$. If ϕ is as in Remark 5.3 we extend it by $d \rightarrow d - X$ where $X \in \mathfrak{h}$ is the unique element such that $[X, X_\alpha] = \phi(\alpha)X_\alpha$ for all roots $\alpha \in \Sigma$. Note that automorphisms of $L(\mathfrak{g})_S$ given by points in $\overline{\mathbf{T}}(k)$ are in the image of $\mathbf{T}(k) \rightarrow \overline{\mathbf{T}}(k)$ and hence they

are generated by unipotent elements. Lastly, if $\phi \in \text{Aut}_k(S)$ is of the form $s \rightarrow as^{-1}$ where $a \in k^\times$ (resp. $s \rightarrow as$) we extend $\tilde{\phi}$ by $\hat{\phi}(d) = -d$ (resp. $\hat{\phi}(d) = d$). We leave it to the reader to verify that in all cases $\hat{\phi}$ preserves the above identity and hence $\hat{\phi}$ is an automorphism of $\widehat{L}(\mathfrak{g})_S$. \square

6.2. Proposition. *One has $\text{Ker } \tau \simeq V$ where $V = \text{Hom}_k(kd, kc)$.*

Proof. We first embed $V \hookrightarrow \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$. Let $v \in V$. Recall that any element $x \in \widehat{L}(\mathfrak{g})_S$ can be written uniquely in the form $x = x' + ad$ where $x' \in \widetilde{L}(\mathfrak{g})_S$ and $a \in k$. We define $\hat{v} : \widehat{L}(\mathfrak{g})_S \rightarrow \widehat{L}(\mathfrak{g})_S$ by $x \rightarrow x + v(ad)$. One checks that \hat{v} is an automorphism of $\widehat{L}(\mathfrak{g})_S$ and thus the required embedding is given by $v \rightarrow \hat{v}$.

Since $\hat{v}(x') = x'$ for all $x' \in \widetilde{L}$ we have $\hat{v} \in \text{Ker } \tau$. Conversely, let $\psi \in \text{Ker } \tau$. Then $\psi(x) = x$ for all $x \in \widetilde{L}(\mathfrak{g})_S$. We need to show that $\psi(d) = ac + bd$ where $a \in k$. Let $\psi(d) = x' + ac + bd$ where $a, b \in k$ and $x' \in L(\mathfrak{g})_S$. Since $[d, X_\alpha]_{\widehat{L}(\mathfrak{g})_S} = 0$ we get

$$[\psi(d), \psi(X_\alpha)]_{\widehat{L}(\mathfrak{g})_S} = 0.$$

Substituting $\psi(d) = x' + ac + bd$ we obtain

$$[x' + ac + bd, X_\alpha]_{\widehat{L}(\mathfrak{g})_S} = 0$$

or $[x', X_\alpha]_{\widetilde{L}(\mathfrak{g})_S} = 0$. Since this is true for all roots $\alpha \in \Sigma$, the element x' commutes with \mathfrak{g} and this can happen if and only if $x' = 0$.

It remains to show that $b = 1$. To see this we can argue similarly by considering the equality

$$[d, X_\alpha \otimes t^{\frac{1}{m}}]_{\widehat{L}(\mathfrak{g})_S} = X_\alpha \otimes t^{\frac{1}{m}}$$

and applying ψ . \square

6.3. Corollary. *The sequence of groups*

$$(6.3.1) \quad 1 \longrightarrow V \longrightarrow \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S) \xrightarrow{\lambda \circ \tau} \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S) \longrightarrow 1$$

is exact.

7. AUTOMORPHISM GROUP OF TWISTED AFFINE KAC-MOODY LIE ALGEBRAS

We keep the notation introduced in §2. In particular, we fix an integer m and a primitive root of unity $\zeta = \zeta_m \in k$ of degree m . Consider the k -automorphism $\zeta^\times : S \rightarrow S$ such that $s \rightarrow \zeta s$ which we view as a k -automorphism of $L(\mathfrak{g})_S$ through the embedding

$$\text{Aut}_k(S) \hookrightarrow \text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S) \simeq \text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S) \rtimes \text{Aut}_k(S)$$

(see Proposition 4.1). As it is explained in §6 we then get the automorphism $\widehat{\zeta}^\times$ (resp. $\widetilde{\zeta}^\times$) of $\widehat{L}(\mathfrak{g})_S$ (resp. $\widetilde{L}(\mathfrak{g})_S$) given by

$$x \otimes s^i + ac + bd \longrightarrow x \otimes \zeta^i s^i + ac + bd$$

where $a, b \in k$ and $x \in \mathfrak{g}$.

Consider now the abstract group $\Gamma = \mathbb{Z}/m\mathbb{Z}$ (which can be identified with $\text{Gal}(S/R)$ as already explained) and define its action on $\widehat{L}(\mathfrak{g})_S$ (resp. $\widetilde{L}(\mathfrak{g})_S, L(\mathfrak{g})_S$) with the use of $\widehat{\zeta}^\times$ (resp. $\widetilde{\zeta}^\times, \zeta^\times$). More precisely, for every $l \in \widehat{L}(\mathfrak{g})_S$ we let $\bar{i}(l) := (\widehat{\zeta}^\times)^i(l)$. Similarly, we define the action of Γ on $\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$ by

$$\bar{i} : \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S) \longrightarrow \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S), \quad x \rightarrow (\widehat{\zeta}^\times)^i x (\widehat{\zeta}^\times)^{-i}.$$

Therefore, $\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$ can be viewed as a Γ -set. Along the same lines one defines the action of Γ on $\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S)$ and $\text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S)$ with the use of ζ^\times . It is easy to see that Γ acts trivially on the subgroup $V \subset \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$ introduced in Proposition 6.2. Thus, (6.3.1) can be viewed as an exact sequence of Γ -groups.

We next choose an element $\pi \in \text{Aut}(\text{Dyn}(\mathbf{G})) \subset \text{Aut}_k(\mathfrak{g})$ of order m (clearly, m can take value 1, 2 or 3 only). Like before, we have the corresponding automorphism $\hat{\pi}$ of $\widehat{L}(\mathfrak{g})_S$ given by

$$x \otimes s^i + ac + bd \longrightarrow \pi(x) \otimes s^i + ac + bd$$

where $a, b \in k$ and $x \in \mathfrak{g}$.

Note that $\widehat{\zeta}^\times \hat{\pi} = \hat{\pi} \widehat{\zeta}^\times$. It then easily follows that the assignment

$$\bar{1} \rightarrow z_{\bar{1}} = \hat{\pi}^{-1} \in \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$$

gives rise to a cocycle $z = (z_{\bar{i}}) \in Z^1(\Gamma, \text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S))$.

This cocycle, in turn, gives rise to a (new) twisted action of Γ on $\widehat{L}(\mathfrak{g})_S$ and $\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S)$. Analogous considerations (with the use of π) are applied to $\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S)$ and $L(\mathfrak{g})_S$. For future reference note that $\hat{\pi}$ commutes with elements in V , hence the twisted action of Γ on V is still trivial. From now on we view (6.3.1) as an exact sequence of Γ -groups, the action of Γ being the twisted action.

7.1. Remark. As we noticed before the invariant subalgebra

$$\mathcal{L} = L(\mathfrak{g}, \pi) = (L(\mathfrak{g})_S)^\Gamma = ((\mathfrak{g} \otimes_k R) \otimes_R S)^\Gamma$$

is a simple Lie algebra over R , a twisted form of a split Lie algebra $\mathfrak{g} \otimes_k R$. The same cohomological formalism also yields that

$$(7.1.1) \quad \text{Aut}_{R\text{-Lie}}(\mathcal{L}) \simeq (\text{Aut}_{S\text{-Lie}}(L(\mathfrak{g})_S))^\Gamma.$$

7.2. Remark. It is worth mentioning that the canonical embedding

$$\iota : (\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S))^\Gamma \hookrightarrow \text{Aut}_{k\text{-Lie}}((L(\mathfrak{g})_S)^\Gamma) = \text{Aut}_{k\text{-Lie}}(\mathcal{L}) \simeq \text{Aut}_{R\text{-Lie}}(\mathcal{L}) \rtimes \text{Aut}_k(R),$$

where the last isomorphism can be established in the same way as in Proposition 4.1, is not necessary surjective in general case. Indeed, one checks that if $m = 3$ then the k -automorphism of R given by $t \rightarrow t^{-1}$ and viewed as an element of $\text{Aut}_{k\text{-Lie}}(\mathcal{L}) \simeq \text{Aut}_{R\text{-Lie}}(\mathcal{L}) \rtimes \text{Aut}_k(R)$ is not in $\text{Im } \iota$. However (7.1.1) implies that the group $\text{Aut}_{R\text{-Lie}}(\mathcal{L})$ is in the image of ι .

7.3. Remark. The k -Lie algebra $\widehat{\mathcal{L}} = (\widehat{L}(\mathfrak{g})_S)^\Gamma$ is a twisted affine Kac–Moody Lie algebra. Conversely, by the Realization Theorem every twisted affine Kac–Moody Lie algebra can be obtained in such a way.

7.4. Lemma. *One has $H^1(\Gamma, V) = 1$.*

Proof. Since Γ is cyclic of order m acting trivially on $V \simeq k$ it follows that

$$Z^1(\Gamma, V) = \{x \in k \mid mx = 0\} = 0$$

as required. \square

The long exact cohomological sequence associated to (6.3.1) together with Lemma 7.4 imply the following.

7.5. Theorem. *The following sequence*

$$1 \longrightarrow V \longrightarrow (\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S))^\Gamma \xrightarrow{\nu} (\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S))^\Gamma \longrightarrow 1$$

is exact. In particular, the group $\text{Aut}_{R\text{-Lie}}(\mathcal{L})$ is in the image of the canonical mapping

$$\text{Aut}_{k\text{-Lie}}(\widehat{\mathcal{L}}) \longrightarrow \text{Aut}_{k\text{-Lie}}(\mathcal{L}) \simeq \text{Aut}_{R\text{-Lie}}(\mathcal{L}) \rtimes \text{Aut}_k(R).$$

Proof. The first assertion is clear. As for the second one, note that as in Remark 7.2 we have the canonical embedding

$$(\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S))^\Gamma \hookrightarrow \text{Aut}_{k\text{-Lie}}((\widehat{L}(\mathfrak{g})_S)^\Gamma) = \text{Aut}_{k\text{-Lie}}(\widehat{\mathcal{L}})$$

and the commutative diagram

$$\begin{array}{ccc} (\text{Aut}_{k\text{-Lie}}(\widehat{L}(\mathfrak{g})_S))^\Gamma & \xrightarrow{\nu} & (\text{Aut}_{k\text{-Lie}}(L(\mathfrak{g})_S))^\Gamma \\ \downarrow & & \downarrow \\ \text{Aut}_{k\text{-Lie}}(\widehat{\mathcal{L}}) & \longrightarrow & \text{Aut}_{k\text{-Lie}}(\mathcal{L}) \end{array}$$

Then surjectivity of ν and Remark 7.2 yield the result. \square

8. SOME PROPERTIES OF AFFINE KAC-MOODY LIE ALGEBRAS

Henceforth we fix a simple finite dimensional Lie algebra \mathfrak{g} and a (diagram) automorphism σ of finite order m . For brevity, we will write $\widehat{\mathcal{L}}$ and $(\widetilde{\mathcal{L}}, \mathcal{L})$ for $(\widehat{L}(\mathfrak{g}, \sigma)$ and $(\widetilde{L}(\mathfrak{g}, \sigma), L(\mathfrak{g}, \sigma))$ respectively.

For all $l_1, l_2 \in \mathcal{L}$ one has

$$(8.0.1) \quad [l_1, l_2] - [l_1, l_2]_{\widehat{\mathcal{L}}} = ac$$

for some scalar $a \in k$. Using (2.0.1) it is also easy to see that for all $y \in \mathcal{L}$ one has

$$(8.0.2) \quad [d, yt^n]_{\widehat{\mathcal{L}}} = mnyt^n + [d, y]_{\widehat{\mathcal{L}}} t^n$$

8.1. Remark. Recall that \mathcal{L} has a natural R -module structure: If $y = x \otimes t^{\frac{p}{m}} \in \mathcal{L}$ then

$$yt := x \otimes t^{\frac{p}{m}+1} = x \otimes t^{\frac{p+m}{m}} \in \mathcal{L}.$$

Therefore since $[d, y]_{\widehat{\mathcal{L}}}$ is contained in \mathcal{L} the expression $[d, y]_{\widehat{\mathcal{L}}} t^m$ is meaningful.

The infinite dimensional Lie algebra $\widehat{\mathcal{L}}$ admits a unique (up to non-zero scalar) invariant nondegenerate bilinear form (\cdot, \cdot) . Its restriction to $\mathcal{L} \subset \widehat{\mathcal{L}}$ is nondegenerate (see [Kac, 7.5.1 and 8.3.8]) and we have

$$(c, c) = (d, d) = 0, \quad 0 \neq (c, d) = \beta \in k^\times$$

and

$$(c, l) = (d, l) = 0 \quad \text{for all } l \in \mathcal{L}.$$

8.2. Remark. It is known that a nondegenerate invariant bilinear form on $\widehat{\mathcal{L}}$ is unique up to nonzero scalar. We may view $\widehat{\mathcal{L}}$ as a subalgebra in the split Kac-Moody Lie algebra $\widehat{L}(\mathfrak{g})_S$. The last one also admits a nondegenerate invariant bilinear form and it is known that its restriction to $\widehat{\mathcal{L}}$ is nondegenerate. Hence this restriction is proportional to the form (\cdot, \cdot) .

Let $\mathfrak{h}_{\bar{0}}$ be a Cartan subalgebra of the Lie algebra $\mathfrak{g}_{\bar{0}}$.

8.3. Lemma. *The centralizer of $\mathfrak{h}_{\bar{0}}$ in \mathfrak{g} is a Cartan subalgebra \mathfrak{h} of \mathfrak{g} .*

Proof. See [Kac, Lemma 8.1]. \square

The algebra $\mathcal{H} = \mathfrak{h}_{\bar{0}} \oplus kc \oplus kd$ plays the role of Cartan subalgebra for $\widehat{\mathcal{L}}$. With respect to \mathcal{H} our algebra $\widehat{\mathcal{L}}$ admits a root space decomposition. The roots are of two types: anisotropic (real) or isotropic (imaginary). This terminology comes from transferring the form to \mathcal{H}^* and computing the “length” of the roots.

The *core* $\widetilde{\mathcal{L}}$ of $\widehat{\mathcal{L}}$ is the subalgebra generated by all the anisotropic roots. In our case we have $\widetilde{\mathcal{L}} = \mathcal{L} \oplus kc$. The correct way to recover \mathcal{L} inside $\widehat{\mathcal{L}}$ is as its core modulo its centre.²

If $\mathfrak{m} \subset \widehat{\mathcal{L}}$ is an abelian subalgebra and $\alpha \in \mathfrak{m}^* = \text{Hom}(\mathfrak{m}, k)$ we denote the corresponding eigenspace in $\widehat{\mathcal{L}}$ (with respect to the adjoint representation of $\widehat{\mathcal{L}}$) by $\widehat{\mathcal{L}}_\alpha$. Thus,

$$\widehat{\mathcal{L}}_\alpha = \{l \in \widehat{\mathcal{L}} \mid [x, l]_{\widehat{\mathcal{L}}} = \alpha(x)l \text{ for all } x \in \mathfrak{m}\}.$$

The subalgebra \mathfrak{m} is called *diagonalizable* in $\widehat{\mathcal{L}}$ if

$$\widehat{\mathcal{L}} = \bigoplus_{\alpha \in \mathfrak{m}^*} \widehat{\mathcal{L}}_\alpha.$$

Every diagonalizable subalgebra of $\mathfrak{m} \subset \widehat{\mathcal{L}}$ is necessarily abelian. We say that \mathfrak{m} is a maximal (abelian) diagonalizable subalgebra (MAD) if it is not properly contained in a larger diagonalizable subalgebra of $\widehat{\mathcal{L}}$.

²In nullity one the core coincides with the derived algebra, but this is not necessarily true in higher nullities.

8.4. **Remark.** Every MAD of $\widehat{\mathcal{L}}$ contains the center kc of $\widehat{\mathcal{L}}$.

8.5. **Example.** The subalgebra \mathcal{H} is a MAD in $\widehat{\mathcal{L}}$ (see [Kac, Theorem 8.5]).

Our aim is to show that an arbitrary maximal diagonalizable subalgebra $\mathfrak{m} \subset \widehat{\mathcal{L}}$ is conjugate to \mathcal{H} under an element of $\text{Aut}_k(\widehat{\mathcal{L}})$. For future reference we record the following facts:

8.6. **Theorem.** (a) *Every diagonalizable subalgebra in \mathcal{L} is contained in a MAD of \mathcal{L} and all MADs of \mathcal{L} are conjugate. More precisely, let \mathbf{G} be the simple simply connected group scheme over R corresponding to \mathcal{L} . Then for any MAD \mathfrak{m} of \mathcal{L} there exists $g \in \mathbf{G}(R)$ such that $\text{Ad}(g)(\mathfrak{m}) = \mathfrak{h}_{\overline{0}}$.*

(b) *There exists a natural bijection between MADs of $\widetilde{\mathcal{L}}$ and MADs of \mathcal{L} . Every diagonalizable subalgebra in $\widetilde{\mathcal{L}}$ is contained in a MAD of $\widetilde{\mathcal{L}}$. All MADs of $\widetilde{\mathcal{L}}$ are conjugate by elements in $\text{Ad}(\mathbf{G}(R)) \subset \text{Aut}_k(\mathcal{L}) \simeq \text{Aut}_k(\widetilde{\mathcal{L}})$.*

(c) *The image of the canonical map $\text{Aut}_k(\widehat{\mathcal{L}}) \rightarrow \text{Aut}_k(\widetilde{\mathcal{L}}) \simeq \text{Aut}_k(\mathcal{L})$ obtained by restriction to the derived subalgebra $\widetilde{\mathcal{L}}$ contains $\text{Aut}_{R\text{-Lie}}(\mathcal{L})$.*

Proof. (a) From the explicit realization of \mathcal{L} one knows that $\mathfrak{h}_{\overline{0}}$ is a MAD of \mathcal{L} . Now (a) follows from [CGP].

(b) The correspondence follows from the fact that every MAD of $\widetilde{\mathcal{L}}$ contains kc . A MAD $\widetilde{\mathfrak{m}}$ of $\widetilde{\mathcal{L}}$ is necessarily of the form $\mathfrak{m} \oplus kc$ for some MAD \mathfrak{m} of \mathcal{L} and conversely. The canonical map $\text{Aut}_k(\widetilde{\mathcal{L}}) \rightarrow \text{Aut}_k(\mathcal{L})$ is an isomorphism by Proposition 5.2.

(c) This was established in Theorem 7.5. □

8.7. **Lemma.** *If $\mathfrak{m} \subset \widehat{\mathcal{L}}$ is a MAD of $\widehat{\mathcal{L}}$ then $\mathfrak{m} \not\subset \widetilde{\mathcal{L}}$.*

Proof. Assume that $\mathfrak{m} \subset \widetilde{\mathcal{L}}$. By Theorem 8.6 (b), there exists a MAD \mathfrak{m}' of $\widetilde{\mathcal{L}}$ containing \mathfrak{m} . Applying again Theorem 8.6 we may assume that up to conjugation by an element of $\text{Aut}_k(\widetilde{\mathcal{L}})$, in fact of $\widehat{\mathbf{G}}(R)$, we have $\mathfrak{m} \subset \mathfrak{m}' = \mathfrak{h}_{\overline{0}} \oplus kc$. Then \mathfrak{m} is a proper subalgebra of the MAD \mathcal{H} of $\widehat{\mathcal{L}}$ and this contradicts the maximality of \mathfrak{m} . □

In the next three sections we are going to prove some preliminary results related to a subalgebra \widehat{A} of the twisted affine Kac-Moody Lie algebra $\widehat{\mathcal{L}}$ which satisfies the following two conditions:

a) *\widehat{A} is of the form $\widehat{A} = A \oplus kc \oplus kd$, where A is an R -subalgebra of \mathcal{L} such that $A \otimes_R K$ is a semisimple Lie algebra over K where $K = k(t)$ is the fraction field of R .*

b) *The restriction to \widehat{A} of the non-degenerate invariant bilinear form $(-, -)$ of $\widehat{\mathcal{L}}$ is non-degenerate.*

In particular, all these results will be valid for $\widehat{A} = \widehat{\mathcal{L}}$.

9. WEIGHTS OF SEMISIMPLE OPERATORS AND THEIR PROPERTIES

Let $x = x' + d \in \widehat{A}$ where $x' \in A$. It induces a k -linear operator

$$ad(x) : \widehat{A} \rightarrow \widehat{A}, \quad y \rightarrow ad(x)(y) = [x, y]_{\widehat{A}}.$$

We say that x is a k -diagonalizable element of \widehat{A} if \widehat{A} has a k -basis consisting of eigenvectors of $ad(x)$. Throughout we assume that $x' \neq 0$ and that x is k -diagonalizable.

For any scalar $w \in k$ we let

$$\widehat{A}_w = \{ y \in \widehat{A} \mid [x, y]_{\widehat{A}} = wy \}.$$

We say that w is a *weight* (= eigenvalue) of $ad(x)$ if $\widehat{A}_w \neq 0$. More generally, if O is a diagonalizable linear operator of a vector space V over k (of main interest to us are the vector spaces \widehat{A} , $\widetilde{A} = A \oplus kc$, A) and if w is its eigenvalue following standard practice we will denote by $V_w \subset V$ the corresponding eigenspace of O .

9.1. Lemma. (a) *If w is a nonzero weight of $ad(x)$ then $\widehat{A}_w \subset \widetilde{A}$.*

(b) $\widehat{A}_0 = \widetilde{A}_0 \oplus \langle x \rangle$.

Proof. Clearly we have $[\widehat{A}, \widehat{A}] \subset \widetilde{A}$ and this implies $ad(x)(\widetilde{A}) \subset \widetilde{A}$. It then follows that the linear operator $ad(x)|_{\widetilde{A}}$ is k -diagonalizable. Let $\widetilde{A} = \bigoplus \widetilde{A}_{w'}$ where the sum is taken over all weights of $ad(x)|_{\widetilde{A}}$. Since $x \in \widehat{A}_0$ and since $\widehat{A} = \langle x \rangle \oplus \widetilde{A}$ we conclude that

$$\widehat{A} = \langle x, \widetilde{A}_0 \rangle \oplus (\bigoplus_{w' \neq 0} \widetilde{A}_{w'}),$$

so that the result follows. \square

The operator $ad(x)|_{\widetilde{A}}$ maps the center $\langle c \rangle = kc$ of \widetilde{A} into itself, hence it induces a linear operator O_x of $A \simeq \widetilde{A}/kc$ which is also k -diagonalizable. The last isomorphism is induced by a natural (projection) mapping $\lambda : \widetilde{A} \rightarrow A$. If $w \neq 0$ the restriction of λ to \widetilde{A}_w is injective (because \widetilde{A}_w does not contain kc). Since $\widetilde{A} = \bigoplus_w \widetilde{A}_w$ it then follows that

$$\lambda|_{\widetilde{A}_w} : \widetilde{A}_w \longrightarrow A_w$$

is an isomorphism for $w \neq 0$. Thus the three linear operators $ad(x)$, $ad(x)|_{\widetilde{A}}$ and O_x have the same nonzero weights.

9.2. Lemma. *Let w be a weight of O_x and let $n \in \mathbb{Z}$. Then $w + mn$ is also a weight of O_x and $A_{w+mn} = t^n A_w$.*

Proof. Assume $y \in A_w \subset A$, hence $O_x(y) = wy$. Let us show that $yt^n \in A_{w+mn}$. We have

$$(9.2.1) \quad O_x(yt^n) = \lambda(ad(x)(yt^n)) = \lambda([x, yt^n]_{\widehat{A}}).$$

Substituting $x = x' + d$ we get

$$[x, yt^n]_{\widehat{A}} = [x', yt^n]_{\widehat{A}} + [d, yt^n]_{\widehat{A}}$$

Applying (8.0.1) and (8.0.2) we get that the right hand side is equal to

$$[x', y] t^n + ac + [d, y]_{\widehat{A}} t^n + mny t^n$$

where $a \in k$ is some scalar. Substituting this into (9.2.1) we get

$$\begin{aligned} O_x(yt^n) &= \lambda([x', y] t^n + ac + [d, y]_{\widehat{A}} t^n + mny t^n) \\ &= [x', y] t^n + \lambda([d, y]_{\widehat{A}} t^n) + mny t^n \end{aligned}$$

By (8.0.1) there exists $b \in k$ such that

$$[x', y] t^n = ([x', y]_{\widehat{A}} + bc) t^n.$$

Here we view $[x', y] t^n$ as an element in \widehat{A} . Therefore

$$\begin{aligned} O_x(yt^n) &= mny t^n + \lambda([x', y]_{\widehat{A}} + bc) t^n + [d, y]_{\widehat{A}} t^n \\ &= mny t^n + \lambda([x, y]_{\widehat{A}} + bc) t^n. \end{aligned}$$

We now note that by construction $[x, y]_{\widehat{A}} + bc$ is contained in $A \subset \widetilde{A}$. Hence

$$\lambda([x, y]_{\widehat{A}} + bc) t^n = \lambda([x, y]_{\widehat{A}} + bc) t^n = \lambda([x, y]_{\widehat{A}}) t^n.$$

Since $\lambda([x, y]_{\widehat{A}}) = O_x(y) = wy$ we finally get

$$O_x(yt^n) = mny t^n + wy t^n = (w + mn)yt^n.$$

Thus we have showed that $A_w t^n \subset A_{w+nm}$. By symmetry $A_{w+nm} t^{-n} \subset A_w$ and we are done. \square

Our next aim is to show that if w is a weight of $ad(x)$ so is $-w$. We remind the reader that \widehat{A} is equipped with the nondegenerate invariant bilinear form $(-, -)$. Hence for all $y, z \in \widehat{A}$ one has

$$(9.2.2) \quad ([x, y]_{\widehat{A}}, z) = -(y, [x, z]_{\widehat{A}}).$$

9.3. Lemma. *If w is a weight of $ad(x)$ then so is $-w$.*

Proof. If $w = 0$ there is nothing to prove. Assume $w \neq 0$. Consider the root space decomposition

$$\widehat{A} = \bigoplus_{w'} \widehat{A}_{w'}.$$

It suffices to show that for any two weights w_1, w_2 of $ad(x)$ such that $w_1 + w_2 \neq 0$ the subspaces \widehat{A}_{w_1} and \widehat{A}_{w_2} are orthogonal to each other. Indeed, the last implies that if $-w$ were not a weight then every element in \widehat{A}_w would be orthogonal to all elements in \widehat{A} , which is impossible.

Let $y \in \widehat{A}_{w_1}$ and $z \in \widehat{A}_{w_2}$. Applying (9.2.2) we have

$$w_1(y, z) = ([x, y]_{\widehat{A}}, z) = -(y, [x, z]_{\widehat{A}}) = -w_2(y, z).$$

Since $w_1 \neq -w_2$ we conclude $(y, z) = 0$. \square

Now we switch our interest to the operator O_x and its weight subspaces. Recall that according to Lemma 9.2 for every weight w of O_x all elements in the set

$$\{w + mn \mid n \in \mathbb{Z}\}$$

are also weights of O_x . We call this set of weights the w -series and denote it by $\{w\}$ (not to be confused with the set whose only element is w . We hope that this abuse of notation will not create any confusion). Recall also that

$$A_{w+mn} = A_w t^n.$$

9.4. Lemma. *Let w be a weight of O_x and let $A_w R$ be the R -span of A_w in A . Then the natural map $\nu : A_w \otimes_k R \rightarrow A_w R$ given by $l \otimes t^n \mapsto l t^n$ is an isomorphism of k -vector spaces.*

Proof. Clearly, the sum $\sum_n A_{w+mn}$ of vector subspaces A_{w+mn} in A is a direct sum. Hence

$$(9.4.1) \quad A_w R = \sum_n A_w t^n = \sum_n A_{w+mn} = \bigoplus_n A_{w+mn}$$

Fix a k -basis $\{e_i\}$ of A_w . Then $\{e_i \otimes t^j\}$ is a k -basis of $A_w \otimes_k R$. Since

$$\nu(e_i \otimes t^n) = e_i t^n \in A_{w+mn}$$

the injectivity of ν easily follows from (9.4.1). The surjectivity is also obvious. \square

Notation: We will denote the R -span $A_w R$ by $A_{\{w\}}$.

By our construction $A_{\{w\}}$ is an R -submodule of A and

$$(9.4.2) \quad A = \bigoplus_w A_{\{w\}}$$

where the sum is taken over fixed representatives of weight series.

9.5. Corollary. $\dim_k A_w < \infty$.

Proof. Indeed, by the above lemma we have

$$\dim_k A_w = \text{rank}_R (A_w \otimes_k R) = \text{rank}_R A_w R = \text{rank}_R A_{\{w\}} \leq \text{rank}_R A < \infty,$$

as required. \square

9.6. Corollary. *There are finitely many weight series.*

Proof. This follows from the fact that A is a free R -module of finite rank. \square

9.7. Lemma. *Let w_1, w_2 be weights of O_x . Then $[A_{w_1}, A_{w_2}] \subset A_{w_1+w_2}$.*

Proof. This is straightforward to check. \square

10. WEIGHT ZERO SUBSPACE

10.1. **Theorem.** $A_0 \neq 0$.

Proof. Assume that $A_0 = 0$. Then, by Lemma 9.2, $A_{mn} = 0$ for all $n \in \mathbb{Z}$. It follows that for any weight w , any integer n and all $y \in A_w$, $z \in A_{-w+mn}$ we have $[y, z] = 0$. Indeed

$$(10.1.1) \quad [A_w, A_{-w+mn}] \subset A_{w+(-w)+mn} = A_{mn} = 0.$$

For $y \in A$ the operator $ad(y) : A \rightarrow A$ may be viewed as a k -operator or as an R -operator. When we deal with the Killing form $\langle -, - \rangle$ on the R -Lie algebra A we will view $ad(y)$ as an R -operator of A .

10.2. **Lemma.** *Let w_1, w_2 be weights of $ad(x)$ such that $\{w_1\} \neq \{-w_2\}$. Then for any integer n and all $y \in A_{w_1}$ and $z \in A_{w_2+mn}$ we have $\langle y, z \rangle = 0$.*

Proof. Let w be a weight of $ad(x)$. By our condition we have $\{w\} \neq \{w + w_1 + w_2\}$. Since $(ad(y) \circ ad(z))(A_{\{w\}}) \subset A_{\{w+w_1+w_2\}}$, in any R -basis of A corresponding to the decomposition (9.4.2) the operator $ad(y) \circ ad(z)$ has zeroes on the diagonal, hence $\text{Tr}(ad(y) \circ ad(z)) = 0$. \square

10.3. **Lemma.** *Let w be a weight of $ad(x)$, n be an integer and let $y \in A_w$. Assume that $ad(y)$ viewed as an R -operator of A is nilpotent. Then for every $z \in A_{-w+mn}$ we have $\langle y, z \rangle = 0$.*

Proof. Indeed, let l be such that $(ad(y))^l = 0$. Since by (10.1.1), $ad(y)$ and $ad(z)$ are commuting operators we have

$$(ad(y) \circ ad(z))^l = (ad(y))^l \circ (ad(z))^l = 0.$$

Therefore $ad(y) \circ ad(z)$ is nilpotent and this implies its trace is zero. \square

Since the Killing form is nondegenerate, it follows immediately from the above two lemmas that for every nonzero element $y \in A_w$ the operator $ad(y)$ is not nilpotent. Recall that by Lemma 9.7 we have $ad(y)(A_{w'}) \subset A_{w+w'}$. Hence taking into consideration Corollary 9.6 we conclude that there exists a weight w' and a positive integer l such that

$$ad(y)(A_{\{w'\}}) \neq 0, (ad(y) \circ ad(y))(A_{\{w'\}}) \neq 0, \dots, (ad(y))^l(A_{\{w'\}}) \neq 0$$

and $(ad(y))^l(A_{\{w'\}}) \subset A_{\{w'\}}$. We may assume that l is the smallest positive integer satisfying these conditions. Then all consecutive scalars

$$(10.3.1) \quad w', w' + w, w' + 2w, \dots, w' + lw$$

are weights of $ad(x)$, $\{w' + iw\} \neq \{w' + (i+1)w\}$ for $i < l$ and $\{w'\} = \{w' + lw\}$. In particular, we automatically get that lw is an integer (divisible by m) which in turn implies that w is a rational number.

Thus, under our assumption $A_0 = 0$ we have proved that all weights of $ad(x)$ are rational numbers. We now choose (in a unique way) representatives w_1, \dots, w_s of all weight series such that $0 < w_i < m$ and up to renumbering we may assume that

$$0 < w_1 < w_2 < \dots < w_s < m.$$

10.4. Remark. Recall that for any weight w_i , the scalar $-w_i$ is also a weight. Since $0 < -w_i + m < m$ the representative of the weight series $\{-w_i\}$ is $m - w_i$. Then the inequality $m - w_i \geq w_1$ implies $m - w_1 \geq w_i$. Hence out of necessity we have $w_s = m - w_1$.

We now apply the observation (10.3.1) to the weight $w = w_1$. Let $w' = w_i$ be as in (10.3.1). Choose the integer $j \geq 0$ such that $w_i + jw_1, w_i + (j+1)w_1$ are weights and $w_i + jw_1 < m$, but $w_i + (j+1)w_1 \geq m$. We note that since m is not a weight of $ad(x)$ we automatically obtain $w_i + (j+1)w_1 > m$. Furthermore, we have $w_i + jw_1 \leq w_s = m - w_1$ (because $w_i + jw_1$ is a weight of $ad(x)$). This implies

$$m < w_i + (j+1)w_1 \leq w_s + w_1 = m - w_1 + w_1 = m$$

– a contradiction that completes the proof of the theorem. \square

11. A LOWER BOUND OF DIMENSIONS OF MADs IN $\widehat{\mathcal{L}}$

11.1. Theorem. *Let $\mathfrak{m} \subset \widehat{\mathcal{L}}$ be a MAD. Then $\dim \mathfrak{m} \geq 3$.*

By Lemma 8.7, \mathfrak{m} contains an element x of the form $x = x' + d$ where $x' \in \mathcal{L}$ and it also contains c . Since x and c generate a subspace of \mathfrak{m} of dimension 2 the statement of the theorem is equivalent to $\langle x, c \rangle \neq \mathfrak{m}$.

Assume the contrary: $\langle x, c \rangle = \mathfrak{m}$. Since \mathfrak{m} is k -diagonalizable we have the weight space decomposition

$$\widehat{\mathcal{L}} = \bigoplus_{\alpha} \widehat{\mathcal{L}}_{\alpha}$$

where the sum is taken over linear mappings $\alpha \in \mathfrak{m}^* = \text{Hom}(\mathfrak{m}, k)$. To find a contradiction we first make some simple observations about the structure of the corresponding eigenspace $\widehat{\mathcal{L}}_0$.

If $\widehat{\mathcal{L}}_{\alpha} \neq 0$, it easily follows that $\alpha(c) = 0$ (because c is in the center of $\widehat{\mathcal{L}}$). Then α is determined uniquely by the value $w = \alpha(x)$ and so instead of $\widehat{\mathcal{L}}_{\alpha}$ we will write $\widehat{\mathcal{L}}_w$.

Recall that by Theorem 10.1, $\mathcal{L}_0 \neq 0$. Our aim is first to show that \mathcal{L}_0 contains a nonzero element y such that the adjoint operator $ad(y)$ of \mathcal{L} is k -diagonalizable. We will next see that y necessarily commutes with x viewed as an element in $\widehat{\mathcal{L}}$ and that it is k -diagonalizable in $\widehat{\mathcal{L}}$ as well. It then follows that the subspace in $\widehat{\mathcal{L}}$ spanned by c, x and y is a commutative k -diagonalizable subalgebra and this contradicts the fact that \mathfrak{m} is a MAD.

11.2. Lemma. *Let $y \in \mathcal{L}$ be nonzero such that $O_x(y) = 0$. Then $[x, y]_{\widehat{\mathcal{L}}} = 0$.*

Proof. Assume that $[x, y]_{\widehat{\mathcal{L}}} = bc \neq 0$. Then

$$(x, [x, y]_{\widehat{\mathcal{L}}}) = (x, bc) = (x' + d, bc) = (d, bc) = \beta b \neq 0.$$

On the other hand, since the form is invariant we get

$$(x, [x, y]_{\widehat{\mathcal{L}}}) = ([x, x]_{\widehat{\mathcal{L}}}, y) = (0, y) = 0$$

– a contradiction which completes the proof. \square

11.3. Lemma. *Assume that $y \in \mathcal{L}_0$ is nonzero and that the adjoint operator $ad(y)$ of \mathcal{L} is k -diagonalizable. Then $ad(y)$ viewed as an operator of $\widehat{\mathcal{L}}$ is also k -diagonalizable.*

Proof. Choose a k -basis $\{e_i\}$ of \mathcal{L} consisting of eigenvectors of $ad(y)$. Thus we have $[y, e_i] = u_i e_i$ where $u_i \in k$ and hence

$$[y, e_i]_{\widehat{\mathcal{L}}} = u_i e_i + b_i c$$

where $b_i \in k$.

Case 1: Suppose first that $u_i \neq 0$. Let

$$\tilde{e}_i = e_i + \frac{b_i}{u_i} \cdot c \in \widehat{\mathcal{L}}.$$

Then we have

$$[y, \tilde{e}_i]_{\widehat{\mathcal{L}}} = [y, e_i]_{\widehat{\mathcal{L}}} = u_i e_i + b_i c = u_i \tilde{e}_i$$

and therefore \tilde{e}_i is an eigenvector of the operator $ad(y) : \widehat{\mathcal{L}} \rightarrow \widehat{\mathcal{L}}$.

Case 2: Let now $u_i = 0$. Then $[y, e_i]_{\widehat{\mathcal{L}}} = b_i c$ and we claim that $b_i = 0$. Indeed, we have

$$(x, [y, e_i]_{\widehat{\mathcal{L}}}) = ([x, y]_{\widehat{\mathcal{L}}}, e_i) = (0, e_i) = 0$$

and on the other hand

$$(x, [y, e_i]_{\widehat{\mathcal{L}}}) = (x, b_i c) = (x' + d, b_i c) = (d, b_i c) = \beta b_i.$$

It follows that $b_i = 0$ and thus $\tilde{e}_i = e_i$ is an eigenvector of $ad(y)$.

Summarizing, replacing e_i by \tilde{e}_i we see that the set $\{\tilde{e}_i\} \cup \{c, x\}$ is a k -basis of $\widehat{\mathcal{L}}$ consisting of eigenvectors of $ad(y)$. \square

11.4. Proposition. *The subalgebra \mathcal{L}_0 contains an element y such that the operator $ad(y) : \mathcal{L} \rightarrow \mathcal{L}$ is k -diagonalizable.*

Proof. We split the proof in three steps.

Step 1: Assume first that there exists $y \in \mathcal{L}_0$ which as an element in $\mathcal{L}_K = \mathcal{L} \otimes_R K$ is semisimple. We claim that our operator $ad(y)$ is k -diagonalizable. Indeed, choose representatives $w_1 = 0, w_2, \dots, w_l$ of the weight series of $ad(x)$. The sets $\mathcal{L}_{w_1}, \dots, \mathcal{L}_{w_l}$ are vector spaces over k of finite dimension, by Lemma 9.5, and they are stable with respect to $ad(y)$ (because $y \in \mathcal{L}_0$). In each k -vector space \mathcal{L}_{w_i} choose a Jordan basis

$$\{e_{ij}, j = 1, \dots, l_i\}$$

of the operator $ad(y)|_{\mathcal{L}_{w_i}}$. Then the set

$$(11.4.1) \quad \{e_{ij}, i = 1, \dots, l, j = 1, \dots, l_i\}$$

is an R -basis of \mathcal{L} , by Lemma 9.4 and the decomposition given in (9.4.2). It follows that the matrix of the operator $ad(y)$ viewed as a K -operator of $\mathcal{L} \otimes_R K$ is a block diagonal matrix whose blocks corresponds to the matrices of $ad(y)|_{\mathcal{L}_{w_i}}$ in the basis $\{e_{ij}\}$. Hence (11.4.1) is a Jordan basis for $ad(y)$ viewed as an operator on $\mathcal{L} \otimes_R K$. Since y is a semisimple element of $\mathcal{L} \otimes_R K$

all matrices of $ad(y)|_{\mathcal{L}_{w_i}}$ are diagonal and this in turn implies that $ad(y)$ is k -diagonalizable operator of \mathcal{L} .

Step 2: We next consider the case when all elements in \mathcal{L}_0 viewed as elements of the R -algebra \mathcal{L} are nilpotent. Then \mathcal{L}_0 , being finite dimensional, is a nilpotent Lie algebra over k . In particular its center is nontrivial since $\mathcal{L}_0 \neq 0$. Let $u \in \mathcal{L}_0$ be a nonzero central element of \mathcal{L}_0 . For any $z \in \mathcal{L}_0$ the operators $ad(u)$ and $ad(z)$ of \mathcal{L} commute. Then $ad(z) \circ ad(u)$ is nilpotent, hence $\langle u, z \rangle = 0$. Furthermore, by Lemma 10.2 $\langle u, z \rangle = 0$ for any $z \in \mathcal{L}_{w_i}$, $w_i \neq 0$. Thus $u \neq 0$ is in the radical of the Killing form of \mathcal{L} – a contradiction.

Step 3: Assume now that \mathcal{L}_0 contains an element y which as an element of \mathcal{L}_K has nontrivial semisimple part y_s . Let us first show that $y_s \in \mathcal{L}_{\{0\}} \otimes_R K$ and then that $y_s \in \mathcal{L}_0$. By Step 1, the last would complete the proof of the proposition.

By decomposition (9.4.2) applied to $A = \mathcal{L}$ we may write y_s as a sum

$$y_s = y_1 + y_2 + \cdots + y_l$$

where $y_i \in \mathcal{L}_{\{w_i\}} \otimes_R K$. In Step 1 we showed that in an appropriate R -basis (11.4.1) of \mathcal{L} the matrix of $ad(y)$ is block diagonal whose blocks correspond to the Jordan matrices of $ad(y)|_{\mathcal{L}_{w_i}} : \mathcal{L}_{w_i} \rightarrow \mathcal{L}_{w_i}$. It follows that the semisimple part of $ad(y)$ is also a block diagonal matrix whose blocks are semisimple parts of $ad(y)|_{\mathcal{L}_{w_i}}$.

Since \mathcal{L}_K is a semisimple Lie algebra over a perfect field we get that $ad(y_s) = ad(y)_s$. Hence for all weights w_i we have

$$(11.4.2) \quad [y_s, \mathcal{L}_{w_i}] \subset \mathcal{L}_{w_i}.$$

On the other hand, for any $u \in \mathcal{L}_{w_i}$ we have

$$ad(y_s)(u) = [y_1, u] + [y_2, u] + \cdots + [y_l, u].$$

Since $[y_j, u] \in \mathcal{L}_{\{w_i+w_j\}} \otimes_R K$, it follows that $ad(y_s)(u) \in \mathcal{L}_{\{w_i\}}$ if and only if $[y_2, u] = \cdots = [y_l, u] = 0$. Since this is true for all i and all $u \in \mathcal{L}_{w_i}$ and since the kernel of the adjoint representation of \mathcal{L}_K is trivial we obtain $y_2 = \cdots = y_l = 0$. Therefore $y_s \in \mathcal{L}_{\{0\}} \otimes_R K$.

It remains to show that $y_s \in \mathcal{L}_0$. We may write y_s in the form

$$y_s = \frac{1}{g(t)}(u_0 \otimes 1 + u_1 \otimes t + \cdots + u_p \otimes t^p)$$

where $u_0, \dots, u_l \in \mathcal{L}_0$ and $g(t) = g_0 + g_1 t + \cdots + g_n t^n$ is a polynomial with coefficients g_0, \dots, g_n in k with $g_n \neq 0$. The above equality can be rewritten in the form

$$(11.4.3) \quad g_0 y_s + g_1 y_s \otimes t + \cdots + g_n y_s \otimes t^n = u_0 \otimes 1 + \cdots + u_p \otimes t^p.$$

Consider an arbitrary index i and let $u \in \mathcal{L}_{w_i}$. Recall that by (11.4.2) we have

$$ad(y_s)(\mathcal{L}_{w_i}) \subset \mathcal{L}_{w_i}.$$

Applying both sides of (11.4.3) to u and comparing \mathcal{L}_{w_i+mn} -components we conclude that $[g_n y_s, u] = [u_n, u]$. Since this is true for all u and all i and

since the adjoint representation of \mathcal{L}_K has trivial kernel we obtain $g_n y_s = u_n$. Since $g_n \neq 0$ we get $y_s = u_n/g_n \in \mathcal{L}_0$. \square

Now we can easily finish the proof of Theorem 11.1. Suppose the contrary. Then $\dim(\mathfrak{m}) < 3$ and hence by Lemma 8.7 we have $\mathfrak{m} = \langle c, x' + d \rangle$ with $x' \in \mathcal{L}$. Consider the operator O_x on \mathcal{L} . By Theorem 10.1 we have $\mathcal{L}_0 \neq 0$. By Propositions 11.4 and 11.3 there exists a nonzero k -diagonalizable element $y \in \mathcal{L}_0$. Clearly, y is not contained in \mathfrak{m} . Furthermore, by Lemma 11.2, y viewed as an element of $\widehat{\mathcal{L}}$ commutes with \mathfrak{m} and by Lemma 11.3 it is k -diagonalizable in $\widehat{\mathcal{L}}$. It follows that the subspace $\mathfrak{m}_1 = \mathfrak{m} \oplus \langle y \rangle$ is an abelian k -diagonalizable subalgebra of $\widehat{\mathcal{L}}$. But this contradicts maximality of \mathfrak{m} .

12. ALL MADs ARE CONJUGATE

12.1. Theorem. *Let $\widehat{\mathbf{G}}(R)$ be the preimage of $\{Ad(g) : g \in \mathbf{G}(R)\}$ under the canonical map $\text{Aut}_k(\widehat{\mathcal{L}}) \rightarrow \text{Aut}_k(\mathcal{L})$. Then all MADs of $\widehat{\mathcal{L}}$ are conjugate under $\widehat{\mathbf{G}}(R)$ to the subalgebra \mathcal{H} in 8.5.*

Proof. Let \mathfrak{m} be a MAD of $\widehat{\mathcal{L}}$. By Lemma 8.7, $\mathfrak{m} \not\subset \widetilde{\mathcal{L}}$. Fix a vector $x = x' + d \in \mathfrak{m}$ where $x' \in \mathcal{L}$ and let $\mathfrak{m}' = \mathfrak{m} \cap \mathcal{L}$. Thus we have $\mathfrak{m} = \langle x, c, \mathfrak{m}' \rangle$. Note that $\mathfrak{m}' \neq 0$, by Theorem 11.1. Furthermore, since \mathfrak{m}' is k -diagonalizable in \mathcal{L} , without loss of generality we may assume that $\mathfrak{m}' \subset \mathfrak{h}_{\overline{0}}$ given that by Theorem 8.6(b) there exists $g \in \mathbf{G}(R)$ such that $Ad(g)(\mathfrak{m}') \subset \mathfrak{h}_{\overline{0}}$ and that by Theorem 7.5 g has lifting to $\text{Aut}_{k-Lie}(\widehat{\mathcal{L}})$.

Consider the weight space decomposition

$$(12.1.1) \quad \mathcal{L} = \bigoplus_i L_{\alpha_i}$$

with respect to the k -diagonalizable subalgebra \mathfrak{m}' of \mathcal{L} where $\alpha_i \in (\mathfrak{m}')^*$ and as usual

$$L_{\alpha_i} = \{z \in \mathcal{L} \mid [t, z] = \alpha_i(t)z \text{ for all } t \in \mathfrak{m}'\}.$$

12.2. Lemma. *L_{α_i} is invariant with respect to the operator O_x .*

Proof. The k -linear operator O_x commutes with $ad(t)$ for all $t \in \mathfrak{m}'$ (because x and \mathfrak{m}' commute in $\widehat{\mathcal{L}}$), so the result follows. \square

12.3. Lemma. *We have $x' \in L_0$.*

Proof. By our construction \mathfrak{m}' is contained in \mathfrak{h}_0 , hence d commutes with the elements of \mathfrak{m}' . But x also commutes with the elements of \mathfrak{m}' and so does $x' = x - d$. \square

$L_0 = C_{\mathcal{L}}(\mathfrak{m}')$, being the Lie algebra of the reductive group scheme $C_{\mathbf{G}}(\mathfrak{m}')$ (see [CGP]), is of the form $L_0 = \mathfrak{z} \oplus A$ where \mathfrak{z} and A are the Lie algebras of the central torus of $C_{\mathbf{G}}(\mathfrak{m}')$ and its semisimple part respectively. Our next goal is to show that $A = 0$.

Suppose this is not true. To get a contradiction we will show that the subset $\widehat{A} = A \oplus kc \oplus kd \subset \widehat{\mathcal{L}}$ is a subalgebra satisfying conditions a) and b)

stated at the end of §8 and that it is stable with respect to $ad(x)$. This, in turn, will allow us to construct an element $y \in A$ which viewed as an element of $\widehat{\mathcal{L}}$ commutes with x and \mathfrak{m}' and is k -diagonalizable. The last, of course, contradicts the maximality of \mathfrak{m} .

Let \mathbf{H} denote the simple simply connected Chevalley-Demazure algebraic k -group corresponding to \mathfrak{g} . Since \mathbf{G} is split over S we have

$$\mathbf{H}_S = \mathbf{H} \times_k S \simeq \mathbf{G}_S = \mathbf{G} \times_R S.$$

Let $C_{\mathfrak{g}}(\mathfrak{m}') = \mathfrak{t} \oplus \mathfrak{r}$ where \mathfrak{t} is the Lie algebra of the central torus of the reductive k -group $C_{\mathbf{H}}(\mathfrak{m}')$ and \mathfrak{r} is the Lie algebra of its semisimple part. Since centralizers commute with base change, we obtain that

$$\mathfrak{t}_S = \mathfrak{t} \otimes_k S = \mathfrak{z} \otimes_R S = \mathfrak{z}_S, \quad \mathfrak{r}_S = \mathfrak{r} \times_k S = A \otimes_R S = A_S.$$

12.4. Lemma. *We have $ad(d)(A) \subset A$ and in particular \widehat{A} is a subalgebra of $\widehat{\mathcal{L}}$.*

Proof. Since \mathfrak{r} consists of “constant” elements we have $[d, \mathfrak{r}]_{\widehat{\mathcal{L}}(\mathfrak{g})_S} = 0$, and this implies that $[d, \mathfrak{r}_S]_{\widehat{\mathcal{L}}(\mathfrak{g})_S} \subset \mathfrak{r}_S$. Also, viewing \mathcal{L} as a subalgebra of $\widehat{\mathcal{L}}(\mathfrak{g})_S$ we have $[d, \mathcal{L}]_{\widehat{\mathcal{L}}} \subset \mathcal{L}$. Furthermore, S/R is faithfully flat, hence $A = A_S \cap \mathcal{L} = \mathfrak{r}_S \cap \mathcal{L}$. Since both subalgebras \mathfrak{r}_S and \mathcal{L} are stable with respect to $ad(d)$, so is their intersection. \square

12.5. Lemma. *The restriction of the nondegenerate invariant bilinear form (\cdot, \cdot) on $\widehat{\mathcal{L}}$ to L_0 is nondegenerate.*

Proof. We mentioned before that the restriction of (\cdot, \cdot) to \mathcal{L} is nondegenerate. Hence in view of decomposition (12.1.1) it suffices to show that for all $a \in L_0$ and $b \in L_{\alpha_i}$ with $\alpha_i \neq 0$ we have $(a, b) = 0$.

Let $l \in \mathfrak{m}'$ be such that $\alpha_i(l) \neq 0$. Using the invariance of (\cdot, \cdot) we get

$$\alpha_i(l)(a, b) = (a, \alpha_i(l)b) = (a, [l, b]) = ([a, l], b) = 0.$$

Hence $(a, b) = 0$ as required. \square

12.6. Lemma. *The restriction of (\cdot, \cdot) to A is nondegenerate.*

Proof. By lemma(12.5) it is enough to show that \mathfrak{z} and A are orthogonal in $\widehat{\mathcal{L}}$. Moreover, viewing \mathfrak{z} and A as subalgebras of the split affine Kac-Moody Lie algebra $\widehat{\mathcal{L}}(\mathfrak{g})_S$ and using Remark 8.2 we conclude that it suffices to verify that $\mathfrak{z}_S = \mathfrak{t}_S$ and $A_S = \mathfrak{r}_S$ are orthogonal in $\widehat{\mathcal{L}}(\mathfrak{g})_S$.

Let $a \in \mathfrak{t}$ and $b \in \mathfrak{r}$. We know that

$$(at^{\frac{i}{m}}, bt^{\frac{j}{m}}) = \langle a, b \rangle \delta_{i+j, 0}$$

where $\langle \cdot, \cdot \rangle$ is a Killing form of \mathfrak{g} . Since \mathfrak{r} is a semisimple algebra we have $\mathfrak{r} = [\mathfrak{r}, \mathfrak{r}]$. It follows that we can write b in the form $b = \sum [a_i, b_i]$ for some $a_i, b_i \in \mathfrak{r}$. Using the facts that \mathfrak{t} and \mathfrak{r} commute and that the Killing form is invariant we have

$$\langle a, b \rangle = \langle a, \sum [a_i, b_i] \rangle = \sum \langle [a, a_i], b_i \rangle = \sum \langle 0, b_i \rangle = 0.$$

Thus $(at^{\frac{i}{m}}, bt^{\frac{j}{m}}) = 0$. \square

12.7. Lemma. *The k -subspace $A \subset \mathcal{L}$ is invariant with respect to O_x .*

Proof. Let $a \in A$. We need to verify that

$$[x, a]_{\widehat{\mathcal{L}}} \in A \oplus kc \subset \widehat{\mathcal{L}}.$$

But $[d, A]_{\widehat{\mathcal{L}}} \subset A + kc$ by Lemma 12.4. We also have

$$[x', A]_{\widehat{\mathcal{L}}} \subset A \oplus kc$$

(because $x' \in L_0$, by Lemma 12.3, and A viewed as a subalgebra in L_0 is an ideal). Since $x = x' + d$ the result follows. \square

According to Lemma 12.3 we can write $x' = x'_0 + x'_1$ where $x'_0 \in \mathfrak{z}$ and $x'_1 \in A$.

12.8. Lemma. *We have $O_x|_A = O_{x'_1+d}|_A$. In particular, the operator $O_{x'_1+d}|_A$ of A is k -diagonalizable.*

Proof. By Lemma 12.7, we have $O_x(A) \subset A$. Since O_x is k -diagonalizable (as an operator of \mathcal{L}), so is the operator $O_x|_A$ of A . Therefore the last assertion of the lemma follows from the first one.

Let now $a \in A$. Using the fact that x'_0 and a commute in \mathcal{L} we have

$$[x', a]_{\widehat{\mathcal{L}}} = [x'_0, a]_{\widehat{\mathcal{L}}} + [x'_1, a]_{\widehat{\mathcal{L}}} = [x'_1, a]_{\widehat{\mathcal{L}}} + bc$$

for some $b \in k$. Thus $O_x(a) = O_{x'_1+d}(a)$. \square

12.9. Lemma. *The operator $ad(x'_1 + d) : \widehat{A} \rightarrow \widehat{A}$ is k -diagonalizable.*

Proof. Since by Lemma 12.8 $O_{x'_1+d}|_A : A \rightarrow A$ is k -diagonalizable we can apply the same arguments as in Lemma 11.3. \square

Now we can produce the required element y . It follows from Lemma 12.6 that the Lie algebra \widehat{A} satisfies all the conditions stated at the end of Section 8. By Lemma 12.9, $ad(x'_1 + d)$ is k -diagonalizable operator of \widehat{A} . Hence arguing as in Theorem 11.1 we see that there exists a nonzero $y \in A$ such that $[y, x'_1 + d]_{\widehat{\mathcal{L}}} = 0$ and $ad(y)$ is a k -diagonalizable operator on \widehat{A} . Then by Lemma 12.8 we have $O_x(y) = O_{x'_1+d}(y) = 0$ and hence, by Lemma 11.2, x and y commute in $\widehat{\mathcal{L}}$.

According to our plan it remains to show that y is k -diagonalizable in $\widehat{\mathcal{L}}$. To see this we need

12.10. Lemma. *Let $z \in \mathfrak{m}'$. Then $[z, y]_{\widehat{\mathcal{L}}} = 0$.*

Proof. Since $y \in A \subset C_{\mathcal{L}}(\mathfrak{m}')$ we have $[z, y]_{\mathcal{L}} = 0$. Then $[z, y]_{\widehat{\mathcal{L}}} = bc$ for some $b \in k$. It follows

$$0 = (0, y) = ([x, z]_{\widehat{\mathcal{L}}}, y) = (x, [z, y]_{\widehat{\mathcal{L}}}) = (x' + d, bc) = (d, bc) = \beta b.$$

This yields $b = 0$ as desired. \square

12.11. **Proposition.** *The operator $ad(y) : \widehat{\mathcal{L}} \rightarrow \widehat{\mathcal{L}}$ is k -diagonalizable.*

Proof. According to Lemma 11.3, it suffices to prove that $ad(y) : \mathcal{L} \rightarrow \mathcal{L}$ is k -diagonalizable. Since y viewed as an element of A is semisimple it is still semisimple viewed as an element of \mathcal{L} . In particular, the R -operator $ad(y) : \mathcal{L} \rightarrow \mathcal{L}$ is also semisimple.

Recall that we have the decomposition of \mathcal{L} into the direct sum of the weight spaces with respect to O_x :

$$\mathcal{L} = \bigoplus_w \mathcal{L}_w = \bigoplus_i \bigoplus_n \mathcal{L}_{w_i+mn} = \bigoplus_i \mathcal{L}_{\{w_i\}}.$$

Since y and x commute in $\widehat{\mathcal{L}}$, for all weights w we have $ad(y)(\mathcal{L}_w) \subset \mathcal{L}_w$. If we choose any k -basis of \mathcal{L}_w it is still an R -basis of $\mathcal{L}_{\{w\}} = \mathcal{L}_w \otimes_k R$ and in this basis the R -operator $ad(y)|_{\mathcal{L}_{\{w\}}}$ and the k -operator $ad(y)|_{\mathcal{L}_w}$ have the same matrices. Since the R -operator $ad(y)|_{\mathcal{L}_{\{w\}}}$ is semisimple, so is $ad(y)|_{\mathcal{L}_w}$, i.e. $ad(y)|_{\mathcal{L}_w}$ is a k -diagonalizable operator. Thus $ad(y) : \mathcal{L} \rightarrow \mathcal{L}$ is k -diagonalizable. \square

Summarizing, assuming $A \neq 0$ we have constructed the k -diagonalizable element

$$y \notin \mathfrak{m} = \langle \mathfrak{m}', x, c \rangle$$

in $\widehat{\mathcal{L}}$ which commutes with \mathfrak{m}' and x in $\widehat{\mathcal{L}}$. Then the subalgebra $\langle \mathfrak{m}, y \rangle$ in $\widehat{\mathcal{L}}$ is commutative and k -diagonalizable which is impossible since \mathfrak{m} is a MAD. Thus A is necessarily trivial and this implies $C_{\mathcal{L}}(\mathfrak{m}')$ is the Lie algebra of the R -torus $C_{\mathfrak{G}}(\mathfrak{m}')$, in particular $C_{\mathcal{L}}(\mathfrak{m}')$ is abelian.

Note that $x' \in C_{\mathcal{L}}(\mathfrak{m}')$, by Lemma 12.3, and that $\mathfrak{h}_{\bar{0}} \subset C_{\mathcal{L}}(\mathfrak{m}')$ (because $\mathfrak{m}' \subset \mathfrak{h}_{\bar{0}}$, by construction). Since $C_{\mathcal{L}}(\mathfrak{m}')$ is abelian and since $x = x' + d$ it follows that $ad(x)(\mathfrak{h}_{\bar{0}}) = 0$. Hence $\langle \mathfrak{h}_{\bar{0}}, x, c \rangle$ is a commutative k -diagonalizable subalgebra in $\widehat{\mathcal{L}}$. But it contains our MAD \mathfrak{m} . Therefore $\mathfrak{m} = \langle \mathfrak{h}_{\bar{0}}, x, c \rangle$. To finish the proof of Theorem 12.1 it now suffices to show that $x' \in \mathfrak{h}_{\bar{0}}$. For that, in turn, we may view x' as an element of $L(\mathfrak{g})_S$ and it suffices to show that $x' \in \mathfrak{h}$ because $\mathfrak{h} \cap \mathcal{L} = \mathfrak{h}_{\bar{0}}$.

12.12. **Lemma.** $x' \in \mathfrak{h}$.

Proof. Consider the root space decomposition of \mathfrak{g} with respect to the Cartan subalgebra \mathfrak{h} :

$$\mathfrak{g} = \mathfrak{h} \oplus \left(\bigoplus_{\alpha \neq 0} \mathfrak{g}_{\alpha} \right).$$

Every k -subspace \mathfrak{g}_{α} has dimension 1. Choose a nonzero elements $X_{\alpha} \in \mathfrak{g}_{\alpha}$. It follows from $\mathfrak{m}' = \mathfrak{h}_{\bar{0}}$ that $C_{L(\mathfrak{g})_S}(\mathfrak{m}') = \mathfrak{h}_S$. Thus $x' \in \mathfrak{h}_S$. Then $\mathfrak{g}_{\alpha} \otimes_k S$ is stable with respect to $ad(x')$ and clearly it is stable with respect to $ad(d)$. Hence it is also stable with respect to O_x .

Arguing as in Lemma 9.2 one can easily see that the operator O_x , viewed as an operator of $L(\mathfrak{g})_S$, is k -diagonalizable. Since $\mathfrak{g}_{\alpha} \otimes_k S$ is stable with

respect to O_x , it is the direct sum of its weight subspaces. Hence

$$\mathfrak{g}_\alpha \otimes_k S = \bigoplus_w (L(\mathfrak{g})_S)_{\{w\}}$$

where $\{w\} = \{w + j \mid j \in \mathbb{Z}\}$ is the weight series corresponding to w . But $\mathfrak{g}_\alpha \otimes_k S$ has rank 1 as an S -module. This implies that in the above decomposition we have only one weight series $\{w\}$ for some weight w of O_x .

We next note that automatically we have $\dim_k (L(\mathfrak{g})_S)_w = 1$. Any its nonzero vector which is a generator of the S -module $\mathfrak{g}_\alpha \otimes_k S$ is of the form $X_\alpha t^{\frac{j}{m}}$. It follows from Lemma 9.2 that $\mathfrak{g}_\alpha = \langle X_\alpha \rangle$ is also a weight subspace of O_x . Thus for every root α we have

$$[x, X_\alpha]_{\widehat{L}(\mathfrak{g})_S} = [x' + d, X_\alpha]_{\widehat{L}(\mathfrak{g})_S} = [x', X_\alpha] = b_\alpha X_\alpha$$

for some scalar $b_\alpha \in k$. Since $x' \in \mathfrak{h}_S$ this can happen if and only if $x' \in \mathfrak{h}$. \square

By the previous lemma we have $x' \in \mathfrak{h}_{\bar{0}}$, hence

$$\mathfrak{m} = \langle \mathfrak{h}_{\bar{0}}, c, d \rangle = \mathcal{H}.$$

The proof of Theorem 12.1 is complete. \square

Acknowledgement: We would like to thank the referee for his/her thorough reading of our work and valuable comments and suggestions.

REFERENCES

- [ABP] B. Allison, S. Berman and A. Pianzola. *Covering Algebras II: Loop algebras of Kac-Moody Lie algebras*, Journal für die reine und angewandte Mathematik (Crelle), **571** (2004), 39-71.
- [BN] G. Benkart and E. Neher. *The centroid of extended affine and root graded Lie algebras*, Journal of Pure and Applied Algebra, **205** (2006), no.1, 117–154.
- [CGP] V. Chernousov, P. Gille, A. Pianzola, *Conjugacy theorems for loop reductive group schemes and Lie algebras*, arXiv: 1109.5236.
- [GP] P. Gille and A. Pianzola, *Torsors, Reductive group schemes and extended affine Lie algebras*, 125pp. (2011). ArXiv:1109.3405v2, to appear in Memoirs of the American Mathematical Society.
- [Kac] V. Kac, *Infinite dimensional Lie algebras*, third edition, Cambridge University Press, Cambridge, 1990.
- [Kmr] S. Kumar, *Kac-Moody groups, their flag varieties and representation theory*, Springer Verlag, 2002.
- [MP] R.V. Moody and A. Pianzola, *Lie algebras with triangular decomposition*, John Wiley, New York, 1995.
- [P1] A. Pianzola, *Locally trivial principal homogeneous spaces and conjugacy theorems for Lie algebras*, J. Algebra, **275** (2004), no. 2, 600–614.
- [P2] A. Pianzola, *Automorphisms of toroidal Lie algebras and their central quotients*, Jour. of Algebra and Applications, **1** (2002), 113–121.
- [PK] D.H. Peterson and V. Kac, *Infinite flag varieties and conjugacy theorems*, Proc. Natl. Acad. Sci. USA, **80** (1983), 1778–1782.
- [SGA3] *Séminaire de Géométrie algébrique de l'I.H.E.S., 1963-1964, schémas en groupes, dirigé par M. Demazure et A. Grothendieck*, Lecture Notes in Math. 151–153, Springer, 1970.
- [Se] J.-P. Serre, *Galois Cohomology*, Springer, 1997.

- [St62] R. Steinberg, *Générateurs, relations et revêtements des groupes algébriques*, Colloque sur la théorie des groupes algébriques (Bruxelles, 1962), Paris: Gauthier-Villars, 1962.
- [St67] R. Steinberg, Lectures on Chevalley Groups, Yale University 1967.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA
T6G 2G1, CANADA
E-mail address: `chernous@math.ualberta.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA
T6G 2G1, CANADA
E-mail address: `uyahorau@math.ualberta.ca`

UMR 8553 DU CNRS, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D'ULM, 75005 PARIS,
FRANCE.
E-mail address: `gille@ens.fr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA
T6G 2G1, CANADA.
CENTRO DE ALTOS ESTUDIOS EN CIENCIA EXACTAS, AVENIDA DE MAYO 866, (1084)
BUENOS AIRES, ARGENTINA.
E-mail address: `a.pianzola@gmail.com`