

Toxicity, repellency and flushing out in *Triatoma infestans* (Hemiptera: Reduviidae) exposed to the repellents DEET and IR3535

Mercedes M.N. Reynoso¹, Emilia A. Seccacini¹, Javier A. Calcagno², Eduardo N. Zerba^{1,3} and Raul A. Alzogaray^{1,3}

¹ UNIDEF, CITEDEF, CONICET, CIPEIN, Villa Martelli, Buenos Aires, Argentina

² Centro de Estudios Biomédicos, Biotecnológicos, Ambientales y de Diagnóstico (CEBBAD), Departamento de Ciencias Naturales y Antropológicas, CONICET, Ciudad Autónoma de Buenos Aires, Argentina

³ Instituto de Investigación e Ingeniería Ambiental (3IA), Universidad Nacional de San Martín (UNSAM), San Martín, Buenos Aires, Argentina

Q1

Please only
ANNOTATE
the proof.
Do not edit
the PDF.
If multiple
authors will
review this PDF,
please return
one file
containing all
corrections.

Submitted 1 February 2017

Accepted 10 April 2017

Published 23 May 2017

Corresponding author

Raul A. Alzogaray,
ralzogaray@hotmail.com

Academic editor

Joao Rocha

Additional Information and
Declarations can be found on
page 9

DOI 10.7717/peerj.3292

© Copyright
2017 Reynoso et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

ABSTRACT

DEET and IR3535 are insect repellents present worldwide in commercial products, which efficacy has been mainly evaluated in mosquitoes. This study compares the toxicological effects and the behavioral responses induced by both repellents on the blood-sucking bug *Triatoma infestans* Klug (Hemiptera: Reduviidae), one of the main vectors of Chagas disease. When applied topically, the Median Lethal Dose (72 h) for DEET was 220.8 $\mu\text{g}/\text{insect}$. Using IR3535, topical application of 250 $\mu\text{g}/\text{insect}$ killed no nymphs. The minimum concentration that produced repellency was the same for both compounds: 1,15 $\mu\text{g}/\text{cm}^2$. The effect of a mixture DEET:IR3535 1:1 was similar to that of their pure components. Flushing out was assessed in a chamber with a shelter containing groups of ten nymphs. The repellents were aerosolized on the shelter and the number of insects leaving it was recorded for 60 min. During that time, 0.006 g/m^3 of the positive control tetramethrin flushed out 76.7% of the nymphs, while 1.76 g/m^3 of DEET or IR3535 flushed out 30 and 0%, respectively. The concentrations required for both compounds to produce toxicity or flushing out are too high to have any practical applications. However, they showed a promisory repellency. Additional research should be done to evaluate their possible use for personal protection against *T. infestans* bites.

Subjects Entomology, Toxicology

Keywords Diethyltoluamide, Ethyl 3-[acetyl(butyl)amino]propanoate, Blood-sucking bugs, Locomotor activity

INTRODUCTION

An insect repellent has been defined as “something that causes insects to make oriented movements away from its source” (*White & Moore, 2015*). The main use given to these substances is for personal protection against the bite of hematophagous insects (*Debboun & Strickman, 2013*). There is a worldwide market of repellent products that contain different active ingredients (*Xue, Muller & Day, 2015*). Most contain N,N-diethyl-3-methylbenzamide (DEET), an active ingredient that has been used for over 60 years all over the world (*White & Moore, 2015*). DEET is a highly efficient repellent for a broad

spectrum of insect species; furthermore, its toxicity in mammals is very slow (EPA, 2000). The United States Environmental Protection Agency (EPA) considers that DEET does not present any risk of concern to human health (Antwi, Shama & Peterson, 2008). However, it can occasionally cause skin reactions and convulsions, especially in children (Osimitz et al., 2010).

The insect repellent ethyl 3-[acetyl(butyl)amino]propanoate (IR3535) was developed by the company Merck in 1975. Compared to DEET, oral or dermal exposure to IR3535 is less toxic and less irritant to mammals (WHO, 2001a; Nentwig, 2003). Other than being very irritating to eyes, no other adverse effects have been reported for IR3535 after more than 30 years of use (Puccetti, 2007). According to the WHO, it is “effective and safe for use in human beings” (WHO, 2001b). In Argentina, DEET and IR3535 are authorized as components in several repellent products (ANMAT, 2012).

Chagas disease, the most severe parasitic disease of the American continent, is caused by the protozoan *Trypanosoma cruzi* (Chagas, 1909) (Lent & Wygodzinsky, 1979). *T. cruzi* is transmitted to humans and other vertebrates by domestic, peridomestic, or sylvatic insects of the Triatominae family (Stevens et al., 2011). The triatomine *Triatoma infestans* (Klug, 1834) is the most important vector of *T. cruzi* in Argentina, Bolivia, Paraguay and Perú (Schofield and Gorla, 2010). In Latin America, Chagas disease affects more than 5.740.000 people (World Health Organization (WHO), 2015).

Different studies have assessed the behavioral response of Chagas disease vectors to synthetic and natural repellents under laboratory conditions (Alzogaray, Fontán & Zerba, 2000; Coelho, De Paula & Spíndola, 2006; Ferrero, González & Chopra, 2006; Abramson, Aldana & Sulbaran, 2007; Mello et al., 2007; Sfara, Zerba & Alzogaray, 2009; López et al., 2011; Pohlit et al., 2011; Avelar-Freitas et al., 2012; Gomes & Favero, 2013). The effects of DEET were barely explored in *T. infestans*, and there are no reports on the effects of IR3535 on triatomines. On the other hand, there is evidence that both repellents have insecticidal activity in house fly and mosquitoes (Pridgeon, Bernier & Becnel, 2009; Swale et al., 2014). Based on this background, the objective of this work was to compare the toxicological and behavioral responses of fifth-instar nymphs of *T. infestans* when exposed to DEET or IR3535.

MATERIALS AND METHODS

Biological material

Fifth-instar nymphs of *T. infestans*, 7–15 days old after last ecdysis, were provided from a laboratory colony maintained by the Centro de Referencia de Vectores (Santa María de Punilla, Córdoba, Argentina). They were kept at 26 ± 2 °C and 60–90% RH until each experiment.

Chemicals

DEET (97%) was purchased from Sigma Aldrich (Buenos Aires, Argentina), IR3535 (99.6%) was a gift from Merck Argentina (Buenos Aires, Argentina), and analytical grade acetone was acquired from Merck Germany (Darmstadt, Germany).

Toxicity bioassays

Six groups of ten nymphs were separated and each was randomly assigned to one of the following treatments: acetone alone (negative control), 31.25, 62.5, 125, 250 or 500 μg of DEET per insect. These concentrations were chosen after a preliminary assay. Treatment were applied using a microsyringe with a dispenser (Hamilton, Reno, NE), and each nymph received 1 μl solution on the abdomen. A similar bioassay was not performed with IR3535 because 500 $\mu\text{g}/\text{insect}$ of this repellent produced no mortality.

Immediately after the treatment, nymphs were placed in plastic container (10 cm high, 8 cm in diameter) closed with a gauze held with a rubber band. The container was maintained in an incubator FOC 225E provided with a thermoregulation system (Velp Scientifica, Usmate, Italy) programmed at 26 ± 2 °C and 60–90% RH. The number of affected nymphs was recorded 72 h after the treatment.

To quantify the toxicity of DEET and IR3535, a circle of filter paper 15 cm in diameter (101 FAST, Hangzhou Xinxing Paper Industry and Co., Ltd., Fuyang, China) was placed within a plastic container (32 cm long, 25 cm wide, and 8 cm high; Colombraro, Buenos Aires, Argentina). The treated nymphs were then carefully dropped in the centre of the paper circle and observed for 1 min. According preliminary observations, control nymphs abandon the paper circle in less than 5 s, following an approximately straight line toward the side of the plastic container. After these observations, a nymph remaining for at least 1 min on the paper circle and showing symptoms of intoxication (difficulty walking or no movement after being gently touched with a soft tweezer) was considered dead. Three independent replicates were made for each assay, and the Median Lethal Dose (*LD50*) was calculated.

Recording equipment

A black and white closed-circuit video camera (VC 1910, Sanyo Electrical Co., Tokyo, Japan) and an image analyser (Videomex V, Columbus Instruments, Columbus, OH) were used to evaluate repellency. The video camera records the movement of the nymph placed in the experimental arena and sends an analogical signal to the image analyser, where it is digitalized. Thus, the nymph appears as a white silhouette (pixels “on”) on the image analyser screen, while the filter paper appears as a black surface (pixels “off”). The Multiple Zone Motion Monitor software compares consecutive frames captured by the camera and records the number of pixels that change from “on” to “off” or vice versa. This software calculates two parameters: (a) Motion (M), the sum of pixels that changed during the assay, and (b) Area (A), the number of pixels that remained “on” (it represents the average area occupied by the nymph).

Repellency bioassays

As a first approach to quantify comparatively the repellence of these compounds, we used a preference test such as is commonly used to evaluate repellent effects on walking insects (Scheffler & Dombrowski, 1992; Aggarwal *et al.*, 2001). A circle of filter paper 110 mm in diameter (101 FAST, Hangzhou Xinxing Paper Industry and Co., Ltd., Fuyang, China) was cut into halves. One half was treated with 0.25 ml of DEET or IR3535 dissolved in acetone, and the other half was treated with 0.25 ml of acetone alone. After the solvent evaporated,

both halves were stuck back together with adhesive tape on the underside, and the circle was placed on a horizontal surface. A glass ring (2.5 cm high, 10 cm in diameter) was used to prevent the insects left the experimental arena. Finally, a nymph was gently deposited on the centre of the arena.

Each nymph was randomly assigned to one of four treatments: solvent alone (control), 0.38, 1.15, 3.43, 10.33, or 31.00 $\mu\text{g}/\text{cm}^2$ of each repellent alone. A mixture of DEET:IR3535 1:1 was also tested at the same concentrations.

The image analyser recorded the nymph movement on each zone of the filter paper for 15 min. Results were used to calculate a Distribution Coefficient (DC) (Moretti, Zerba & Alzogaray, 2013):

$$DC = (AT - At)/AT.$$

AT is the area occupied by the nymph throughout the assay, At is the area occupied by the nymph in the treated zone of the experimental arena. Values of DC vary between 0 (maximum attraction to the treated zone) and 1 (maximum repellence). Values close to 0.5 indicate that the insect spent the approximately the same amount of time on each zone.

The experimental arena was illuminated with a cold light lamp (22 watts; Luxa, Shanghai, China) located at the zenith. Temperature varied between 24 and 28 °C. Each assay was repeated four times. Replicates were carried out on different days with newly prepared solutions, and each insect was used only once.

Flushing out bioassays

Bioassays were performed inside a glass chamber (70 × 70 × 70 cm), illuminated by two cold light tubes of 20 watts each (Osram, Buenos Aires, Argentina) placed externally at the upper rear corner (Fig. 1). The front panel had a hole (5 cm in diameter) through which the flushing out agents were aerosolized into the chamber. Room temperature was maintained at 26 ± 2 °C. A black cardboard shaped as a triangular hollow prism (3 cm × 15 cm high), with its two ends opened, was located vertically inside the glass chamber (5 cm from the back wall and equidistant from the lateral walls). Ten fifth instar nymphs were gently released inside the black cardboard refuge and were allowed 15 min of familiarization. Following this, 1 ml of repellent in acetone (150, 300 or 600 mg/ml, equivalent to 0.44, 0.88, and 1.76 g/m^3 , respectively) was aerosolized through the front hole of the chamber using a glass sprayer. Compressed nitrogen was used as the carrier (3.5–3.8 psi). After the treatment, the hole in the chamber was sealed with a rubber stopper, and the number of insects leaving the refuge was recorded every 5 min during 1 h. One ml of acetone alone was aerosolized as a negative control and a solution of tetramethrin (0.002 mg/ml, equivalent to 0.006 g/m^3) was applied as a positive control. Three independent replicates were performed for each treatment.

Statistical analysis

LD_{50} values with their respective 95% Confidence Limits were calculated using the PoloPlus 2.0 programme (LeOra Software, 2002). Results from the repellency flushing out bioassays were analysed using one-way ANOVA, followed by Tukey's *post hoc* comparisons when $P < 0.05$. Results of repellency were also used to calculate linear regressions.




Figure 1 Glass chamber used for flushing out bioassays. 1, Exhaust fan (it is used to exhaust the contaminated air out of the glass chamber after each assay); 2, hole sealed with a rubber stopper; 3, front panel; 4, black cardboard triangular hollow prism (insects refuge); 5, two cold light tubes.

Table 1 Toxicity of DEET on fifth instar nymphs of *Triatoma infestans*.

	LD50 ^a (μg/insect) (95% CL) ^b	N	Slope ± SE	Chi-square
DEET	220.8 (167.8–313.0)	180	1.9 ± 0.3	1.99
IR3535	>500.0 ^c	–	–	–

Notes.

^aMedian Lethal Dose at 72 h.

^b95% Confidence Limit.

^cNo mortality was observed when this dose was applied.

RESULTS

Topical application of DEET and IR3535 showed very low toxicity on fifth instar nymphs of *T. infestans* (Table 1). The LD50 at 72 h for DEET was 220.8 μg/insect. No mortality was observed after topical application of 500 μg/insect of IR3535.

The repellent effect of pure and mixed solutions of DEET and IR3535, applied at concentrations ranging between 0.38 and 31.00 μg/cm², increased as a linear function of log concentration (Fig. 2 and Table 2). The values of r^2 varied between 0.922 and 0.963,




Figure 2 Repellency of pure and mixed DEET and IR3535 in fifth instar nymphs of *Triatoma infestans*. Distribution Coefficient = $(AT - At)/AT$, where AT is the total area occupied by the nymph on the arena, and At is the area occupied by the nymph on the treated zone throughout the experiment.

Table 2 Linear regressions for the independent and joint repellency of DEET and IR3535 on fifth instar nymphs of *Triatoma infestans*.

DEET:IR3535	Regression	r^2	F	df	P
1:0	$RC = 0.635 + 0.242 \log C$	0.963	77.920	1, 3	0.003
0:1	$RC = 0.683 + 0.193 \log C$	0.951	57.779	1, 3	0.005
1:1	$RC = 0.622 + 0.199 \log C$	0.922	35.613	1, 3	0.009

Notes.

RC , Response Coefficient = $(AT - At)/AT$, where AT is the total area occupied by the nymph on the arena, and At is the area occupied by nymphs on the treated zone throughout the experiment; $\log C$, log of concentration.

indicating a good fit to the model in all cases (Table 2). The minimum concentration that was significantly different from control (i.e., the minimum concentration that produced repellency) was the same for both substances: $1.15 \mu\text{g}/\text{cm}^2$ ($p < 0.05$). The effect of the mixture 1:1 was similar to the effects of their separate components (Table 3). In other words, neither synergy nor antagonism was observed.

No flushed out nymphs were observed when acetone or IR3535 alone were aerosolized on the insect refuges. Tetramethrin and DEET produced a significant flushing out ($F = 26.51$; $df = 3, 8$; $p < 0.001$) (Fig. 3). The positive control tetramethrin flushed out 76.7% of nymphs when applied at $0.006 \text{ g}/\text{m}^3$. Flushing out by DEET increased as the concentration increased. However, despite the high concentrations of this repellent used (0.44 – $1.76 \text{ g}/\text{m}^3$), none of them exceeded 40% of flushing out during the experimental time. IR3535 did not flushed out any nymph even at $1.76 \text{ g}/\text{m}^3$.

Table 3 Statistical analysis of different concentrations of pure or mixed DEET and IR3535.

Concentration ($\mu\text{g}/\text{cm}^2$)	DC ^a			F	df	P
	DEET:IR3535					
	1:0	0:1	1:1			
0.38	0.54a (0.06)	0.61a (0.03)	0.52a (0.04)	0.840	2, 9	0.463
1.15	0.64a (0.06)	0.67a (0.07)	0.70ab (0.06)	0.198	2, 9	0.824
3.4	0.74ab (0.06)	0.78ab (0.07)	0.68ab (0.04)	0.754	2, 9	0.498
10.3	0.94bc (0.03)	0.93b (0.03)	0.80bc (0.06)	3.271	2, 9	0.086
31	0.96c (0.02)	0.94b (0.03)	0.92c (0.03)	0.274	2, 9	0.767
F	24.0	16.1	11.7			
df	5, 18	5, 18	5, 18			
P	<0.001	<0.001	<0.001			

Notes.

^aDistribution Coefficient = $(AT - At)/AT$, where AT is the total area occupied by the nymph on the arena, and At is the area occupied by nymphs on the treated zone throughout the experiment. In each column, values followed by the same lowercase letter are not significantly different ($P > 0.05$). Statistical parameters are from one-way ANOVA.




Figure 3 Flushing out in fifth instar nymphs of *Triatoma infestans* exposed to aerosolized DEET. In the legend, values in brackets are expressed in units of g/m^3 . Different letters indicate significant differences ($P < 0.05$) (IR3535 produced 0% of flushing out in all replicates, so it was not included in the ANOVA). All replicates of IR3535 produced 0% of flushing out.

DISCUSSION

In this work, it was studied for the first time in *T. infestans*: (a) DEET and IR3535 toxicity, (b) the behavioral response to IR3535 alone or mixed with DEET, and (c) the flushing out effect of both compounds.

There are very few studies on the insecticidal properties of DEET and IR3535. In the house fly and the mosquitoes *Aedes aegypti* (Linnaeus, 1782). and *Anopheles gambiae* (Giles, 1902), DEET showed moderate toxicity when a few micrograms were applied topically (Swale *et al.*, 2014). It was also slightly more toxic than IR3535 in several mosquito species (Pridgeon, Bernier & Becnel, 2009).

Here, the LD50 (72 h) for DEET was 220.8 $\mu\text{g}/\text{insect}$. This is an extremely low toxicity when compared, for example, to deltamethrin, a pyrethroid widely used for controlling *T. infestans*. The Median Lethal Dose (72 h) of this insecticide in fifth instar nymphs is 1.27 ng/insect (De Oliveira Filho, 1999). Toxicity of DEET in triatomines was even lower when applied as films on filter paper on *T. rubida* (Uhler, 1894) (Terriquez *et al.*, 2013). On the other hand, the topical application of a high concentration of IR3535 produced no toxic effects on the nymphs. The same result was reported in *R. prolixus* (Alzogaray, 2016).

The primary site of action by which this repellent exerts its insecticidal activity has not yet been identified. DEET inhibits house fly and mosquito acetylcholinesterase activity, but only at very high concentrations (Corbel *et al.*, 2009). Neurophysiological studies suggest that the octopaminergic receptor of insects might be its target (Swale *et al.*, 2014).

Several reasons related to toxicokinetic and toxicodynamic processes could be the cause of the low toxicity of DEET and IR3535 in triatomines. For example, a low rate of cuticular penetration or a high rate of biotransformation might explain this characteristic. In *R. prolixus*, mixed function microsomal oxidases could be involved in the biotransformation of DEET because when these enzymes are inhibited with pyperonil butoxide, the toxicity of DEET is doubled (Alzogaray, 2016).

The repellent activity of these two compounds have been mainly studied in mosquitoes. IR3535 resulted as efficient as DEET in *Aedes* and *Culex* spp., but less efficient in *Anopheles* (Barnard *et al.*, 2002; Fradin & Day, 2002; Barnard & Xue, 2004; Cilek, Petersen & Hallmon, 2004; N'Guessan *et al.*, 2006). In the present work, DEET and IR3535 were equally repellent to fifth instar nymphs of *T. infestans*, presenting the same minimum concentration that produced repellency (1.15 $\mu\text{g}/\text{cm}^2$).

Pyrethroids and some botanical monoterpenes induce a non-directional increase in the locomotor activity of insects (Gammon, 1978; Alzogaray, Fontán & Zerba, 1997; Moretti, Zerba & Alzogaray, 2013). If the exposed insects are hidden in a shelter, they leave it by chance. This phenomenon is called flushing-out and is exploited to detect the presence of triatomines (Pinchin, De Oliveira Filho & Pereira, 1980). In Argentinian rural areas where Chagas disease is endemic, sanitary agents use aerosolized tetramethrin to flush out *T. infestans* from their shelters (Gürtler *et al.*, 1993). Flushing out allows determining whether a domicile is infested with triatomines; it is also used to evaluate the efficacy of an insecticide treatment and study the reinfestation of treated houses (Gürtler *et al.*, 2001). In the last years, resistance to pyrethroids has been reported in *T. infestans* populations from Argentina and Bolivia (Picollo *et al.*, 2005; Roca-Acevedo, Picollo & Santo-Orihuela, 2013).

The individuals from these populations are resistant to both knock down and hyperactivation produced by pyrethroids (Sfara, Zerba & Alzogaray, 2006). It is therefore highly important to identify alternative flushing out agents. Among natural compounds, isobutyric acid, 3-pentanol, 3-methyl-1-butanol (Minoli et al., 2013), and several monoterpenes (Moretti, Zerba & Alzogaray, 2015) showed flushing out activity on triatomines.

In the present work, DEET showed a very weak flushing out capacity compared to tetramethrin, a pyrethroid usually used by sanitary agents in Argentina to flush out triatomines. IR3535 produced no flushing out at all. Considering that hyperactivity is a symptom of intoxication, these results could be considered a consequence of the very low toxicity of these compounds in *T. infestans*.

Regrettably, the concentrations of DEET and IR3535 needed required to produce toxicity or flushing out in these species seem too high to have any practical applications. However, both compounds showed a similar and promisory repellency. Additional research should be done to evaluate the possible use of these compounds for personal protection against *T. infestans* bites. In particular, it may be worth to look for synergistic interactions with other compounds; for example, the botanical monoterpene eucalyptol, which showed repellent and insecticidal activity in *T. infestans* (Moretti, Zerba & Alzogaray, 2013).

ACKNOWLEDGEMENTS

The authors are grateful to Catriona Kirkwood for correcting the English style of this article, to Merck Argentina for donating IR3535, and to the two reviewers whose valuable comments and suggestions allowed improving this article.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work received financial support from Argentina's ANPCyT (PICT 2012-1471) and from Argentina's CONICET (PIP 2013-0238). JAC, ENZ and RAA are members of the Carrera del Investigador Científico y Tecnológico from CONICET. MMNR is a fellowship holder of the Agencia Nacional de Promoción Científica y Tecnológica from Argentina (ANPCyT). EAS is a member of the Carrera del Personal de Apoyo del Consejo Nacional de Investigaciones Científicas y Técnicas from Argentina (CONICET). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Argentina's ANPCyT: PICT 2012-1471.

Argentina's CONICET: PIP 2013-0238.

Carrera del Investigador Científico y Tecnológico.

Agencia Nacional de Promoción Científica y Tecnológica.

Carrera del Personal de Apoyo del Consejo Nacional de Investigaciones Científicas y Técnicas.

All supplemental information will be made available for download exactly as they were supplied. This link to the SI will only work when the article is published.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Mercedes M.N. Reynoso performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Emilia A. Seccacini performed the experiments, reviewed drafts of the paper.
- Javier A. Calcagno analyzed the data, reviewed drafts of the paper.
- Eduardo N. Zerba conceived and designed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Raul A. Alzogaray conceived and designed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as a [Data S1](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3292#supplemental-information>.

REFERENCES

- Abramson CI, Aldana E, Sulbaran E. 2007.** Exposure to citral, cinnamon and ruda disrupts the life cycle of a vector of Chagas Disease. *American Journal of Environmental Sciences* 3:7–8 DOI 10.3844/ajessp.2007.7.8.
- Administración Nacional de Medicamentos, Alimentos y Tecnología Médica (ANMAT). 2012.** Registro de repelentes de insectos inscriptos (actualizado al 18/06/2012) ANMAT, Buenos Aires. Available at http://www.anmat.gov.ar/listados/Listado_repelentes_jun_2012pdf.
- Aggarwal KK, Tripathi AK, Prajapati V, Kumar S. 2001.** Toxicity of 1,8-cineole towards three species of stored product coleopterans. *Insect Science and its Applications* 21:155–160.
- Alzogaray RA. 2016.** Behavioral and toxicological responses of *Rhodnius prolixus* (Hemiptera: Reduviidae) to the insect repellents DEET and IR3535. *Journal of Medical Entomology* 53:387–393 DOI 10.1093/jme/tjv194.
- Alzogaray RA, Fontán A, Zerba EN. 1997.** Evaluation of hyperactivity produced by pyrethroid treatment on nymph III of *Triatoma infestans* (Hemiptera: Reduviidae). *Archives of Insect Biochemistry and Physiology* 35:323–333 DOI 10.1002/(SICI)1520-6327(199705)35:3<323::AID-ARCH6>3.0.CO;2-U.
- Alzogaray RA, Fontán A, Zerba EN. 2000.** Repellency of deet to nymphs of *Triatoma infestans*. *Medical and Veterinary Entomology* 14:6–10 DOI 10.1046/j.1365-2915.2000.00213.x.

- Antwi FB, Shama LM, Peterson RK. 2008.** Risk assessments for the insect repellents DEET and picaridin. *Regulatory Toxicology and Pharmacology* **51**:31–36 DOI [10.1016/j.yrtph.2008.03.002](https://doi.org/10.1016/j.yrtph.2008.03.002).
- Avelar-Freitas BA, Dias JVL, Tibães GG, Rocha CA, Bahia-Oliveira GH, Graef CFF, Pires HHR. 2012.** The effect of *Ageratum fastigiatum* extract on *Rhodnius nasutus*, vector of Chagas disease. *Brazilian Journal of Pharmacognosy* **23**:366–369 DOI [10.1590/S0102-695X2012005000139](https://doi.org/10.1590/S0102-695X2012005000139).
- Barnard DR, Bernier UR, Posey KH, Xue D. 2002.** Repellency of IR3535, KBR3023, *para*-menthane-3,8-diol, and Deet to black salt marsh mosquitoes (Diptera: Culicidae) in the Everglades National Park. *Journal of Medical Entomology* **39**:895–899 DOI [10.1603/0022-2585-39.6.895](https://doi.org/10.1603/0022-2585-39.6.895).
- Barnard DR, Xue RD. 2004.** Laboratory evaluation of mosquito repellents against *Aedes albopictus*, *Culex nigripalpus*, and *Ochlerotatus triseriatus* (Diptera: Culicidae). *Journal of Medical Entomology* **41**:726–730 DOI [10.1603/0022-2585-41.4.726](https://doi.org/10.1603/0022-2585-41.4.726).
- Cilek JE, Petersen JL, Hallmon CF. 2004.** Comparative efficacy of IR3535 and DEET as repellents against adult *Aedes aegypti* and *Culex quinquefasciatus*. *Journal of the American Mosquito Control Association* **20**:299–304.
- Coelho AAM, De Paula JE, Spíndola LS. 2006.** Insecticidal activity on cerrado plants extracts on *Rhodnius milesi* Carcavallo, Rocha, Garvão and Jurberg (Hemiptera: Reduviidae), under laboratory conditions. *Neotropical Entomology* **35**:133–138 DOI [10.1590/S1519-566X2006000100018](https://doi.org/10.1590/S1519-566X2006000100018).
- Corbel V, Stankiewicz M, Pennetier C, Fournier D, Stojan J, Girard E, Dimitrov M, Molgó J, Hougard JM, Lapied B. 2009.** Evidence for inhibition of cholinesterases in insect and mammalian nervous system by the insect repellent DEET. *BMC Biology* **7**:47–57 DOI [10.1186/1741-7007-7-47](https://doi.org/10.1186/1741-7007-7-47).
- De Oliveira Filho AM. 1999.** Differences of susceptibility of five triatomine species to pyrethroid insecticides—implications for Chagas disease vector control. *Memorias do Instituto Oswaldo Cruz* **94**:425–428.
- Debboun M, Strickman D. 2013.** Insect repellents and associated personal protection for a reduction in human disease. *Medical and Veterinary Entomology* **27**:1–9 DOI [10.1111/j.1365-2915.2012.01020.x](https://doi.org/10.1111/j.1365-2915.2012.01020.x).
- Environmental Protection Agency (EPA). 2000.** *N,N-Diethyl-meta-toluamide (DEET)*. Risk characterization document. Sacramento: EPA.
- Ferrero A, González J, Chopa C. 2006.** Biological activity of *Schinus molle* on *Triatoma infestans*. *Fitoterapia* **77**:381–383 DOI [10.1016/j.fitote.2006.03.004](https://doi.org/10.1016/j.fitote.2006.03.004).
- Fradin MS, Day JF. 2002.** Comparative efficacy of insect repellents against mosquito bites. *New England Journal of Medicine* **347**:13–18 DOI [10.1056/NEJMoa011699](https://doi.org/10.1056/NEJMoa011699).
- Gammon DW. 1978.** Neural effects of allethrin on the free walking cockroach *Periplaneta americana*: and investigation using defined doses at 15 and 32 °C. *Journal of Pesticide Science* **9**:79–81 DOI [10.1002/ps.2780090115](https://doi.org/10.1002/ps.2780090115).
- Gomes SP, Favero S. 2013.** Assessment of the insecticidal potential of *Eucalyptus urograndis* essential oil against *Rhodnius neglectus* Lent (Hemiptera: Reduviidae). *Neotropical Entomology* **42**:431–435 DOI [10.1007/s13744-013-0144-2](https://doi.org/10.1007/s13744-013-0144-2).

- Gürtler RE, Schweigmann NJ, Cécere MC, Chuit R, Wisnivesky-Colli C. 1993.** Comparison of two sampling methods for domestic populations of *Triatoma infestans* in north-west Argentina. *Medical and Veterinary Entomology* 7:238–242 DOI [10.1111/j.1365-2915.1993.tb00683.x](https://doi.org/10.1111/j.1365-2915.1993.tb00683.x).
- Gürtler RE, Vázquez Prokopec GM, Ceballos LA, Lund Petersen C, Salomón OD. 2001.** Comparison between two artificial shelter units and timed manual collections for detecting peridomestic *Triatoma infestans* (Hemiptera: Reduviidae) in rural northwestern Argentina. *Journal of Medical Entomology* 38:429–436 DOI [10.1603/0022-2585-38.3.429](https://doi.org/10.1603/0022-2585-38.3.429).
- Lent H, Wygodzinsky P. 1979.** Revision of the triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas' disease. *Bulletin of the American Museum of Natural History* 163:125–520.
- LeOra Software 2002.** POLO PC: a user's guide for probit or logit analysis. Berkeley: LeOra Software.
- López B, López M, Aragón L, Tereschuk M, Slaris A, Feresin G, Zygadlo J, Tapia AA. 2011.** Composition and anti-insect activity of essential oils from *Tagetes L.* species (Asteraceae, Helenieae) on *Ceratitis capitata* Wiederman and *Triatoma infestans* Klug. *Journal of Agricultural and Food Chemistry* 59:5286–5292 DOI [10.1021/jf104966b](https://doi.org/10.1021/jf104966b).
- Mello C, Uzeda C, Bernardino M, Mendonça-Lopes D, Kelecom A, Fevereiro P, Guerra M, Oliveira A, Rocha L, González M. 2007.** Effects of the essential oil obtained from *Pilocarpus spicatus* Saint-Hilarie (Rutaceae) on the development of *Rhodnius prolixus* nymphae. *Brazilian Journal of Pharmacognosy* 17:514–520 DOI [10.1590/S0102-695X2007000400007](https://doi.org/10.1590/S0102-695X2007000400007).
- Ministerio de Salud de la Nación. 2014.** Programa nacional de Chagas: diagnóstico de situación. Buenos Aires: Ministerio de Salud de la Nación.
- Minoli S, Palottini F, Crespo JG, Manrique G. 2013.** Dislodgement effect of natural semiochemicals released by disturbed triatomines: a possible alternative monitoring tool. *Journal of Vector Ecology* 38:353–360 DOI [10.1111/j.1948-7134.2013.12051.x](https://doi.org/10.1111/j.1948-7134.2013.12051.x).
- Moretti A, Zerba EN, Alzogaray RA. 2013.** Behavioral and toxicological responses of *Rhodnius prolixus* and *Triatoma infestans* (Hemiptera: Reduviidae) to ten monoterpene alcohols. *Journal of Medical Entomology* 50:1046–1054 DOI [10.1603/ME12248](https://doi.org/10.1603/ME12248).
- Moretti A, Zerba EN, Alzogaray RA. 2015.** Lethal and sublethal effects of eucalyptol on *Triatoma infestans* and *Rhodnius prolixus*, vectors of Chagas disease. *Entomologia Experimentalis et Applicata* 154:62–70 DOI [10.1111/eea.12256](https://doi.org/10.1111/eea.12256).
- Nentwig G. 2003.** Use of repellents as prophylactic agents. *Parasitology Research* 1:S40–S48 DOI [10.1007/s00436-002-0755-8](https://doi.org/10.1007/s00436-002-0755-8).
- N'Guessan R, Rowland M, Traore Lamizana M, Kesse NB, Carnevale P. 2006.** Evaluation of synthetic repellents on mosquito nets in experimental huts against insecticide-resistant *Anopheles gambiae* and *Culex quinquefasciatus* mosquitoes. *Tropical Medicine & International Health* 100:1091–1097 DOI [10.1016/j.trstmh.2006.04.005](https://doi.org/10.1016/j.trstmh.2006.04.005).

- Osimitz TG, Murphy JV, Fell LA, Page B. 2010.** Adverse events associated with the use of insect repellents containing N,N-diethyl-m-toluamide (DEET). *Regulatory Toxicology and Pharmacology* **56**:93–99 DOI [10.1016/j.yrtph.2009.09.004](https://doi.org/10.1016/j.yrtph.2009.09.004).
- Piccolo MI, Vassena C, Orihuela P, Barrios S, Zaidemberg M, Zerba EN. 2005.** High resistance to pyrethroid insecticides associated to the ineffectiveness of field treatments in *Triatoma infestans* (Hemiptera, Reduviidae) from the north of Argentina. *Journal of Medical Entomology* **42**:637–642 DOI [10.1093/jmedent/42.4.637](https://doi.org/10.1093/jmedent/42.4.637).
- Pinchin R, De Oliveira Filho AM, Pereira ACB. 1980.** The flushing-out activity of pyrethrum and synthetic pyrethroids on *Panstrongilus megistus*, a vector of Chagas' disease. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **75**:691–694 DOI [10.1016/0035-9203\(80\)90207-2](https://doi.org/10.1016/0035-9203(80)90207-2).
- Pohlit AM, Ribeiro Rezende A, Lopes Baldin EL, Peoporine Lopes N, Ferreira de Andrade Neto V. 2011.** Plant extracts, isolated phytochemicals, and plant-derived agents which are lethal to arthropod vectors of human tropical diseases—a review. *Planta Medica* **77**:618–630 DOI [10.1055/s-0030-1270949](https://doi.org/10.1055/s-0030-1270949).
- Pridgeon JW, Bernier UR, Becnel JJ. 2009.** Toxicity comparison of eight repellents against four species of female mosquitoes. *Journal of the American Mosquito Control Association* **25**:168–173 DOI [10.2987/08-5837.1](https://doi.org/10.2987/08-5837.1).
- Puccetti G. 2007.** IR3535 (ethyl butylacetylaminopropionate). In: Debboun M, Frances SP, Strickman D, eds. *Insect repellent*. Boca Raton: CRC Press, 353–360.
- Roca-Acevedo G, Piccolo MI, Santo-Orihuela P. 2013.** Expression of insecticide resistance in immature life stages of *Triatoma infestans* (Hemiptera: Reduviidae). *Journal of Medical Entomology* **50**:816–818 DOI [10.1603/ME12116](https://doi.org/10.1603/ME12116).
- Scheffler I, Dombrowski M. 1992.** Behavioural responses of German cockroaches (*Blattella germanica* L.) induced by plant repellents. In: Otto D, Weber B, eds. *Insecticides: mechanism of action and resistance*. Andover: Intercept, 107–116.
- Sfara V, Zerba E, Alzogaray RA. 2006.** Toxicity of pyrethroids and repellency of diethyltoluamide in two deltamethrin resistant colonies of *Triatoma infestans* Klug 1834 (Hemiptera: Reduviidae). *Memorias Do Instituto Oswaldo Cruz* **101**:89–94 DOI [10.1590/S0074-02762006000100017](https://doi.org/10.1590/S0074-02762006000100017).
- Sfara V, Zerba EN, Alzogaray RA. 2009.** Fumigant insecticidal activity and repellent effect of five essential oils and seven monoterpenes in first instar nymphs of *Rhodnius prolixus*. *Journal of Medical Entomology* **46**:511–515 DOI [10.1603/033.046.0315](https://doi.org/10.1603/033.046.0315).
- Stevens L, Dorn PL, Schmidt JO, Klotz JH, Lucero D, Klotz SA. 2011.** Kissing bugs: vectors of Chagas. *Advances in Parasitology* **75**:169–192 DOI [10.1016/B978-0-12-385863-4.00008-3](https://doi.org/10.1016/B978-0-12-385863-4.00008-3).
- Swale DR, Sun B, Tong F, Bloomquist JR. 2014.** Neurotoxicity and mode of action of N,N-diethyl-meta-toluamide (DEET). *PLOS ONE* **9**:e103713 DOI [10.1371/journal.pone.0103713](https://doi.org/10.1371/journal.pone.0103713).
- Terriquez JA, Klotz SA, Meister EA, Klotz JH, Schmidt JO. 2013.** Repellency of DEET, picaridin, and three essential oils to *Triatoma rubida* (Hemiptera: Reduviidae: Triatominae). *Journal of Medical Entomology* **50**:664–667 DOI [10.1603/ME12282](https://doi.org/10.1603/ME12282).

- White GB, Moore SJ. 2015.** Terminology of insect repellents. In: Debboun M, Frances SP, Strickman D, eds. *Insect repellent, handbook*. 2nd edition. Boca Raton: CRC Press, 3–30.
- World Health Organization (WHO). 2001a.** *IR3535. Interim specification WHO/IS/TC/667/2001*. Geneva: WHO.
- World Health Organization (WHO). 2001b.** Review of IR3535; KBR 3023; (RS)-methoprene 20% EC, pyriproxyfen 0.5% GR; and lambda-cyhalothrin 2.5% CS. Report of the fourth WHOPES working group meeting. WHO, Geneva, Switzerland.
- World Health Organization (WHO). 2015.** Chagas disease in Latin America: an epidemiological update based on 2010 estimates. *Weekly Epidemiological Record* **90**:33–44.
- Xue R, Muller GC, Day JF. 2015.** Commercially available insect repellents and criteria for their use. In: Debboun M, Frances SP, Strickman D, eds. *Insect repellent, handbook*. 2nd edition. Boca Raton: CRC Press, 3–30.

Author Queries

Journal: PEERJ

Article id: 3292

Author: Reynoso et al.

Title: Toxicity, repellency and flushing out in *Triatoma infestans* (Hemiptera: Reduviidae) exposed to the repellents DEET and IR3535

Q1 (Page 1)

Please check all author names, initials, and affiliations to confirm they are accurate.

Uncited Reference(s)

This section includes references in the reference list but are not cited in the body of the text. Please indicate where each reference should be cited in the text or, alternatively, indicate that it should be deleted. Any reference not cited in the text but required as a resource must instead be placed in a “Further Reading” section, please identify any such references in the proof.

[Ministerio de Salud de la Nación \(2014\)](#)