



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: July 3, 2014

ACCEPTED: August 31, 2014

PUBLISHED: September 23, 2014

Anisotropic scale invariant spacetimes and black holes in Zwei-Dreibein Gravity

A.F. Goya

*Universidad de Buenos Aires FCEN-UBA and IFIBA-CONICET,
Ciudad Universitaria, Pabellón I, 1428, Buenos Aires, Argentina*

E-mail: af.goya@df.uba.ar

ABSTRACT: We show that Zwei-Dreibein Gravity (ZDG), a bigravity theory recently proposed by Bergshoeff, de Haan, Hohm, Merbis, and Townsend in ref. [1], admits exact solutions with anisotropic scale invariance. These type of geometries are the three-dimensional analogues of the spacetimes which were proposed as gravity duals for condensed matter systems. In particular, we find Schrödinger invariant spaces as well as Lifshitz spaces with arbitrary dynamical exponent z . We also find black holes that are asymptotically Lifshitz with $z = 3$, showing that these (non-constant curvature) solutions of New Massive Gravity (NMG) are persistent after the introduction of the infinite tower of higher-curvature terms of ZDG, provided a renormalization of the parameters. Black holes in asymptotically warped Anti-de Sitter spaces are also found. Interestingly, in almost all the geometries studied in this work, the metric associated with the second dreibein turns out to be equivalent, up to a constant global factor, to the first one. This phenomenon has been previously observed in other bigravity theories in asymptotically flat and asymptotically Anti-de Sitter backgrounds. However, for the particular case of the $z = 3$ Lifshitz black hole, here we found that the second metric corresponds to a different black hole that coincides with the former only in the asymptotic region. In fact, we find a new family of $z = 3$ black holes that corresponds to a one-parameter deformation of the NMG solution.

KEYWORDS: Holography and condensed matter physics (AdS/CMT), Black Holes, Classical Theories of Gravity

ARXIV EPRINT: [1406.4771](https://arxiv.org/abs/1406.4771)

Contents

1	Introduction	1
2	Review of New Massive Gravity	3
2.1	Lifshitz spacetimes in NMG	3
2.2	Lifshitz black hole in NMG	4
2.3	Schrödinger spacetimes in NMG	4
2.4	Warped Anti-de Sitter spacetimes in NMG	4
3	Zwei-Dreibein Gravity	5
3.1	A relation between $g_{1\mu\nu}$ and $g_{2\mu\nu}$	7
4	New exact solutions of Zwei-Dreibein Gravity	8
4.1	Lifshitz spacetimes in ZDG	9
4.2	Lifshitz black holes in ZDG	10
4.3	Schrödinger spacetimes in ZDG	11
4.4	Warped Anti-de Sitter spacetimes in ZDG	11
5	Conclusions	13

1 Introduction

The idea that a theory of quantum gravity must be ‘holographic’, in the sense that the relevant degrees of freedom must be ‘on the surface’, was surrounding the theoretical physics community for several years [2, 3]. The first concise and most successful realization of holographic ideas was given by Maldacena in 1997 [4], relating type IIB string theory in five-dimensional Anti-de Sitter space (AdS) with $\mathcal{N} = 4$ Super Yang-Mills theory in four-dimensional Minkowski space. Since then, it is commonly believed that this gravity/gauge duality [5, 6] is an intrinsic property of any sensible candidate for quantum gravity, so it must hold not only for string theory but for any gravity theory that is expected to be well defined beyond the classical limit. Therefore, this relation must establish a connection between any complete gravity theory on AdS_{d+1} and a conformal field theory (CFT) in d -dimensional Minkowski space.

Since 2008, there have been attempts to extend AdS/CFT in order to apply holography to non-relativistic field theories scenarios, like superconductors [7] and other problems in condensed matter [8]. These proposals to generalize AdS/CFT had even involved non-AdS bulk geometries; such is the case of spacetimes having the same symmetries as the Schrödinger group with anisotropic scale invariance, which were conjectured to be gravity duals of strongly coupled condensed matter systems [9, 10]; other example is the so-called

Lifshitz spacetimes, which were proposed as gravity duals for Lifshitz fixed points [11]. The question then arised as to what kind of gravity theory would admit these anisotropic scale invariant spaces as exact solutions. Notably, a three-dimensional higher-derivative gravity theory, dubbed as New Massive Gravity (NMG) [12, 13], was shown to contain in its spectrum both Schrödinger spaces [14] and Lifshitz spaces [15] as exact solutions. In addition, the theory also contains Lifshitz black holes [15], which would be dual to finite temperature systems of these type. This invites to consider NMG in the context of condensed matter applications of holography, at least as a toy model. On the other hand, NMG also admits as exact solutions other interesting non-AdS spaces that could be relevant for holography, such as Warped Anti-de Sitter (WAdS) geometries, which are stretched or squashed deformation of AdS_3 , and which could provide another interesting playground to study extensions of the AdS/CFT correspondence. The isometry group of WAdS_3 is $\text{SL}(2, \mathbb{R}) \times \text{U}(1)$ and provides enough information to build a boundary field theory [16]. This kind of spacetimes and the black holes that asymptote to them are solutions of NMG as well [17]. All this convinces ourselves that three-dimensional massive gravity seems to be a good arena to study different non-AdS extensions of holography.

Very recently, a new 3D gravity model with very interesting properties, called Zwei-Dreibein Gravity (ZDG), was proposed [1]. It was shown in [18] that this bigravity theory can be viewed as a higher-derivative gravity theory of which NMG is a particular limit. Some indications were found in [18] that ZDG at some particular points of the parameter space can be interpreted as a gravity dual of the Logarithmic Conformal Field Theories (LCFT) [19], which are also relevant in the context of condensed matter. Hence, a natural question is whether ZDG can also be used in non-relativistic extensions of the AdS/CFT correspondence. In order to answer this we should verify whether Lifshitz, Schrödinger and Warped AdS spacetimes are solutions of ZDG as well. This would give a hint that ZDG could provide another sensible model to test non-relativistic holography.

In this paper we will show that, indeed, the recently proposed ZDG theory does admit all those spaces as exact solutions. We will construct explicit solutions of ZDG representing Schrödinger spacetimes as well as black holes that asymptote to both Lifshitz and WAdS spacetimes. An interesting feature observed in other bigravity theories considered in the literature is that, when a black hole solution is considered, the two ‘metrics’ of the theory are related each other by coordinate transformations and a global rescaling. Here, we will show that this sort of ‘duality’ also appears in ZDG for some solutions. However, for the asymptotically Lifshitz black hole, the second ‘metric’ represents a new Lifshitz black hole that coincides with the first one only in the asymptotic region. In fact, here we find a new one-parameter family of Lifshitz black holes.

The paper is organized as follows: in section 2, we will review NMG theory and its non-AdS solutions. In sections 3 and 4, we will present ZDG theory and explicitly show how the non-AdS geometries of NMG persists in this three-dimensional bigravity theory, and we will give the explicit relation between the two spin-two fields. Finally in section 5, we will conclude with some final remarks and outlook.

2 Review of New Massive Gravity

In this section, we will review the important aspects of New Massive Gravity (NMG) [12, 13] and the relevant solutions of this theory for this work.

NMG is a higher-derivative gravity theory that at linearized level is equivalent to a Fierz-Pauli theory for a massive spin 2 field. Its action is given by the Einstein-Hilbert term (with a sign parameter $\varsigma = \pm 1$), a cosmological constant term, and a particular term quadratic in the curvature; namely

$$S_{NMG} = \frac{M_P}{2} \int d^3x \sqrt{-g} \left(\varsigma R - 2\Lambda + \frac{1}{m^2} \left(R_{\mu\nu} R^{\mu\nu} - \frac{3}{8} R^2 \right) \right),$$

where the free parameters of the theory are the Planck mass M_P , the bare cosmological constant Λ , and the mass of the gravitons m^2 . The specific choice of the relative coefficient $-3/8$ between the squared curvature terms is necessary in order to eliminate a scalar ghost mode from the spectrum.

An interesting feature of NMG is that it contains a large variety of solutions. In particular, due to its higher-derivative nature, it is possible to circumvent the obstructions of Birkhoff-like theorems and find different static spherically symmetric solutions. The spectrum of NMG contains hairy black holes, wormholes, gravitational solitons, and kinks [20]. It is also possible to find asymptotically AdS₃ spaces with weaker boundary conditions relative to the Brown-Henneaux ones [21, 22]. Remarkably, this theory also supports asymptotically dS₃ black holes [13, 20], unlike three-dimensional General Relativity. Besides, as we said before, NMG provides a playground for non-relativistic holography since it contains solutions whose isometries realizes non-relativistic symmetries. Let us review some of these solutions below.

2.1 Lifshitz spacetimes in NMG

Lifshitz spacetimes have an anisotropic scale invariance under

$$t \rightarrow \lambda^z t, \quad x \rightarrow \lambda x.$$

As we said before, geometries that realize this kind of symmetries were conjectured to be gravity duals of condensed matter systems of Lifshitz type [11]. The so-called Lifshitz spacetimes in $d + 1$ dimensions are given by

$$ds^2 = -\frac{r^{2z}}{\ell^{2z}} dt^2 + \frac{\ell^2}{r^2} dr^2 + \frac{r^2}{\ell^2} d\vec{x} \cdot d\vec{x}, \tag{2.1}$$

where \vec{x} is a $(d - 1)$ -dimensional spacelike vector. It is worth noticing that for $z = 1$ the spacetime (2.1) correspond to AdS _{$d+1$} . The three-dimensional version of (2.1) is an exact solution of NMG [15]. The curvature radius ℓ and the mass parameter m^2 are related with the parameter z by

$$m^2 \ell^2 = \frac{1}{2\varsigma} (z^2 - 3z + 1), \quad \Lambda \ell^2 = -\frac{\varsigma}{2} (z^2 + z + 1). \tag{2.2}$$

Notably, NMG also contains asymptotically Lifshitz black holes as we will see below.

2.2 Lifshitz black hole in NMG

In order to study condensed matter theories at finite temperature it is necessary to find black holes that coincide with the Lifshitz space in the asymptotic region. In [15] a black hole was found that is asymptotically Lifshitz with $z = 3$. It was recently shown that this is the only asymptotically Lifshitz black hole with $z \neq 1$ in the theory [23].¹ The metric of this three-dimensional $z = 3$ Lifshitz black hole is given by the following line element

$$ds^2 = -\frac{r^6}{\ell^6} \left(1 - \frac{M\ell^2}{r^2}\right) dt^2 + \frac{r^2}{\ell^2} dx^2 + \left(\frac{r^2}{\ell^2} - M\right)^{-1} dr^2.$$

The metric above solves the equations of motion of NMG provided

$$m^2\ell^2 = \frac{1}{2\zeta}, \quad \Lambda\ell^2 = -\frac{13\zeta}{2}.$$

The above conditions are just (2.2) evaluated at $z = 3$. We will see that this solution also persists in Zwei-Dreibein Gravity (ZDG) [1].

2.3 Schrödinger spacetimes in NMG

Other interesting geometries for non-AdS holography applications are Schrödinger spaces. These spaces realize geometrically the non-relativistic conformal group, also known as the Schrödinger group. These spacetimes are particular cases of AdS waves [14] and are also equivalent to null WAdS spaces, which we will discuss in the next subsection. The Schrödinger line element is given by

$$ds^2 = \frac{\ell^2}{y^2} \left(-\frac{\ell^{2\nu}}{y^{2\nu}} du^2 - 2dudv + dy^2 \right),$$

and happens to be a solution of NMG if

$$m^2\ell^2 = -2\zeta(1 + 8\nu(\nu + 1)), \quad \Lambda = \frac{1 - 4m^2\ell^2\zeta}{4m^4\ell^4}.$$

Next, let us discuss the WAdS spaces.

2.4 Warped Anti-de Sitter spacetimes in NMG

Warped Anti-de Sitter space can be obtained as a Hopf fibration of \mathbb{R} over AdS_2 and multiplying the fiber by a constant warp factor. In a convenient system of coordinates the line element reads

$$ds^2 = \frac{1}{4} \left(-\cosh^2 x d\tau^2 + dx^2 + \frac{4\nu^2}{\nu^2 + 3} (dy + \sinh x d\tau)^2 \right).$$

For $\nu = 1$ this line element is equivalent to AdS_3 in a rotating frame. There are different types of deformations of AdS_3 producing timelike, spacelike, and null WAdS_3 spaces. As we said before, the Schrödinger space correspond to the null WAdS_3 . It is interesting to

¹At least for spacetimes that can be written as a Kerr-Schild deformation of a Lifshitz space.

notice that the timelike version of WAdS₃ actually corresponds to the three-dimensional part of the Gödel solution of General Relativity [24].² The Gödel spacetime can be written as a direct product $\Sigma_3 \times \mathbb{R}$, and this Σ_3 piece is precisely the timelike WAdS₃ geometry. In this work we will focus on the spacelike WAdS₃. Notably, there are exact solutions that represent black holes that are locally and asymptotically equivalent to spacelike WAdS₃ spaces in the same way as how the BTZ black hole is locally and asymptotically equivalent to AdS₃. The WAdS₃ black holes are given by

$$ds^2 = -N^2 dt^2 + \rho^2 (d\varphi + N^\varphi dt)^2 + \frac{\ell^2 dr^2}{4\rho^2 N^2}, \quad (2.3a)$$

with

$$\begin{aligned} \rho^2 &= \frac{r}{4} \left(3(\nu^2 - 1)r + (\nu^2 + 3)(r_+ + r_-) - 4\nu\sqrt{(\nu^2 + 3)r_+ r_-} \right), \\ N^2 &= \frac{(\nu^2 + 3)(r - r_+)(r - r_-)}{4\rho^2}, \\ N^\varphi &= \frac{2\nu r - \sqrt{(\nu^2 + 3)r_+ r_-}}{2\rho^2}. \end{aligned} \quad (2.3b)$$

In the context of parity even massive gravity theories, these black holes were first discussed by Clément in ref. [17]. They are solutions of NMG if m^2 and Λ satisfy the following relations

$$m^2 \ell^2 = -\frac{(20\nu^2 - 3)\varsigma}{2}, \quad \Lambda \ell^2 = \frac{\varsigma(9 - 48\nu^2 + 4\nu^4)}{2(20\nu^2 - 3)}.$$

The WAdS₃ black holes of NMG were studied in detail in ref. [25]; see also [26].

3 Zwei-Dreibein Gravity

Now, let us describe the main aspects of the Zwei-Dreibein Gravity theory (ZDG) proposed in ref. [1].

ZDG is a bigravity theory whose dynamical fields are the two dreibein 1-forms $e_I^a = e_{I\mu}^a dx^\mu$ and the two spin-connection 1-forms $\omega_I^a = \omega_{I\mu}^a dx^\mu$. While the subindex $I = 1, 2$ identifies each of the two spin-2 fields, the superindex $a = \hat{1}, \hat{2}, \hat{3}$ refers to the coordinates in the tangent bundle. Index $\mu = 0, 1, 2$ labels spacetime coordinates. The Lagrangian of the theory is given by

$$\begin{aligned} L_{ZDG} = -\mathcal{M}_P \left\{ \sigma e_{1a} R_1^a + e_{2a} R_2^a + \frac{1}{6} m^2 \epsilon_{abc} \left(\alpha_1 e_1^a e_1^b e_1^c + \alpha_2 m^2 e_2^a e_2^b e_2^c \right) \right. \\ \left. - \frac{1}{2} m^2 \epsilon_{abc} \left(\beta_1 e_1^a e_1^b e_2^c + \beta_2 e_1^a e_2^b e_2^c \right) \right\}, \end{aligned} \quad (3.1)$$

where we suppressed the wedge symbol \wedge and where all the products between forms must be understood as exterior products.

²The author thanks Geoffrey Compère and Stéphane Detournay for clarifying comments about this subject.

The curvature and the torsion 2-forms are given by the following expressions

$$R_I^a = d\omega_I^a + \frac{1}{2}\epsilon^{abc}\omega_{Ib}\omega_{Ic}, \quad T_I^a = \mathcal{D}_I e_I^a \equiv de_I^a + \epsilon^{abc}\omega_{Ib}e_{Ic},$$

where the usual rank-2 valued 2-forms have been replaced by its dual tensors, e.g. $\mathcal{T}^a = (1/2)\epsilon^a_{bc} \mathcal{T}^{bc}$.

The independent parameters in (3.1) are the two cosmological constant parameters α_I , the two interaction constants β_I , and the Planck mass \mathcal{M}_P . The parameter m^2 is a convenient but redundant mass parameter, and $\sigma = \pm 1$ is a sign parameter.

For arbitrary values of the parameters β_1 and β_2 the theory contains three bulk degrees of freedom, two corresponding to the helicity ± 2 modes of a massive graviton. The remaining degree of freedom is potentially a ghost mode. One way to avoid this presumably ghost mode is by restricting the parameter space of the theory, for example, choosing $\beta_2 = 0$, and assuming that e_1 is invertible [1, 27–29].³

Therefore, the equations of motion for e_1^a, e_2^a , and ω_I^a , with $\beta_2 = 0$, and with an invertible e_1 , derived from the Lagrangian density (3.1) are given by:

$$0 = \sigma R_1^a + \frac{1}{2}m^2\epsilon^a_{bc} \left(\alpha_1 e_1^b e_1^c - 2\beta_1 e_1^b e_2^c \right), \tag{3.2a}$$

$$0 = R_2^a + \frac{1}{2}m^2\epsilon^a_{bc} \left(\alpha_2 e_2^b e_2^c - \beta_1 e_1^b e_1^c \right), \tag{3.2b}$$

$$0 = T_I^a, \tag{3.2c}$$

respectively.

Notice that the curvature and torsion 2-forms satisfy the Bianchi identities

$$\mathcal{D}_I R_I^a = 0, \quad \mathcal{D}_I T_I^a = \epsilon^{abc} R_{Ib} e_{Ic}.$$

The kinetic terms of e_1 and e_2 in (3.1) are invariant under their respective diffeomorphisms and local Lorentz transformations. Notice that the presence of the interaction term breaks the symmetries to their diagonal subgroup, given by the identification of the gauge parameters of each group.

An interesting remark is that ZDG can be understood as a theory with a single dreibein that, however, contains an infinite number of higher-derivative terms [18]. The fact that bigravity theories can be viewed as higher-derivative theories of one metric field was first noticed in [30]. A similar procedure was followed in [18] for the ZDG case. To see this, first we observe that we can solve the equations of motion (3.2a) to obtain an expression for e_2^a in terms of e_1^a . Then, using the property $\epsilon^{\rho\sigma\tau} R_{1\sigma\tau}^a = \det(e_1) e_1^{\sigma a} G_{1\rho}^\sigma$, we obtain the following expression for e_2^a :

$$e_{2\mu}^a = \frac{\alpha_1}{2\beta_1} e_{1\mu}^a + \frac{\sigma}{m^2\beta_1} S_{1\mu}^a, \tag{3.3}$$

³The most general way to ensure that ZDG is a ghost free theory is to assume the invertibility of the linear combination $\beta_1 e_1^a + \beta_2 e_2^a$ (for more details see [29]). The choice made in this work (e_1 invertible and $\beta_2=0$) is a particular case of the former one.

where $S_{1\mu}{}^a \equiv S_{1\mu\nu}e_1^{\nu a}$ and where $S_{1\mu\nu} = R_{1\mu\nu} - \frac{1}{4}R_1g_{1\mu\nu}$ is the Schouten tensor associated with the metric $g_{1\mu\nu} \equiv e_{1\mu}{}^a e_{1\nu}{}^b \eta_{ab}$. Here, we identify $g_{1\mu\nu}$ with the physical metric since, as we said before, we have assumed e_1 to be invertible. The ‘auxiliary field’ e_2 represents the higher-derivative content of the theory. The dreibein e_2 is auxiliary in the sense that it can be solved for algebraically from the equations of motion (3.2a), in the same way as in the second derivative formulation of NMG [31].

It is worth pointing out that the resulting higher-derivative equations of motion for g_1 up to order $1/m^2$ can be integrated to an action if the parameters of the theory satisfy a relation consistent with the NMG limit (see appendix B of [18] for more details). This does not mean that ZDG with generic values of its coupling constant is a higher-derivative completion of NMG. However, we remark that the resulting action at order $1/m^2$ is precisely the NMG action. In general, the interpretation of ZDG as a higher-derivative gravity theory of one metric, only holds at the level of the equations of motion without matter fields.

3.1 A relation between $g_{1\mu\nu}$ and $g_{2\mu\nu}$

Besides the relation (3.3) between e_{1a}^μ and e_{2a}^μ and the corresponding metrics $g_{1\mu\nu} = e_{1\mu}{}^a e_{1\nu}{}^b \eta_{ab}$ and $g_{2\mu\nu} = e_{2\mu}{}^a e_{2\nu}{}^b \eta_{ab}$, one can wonder whether for a particular class of solutions, there exists a coordinate transformation that maps g_2 into g_1 and viceversa.⁴ It is worth mentioning that here and in what follows we will refer to the g_2 field as a ‘metric’, as a consequence of the form that it takes that resembles a metric. Since we are only assuming the invertibility of e_1 , g_1 is the physical metric.⁵ Whether both spin-two fields have a clear geometric interpretation is not yet understood.

In this subsection we will explore whether both ‘metrics’ g_1 and g_2 are related by coordinate/scale transformations. This sort of ‘duality’ between the two spin-two fields were found in the so-called *f-g theory* [33], where the two metrics ($g_{\mu\nu}$ and $f_{\mu\nu}$) represent the Schwarzschild-(A)dS solution at the same time [34]. In [35], hairy black hole solutions were found that asymptote (A)dS in the Hassan-Rosen bigravity theory [36] and a similar phenomenon happens there. All these are examples of four dimensional bigravity theories; however, there also exists an example of this type in three dimensions: in [32] the three dimensional version of the *f-g theory* was studied and it was found asymptotically AdS₃ black hole solutions for the two metrics. Therefore, it is natural to ask whether ZDG has this ‘duality’ as well. In the particular case in which g_1 is a maximally symmetric space, provided $R_{1\mu\nu} = 2\Lambda g_{1\mu\nu}$, using the expression (3.3) it becomes clear that g_2 will be proportional to g_1 . This is also true for the Bañados-Teitelboim-Zanelli (BTZ) black hole [37], since it is locally equivalent to AdS₃. More precisely, if the metric associated to e_1 is given by

$$ds_1^2 = - \left(\frac{r^2}{\ell^2} - M \right) dt^2 + 2J dt d\varphi + r^2 d\varphi^2 + \left(\frac{r^2}{\ell^2} - M + \frac{J}{4r^2} \right)^{-1} dr^2,$$

⁴The author thanks Gastón Giribet for draw his attention to this point, specially to the reference [32].

⁵In the footnote 4 of ref. [18] the authors said that it is possible to find the inverse of e_2 as a power expansion in $1/m^2$. On the other hand, even when the invertibility on the dreibein e_2 is not imposed as a necessary condition, notice that the explicit solutions we found here for the two spin-two fields are manifestly invertible.

then, the ‘metric’ associated to e_2 is given by a constant conformal factor times the BTZ black hole metric

$$ds_2^2 = \frac{1}{4m^2\ell^4\beta_1^2} \left\{ - \left(\frac{r^2}{\ell^2} - M \right) dt^2 + 2Jdt d\varphi + r^2 d\varphi^2 + \left(\frac{r^2}{\ell^2} - M + \frac{J}{4r^2} \right)^{-1} dr^2 \right\}.$$

The same happens for the AdS waves studied in [18]. In the latter case, the metric associated to e_1 is given by

$$ds_1^2 = \frac{\ell^2}{y^2} \left\{ -f(u, y) du^2 - 2dudv + dy^2 \right\},$$

where the wave profile $f(u, y)$ is polynomial in y

$$f(u, y) = f_0(u) + f_2(u) \left(\frac{y}{\ell} \right)^2 + f_+ \left(\frac{y}{\ell} \right)^{n_+} + f_-(u) \left(\frac{y}{\ell} \right)^{n_-}, \quad (3.4)$$

and $\{0, 2, n_+, n_-\}$ are the roots of the characteristic polynomial obtained from the equations of motion (3.2a)–(3.2b).⁶

On the other hand, the ‘metric’ of e_2 is given by

$$ds_2^2 = \frac{(m^2\ell^2\alpha_1 - \sigma)^2 \ell^2}{4m^2\ell^4\beta_1^2} \frac{\ell^2}{y^2} \left\{ -F(u, y) du^2 - 2dudv + dy^2 \right\},$$

where

$$F(u, y) = (1 - 2\sigma y(y\partial_y^2 - \partial_y)) f(u, y),$$

is identified with the g_2 wave profile. Here we will assume that $m^2\ell^2\alpha_1 \neq \sigma$.

Therefore, we observe that in ZDG we have a similar phenomenon: for some solutions, spin-two fields g_1 and g_2 represent the same type of geometry, what we mean is that when one dreibein field solution corresponds to a given geometry, the other dreibein field solution is given by the same geometry up to parameter redefinitions and diffeomorphisms. It would be interesting to find out if the same ‘duality’ holds in the case of non-AdS geometries. As far as we know, all the known solutions for which such a ‘duality’ between metrics holds, either in three or four spacetime dimensions, are asymptotically flat, dS or AdS geometries. Here we will show that, indeed, ZDG bigravity theory, possesses non-AdS solutions which exhibit this sort of ‘duality’ between the two dynamical fields. This suggests that this phenomenon is much more universal of what we thought and it is not necessarily related to the simplicity of asymptotically constant curvature solutions.

4 New exact solutions of Zwei-Dreibein Gravity

In this section we will construct explicit non-AdS exact solutions of ZDG.

⁶The special cases where the roots n_{\pm} degenerate were studied in detail in [14] for the NMG case, and in [18] for the ZDG case.

4.1 Lifshitz spacetimes in ZDG

Let us begin with Lifshitz type solutions. The expression (3.3) gives us a hint to propose an ansatz to find a Lifshitz solution of ZDG. It is convenient to rescale the parameters α_1 , α_2 , and β_1 by defining $\alpha \equiv m^2 \ell^2 \alpha_1$, $\mathcal{A} \equiv m^2 \ell^2 \alpha_2$, and $\beta \equiv m^2 \ell^2 \beta_1$. The dreibein e_1 will be related to the Lifshitz metric in the usual way $g_{1\mu\nu} = e_{1\mu}{}^a \eta_{ab} e_{1\nu}{}^b$; and using (3.3) the ansatz for e_2^a is

$$\begin{aligned} e_2^{\hat{1}} &= \frac{\alpha - \sigma(z^2 + z - 1)}{2\beta} \left(\frac{r}{\ell}\right)^z dt, \\ e_2^{\hat{2}} &= \frac{\alpha + \sigma(z^2 - z - 1)}{2\beta} \left(\frac{r}{\ell}\right) dx, \\ e_2^{\hat{3}} &= \frac{\alpha - \sigma(z^2 - z + 1)}{2\beta} \left(\frac{\ell}{r}\right) dr. \end{aligned} \tag{4.1}$$

Three-dimensional Lifshitz spacetime is a solution of the ZDG equations of motion (3.2a)–(3.2b) provided \mathcal{A} and β satisfy the following relations

$$\begin{aligned} \mathcal{A} &= -\frac{(\alpha(\alpha - 2\sigma z) - (z^2 + z - 1)(z^2 - z - 1))^2 (\alpha - \sigma(z^2 + z + 1))}{2\sigma(\alpha - \sigma(z^2 - z + 1))^4}, \\ \beta &= -\frac{\alpha(\alpha - 2\sigma z) - (z^2 + z - 1)(z^2 - z - 1)}{2\sigma(\alpha - \sigma(z^2 - z + 1))}, \end{aligned} \tag{4.2}$$

or, equivalently,

$$\frac{\mathcal{A}}{\beta} = \frac{(\alpha(\alpha - 2z\sigma) - (z^2 + z - 1)(z^2 - z - 1)) (\alpha - \sigma(z^2 + z + 1))}{(\alpha - \sigma(z^2 - z + 1))^3}, \tag{4.3}$$

assuming in particular $\alpha\sigma \neq z^2 - z + 1$.

As we said in the previous section, it would be interesting to investigate whether there exists non-AdS solutions to bigravity theories like ZDG that exhibit the mentioned ‘duality’ between the two spin-two fields. To give a first answer to this question we can observe that the ‘metric’ associated with (4.1) can be put in the following form

$$ds_2^2 = \Omega^2 \left(-\frac{r^{2z}}{\ell^{2z}} d\tilde{t}^2 + \frac{r^2}{\ell^2} d\tilde{x}^2 + \frac{\ell^2}{r^2} dr^2 \right),$$

where we have (re-)defined

$$\begin{aligned} \Omega^2 &= \frac{(\alpha - \sigma(z^2 - z + 1))^2}{4\beta^2}, \quad \tilde{t} = \frac{\alpha - \sigma(z^2 + z - 1)}{\alpha - \sigma(z^2 - z + 1)}, \\ \tilde{x} &= \frac{\alpha - \sigma(z^2 - z - 1)}{\alpha - \sigma(z^2 - z + 1)}. \end{aligned}$$

Thus, g_2 also represents a Lifshitz spacetime with the same dynamical exponent z , that is a global rescaling of g_1 , assuming additionally $\alpha \neq \sigma(z^2 \pm z - 1)$.

4.2 Lifshitz black holes in ZDG

Remarkably, for the special case $z = 3$, ZDG also admits a static circularly symmetric black hole solution that asymptotes Lifshitz space. This shows that the Lifshitz black hole of NMG

$$ds^2 = -\frac{r^6}{\ell^6} \left(1 - \frac{M\ell^2}{r^2}\right) dt^2 + \frac{r^2}{\ell^2} dx^2 + \left(\frac{r^2}{\ell^2} - M\right)^{-1} dr^2, \quad (4.4)$$

persists as an exact solution after the infinite set of higher-curvature terms of ZDG were introduced.⁷ This is notable due to the fact that Lifshitz black holes do not exhibit constant curvature invariants.

To prove that the black hole metric (4.4) is solution of ZDG, it is convenient to consider the following ansatz for e_2

$$\begin{aligned} e_2^{\hat{1}} &= \frac{(\alpha - 11\sigma)r^2 + 4M\ell^2\sigma}{2\beta r^2} \left(\frac{r}{\ell}\right)^3 \sqrt{1 - \frac{M\ell^2}{r^2}} dt, \\ e_2^{\hat{2}} &= \frac{\alpha + 5\sigma}{2\beta} \frac{r}{\ell} dx, \\ e_2^{\hat{3}} &= \frac{\alpha - 7\sigma}{2\beta} \left(\frac{r^2}{\ell^2} - M\right)^{-1/2} dr. \end{aligned} \quad (4.5)$$

The three-dimensional Lifshitz black hole of [15] solves the ZDG equations of motion if \mathcal{A} and β satisfy the conditions (4.3) evaluated at $z = 3$; namely

$$\mathcal{A} = -\frac{(\alpha(\alpha - 6\sigma) - 55)^2(\alpha - 13\sigma)}{2\sigma(\alpha - 7\sigma)^4}, \quad \beta = -\frac{\alpha(\alpha - 6\sigma) - 55}{2\sigma(\alpha - 7\sigma)}, \quad (4.6)$$

or, equivalently,

$$\frac{\mathcal{A}}{\beta} = \frac{(\alpha(\alpha - 6\sigma) - 55)(\alpha - 13\sigma)}{(\alpha - 7\sigma)^3}. \quad (4.7)$$

Here we are assuming $\alpha \neq 11\sigma$, $\alpha \neq 7\sigma$, and $\alpha \neq -5\sigma$ for (4.5)–(4.6) to be well defined.

We can see that the ‘metric’ associated with the second dreibein (4.5) can be rewritten as a global factor times an asymptotically $z = 3$ Lifshitz black hole. Actually, the line element of g_2 reads

$$ds_2^2 = \Omega^2 \left\{ -\left(\frac{r^2}{\ell^2} - M\right) \left(\frac{r^2}{\ell^2} + \mu\right)^2 d\tilde{t}^2 + \frac{r^2}{\ell^2} d\tilde{x}^2 + \frac{dr^2}{\left(\frac{r^2}{\ell^2} - M\right)} \right\}, \quad (4.8)$$

where we (re-)defined

$$\Omega^2 = \frac{(\alpha - 7\sigma)^2}{4\beta^2}, \quad \mu = \frac{4M\sigma}{\alpha - 11\sigma}, \quad \tilde{t} = \frac{\alpha - 11\sigma}{\alpha - 7\sigma}, \quad \tilde{x} = \frac{\alpha - 5\sigma}{\alpha - 7\sigma}. \quad (4.9)$$

The parameter μ is positive provided $\text{sign}(M\sigma)(\alpha - 11\sigma) > 0$.

The ‘spacetime’ described by the line element (4.8) has non-constant curvature invariants with the same divergences structure than (4.4). However, it is important to remark

⁷As we said in section 3, ZDG is not a higher-curvature completion of NMG.

that black hole (4.8) is different from the one found in [15], representing a one-parameter deformation of the latter. The presence of the parameter μ makes the black hole (4.8) not locally equivalent to (4.4), and it may also contribute to global properties of the geometry like the value of its mass. Both black holes share, however, the same asymptotic behaviour, approaching $z = 3$ Lifshitz space at large r .

Notice that, provided both M and μ are positive, black hole (4.8) has positive temperature, given by

$$T = \frac{\sqrt{M}(M + \mu)}{2\pi\ell}.$$

It remains an open problem how to define conserved charges in this setup, in order to study, for example, the thermodynamics of the new black hole (4.8).

4.3 Schrödinger spacetimes in ZDG

Now, let us consider other anisotropic spaces, the Schrödinger spacetimes. These are a special case of the AdS-waves studied in [18], for which the wave profile functions (3.4) takes the values $f_+(u) = 0$, $f_-(u) = f_0$, and $n_- = -2\nu$. The dreibein e_1 in this case is given by

$$e_1^{\hat{1}} = \frac{\ell}{y} \left(\frac{\ell^\nu}{y^\nu} du + \frac{y^\nu}{\ell^\nu} dv \right), \quad e_1^{\hat{2}} = \frac{\ell}{y} \frac{y^\nu}{\ell^\nu} dv, \quad e_1^{\hat{3}} = \frac{\ell}{y} dy,$$

while e_2 is given by

$$\begin{aligned} e_2^{\hat{1}} &= \frac{\ell}{y} \left[\left(\gamma - \frac{2\sigma\nu(1+\nu)}{\beta} \right) \frac{\ell^\nu}{y^\nu} du + \gamma \frac{y^\nu}{\ell^\nu} dv \right], \\ e_2^{\hat{2}} &= \frac{\ell}{y} \left[-\frac{2\sigma(1+\nu)}{\beta} \frac{\ell^\nu}{y^\nu} du + \gamma \frac{y^\nu}{\ell^\nu} dv \right], \\ e_2^{\hat{3}} &= \frac{\ell\gamma}{y} dy. \end{aligned}$$

For this geometry to be a solution of ZDG, the parameters of the theory must satisfy the following relations

$$\alpha = -\sigma(2\beta + 1 - 8\nu - 8\nu^2), \quad \mathcal{A} = -\frac{\beta^2(\beta - 1)}{(\beta - 4\nu(1 + \nu))^2}, \quad \gamma = -\frac{\sigma(\beta - 4\nu(1 + \nu))}{\beta}.$$

Since the Schrödinger spaces are particular cases of the AdS waves, the spin-two field g_2 is a global rescaling of g_1 as we discussed in section 3.1. To avoid repetition we address the reader's attention to ref. [18] for details.

4.4 Warped Anti-de Sitter spacetimes in ZDG

We already mentioned that a special case of Schrödinger spaces is the so-called null WAdS geometry. This is therefore a solution of ZDG as well. Besides, the theory admits also spacelike and timelike WAdS spaces. Interestingly, the spacelike stretched WAdS as well as the black hole that is locally equivalent (and asymptote) to it, namely (2.3) solves the

ZDG equations of motion (3.2a)–(3.2b). So, the first dreibein has the form

$$\begin{aligned} e_1^{\hat{1}} &= dt + \left(\nu r - \frac{1}{2} \sqrt{(\nu^2 + 3)r_+ r_-} \right) d\varphi, \\ e_1^{\hat{2}} &= \frac{1}{2} \sqrt{(\nu^2 + 3)(r - r_+)(r - r_-)} d\varphi, \\ e_1^{\hat{3}} &= \frac{\ell}{\sqrt{(\nu^2 + 3)(r - r_+)(r - r_-)}} dr, \end{aligned} \tag{4.10}$$

while the second dreibein has the form

$$\begin{aligned} e_2^{\hat{1}} &= \frac{\alpha - \sigma(4\nu^2 - 3)}{4\beta} \left[2dt + \left(2\nu r - \sqrt{(\nu^2 + 3)r_+ r_-} \right) d\varphi \right], \\ e_2^{\hat{2}} &= \frac{\alpha + \sigma(2\nu^2 - 3)}{4\beta} \sqrt{(\nu^2 + 3)(r - r_+)(r - r_-)} d\varphi, \\ e_2^{\hat{3}} &= \frac{\ell}{2\beta} \frac{\alpha + \sigma(2\nu^2 - 3)}{\sqrt{(\nu^2 + 3)(r - r_+)(r - r_-)}} dr. \end{aligned} \tag{4.11}$$

For these black hole geometries to be solutions of ZDG, the parameters α , \mathcal{A} , and β must satisfy

$$\begin{aligned} \frac{\mathcal{A}}{\beta} &= \frac{(\alpha - \sigma(4\nu^2 - 3))}{(\alpha + \sigma(2\nu^2 - 3))^6} (20\nu^4 - 12\sigma(\alpha + \sigma)\nu^2 + (\alpha - 3\sigma)^2) \\ &\quad \times (8\sigma\nu^6 + 12(\alpha - 3\sigma)\nu^4 - 12\sigma(\alpha^2 - 3)\nu^2 + (\alpha - 3\sigma)^3). \end{aligned} \tag{4.12}$$

The ‘metric’ associated to the second dreibein of the spacelike WAdS black hole (4.11) can be put in the following form

$$\begin{aligned} ds_2^2 &= \Omega^2 \left\{ d\tilde{t}^2 + 2 \left(\nu r - \frac{1}{2} \sqrt{(\nu^2 + 3)r_+ r_-} \right) d\tilde{t} d\tilde{\varphi} + \frac{\ell^2 dr^2}{(\nu^2 + 3)(r - r_+)(r - r_-)} \right. \\ &\quad \left. + \frac{r}{4} \left(3(\nu^2 - 1)r + (\nu^2 + 3)(r_+ + r_-) - 4\nu \sqrt{(\nu^2 + 3)r_+ r_-} \right) d\tilde{\varphi}^2 \right\} \end{aligned} \tag{4.13}$$

where we (re-)defined

$$\Omega^2 = \frac{9(\nu^2 - 1)^2}{4\beta^2}, \quad \alpha = \nu^2 \sigma, \quad \tilde{t} = -t, \quad \tilde{\varphi} = -\varphi, \tag{4.14}$$

and where the coupling constants take the values

$$\mathcal{A} = -\frac{(\nu^2 - 3)}{2}, \quad \beta = \frac{3(\nu^2 - 1)}{2}.$$

It is worth mentioning that the ‘spacetime’ described by (4.13) does not differ globally from (2.3) since the parameter rescaling given by (4.14) does not introduce any angular deficit.

5 Conclusions

In this paper we have constructed explicit solutions of the recently proposed Zwei-Dreibein Gravity (ZDG) theory, which is a consistent bigravity theory in three dimensions. Among the exact solutions of this theory we have found geometries with anisotropic scale invariance with and without Galilean invariance. These are three-dimensional analogues of the Schrödinger and Lifshitz spaces, respectively, which were conjectured to be dual to strongly coupled condensed matter systems. We have also found static circularly symmetric black holes that asymptote Lifshitz space, which is remarkable as they represent non-constant curvature solutions of New Massive Gravity that persist after the introduction of the infinite set of higher-curvature terms of ZDG. Warped Anti-de Sitter black holes were also derived. For the Lifshitz, Schrödinger and Warped Anti-de Sitter spacetimes, the two spin-two fields of the theory are related each other by a global rescaling and a coordinate transformation. For the Lifshitz black hole, in contrast, the second ‘metric’ represents a new Lifshitz black hole that asymptote $z = 3$ Lifshitz geometry, representing a new family of one-parameter Lifshitz black holes. Whether the two dynamical fields of the theory have a clear geometric interpretation remains as an open problem. In addition, it would be interesting to find out whether ZDG contains in its spectrum new anisotropic scale invariance solutions, like Lifshitz black holes with $z \neq 3$. An open problem is how to define conserved charges for this theory; finding a method to do so would be important in order to study, for example, the thermodynamics of the bigravity black holes. The results of this paper give additional motivations to study three-dimensional massive (bi)gravity theories as toy models to study AdS/CFT holography and its ramifications.

Acknowledgments

The work of A.F.G. is supported by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and by Universidad de Buenos Aires (UBA). A.F.G. thanks G. Giribet and W. Merbis for useful comments and remarks on the manuscript, and to E. Bergshoeff and J. Rossell for discussions on Zwei-Dreibein Gravity. A.F.G. would like to thank the hospitality of Centre for Theoretical Physics of the University of Groningen where part of this work was done.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E.A. Bergshoeff, S. de Haan, O. Hohm, W. Merbis and P.K. Townsend, *Zwei-Dreibein Gravity: a two-frame-field model of 3D massive gravity*, *Phys. Rev. Lett.* **111** (2013) 111102 [[arXiv:1307.2774](https://arxiv.org/abs/1307.2774)] [[INSPIRE](#)].
- [2] G. 't Hooft, *Dimensional reduction in quantum gravity*, [gr-qc/9310026](https://arxiv.org/abs/gr-qc/9310026) [[INSPIRE](#)].

- [3] L. Susskind, *The world as a hologram*, *J. Math. Phys.* **36** (1995) 6377 [[hep-th/9409089](#)] [[INSPIRE](#)].
- [4] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)] [[INSPIRE](#)].
- [5] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[INSPIRE](#)].
- [6] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [7] S.A. Hartnoll, C.P. Herzog and G.T. Horowitz, *Building a holographic superconductor*, *Phys. Rev. Lett.* **101** (2008) 031601 [[arXiv:0803.3295](#)] [[INSPIRE](#)].
- [8] S.A. Hartnoll, *Lectures on holographic methods for condensed matter physics*, *Class. Quant. Grav.* **26** (2009) 224002 [[arXiv:0903.3246](#)] [[INSPIRE](#)].
- [9] D.T. Son, *Toward an AdS/cold atoms correspondence: A Geometric realization of the Schrödinger symmetry*, *Phys. Rev. D* **78** (2008) 046003 [[arXiv:0804.3972](#)] [[INSPIRE](#)].
- [10] K. Balasubramanian and J. McGreevy, *Gravity duals for non-relativistic CFTs*, *Phys. Rev. Lett.* **101** (2008) 061601 [[arXiv:0804.4053](#)] [[INSPIRE](#)].
- [11] S. Kachru, X. Liu and M. Mulligan, *Gravity duals of Lifshitz-like fixed points*, *Phys. Rev. D* **78** (2008) 106005 [[arXiv:0808.1725](#)] [[INSPIRE](#)].
- [12] E.A. Bergshoeff, O. Hohm and P.K. Townsend, *Massive gravity in three dimensions*, *Phys. Rev. Lett.* **102** (2009) 201301 [[arXiv:0901.1766](#)] [[INSPIRE](#)].
- [13] E.A. Bergshoeff, O. Hohm and P.K. Townsend, *More on massive 3D gravity*, *Phys. Rev. D* **79** (2009) 124042 [[arXiv:0905.1259](#)] [[INSPIRE](#)].
- [14] E. Ayon-Beato, G. Giribet and M. Hassaine, *Bending AdS waves with new massive gravity*, *JHEP* **05** (2009) 029 [[arXiv:0904.0668](#)] [[INSPIRE](#)].
- [15] E. Ayon-Beato, A. Garbarz, G. Giribet and M. Hassaine, *Lifshitz black hole in three dimensions*, *Phys. Rev. D* **80** (2009) 104029 [[arXiv:0909.1347](#)] [[INSPIRE](#)].
- [16] S. Detournay, T. Hartman and D.M. Hofman, *Warped conformal field theory*, *Phys. Rev. D* **86** (2012) 124018 [[arXiv:1210.0539](#)] [[INSPIRE](#)].
- [17] G. Clement, *Warped AdS₃ black holes in new massive gravity*, *Class. Quant. Grav.* **26** (2009) 105015 [[arXiv:0902.4634](#)] [[INSPIRE](#)].
- [18] E.A. Bergshoeff, A.F. Goya, W. Merbis and J. Rosseel, *Logarithmic AdS waves and Zwei-Dreibein Gravity*, *JHEP* **04** (2014) 012 [[arXiv:1401.5386](#)] [[INSPIRE](#)].
- [19] V. Gurarie, *Logarithmic operators in conformal field theory*, *Nucl. Phys. B* **410** (1993) 535 [[hep-th/9303160](#)] [[INSPIRE](#)].
- [20] J. Oliva, D. Tempo and R. Troncoso, *Three-dimensional black holes, gravitational solitons, kinks and wormholes for BHT massive gravity*, *JHEP* **07** (2009) 011 [[arXiv:0905.1545](#)] [[INSPIRE](#)].
- [21] G. Giribet and M. Leston, *Boundary stress tensor and counterterms for weakened AdS₃ asymptotic in new massive gravity*, *JHEP* **09** (2010) 070 [[arXiv:1006.3349](#)] [[INSPIRE](#)].
- [22] G. Giribet, A. Goya and M. Leston, *Boundary stress tensor and asymptotically AdS₃ non-Einstein spaces at the chiral point*, *Phys. Rev. D* **84** (2011) 066003 [[arXiv:1108.0400](#)] [[INSPIRE](#)].

- [23] E. Ayón-Beato, M. Hassaine and M.M. Juárez-Aubry, *Towards the uniqueness of Lifshitz black holes and solitons in New Massive Gravity*, *Phys. Rev. D* **90** (2014) 044026 [[arXiv:1406.1588](#)] [[INSPIRE](#)].
- [24] K. Gödel, *An example of a new type of cosmological solutions of Einstein's field equations of gravitation*, *Rev. Mod. Phys.* **21** (1949) 447.
- [25] E. Tonni, *Warped black holes in 3D general massive gravity*, *JHEP* **08** (2010) 070 [[arXiv:1006.3489](#)] [[INSPIRE](#)].
- [26] G. Giribet and A. Goya, *The Brown-York mass of black holes in warped Anti-de Sitter space*, *JHEP* **03** (2013) 130 [[arXiv:1212.2100](#)] [[INSPIRE](#)].
- [27] C. Deffayet, J. Mourad and G. Zahariade, *Covariant constraints in ghost free massive gravity*, *JCAP* **01** (2013) 032 [[arXiv:1207.6338](#)] [[INSPIRE](#)].
- [28] M. Bañados, C. Deffayet and M. Pino, *The Boulware-Deser mode in 3D first-order massive gravity*, *Phys. Rev. D* **88** (2013) 124016 [[arXiv:1310.3249](#)] [[INSPIRE](#)].
- [29] E.A. Bergshoeff, O. Hohm, W. Merbis, A.J. Routh and P.K. Townsend, *The hamiltonian form of three-dimensional Chern-Simons-like gravity models*, [arXiv:1402.1688](#) [[INSPIRE](#)].
- [30] S.F. Hassan, A. Schmidt-May and M. von Strauss, *Higher derivative gravity and conformal gravity from bimetric and partially massless bimetric theory*, [arXiv:1303.6940](#) [[INSPIRE](#)].
- [31] O. Hohm and E. Tonni, *A boundary stress tensor for higher-derivative gravity in AdS and Lifshitz backgrounds*, *JHEP* **04** (2010) 093 [[arXiv:1001.3598](#)] [[INSPIRE](#)].
- [32] M. Bañados and S. Theisen, *Three-dimensional massive gravity and the bigravity black hole*, *JHEP* **11** (2009) 033 [[arXiv:0909.1163](#)] [[INSPIRE](#)].
- [33] C.J. Isham, A. Salam and J.A. Strathdee, *F-dominance of gravity*, *Phys. Rev. D* **3** (1971) 867 [[INSPIRE](#)].
- [34] C.J. Isham and D. Storey, *Exact spherically symmetric classical solutions for the F-G theory of gravity*, *Phys. Rev. D* **18** (1978) 1047 [[INSPIRE](#)].
- [35] M.S. Volkov, *Hairy black holes in theories with massive gravitons*, [arXiv:1405.1742](#) [[INSPIRE](#)].
- [36] S.F. Hassan and R.A. Rosen, *Bimetric Gravity from Ghost-free Massive Gravity*, *JHEP* **02** (2012) 126 [[arXiv:1109.3515](#)] [[INSPIRE](#)].
- [37] M. Bañados, C. Teitelboim and J. Zanelli, *The black hole in three-dimensional space-time*, *Phys. Rev. Lett.* **69** (1992) 1849 [[hep-th/9204099](#)] [[INSPIRE](#)].