

Publicación Especial N° 13



ASOCIACIÓN PALEONTOLÓGICA ARGENTINA

CONODONTS FROM THE ANDES

Proceedings of the 3rd International Conodont Symposium
&
Regional Field Meeting of the IGCP project 591

Edited by
Guillermo L. Albanesi and Gladys Ortega

BUENOS AIRES
2013

MIDDLE ORDOVICIAN CONODONTS AND GRAPTOLITES AT LOS CAUQUENES RANGE, CENTRAL PRECORDILLERA OF SAN JUAN, ARGENTINA



VOLDMAN, G. G.^{1,2}, ORTEGA, G.¹ AND ALBANESI, G. L.^{1,2}

¹ CONICET-Museo de Paleontología, CIGEA, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, X5000JJC, Córdoba, Argentina; gvoldman@efn.uncor.edu, gcortega@arnet.com.ar, galbanes@com.uncor.edu

² CICTERRA-CONICET, Córdoba, Argentina.

Keywords: Conodonts. Graptolites. Biostratigraphy. Middle Ordovician. Precordillera. Argentina.

THE Ordovician System is superbly represented in the Precordillera of western Argentina, at the Andean foothills. During the Middle Ordovician, an important paleogeographical rearrangement of depocenters and source areas took place associated with the demise of the Eopaleozoic carbonate platform of the Precordillera. This critical interval is recorded in the basin through widespread deposition of black shales over the fossiliferous limestones of the San Juan Formation, punctuated by local deposition of olistostromes, debris flows, conglomerates, and turbidites (*e.g.*, Astini *et al.*, 1995; Keller, 1999). Fossil dating of the lowest black shales and equivalent units indicates a diachronous deposition from north to south in the Precordillera basin (Hünicken, 1985; Albanesi *et al.*, 1998).

The occurrence of Ordovician clastic rocks in the Jáchal area is known since the study of Borrello and Gareca (1951), who recorded *Nemagraptus gracilis* (J. Hall) in the Ordovician upper shaly succession that is exposed in the western flank of the Cerro Viejo, a prominent mountain situated midway between Huaco and Jáchal city. Harrington (in Harrington and Leanza, 1957) distinguished two formations in the lower and upper parts of this succession, the Cerro Viejo Shales and Los Azules Shales, respectively. Furque (1979) referred all the Ordovician clastic sequence to the Los Azules Formation, equivalent to the Gualcamayo Formation elsewhere. Later, Ortega (1987) refined the biostratigraphy of the Los Azules Formation by determining a lower member, made up of dark claystones and siltstones interbedded with K-bentonites (lower Darriwilian), a middle member that consists of black shales and siltstones (upper Darriwilian), and an upper member of yellowish calcareous silt-

stones and calcarenites (upper Sandbian). A hiatus covering the highest Darriwilian and the lowest Sandbian is recorded between the middle and upper members.

The Los Azules Formation is arranged in small isolated outcrops aligned in narrow elongated belts with meridian orientation. The best exposures of the Los Azules Formation are located at the foothill of the Cerro Viejo, between 30° 11' 40" and 30° 15' 30" S Latitude, and 68° 34' 30" and 68° 35' 20" W Longitude, uplifted by the Niquivil Thrust (*e.g.*, Ortega *et al.*, 2007). It also crops out along western thrusts in the less studied successions of the La Chilca, Mogotes Azules, and the Aguada de Los Azulejitos on the northernmost Talacasto Range.

The purpose of this paper is to describe for the first time a restricted stratigraphic exposure that bear a significant conodont and graptolite fauna of the upper section of the San Juan Formation and the overlying the Los Azules Formation (Fig. 1), which is currently under study. The stratigraphic section is located at the Los Cauquenes Range, a parallel thrust belt to the Cerro Viejo de Huaco thrust. The latter thrust belt is 4 km to the east, where the Los Azules Creek, the type section of the homonymous formation, is located.

At the Oculca Creek, the upper levels of the San Juan Formation are characterized by nodular limestones, which are highly fossiliferous. These are roofed by a hardground surface, rich in nautiloids at the contact with the Los Azules Formation, which is interpreted as a marine flooding surface (Astini, 1994). The Los Azules Formation presents 226 m in thickness at the Oculca Creek, with the lower, middle and upper members well developed. It is unconformably overlain through angular

discordance by Holocene alluvial fan deposits. The Los Azules Formation starts with dark-brown argillites with yellowish alteration and conchoidal fracture interbedded with K-bentonites. The middle member consists of black-grey shales and siltstones, and the upper member is made up of calcareous siltstones of dark colour and yellow alterations, with scarce mudstones strata intercalated, some of them reaching up to 1 m.

In order to define the age of aforementioned strata by conodont dating, we took 3 rock samples from the uppermost levels

of the San Juan Formation (4.5 kg) and 2 rock samples from the base of the Los Azules Formation (3 kg). The samples are condensed limestone deposits, which after digesting in 10% acetic acid following the conventional procedures, produced over 2200 conodont elements. The collected specimens are well preserved and exhibit a CAI 2 (Epstein *et al.*, 1977). These are stored in the Museo de Paleontología, Universidad Nacional de Córdoba, under repository code CORD-MP.

The conodont collection exhibits a high species diversity (Fig. 2), including *Ansella jemtlandica* (Löfgren), *Baltoniodus* sp.,




Figure 1. Geological map with the location of the Oculita Creek, Central Precordillera, San Juan Province, Argentina (modified from Furque, 1979, Furque *et al.*, 1998).



Figure 2. Darriwilian (*Yangtzeplacognathus crassus* Zone, Da2) conodonts from the Oculita Creek, Central Precordillera, San Juan Province, Argentina. **1, 7**, *Drepanodus arcuatus* Pander. **1**, Pb element, CORD-MP28039, **7**, Sa element, CORD-MP28045; **2**, *Cornuodus longibasis* (Lindström), Sd element, CORD-MP 28040; **3**, *Ansella jemtländica* (Löfgren), M element, CORD-MP 28041; **4**, *Protopanderodus calceatus* Bagnoli and Stouge, P element, CORD-MP 28042; **5, 12, 15**, *Yangtzeplacognathus crassus* (Chen and Zhang). **5**, Pb element, antero-lateral view, CORD-MP28043, **12**, Sa element, upper view, CORD-MP 28050, **15**, Pa element, upper view, CORD-MP28053; **6**, *Parapaltodus simplicissimus* Stouge, M element, CORD-MP 28044; **8–9**, *Periodon* n. sp. **8**, M element, CORD-MP 28046, **9**, Pa element, CORD-MP 28047; **10**, *Semiacontiodus potrerillensis* Albanesi, long based coniform, CORD-MP 28048; **11**, *Parioistodus horridus* (Barnes and Poplawski), Sc element, CORD-MP28049; **13**, *Microzarkodina hagetiana* Stouge and Bagnoli, M element, CORD-MP28051; **14**, *Histiodela holodentata* Ethington and Clark, Pa element, CORD-MP28052; **16**, Gen. et sp. indet., CORD-MP28054; **17**, *Pteracniodus cryptodens* (Mound), M element, CORD-MP28055; **18**, *Periodon macrodentatus* (Graves and Ellison), M element, CORD-MP28056; **19**, *Erraticodon alternans* (Hadding), Pb element, CORD-MP28057; **20**, *Drepanoistodus bellburnensis* Stouge, M element, CORD-MP28058; **21**, *Fahraesodus marathonsis* (Bradshaw), Sa element, CORD-MP 28059; **22**, *Drepanoistodus costatus* (Abaimova), Sa element, CORD-MP 28060; **23**, *Costiconus costatus* (Dzik), Sb element, CORD-MP 28061; **24**, “*Bryantodina*” aff. *typicalis* Stauffer, Pa element, CORD-MP 28062; **25**, *Drepanoistodus tablepointensis* Stouge, M element, CORD-MP 28063. All illustrated specimens proceed from sample TFSJ1° hrdgrnd. All elements show inner-lateral views except where indicated. Scale bar: 0.1 mm.

“*Bryantodina*” aff. *typicalis* Stauffer, *Cornuodus longibasis* (Lindström), *Costiconus costatus* (Dzik), *Drepanodus arcuatus* Pander, *D. reclinatus* (Lindström), *Drepanoistodus bellburnensis* Stouge, *D. costatus* (Abaimova), *D. forceps* (Lindström), *D. tablepointensis* Stouge, *Erraticodon alternans* (Hadding), *Fabraesodus marathoniensis* (Bradshaw), *Histiodella holodentata* Ethington and Clark, *Microzarkodina hagetiana* Stouge and Bagnoli, *Parapaltodus simplicissimus* Stouge, *Paroistodus originalis* (Sergeeva), *Paroistodus horridus primus* Albanesi, *P. h. secundus* Albanesi, *P. horridus horridus* (Barnes and Poplawski), *Periodon macrodentatus* (Graves and Ellison), *Protopanderodus calceatus* Bagnoli and Stouge, *Pteracantiodus cryptodens* (Mound), *Scolopodus striatus* Pander, *Semiacantiodus potrerillensis* Albanesi and the index species *Yangtzeplacognathus crassus* (Chen and Zhang). It also includes a new species of *Periodon* and a Gen. et sp. indet. This conodont association allows constraining the uppermost levels of the San Juan Formation and the lower levels of the Los Azules Formation to the *Yangtzeplacognathus crassus* Zone of early middle Darriwilian age (Da2) (Fig. 3), following the Baltoscandic and Chinese schemes (Zhang, 1998; Löfgren and Zhang, 2003). This is consistent with the timing of the drowning of the San Juan Formation in the nearby Cerro Viejo de Huaco (Ortega *et al.*, 2007), as well as Los Blanquitos and La Trampa ranges, central area of the Precordillera (Albanesi *et al.*, in press). In Laurentia, this interval correlates with *Histiodella holodentata* Subzone of the *Periodon macrodentatus* Zone (Stouge, 2012). The studied conodont assemblage is accompanied by sponge

spicules, brachiopods, foraminifers and gastropods.

On the other hand, the argillites of the lower member of the Los Azules Formation contain a rich graptolite association referable to the early middle Darriwilian *Levisograptus dentatus* Zone. It is composed of *Tetragraptus quadribrachiatus* (J. Hall), *Pseudotrigranograptus ensiformis* (J. Hall), *Xiphograptus lofuensis* Lee, *Acrograptus* sp., *Holmograptus bovis* Williams and Stevens, *Isograptus* cf. *victoriae divergens* Harris, *Parisograptus caduceus* (Salter), *Arienigraptus* sp., *Paraglossograptus tentaculatus* (J. Hall), *Levisograptus austrodentatus* (Harris and Keble), *L.* cf. *dentatus* (Brongniart), *L. sinicus* (Mu and Lee) and *L. primus* (Legg). The stratigraphic contact of the lower member with the black-grey shales and siltstones of the middle member is partially splintered and covered. The latter member hosts a significant graptolite fauna corresponding to the *Pterograptus elegans* Zone, just above the deformed levels. The overlying levels are dominated by biserial graptolites that represent the *Hustedograptus teretiusculus* Zone, following Ortega (1987). This way, the middle member is late Darriwilian in age, yet the temporal definition of its lower limit is hampered by the pervasive deformation of the stratigraphic levels. The upper member of the Los Azules Formation includes graptolites of the *Climacograptus bicornis* Zone (late Sandbian) characterized by the presence of the nominal taxon associated with *Nemagraptus gracilis*, *Cryptograptus tricornis* (Carruthers), *Glossograptus ciliatus* (Emmons) and a variety of dicranograptids, and biserial graptolites.

Global Series	G R A P T O L I T E S										CONODONTS				
	Australia		N. America		China		Great Britain		Baltoscandia		Argentine Precordillera		Baltoscandia		
MIDDLE ORDOVICIAN	LATE ORDOV. Sandbian	Gisbornian	calcaratus	americanus	wilsoni	Caradoc	foliaceus	foliaceus	bicornis	tvaerensis	tvaerensis	tvaerensis	tvaerensis		
				bicornis										bicornis	
		Darrwilian	Da4	riddellensis	teretiusculus	teretiusculus	Lianvirn	Li	teretiusculus	vagus	teretiusculus	anserinus	anserinus	anserinus	anserinus
					elegans	jiangxiensis									
			Da3	decoratus	fasciculatus	elegans	Ab	murchisoni	Uh	elegans	elegans	pseudoplanus	pseudoplanus		
					lentus	fasciculatus	As							lentus	
	Da2	intersitus	dentatus	ellesae	austro-	Aren.	Fe	hirundo	Öland	dentatus	crassus	crassus	variabilis	variabilis	
				austro-	sinicus										dentatus
	Da1	austro-	austro-	austro-	austro-	Öland	V. J. Kunda	Uh	austro-	austro-	austro-	austro-	austro-	austro-	

Figure 3. Biostratigraphic chart for conodonts and graptolites of the Argentine Precordillera and correlation with reference schemes [modified from Webby *et al.* (2004), Ortega *et al.* (2007), and Maletz *et al.* (2009)].

ACKNOWLEDGMENTS

This study was funded by CONICET (Resolution 4541-12) and SECyT, Universidad Nacional de Córdoba. The present paper is a contribution to the IGCP Project 591.

REFERENCES

- Albanesi, G.L., Hünicken, M.A. and Barnes, C.R. 1998. Biostratigrafía de conodontes de las secuencias ordovícicas del Cerro Potrerillo, Precordillera Central de San Juan, R. Argentina. *Actas de la Academia Nacional de Ciencias, Córdoba* 12: 7–72.
- Albanesi, G.L., Bergström, S.M., Serra, F., Feltes, N.A., Voldman, G.G. and Ortega, G. (in press). Darriwilian (Middle Ordovician) 13C chemostratigraphy in the Precordillera of Argentina: Documentation of the middle Darriwilian Isotope Carbon Excursion (MDICE) and its use for intercontinental correlation. *Palaeogeography, Palaeoclimatology, Palaeoecology*.
- Astini, R.A. 1994. Análisis secuencial y paleoambientes de las pelitas negras (aloformación Gualcamayo) que suprayacen a las sucesiones carbonáticas eo-ordovícicas en la Precordillera Argentina. *Revista de la Asociación Geológica Argentina* 49: 71–84.
- Astini, R.A., Benedetto, J.L. and Vaccari, N.E. 1995. The early Paleozoic evolution of the Argentine Precordillera as a Laurentian rifted, drifted and collided terrane: A geodynamic model. *Geological Society of America Bulletin* 107: 253–273.
- Borrello, A.V. and Gareca, P.G. 1951. Sobre la presencia de “*Nemagraptus gracilis*” (Hall) en el Ordovícico del norte de San Juan. *Revista de la Asociación Geológica Argentina* 6: 187–193.
- Epstein, A.G., Epstein, J.B. and Harris, L.D. 1977. Conodont color alteration - An index to organic metamorphism. *Geological Survey Professional Paper* 995: 1–27.
- Furque, G. 1979. Descripción Geológica de la Hoja 16c Jáchal. *Servicio Geológico Nacional, Boletín* 164: 1–79.
- Furque, G., González, P. and Caballé, M. 1998. Descripción de la Hoja Geológica 3169-II, San José de Jáchal (Provincias de San Juan y La Rioja). *Servicio Geológico Minero Argentino, Boletín* 259: 1–150.
- Harrington, H.J. and Leanza, A.A. 1957. *Ordovician trilobites of Argentina*. University of Kansas Press, Department of Geology, Special Publication 1, Lawrence, 276 p.
- Hünicken, M.A. 1985. Lower Ordovician conodont biostratigraphy in Argentina. *Boletín Academia Nacional de Ciencias, Córdoba* 56: 309–321.
- Keller, M. 1999. Argentine Precordillera: sedimentary and plate tectonic history of a Laurentian crustal fragment in South America. *Geological Society of America, Special Paper* 341: 1–130.
- Löfgren, A. and Zhang, J. 2003. Element association and morphology in some Middle Ordovician platform-equipped conodonts. *Journal of Paleontology* 77: 721–737.
- Maletz, J., Albanesi, G.L. and Voldman, G.G., 2009. Lower Darriwilian radiolarians from the Argentine Precordillera. *Geobios* 42: 53–61.
- Ortega, G. 1987. [Las graptofaunas y los conodontes de la Formación Los Azules, Cerro Viejo, zona de Huaco, Departamento Jáchal, San Juan. Doctoral Thesis, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba. Unpublished]: 1–210.
- Ortega, G., Albanesi, G.L. and Frigerio, S.E. 2007. Graptolite–conodont biostratigraphy and biofacies of the Middle Ordovician Cerro Viejo succession, San Juan Precordillera, Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology* 245: 245–263.
- Stouge, S. 2012. Middle Ordovician (late Dapingian–Darriwilian) conodonts from the Cow Head Group and Lower Head Formation, western Newfoundland, Canada. *Canadian Journal of Earth Sciences* 49: 59–90.
- Webby, B.D., Cooper, R.A., Bergström, S.M., and Paris, F. 2004. Stratigraphic framework and time slices. In: B.D. Webby, F. Paris, M.L. Droser, and I.G. Percival (Eds.), *The Great Ordovician Biodiversification Event*. Columbia University Press, New York, p. 41–47.
- Zhang, J. 1998. Conodonts from the Guniutan Formation (Llanvirnian) in Hubei and Hunan Provinces, south-central China. *Stockholm Contributions in Geology* 46: 1–161.