

ISCR2 and IS26: Two Insertion Sequences Highly Dispersed among *Acinetobacter* spp. Clinical Strains

Abstract

The aim of this work is to study the dispersion of two ISs, ISCR2 and IS26, which are known to contribute in the acquisition of resistance determinants and genome plasticity, among a collection of hundred and seventy-four *Acinetobacter* spp. clinical isolates. PCR amplification reactions using total DNA were performed to search ISCR2, IS26, and different antibiotic resistance genes (*tet(B)*, *aphA6*, *sul2*, *floR*, *dfr9*) previously described in the genetic context of ISCR2.

Among *A. baumannii*, positive amplification for ISCR2 was obtained in 66% of the included isolates and most of them (93%) were positive for IS26 amplification. The platform *tet(B)::ISCR2* was found in 57% of the ISCR2 positive isolates and only 14 gave negative amplification for *tet(B)*. In these isolates positive amplifications of genes that were previously described associated to ISCR2 -such as, *aphA6*, *sul2*, *floR*, and *dfr9*- were observed. When searching these ISs in the *non-baumannii* *Acinetobacter* studied-collection, two isolates were positive for ISCR2 and two isolates for IS26. No positive results were obtained for *tet(B)*, but the presence of *sul2*, *floR* and *dfr9* was found. Our results exposed a great dispersion of ISCR2 as well as IS26 among *Acinetobacter* spp. clinical isolates reinforcing the idea that the ISs play a crucial role in the plasticity and evolution towards drug resistance in this genus.

Keywords: *Acinetobacter* spp.; ISCR2; IS26; Insertion sequences; Minimum inhibitory concentration

Short Communication

Volume 4 Issue 1 - 2017

Montaña S¹, Almuzara M², Pennini M³,
Sucari A³, Centrón D¹, Vay C² and María
Soledad Ramírez^{4*}

¹Instituto de Microbiología y Parasitología Médica, Argentina

²Laboratorio de Bacteriología, Argentina

³Unidad Microbiología, Argentina

⁴Department of Biological Science, California State University Fullerton, USA

*Corresponding author: María Soledad Ramírez, Assistant Professor, Department of Biological Science California State University Fullerton, Fullerton, CA, USA, Email: msramirez@fullerton.edu

Received: December 26, 2016 | Published: January 26, 2017

Introduction

Acinetobacter are gram-negative, *coccobacilli*, non-fermenting aerobes and can cause a variety of nosocomial infections [1-4]. Currently forty-one distinctive *Acinetobacter* species with unique names and characteristics were described [3, (<http://www.bacterio.net/a/acinetobacter.html>)]. Among this genus, *A. baumannii* is responsible for the majority of nosocomial infections and has an intrinsic ability to acquire and to develop antibiotic resistance determinants to all available antibiotics to treat it [5]. Patient outcomes associated with extensively drug-resistant *A. baumannii* infections can be deadly. The CDC reports 500 deaths annually due to infections with multidrug-resistant *Acinetobacter* (CDC). Recently due to advances in technologies used to identify bacterial species, other members of the *Acinetobacter* genus have been recovered in the clinical setting and have demonstrated resistance to different antibiotics [3,6].

Insertion sequences (ISs) are important for the acquisition and dissemination of antimicrobial resistance determinants, and can also contribute to resistance by insertional inactivation of proteins, such as transcriptional regulators or outer membrane protein [6,7]. More than 30 different types of ISs have been reported in *Acinetobacter* spp. [8] supporting the idea that ISs play a crucial role in the evolution of this genus and that they have contributed to the development of the multidrug-resistant phenotype observed in this genus.

Among the ISs described in *A. baumannii*, most of the studies referred to ISAba1, ISAba125 and ISAba3 since they are associated

with *bla*_{OXA-like} carbapenemases [7]. Among other ISs that have been not deeply studied, we found ISCR2. ISCR are recognized as powerful elements that can capture and mobilize antibiotic resistance genes, as well as, array extended clusters of antibiotic resistance genes on plasmids or on chromosomes [9]. Twenty-one members of the ISCR family have been described [10] and most of them are related with antibiotic resistance [11]. ISCR2 constitutes the second group of ISCR elements and it has been found in many species such as *Vibrio cholerae*, *Shigella flexneri*, *Salmonella enterica*, *Escherichia coli*, *Klebsiella pneumoniae*, *Aeromonas salmonicida*, *Pasteurella piscicida*. ISCR2 have been described on numerous plasmids carrying trimethoprim, tetracycline, chloramphenicol and sulphonamides resistance gene [9,12] and also within resistance island [13-17].

Another important IS is IS26, which was recognized to play an important role in the acquisition and dissemination of genes that confer resistance to many different classes of antibiotics. It has been reported that IS26 plays a relevant role in the genomic plasticity observed in *A. baumannii*. They are involved in building transposons carrying different resistance genes in the fusion of additional transposons into resistance regions and also as a contributor of the variability observed among the genomic resistance island [18].

While conducting a surveillance exploring the dispersion of tetracycline resistance determinants among extensively-drug resistance (XDR) *A. baumannii* strains, we found a high prevalence of the genetic platform *tet(B)::ISCR2* among minocycline resistant strains [19]. ISCR2 has been also described in *Acinetobacter*

spp. associated with *bla*_{VEB1}, suggesting its involvement in the acquisition and the mobilization of a β -lactamase [1]. Moreover, it was found as part of transposon and in *AbaR*-like resistant island close to tetracycline and aminoglycosides resistance genes [13-15]. The aim of the present work was to further explore the dispersion of this particular insertion sequence (ISCR2) among a collection of hundred and seventy-five XDR *Acinetobacter* spp. clinical isolates in order to expose its prevalence and its potential role as an important IS among XDR isolates. Moreover, we also explore the presence IS26 since it is known to have an important role in the plasticity of this species.

A total of 174 *Acinetobacter* spp. isolates were recovered from a variety of clinical sites and samples from individual patients including blood, urine, and respiratory tract, among others, from 12 different hospitals during 2010-2014. One hundred and sixty-four were *A. baumannii*, while the remaining ten were a novel species, *Acinetobacter* spp. A47 strain [2], *A. pittii*, *A. radioresistans*, *A. johnsonii*, *A. ursingii*, *A. guilloiae*, *A. lwoffii*, *A. haemolyticus*, *A. soli* and *A. junii*.

The species of the isolates were confirmed by MALDI-TOF MS (Bruker Daltonik, Bremen, Germany) and *rpoB* amplification and sequencing. The resistance profiles of the isolates to ampicillin, ampicillin/sulbactam, piperacillin/tazobactam, cefalotin, ceftazidime, cefepime, imipenem, meropenem, amikacin, gentamicin, nalidixic acid, ciprofloxacin, nitrofurantoin, colistin and trimethoprim/sulfamethoxazole were determined by disk diffusion according to Clinical Laboratory Standards Institute (CLSI) guidelines or using the VITEK 2 System (bioMérieux, Marcy, L'Étoile, France) employing the panel AST-082 (GNS susceptibility

card) and the minimum inhibitory concentration (MIC) results were interpreted using the CLSI categories.

We extracted total DNA and used it to perform PCR amplification reactions according to the manufacturer's instructions (Promega, Madison, Wisconsin). Specific primers for ISCR2 (ISCR2F: AAGAATTTCTCCAATGCGGG and ISCR2R: GCGGCTCCTTTCCGACAAC) and IS26 (IS26F: GCTGGCTGAACGCGGAG and IS26R: ATACCTTTGATGGTGGC) were used. Moreover, we searched different antibiotic resistance genes (*tet(B)*, *aphA6*, *sul2*, *floR*, *dfr9*) previously described in the genetic context of ISCR2 and we confirm its association in the required cases [9].

To confirm the association of antibiotic resistance genes (*tet(B)*, *aphA6*, *sul2*, *floR*, *dfr9*) with ISCR2 several PCR products were sequenced after purifying the DNA by using the Wizard SV Gel and PCR clean-up System kit according to the manufacturer's directions (Promega, USA). Sequencing was performed on both DNA strands, using an ABI Prism 3100 Bio Analyzer equipment. The nucleotide sequences were analyzed using the Blast V2.0 software (<http://www.ncbi.nlm.nih.gov/BLAST/>).

Conclusion

All *A. baumannii* isolates were categorized as extreme drug-resistant (XDR) according to Magiorakos et al. [20] being resistant to carbapenems and all antibiotics tested except colistin, and in some cases minocycline. Among *non-baumannii* *Acinetobacter* isolates, six were categorized as multi drug-resistant (MDR), and three were susceptible to all antibiotics tested (Table 1).

Table 1: Minimum inhibitory concentrations (MICs) of antimicrobial agents in non-*baumannii* *Acinetobacter* isolates.

Isolate	MIC (mg/L) ^a																
	AMP	SAM	TZP	CEP	FOX	CTX	CAZ	CEF	IPM	MEM	AMK	GEN	NAL	CIP	NIT	COL	SXT
47	≤2	≤2	≤4	≥64	≤4	≤1	≤1	≤1	≤1	≤0.25	≤2	≤1	4	≤0.25	≥512	≤0.5	≤20
23	≥32	≤2	16	≥64	ND	8	4	2	4	≥16	16	≤1	≤2	≤0.25	≥512	≤0.5	160
350	16	≤2	≥128	≥64	ND	32	≥64	8	≤0.25	0.5	≤2	≤1	16	≤0.25	≥512	≤0.5	≤20
306	≤2	≤2	≤4	≤2	≤4	≤1	≤1	≤1	≤1	≤0.25	≤2	≤1	8	≤0.25	32	2	≤20
7	8	≤2	8	≥64	ND	8	8	2	≤0.25	≤0.25	≤2	≤1	4	≤0.25	256	≤0.5	≤20
181	>16	>16/8	>64/4	ND	>16	ND	>16	>16	>8	>8	≤8	≤2	≤8	≤0.125	>128	≤1	1/19
407	>16	≤4/2	≤4/4	ND	>16	ND	16	2	≤1	≤1	≤8	≤2	≤8	0.5	>128	≤1	≤0.5/9.5
432	≤4	≤4/2	≤4/4	ND	8	ND	8	≤1	≤1	≤1	≤8	≤2	≤8	≤0.125	>128	≤1	1/19
476	≤4	≤4/2	≤4/4	ND	>16	ND	4	4	≤1	≤1	≤8	≤2	≤8	≤0.125	>128	≤1	>2/38

^aBold indicates resistance.

AMP: Ampicillin; SAM: Ampicillin/Sulbactam; TZP: Piperacillin/Tazobactam; CEP: Cefalotin; FOX: Cefoxitin; CTX: Cefotaxime; CAZ: Ceftazidime; CEF: Cefepime; IPM: Imipenem; MEM: Meropenem; AMK: Amikacin; GEN: Gentamicin; NAL: Nalidixic Acid; CIP: Ciprofloxacin; NIT: Nitrofurantoin; COL: Colistin; SXT: trimethoprim/sulfamethoxazole; ND, no determined.

Surprisingly, in the *A. baumannii* isolates we obtained positive results for the amplification of ISCR2 in 108 out of 164 (66%). The distribution of IS26 was also high among our isolates since 93% (153/164) of the isolates resulted positive for the amplification of this IS. This last result confirms the previous suggested role of IS26 as a ubiquitous and highly disperse IS [21]. Among the 164 *A. baumannii* isolates included in this study 102 of the isolates were harboring both IS. As we previously found a high association of ISCR2 with *tet(B)* in our *A. baumannii* population, we decided to investigate the presence of the previously reported platform *tet(B)::ISCR2* [19]. We obtained positive results for the association of *tet(B)* with the ISCR2 element in 94 *A. baumannii* isolates (Figure 1). Moreover, 14 of the isolates that were positive for ISCR2 and IS26 gave negative amplification for *tet(B)*. In these 14 isolates we investigated for the presence of other genes (*aphA6*, *sul2*, *floR*, *dfr9*) that were previously described associated to ISCR2. We found that all this isolates were positive for *sul2*, and in most (13/14) of the cases other genes were also present. We found that four isolates were positive for *sul2* and *dfr9*, five were positive for *floR* apart of *sul2*, one was positive for *aphA6*, *sul2* and *dfr9* and another was positive for *floR*, *sul2* and *dfr9* (Figure 1). We were able to determine the association of ISCR2 with genes different than *tet(B)* only in one isolate, we were able to link ISCR2

with *floR*. Moreover, among ISCR2 negative isolates, we found three *tet(B)* positive isolates that were also IS26 negative.

On the other hand, in the *non-baumannii Acinetobacter* isolates we obtained positives results for ISCR2 in two isolates, one of which was positive for *sul2* and IS26 while the other was positive for *sul2* and *floR*. Positives results for IS26 were found in two isolates, for *sul2* in five isolates, for *aphA6* in two isolates, and for *floR* in 1 isolate. No positive results were found for *tet(B)*.

This study shows a great dispersion of ISCR2 as well as IS26 among *Acinetobacter* spp. clinical isolates reinforcing the idea that ISs plays a crucial role in the plasticity and evolution towards drug resistance in this genus. The association of ISCR2 with *tet(B)* is the predominant platform among the *A. baumannii* isolates studied. However, ISCR2 is also present in other *Acinetobacter* spp. isolates where *tet(B)* is not present. We plan to perform further studies to find out the genetic environment of the ISCR2 positive isolates where *tet(B)* is not present. The high prevalence of ISCR2 and IS26 in our *Acinetobacter* spp. isolates allow us to suggest that this element is successfully been maintained in this species, giving a tool that could be involved in the acquisition and dissemination of resistance traits.




Figure 1: Diagram showing the dispersion of ISCR2, IS26 and the most prevalent genes associated to ISCR2 among *A. baumannii* and *A. non baumannii* isolates.

Acknowledgment

M.S.R. and D.C. are members of the career investigator of CONICET, Argentina. S.M. has a Doctoral Fellowship from CONICET. This study was supported by grants UBACyT and PICT 0120 to MSR, Buenos Aires, Argentina.

References

1. Poirel L, Mugnier PD, Toleman MA, Walsh TR, Rapoport MJ, et al. (2009) ISCR2, another vehicle for bla (VEB) gene acquisition. *Antimicrob Agents Chemother* 53(11): 4940-4943.
2. Almuzara M, Traglia GM, Krizova L, Barberis C, Montaña S, et al. (2015) A taxonomically unique *Acinetobacter* strain with proteolytic and hemolytic activities recovered from a patient with a soft tissue injury. *J Clin Microbiol* 53(1): 349-351.
3. Touchon M, Cury J, Yoon EJ, Krizova L, Cerqueira GC, et al. (2014) The genomic diversification of the whole *Acinetobacter* genus: origins, mechanisms, and consequences. *Genome Biol Evol* 8(11): 2866-2882.
4. Turton JF, Shah J, Ozongwu C, Pike R (2010) Incidence of *Acinetobacter* species other than *A. baumannii* among clinical isolates of *Acinetobacter*: evidence for emerging species. *J Clin Microbiol* 48(4): 1445-1449.
5. Peleg AY, de Breij A, Adams MD, Cerqueira GM, Mocali S, et al. (2012) The success of *Acinetobacter* species: genetic, metabolic and virulence attributes. *PLoS One* 7(10): e46984.
6. Roca I, Espinal P, Vila-Farrés X, Vila J (2012) The *Acinetobacter baumannii* oxymoron: commensal hospital dweller turned pan-drug-resistant menace. *Front Microbiol* 3: 148.
7. Mussi MA, Limansky AS, Viale AM (2005) Acquisition of resistance to carbapenems in multidrug-resistant clinical strains of *Acinetobacter baumannii*: natural insertional inactivation of a gene encoding a member of a novel family of beta-barrel outer membrane proteins. *Antimicrob Agents Chemother* 49(4): 1432-1440.
8. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M (2006) IS finder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 34(Database issue): D32-D36.
9. Toleman MA, Bennett PM, Walsh TR (2006) ISCR Elements: novel gene-capturing systems of the 21st century? *Microbiol Mol Biol Rev* 70(2): 296-316.
10. Schleinitz KM, Vallaeyts T, Kleinstaub S (2010) Structural characterization of ISCR8, ISCR22, and ISCR23, Subgroups of IS91-Like Insertion Elements. *Antimicrob Agents Chemother* 54(10): 4321-4328.
11. Boerlin P, Reid Smith RJ (2008) Antimicrobial resistance: its emergence and transmission. *Anim Health Res Rev* 9(2): 115-126.
12. Toleman MA, Bennett PM, Walsh TR (2006) Common regions e.g. orf513 and antibiotic resistance: IS91-like elements evolving complex class 1 integrons. *J Antimicrob Chemother* 58(1): 1-6.
13. Nigro SJ, Post V, Hall RM (2011) The multiresistant *Acinetobacter baumannii* European clone I type strain RUH875 (A297) carries a genomic antibiotic resistance island AbaR21, plasmid pRAY and a cluster containing ISAbal-sul2-CR2-strB-strA. *J Antimicrob Chemother* 66(8): 1928-1930.
14. Seputiene V, Povilonis J, Suziedeliene E (2012) Novel variants of AbaR resistance islands with a common backbone in *Acinetobacter baumannii* isolates of European clone II. *Antimicrob Agents Chemother* 56(4): 1969-1973.
15. Nigro SJ, Hall RM (2011) GIsul2, a genomic island carrying the sul2 sulphonamide resistance gene and the small mobile element CR2 found in the *Enterobacter cloacae* subspecies *cloacae* type strain ATCC 13047 from 1890, *Shigella flexneri* ATCC 700930 from 1954 and *Acinetobacter baumannii* ATCC 17978 from 1951. *J Antimicrob Chemother* 66(9): 2175-2176.
16. Nigro SJ, Hall RM (2012) Tn6167, an antibiotic resistance island in an Australian carbapenem-resistant *Acinetobacter baumannii* GC2, ST92 isolate. *J Antimicrob Chemother* 67(6): 1342-1346.
17. Toleman MA, Walsh TR (2011) Combinatorial events of insertion sequences and ICE in Gram-negative bacteria. *FEMS Microbiol Rev* 35(5): 912-935.
18. Nigro SJ, Farrugia DN, Paulsen IT, Hall RM (2013) A novel family of genomic resistance islands, AbGRI2, contributing to aminoglycoside resistance in *Acinetobacter baumannii* isolates belonging to global clone 2. *J Antimicrob Chemother* 68(3): 554-557.
19. Vilacoba E, Almuzara M, Gulone L, Traglia GM, Figueroa SA, et al. (2013) Emergence and spread of plasmid-borne tet(B)::ISCR2 in minocycline-resistant *Acinetobacter baumannii* Isolates. *Antimicrob Agents Chemother* 57(1): 651-654.
20. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, et al. (2012) Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clinical Microbiology and Infection* 18(3): 268-281.
21. He S, Hickman AB, Varani AM, Siguier P, Chandler M, et al. (2015) Insertion Sequence IS26 reorganizes plasmids in clinically isolated multidrug-resistant bacteria by replicative transposition. *Mbio* 6(3): e00762.