

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

# **The Regioselective Synthesis of Deuterated 4-Alkyl- $\gamma$ -Lactones.**

A thesis presented in partial fulfilment of the requirements  
for the degree of

**Master of Science in Chemistry**

At Massey University.

**Jo-Anna Hislop**

2001

## Corrections

p. 3, para 1, line 7	predominantly (not predominately)
p. 14, para 2, line 3	radical cations should read radical anions
p. 18, para 2, line 5	propargylic (not propagylic)
p. 22, para 1, line 1	propargylic (not propagylic)
p. 20, ref 24	Nevada (not Nervada)
p. 61, ref 1	Nevada (not Nervada)
p. 21, ref 47	The second author's surname is Tramontano
p. 26, para 2, line 6	rationale (not rational)
p. 29, line 6	integration (not intergration)
p. 37, para 2, line 1	replacing water with benzene (not replacing water for benzene)
p. 42, para 2, line 4	stoichiometric (not stiochiometric)
p. 44, para 3, line 6	reaction's (not reactions)
p. 47, para 2, line 1	1- <sup>2</sup> H-decyne is compound (3.20) – not (3.19)
p. 48, line 7	<sup>1</sup> H (superscript 1)
p. 50, para 2, line 14	confirmation (not conformation)
p. 51, para 2, line 3	from the BuLi solution (not for)
p. 56, para 1, line 9	from (not form)
p. 56, para 2, line 3	affected (not effected)
p. 58, para 2, line 1	from should be deleted from this sentence
p. 58, para 4, line 3	unfavourably with (not unfavourable to)
p. 62, title	Comparative (not Comparitive)
p. 69, line 5	equilibria (not equilibra)
p. 70, para 2, line 1	be should be deleted from this sentence
p. 83, para 2, line 6	hexane (not hesane)
p. 87	the footnote belongs on page 86

### *Abstract.*

$\gamma$ -Lactones are important flavour compounds that occur naturally in foodstuffs such as fruit and dairy products. Their presence, or absence, has a considerable influence on the perceived quality of these products. It is therefore important to be able to accurately measure the concentrations of certain  $\gamma$ -lactones. Deuterium labelled compounds offer the possibility of achieving this through stable isotope dilution assays (SIDA). To achieve maximum sensitivity in SIDA of  $\gamma$ -lactones requires two to four deuterium atoms to be placed regioselectively within the lactone ring, where they are retained in the base peak of the labelled analogue upon electron ionisation mass spectroscopy. To this end, two syntheses of regioselectively deuterium ring labelled  $\gamma$ -lactones have been developed.

Initially, three tetradeuterated 2,2,3,3- $^2\text{H}_4$ - $\gamma$ -lactones were prepared. The key step in these syntheses involved the reduction of a doubly protected hydroxy acetylenic acid with deuterium gas in the presence of Wilkinson's catalyst. 2,2,3,3- $^2\text{H}_4$ - $\gamma$ -Dodecalactone was prepared in 25% overall yield with 95% deuterium incorporation, 2,2,3,3- $^2\text{H}_4$ - $\gamma$ -decalactone in 46% yield with 89% deuterium incorporation and 2,2,3,3- $^2\text{H}_4$ - $\gamma$ -octalactone in 36% yield with 90% deuterium incorporation. While higher catalyst loadings (10 mol%) resulted in shorter reduction times, a higher degree of deuterium incorporation was achieved at lower catalyst loadings (5 mol%).

A second, one-pot radical synthesis of ring-labelled  $\gamma$ -lactones was also developed which involved the addition of a two-carbon acetoxy radical to an appropriate 1,1,2-deuterated 1-alkene. This synthesis produced trideuterated  $\gamma$ -lactones without deuterium in the potentially exchangeable position  $\alpha$  to the carbonyl moiety. Extensive GC optimisation for this process was undertaken using unlabelled precursors. A combination of 1-alkene (1.5 mmol), 2-iodoacetamide (0.5 mmol), 1,1'-azo-*bis*-cyclohexanecarbonitrile (radical initiator) (1.0 mmol) and water (50 mmol) in benzene (10 mL) was found to be optimal. The synthesis of 1,1,2- $^2\text{H}_3$ -1-decene, 1,1,2- $^2\text{H}_3$ -1-octene and 1,1,2- $^2\text{H}_3$ -1-hexene was pursued via the reduction of appropriate 1- $^2\text{H}$ -1-alkynes with deuterium gas over Lindlar's

catalyst. This apparently simple transformation, proved difficult due to the volatility of the target 1,1,2-<sup>2</sup>H<sub>3</sub>-1-alkenes. Nonetheless, under the optimised radical conditions, 3,3,4-<sup>2</sup>H<sub>3</sub>- $\gamma$ -dodecalactone was prepared in a 69% isolated yield from 1,1,2-<sup>2</sup>H<sub>3</sub>-1-decene with 96% deuterium incorporation. 3,3,4-<sup>2</sup>H<sub>3</sub>- $\gamma$ -Octalactone was prepared in 17% yield from 1,1,2-<sup>2</sup>H<sub>3</sub>-1-hexene with 92% deuterium incorporation. It is suggested the efficiency of the radical lactonisation process was dependant upon the purity of the 1,1,2-<sup>2</sup>H<sub>3</sub>-1-alkene as under the optimised conditions, unlabelled  $\gamma$ -decalactone and  $\gamma$ -octalactone were prepared in 91% and 94% isolated yields respectively.

Application of the radical strategy for the synthesis of 3,3,4-<sup>2</sup>H<sub>3</sub>-(Z)-6-dodecen- $\gamma$ -lactone was envisaged but the required deuterated precursor was not accessible. The alternative dideuterated 6,7-<sup>2</sup>H<sub>2</sub>-(Z)-6-dodecen- $\gamma$ -lactone was, however, prepared in 75% yield with 96% deuterium incorporation via the partial reduction of 6-dodecyn- $\gamma$ -lactone with deuterium gas over Lindlar's catalyst.

The deuterated lactones prepared in this thesis have since proved of value as internal standards for SIDA.

### *Acknowledgements.*

Firstly a huge thank you to my supervisors;

**Dr. Simon Fielder** for your endless ideas, encouragement, guidance and support throughout the course of my research and writing. Thanks also for all the time you invested and for the “cooking beer”; you’re well dodgy.

**Dr. Dave Harding** for actually convincing me organic chemistry is interesting and for your patience, willingness to help in any way possible and listening to me whinge. Special thanks for the chocolate; it helped lots.

**Dr. Daryl Rowan** for the valued scientific advice and for more often than not lending a new point of view.

Many thanks also to HortResearch, for allowing me the opportunity to study an interesting research area, and for providing the support and resources to make this project possible.

Big thank you to Giovanna (G), where would I be without you and our morning lambington chats? Justin (I can hear a cat!), Rekha for helping me settle into the lab, Paul and Jim. You have all been of great help somehow and contributed to the fun I’ve had both in and out of Massey.

Thanks to members of the porphyrin group; Gav, Wayne and Esther for the odd piece of advice, chemical and help. Also greatly appreciated is the expertise and assistance of Pat Edwards (NMR), John Allen (MS) and Marty Hunt and Adam Matich (GCMS and all sorts of other bits and pieces when required).

Finally my family; Mum, Dad, Mark, Nathan and my Grandparents. Your unconditional support (both financially and emotionally), enthusiasm and encouragement for my studies, has without a doubt been the most significant contribution of all. Thank you.

*Simon and Dave.*

Simon and Dave are not what they seem,  
Their dull appearance is part of their scheme.  
I know of their plans. I know their techniques.  
My supervisors are outer space alien freaks!

They landed on earth in spaceships humongous.  
Posing as chemists they now walk among us.  
My supervisors deny this, but I know the truth.  
They're here to enslave me and spoil my youth.

Early each morning as the sun rises,  
Simon and Dave put on their earthling disguises.  
I knew right away their masks weren't legit.  
Their faces are lined – they sag and don't fit.

The earth's gravity makes them sluggish and slow.  
They say not to run wherever I go.  
They live by the clock. They're slaves to routines.  
They work the year 'round. They're almost machines.

They deny that coffee breaks have much worth.  
They cannot be human. They're not of this earth.  
I cannot escape their alien gaze,  
And they're warping my mind with their alien ways.  
For sinister plots, this ones a gem.  
They're trying to turn me into on of them!

Poem modified from William B. Watterson II.

*Table of Contents.*

	Page
Title page.	i
Abstract.	ii
Acknowledgements.	iv
Poem: Simon and Dave.	v
Table of contents.	vi
Abbreviations.	x
<b>Chapter 1: Introduction.</b>	
1.1 General Introduction.	1
1.2 Measurement Techniques.	3
1.2.1 Isotope Dilution Assays (IDAs).	4
1.2.2 Mass Spectral Requirements.	5
1.2.3 Effects of Deuterium on the NMR Spectra.	7
1.2.3.1 $^1\text{H}$ NMR.	7
1.2.3.2 $^{13}\text{C}$ NMR.	8
1.3 Synthetic routes to $\gamma$ -Lactones.	9
1.3.1 Incorporation of Deuterium via a Radical Strategy.	11
1.3.2 Incorporation of Deuterium via a Reductive Strategy.	15
1.4 Conclusions.	18
References.	19



**Chapter 2: *Synthesis of 2,2,3,3-<sup>2</sup>H<sub>4</sub>-4-Alkyl- $\gamma$ -Lactones.***

2.1	Introduction.	22
2.2	Synthesis of 4-Hydroxydodecanoic Acid.	24
2.3	Synthesis of 2,2,3,3- <sup>2</sup> H <sub>4</sub> - $\gamma$ -Dodecalactone.	25
2.4	Synthesis of 2,2,3,3- <sup>2</sup> H <sub>4</sub> - $\gamma$ -Decalactone.	27
2.5	Synthesis of 2,2,3,3- <sup>2</sup> H <sub>4</sub> - $\gamma$ -Octalactone.	29
2.6	Conclusions.	32
	References.	34

**Chapter 3: *Free Radical Synthesis of 3,3,4-<sup>2</sup>H<sub>3</sub>-4-Alkyl- $\gamma$ -Lactones and a Comparative  $\gamma$ -Lactone Synthesis.***

3.1	Introduction.	35
3.2	The Halogen Transfer Reaction as a route to 3,3,4- <sup>2</sup> H <sub>3</sub> -4-Alkyl- $\gamma$ -Lactones.	36
3.3	Optimisation Reactions.	39
	3.3.1 Reaction Conditions.	39
	3.3.2 First Optimisation Series.	40
	3.3.3 Second Optimisation Series.	41
	3.3.4 Third Optimisation Series.	42
	3.3.5 Final Optimisations.	45
3.4	Deuterium Incorporation.	46
	3.4.1 Synthesis of 1,1,2- <sup>2</sup> H <sub>3</sub> -Alkenes.	47
	3.4.1.1 1,1,2- <sup>2</sup> H <sub>3</sub> -Decene.	47
	3.4.1.2 1,1,2- <sup>2</sup> H <sub>3</sub> -Hexene via 1- <sup>2</sup> H-Hexyne.	48
	3.4.1.2.1 Tetraethyleneglycol Dimethyl Ether as Solvent.	49
	3.4.1.2.2 Undecane as Solvent.	49
	3.4.1.2.2 Tetradecane as Solvent.	50

3.4.1.3	1,1,2- <sup>2</sup> H <sub>3</sub> -Octene.	51
3.4.1.3.1	Method A: n-Butyl lithium as base.	51
3.4.1.3.2	Method B: Sodium hydride as base.	52
3.5	3,3,4- <sup>2</sup> H <sub>3</sub> -4-Alkyl- $\gamma$ -Lactone synthesis via the radical method.	53
3.5.1	3,3,4- <sup>2</sup> H <sub>3</sub> - $\gamma$ -Dodecalactone.	54
3.5.2	3,3,4- <sup>2</sup> H <sub>3</sub> - $\gamma$ -Octalactone.	55
3.5.3	Unlabelled $\gamma$ -Octalactone.	56
3.5.4	Attempted Synthesis of 3,3,4- <sup>2</sup> H <sub>3</sub> - $\gamma$ -Decalactone.	57
3.6	Comparative Procedure: Mn (III) Chemistry.	58
3.7	Conclusions.	60
	References.	61

**Chapter 4: Approaches to 3,3,4-<sup>2</sup>H<sub>3</sub>-(Z)-6-Dodecen- $\gamma$ -lactone and a Comparative  $\gamma$ -Lactone Synthesis.**

4.1	Introduction.	62
4.2	Synthetic approaches to (Z)-6-Dodecen- $\gamma$ -lactone.	63
4.3	Synthesis of (Z)-6-Dodecen- $\gamma$ -lactone.	65
4.3.1	Synthesis of 1-Decen-4-yne.	66
4.3.2	Synthesis of 6-Dodecyn- $\gamma$ -lactone via the Optimised Radical Method.	66
4.3.3	Synthesis of (Z)-6-Dodecen- $\gamma$ -lactone.	67
4.3.4	“Formal Synthesis” of 3,3,4- <sup>2</sup> H <sub>3</sub> -(Z)-6-Dodecen- $\gamma$ -lactone.	68
4.3.5	Synthesis of 6,7- <sup>2</sup> H <sub>2</sub> -(Z)-6-Dodecen- $\gamma$ -lactone.	69
4.4	Comparitive procedure: Mn(III) chemistry.	70
4.5	Conclusions.	71
	References.	72

**Chapter 5: *Experimental.***

5.1	General Experimental.	73
5.2	Experimental for Chapter 2.	75
5.3	Experimental for Chapter 3.	86
5.4	Experimental for Chapter 4.	101
	References.	105

***Appendicies.***

Appendix 3.1	106
--------------	-----

*Abbreviations.*

ACCN	1,1'-azo-bis-cyclohexanecarbonitrile
ACPA	4,4'-azo- <i>bis</i> -4-cyanopentanoic acid
AIBN	2,2'-azo- <i>bis</i> -isobutyronitrile
b.p.	boiling point
br.	broad
BuLi	<i>n</i> -butyl lithium
cat.	catalytic
CI	chemical ionisation
cm <sup>-1</sup>	wave number
°C	degrees Celsius
d	doublet
DMF	dimethylformamide
δ	chemical shift
EI	electron impact
eq.	molar equivalent(s)
eV	electron volts
GC	gas chromatography
GCMS	gas chromatography-mass spectrometry
hr(s)	hour(s)
Hz	Hertz
i.d.	internal diameter
IDA	isotope dilution assay
IR	infrared
J	coupling constant
LDA	lithium diisopropylamide
m	multiplet
min(s)	minute(s)
mmHg	millimetres of mercury

mmol	millimole
mol	mole
MS	mass spectrometry
NMR	nuclear magnetic resonance
$v_{\max}$	absorption maxima (IR)
PhH	benzene
ppm	parts per million
p.s.i.	pounds per square inch
q	quartet
quint	quintet
r.p.m.	revolutions per minute
rt	room temperature
SIDA	stable isotope dilution assay
t	triplet
TBDMS	<i>tert</i> -butyldimethylsilyl
TBDMSCl	<i>tert</i> -butyldimethylsilyl chloride
THF	tetrahydrofuran
tlc	thin layer chromatography
TMEDA	N,N,N',N'-tetramethylethylenediamine