

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Influential Factors in Nectar Composition
and Yield
in
*Leptospermum scoparium***

**A thesis presented in partial fulfilment of the
requirements for the degree of
Doctor of Philosophy in Plant Science**



**Institute of Agriculture and the Environment
College of Sciences, Massey University
Palmerston North, New Zealand**

Elizabeth Mary Nickless

2015

ABSTRACT

Leptospermum scoparium (Mānuka) is the plant nectar source for medically bioactive honey, commercially marketed in New Zealand as Unique Mānuka Factor honey (UMF-honey). Methylglyoxal (MGO) is the unique bioactive component of UMF honey with Mānuka nectar containing significant amounts of the carbohydrate dihydroxyacetone (DHA), the chemical precursor for MGO. Anecdotal evidence and recently published data from nectar samples collected from various cultivars in natural sites or botanical gardens has indicated that the DHA and overall composition of *L. scoparium* nectar varies according to cultivar. The source of this variation is not clearly understood and although there is considerable literature on climatic and genetic influences on nectar composition and yield within various other plant species, there is little published work available on the influence of genetic and environmental factors on the composition and yield of nectar in *L. scoparium*.

Of value to the commercial UMF honey industry in New Zealand is the ability to assess cultivars from breeding programs for the best potential to increase overall UMF honey yield. Predictive modelling of yields is invaluable to the developing honey industry to allow assessment of environmental influences that may affect overall yield along with seasonal influences on nectar production in Mānuka. The research in this thesis establishes the effect of various parameters on overall DHA yield from Mānuka and the beginnings of modelling influencing environmental factors.

To determine influences on dihydroxyacetone (DHA) concentration and yield in the nectar of *L. scoparium* a number of studies were carried out. Methodologies for the collection and analysis of nectar were established. Ten different cultivars of *L. scoparium*

with a range of genetic parentage were studied in controlled glasshouse conditions to assess phenotypic variability in terms of nectar composition and yield as well as plant growth and flowering amongst these cultivars. Significant differences in plant growth and flowering habits were observed amongst the ten cultivars, significant differences in nectar yield and nectar composition with regard to DHA yield were also observed. DHA yields ranged from 2714-7459 mg of DHA/kilogram normalised to 80 °BRIX, with total nectar sugar yields ranging between 0.7 and 4.8 mg amongst the ten cultivars studied. Preliminary research into the effect of temperature, radiation and humidity on nectar composition and yield were also undertaken.

Effects of soil composition on these same parameters were researched with a subset of three of the ten cultivars grown on ten different soil types. Plant relative growth rates, dry weights and total plant height were measured throughout a 15 month glasshouse trial. Plant growth, flowering phenology, floral density, nectar yield and DHA composition data was gathered. Soils were analysed for various macronutrient and micronutrient levels and these parameters were modelled against plant data to determine which soil components were influencing plant parameters of interest. Soil type was shown to have no significant effect on DHA concentrations in nectar but results did show that soil type had a significant effect on flowering density amongst the three *L. scoparium* cultivars studied in the trial. Results from regression analysis of soil chemistry against measured plant parameters indicate that a fertiliser regime has the potential to increase nectar yields due to increased flower numbers. Multivariate analysis using partial least squares regression of soil composition data against plant parameters of value showed that soil components; phosphorus, sulphate, ferric and chloride were commonly shown to influence plant parameters measured.

Analytical spectroscopy was investigated as a method to chemotype *L. scoparium* cultivars and also as a method for quantifying nectar components sucrose, glucose, fructose and DHA.

Nectar composition was analysed using high pressure liquid chromatography (HPLC) and compared with fourier transform Raman spectroscopy (FT-Raman) and attenuated total reflectance infrared spectroscopy (ATR-FTIR) analytical spectroscopy methods.

FT-Raman spectroscopy was shown to be useful in chemotyping cultivars and in addition proved to be a useful analytical method to predict DHA yield using leaf material from *L. scoparium* plants from the ten cultivars. FT-Raman and ATR-FTIR proved to be relatively accurate techniques to quantify *L. scoparium* nectar components DHA, fructose, glucose and sucrose, compared with HPLC methods which use extensive preparation techniques. R-squared values were very good for all nectar components measured excepting the sucrose model at $R^2 = 0.77$. The R^2 for the FT-Raman predictions of DHA against HPLC data are very good at 0.85. FTIR prediction data against HPLC data was also good at 0.86 R^2 . Overall an accurate model is possible for quantifying DHA concentrations in nectar using both FTIR-ATR and FT-Raman spectroscopy.

Overall results show that various factors need to be considered when assessing plants for commercial use in the (UMF) Mānuka honey industry within New Zealand. Due to their large impact on overall nectar yield; floral density and plant growth rate parameters are the two key factors of value for commercial assessment of Mānuka cultivars. This research also highlights the importance of assessing not just DHA concentration in deducing cultivar value, but overall nectar yield. These key features

must be explored when assessing *L. scoparium* plants within breeding programs, prior to selection for large-scale field production of high UMF Mānuka honey.

Acknowledgements

This work was supported by Primary Growth Partnership (PGP) funding (Ministry of Primary Industries, New Zealand) awarded to Mānuka Research Partnership (NZ) Limited (MRPL), and a scholarship from Callaghan Innovation (NZ) to L.M.N. [Contract: MARP1001].

Supervision and staff support from the Institute of Agriculture & Environment, Massey University; Palmerston North, New Zealand.

The Primary Growth Partnership (PGP) scheme brings the resources of many people together into one research program. And I feel privileged to have worked on my thesis under this collaborative scheme. The PGP scheme brings an extra focus to the work and the connection and interaction with the commercial partners (MRPL) in the scheme has been an invaluable and enjoyable experience.

There are many people, Dr Steve Holroyd, Professor Keith Gordon, Dr Chris Anderson, Dr Jason Wargent, Dr Jonathan Stephens, Professor Michael McManus, Dr James Milner and Associate Professor Alastair Robertson, who my sincere thanks go to for their scientific input and support for the research in this thesis. My special thanks and appreciation go to Georgie Hamilton for the technical support she has given me, along with our many enjoyable discussions on various other topics of interest while we worked on the Mānuka program. Additional thanks and much appreciation go to my supervisor Dr Steve Holroyd whose unfailing support encouraged me to attempt this thesis.

There are many others who I owe a debt of gratitude for the help and support with the experimental work in this thesis in particular the staff at the plant growth unit at Massey

University, Lindsey, Leslie and Steve and in addition Chris Rawlingson for his help with the analytical work on the HPLC and Geoff Smith at Otago University for his help with using the Raman equipment. And my employer: Fonterra Research and Development Centre (FRDC) Palmerston North who have allowed me the time from work to work on my thesis, and especially the support from FRDC statisticians Barbara Kuhn-Sherlock and Maree Luckman.

Last thanks go to my family and the support of my partner Daryl and son Henry who have suffered my lack of attention when my evenings were spent focussed on writing and my thoughts often elsewhere focussed on data analysis. Special thanks to my son Henry who helped me measure plants, collect leaves and chop up plants when I needed his help and entertained me with his wonderful wit, humour and patience along the way.

For myself, I have greatly enjoyed working on this thesis, it has been a very enjoyable experience and I have learnt an immense amount along the way and have really enjoyed getting to know all those involved and have greatly appreciated the input and knowledge and experience shared with me.

Thank you!!

Table of Contents

Contents

Title: Influential Factors in Nectar Composition and Yield in *Leptospermum scoparium*

ABSTRACT	i
Acknowledgements	v
Table of Contents	vii
Contents	vii
List of Tables	xii
List of Figures.....	xiv
List of Abbreviations and Symbols	xix
Chapter 1: Introduction	1
1.1 Project Overview	1
1.2 Ministry of Primary Industries Primary Growth Partnership	3
1.3 Research Objectives	5
1.4 Thesis Outline.....	5
1.5 References.....	8
Chapter 2: Literature Review	9
2.1 Overview of <i>Leptospermum scoparium</i>	9
2.2 Overview of Conditions Affecting Vegetative Growth.....	10
2.3 Overview of Commercial Uses of Mānuka	13
2.4 Bioactive Constituents of Mānuka Honey.....	15
2.4.1 Methylglyoxal and Dihydroxyacetone	15
2.5 Nectar Chemistry	18
2.5.1 Nectar Composition	19
2.6 Review of Secondary Metabolism and Metabolites in Plants	20
2.6.1 Metabolomics	21
2.7 Overview of Raman and FTIR Spectroscopic Applications.....	23
2.7.1 FTIR Spectroscopy	24
2.7.2 Raman Spectroscopy.....	26
2.7.3 Application of FTIR and Raman in Analysing Carbohydrates in Plants	28
2.8 Specific Research Objectives.....	29
2.9 References.....	31

Chapter 3: Glasshouse experimental methods	39
3.1 Overview.....	39
3.2 Introduction.....	39
3.2.1 Genotypes	39
3.2.2 Soil Composition.....	40
3.2.3 Experimental Design.....	41
3.3 Plant Breeding History.....	42
3.4 Glasshouse Plant Conditions Experiment 1: Genotype Comparisons.....	49
3.5 Glasshouse Experimental Design	52
3.6 Glasshouse Experiment 2: Soils.....	54
3.6.1 Soil Analysis	58
3.6.2 Soil Chemistry.....	59
3.7 References	62
Chapter 4: Chemometrics: An outline of the development of the chemometric workflow applied in the analysis of Mānuka plant material	63
4.1 Overview.....	63
4.2 Introduction.....	63
4.3 Workflow and Methods	65
4.3.1 Preprocessing Spectral Data.....	65
4.3.2 Baseline Adjustments.....	69
4.3.3 Scatter Correction	71
4.3.4 Standard Normal Variate (SNV).....	71
4.3.5 Detection of Outliers in Spectral Data	73
4.3.6 Use of Derivative Spectra	74
4.4 Multivariate Data Analysis as a Tool to Explore Data	75
4.4.1 Principal Component Analysis (PCA)	75
4.4.2 Analysing Loading Results	81
4.4.3 Explained Variance	84
4.4.4 Prediction Modelling Using Partial Least Squares (PLS).....	85
4.5 Summary.....	91
4.6 References	92
Chapter 5: Genetic and environmental influences on nectar composition and yield in <i>Leptospermum scoparium</i> (Mānuka)	95
5.1 Abstract	96
5.2 Introduction.....	97
5.3 Materials and Methods	99

5.3.1 Plant Material and Growth Conditions.	99
5.3.2 Nectar Collection.....	100
5.3.3 High Pressure Liquid Chromatography (HPLC) Conditions for DHA Analysis.....	101
5.3.4 Environmental Data.	103
5.4 Results and Discussion	104
5.4.1 Floral Density as a Component of DHA Yield.	109
5.4.2 Variability in Plant Growth.....	112
5.4.3 Environmental Influences on Nectar Production.	115
5.5 Acknowledgements.....	120
5.6 References.....	121
Chapter 6: Soil type influences floral density and growth in Mānuka (<i>Leptospermum scoparium</i>).....	125
6.1 Abstract	126
6.2 Introduction	127
6.3 Materials and Methods.....	129
6.3.1 Soil collection and experimental design	129
6.3.2 Nectar Collection.....	132
6.3.3 Analysis of Dihydroxyacetone.....	133
6.3.4 HPLC Conditions.	133
6.3.5 Soil Analysis.....	134
6.3.6 Plant Growth Measurement Methods.....	135
6.3.7 Soil Composition	135
6.3.8 Plant data.	136
6.4 Results and discussion.....	137
6.4.1 Soil Chemistry.....	137
6.4.2 Plant Response to Soil Chemistry	140
6.5 Discussion.....	155
6.6 Acknowledgements.....	157
6.7 References.....	158
Chapter 7: Analytical FT-Raman spectroscopy to chemotype <i>Leptospermum scoparium</i> and generate predictive models for screening for dihydroxyacetone levels in floral nectar	163
7.1 Acknowledgements.....	164
7.2 Abstract	164
7.3 Introduction	165

7.4 Methods	167
7.4.1 Experimental Design.....	167
7.4.2 Multivariate Analysis of the Spectral Data.....	171
7.5 Results and discussion	171
7.5.1 Multivariate Analysis of Spectral Data from <i>L. scoparium</i> Genotypes.	171
7.5.2 Multivariate Partial Least Squares (PLS) Regression.....	178
7.6 Conclusion	180
7.7 References.....	181
Chapter 8: Analytical method development using FTIR-ATR and FT-Raman spectroscopy to assay the carbohydrates; fructose, sucrose, glucose and dihydroxyacetone, in <i>Leptospermum scoparium</i> nectar	187
8.1 Abstract	188
8.2 Introduction.....	189
8.3 Experimental Methods.....	191
8.3.1. HPLC conditions for DHA analysis.	192
8.4. Spectroscopic methods	196
8.5 Results and Discussion	199
8.5.1 Regression Analysis.	200
8.6 Conclusion	203
8.7 Acknowledgements	203
8.8 References.....	204
Chapter 9: Discussion and Future Work.....	209
9.1 Introduction.....	209
9.2 Research Objectives	210
9.3 Discussion	211
9.3.1. Research of available literature to establish standard methodologies for nectar collection and decide on best practises for experimental design for assessing nectar composition, plant growth, flowering and overall nectar yield in <i>L. scoparium</i> (Chapters 3 and 5).	211
9.3.2 Identification of the variance of specific nectar and growth parameters within several cultivars of <i>L. scoparium</i> to establish the potential of breeding programmes for increasing UMF honey yield from <i>L. scoparium</i> . Model the effects of a range of climatic conditions on nectar composition and yield in several cultivars of <i>L. scoparium</i> to allow identification of localities where climatic factors may be beneficial or detrimental to overall UMF honey yield (Chapter 5 and 6).....	212
9.3.3 Identification the effect of the influence of the environment in terms of soil composition on nectar, plant growth and flowering parameters to	

allow decisions to be made regarding best soil sites to establish <i>L. scoparium</i> plantations. (Chapters 5 and 6).....	213
9.3.4 Research into the analytical development of FT-Raman spectroscopy to chemotype cultivars of <i>L. scoparium</i> and investigate the potential for linking metabolites from chemotype profiles produced by FT-Raman data to UMF potential in <i>L. scoparium</i> and test the capability of FT-Raman and ATR-FTIR spectroscopy along with chemometric modelling techniques to quantify nectar components of <i>L. scoparium</i> (Chapters 4, 7 and 8).	216
9.4 Recommendations for Future Work	218
Appendix A	220

List of Tables

Table 2.1. This table shows the classification of plant metabolites into primary and secondary groups.....	21
Table 3.1. <i>Leptospermum scoparium</i> cultivar parentage information	43
Table 3.2. Chemical composition of Long-Term Fertiliser	49
Table 3.3. Chemical composition of Short-Term Fertiliser	50
Table 3.4. Ingredients of Dolomite	50
Table 3.5. Peters Liquid feed formulations.....	52
Table 3.6. Soil Collection Sites with information on soil types with rationale for choosing soils	60
Table 3.7. Soil composition data from site collected soils.....	61
Table 4.1. Example of a data matrix X . Individual objects or samples denoted by n rows and the variables denoted by X columns.....	76
Table 5.1. Data ranges of environmental parameters logged at the time of nectar collection.	107
Table 5.2. Table of means of the main sugars; sucrose, glucose and fructose and DHA in <i>L. scoparium</i> nectar normalised to 80°BRIX.	110
Table 5.3. Potential DHA yield for each cultivar calculated as the product of μg of DHA per flower.....	111
Table 5.4. Growth data parameters measured over 15 months for each cultivar in the glasshouse experiment.	113
Table 5.5. Comparison of the final biomass data between the 10 cultivars	114
Table 5.6. Proposed estimate of potential nectar yield and growth i.e. overall cultivar rating.....	115
Table 5.7. Data ranges of environmental parameters logged at the time of nectar collection.....	116
Table 5.8. Partial Least Squares Regression (PLS) analysis of climatic influences on nectar production and composition.	119

Table 6.1. Description of soil type with codes Soil 1 (S1) to Soil 10 (S10) used for the experimental analysis.	131
Table 6.2. Average macronutrient and micronutrient concentration in the ten soils used for the glasshouse trial.....	139
Table 6.3. Averaged data is displayed for plant response parameters for Cultivar R.	144
Table 6.4. Averaged data is displayed for plant response parameters for Cultivar G.	145
Table 6.5. Averaged data is displayed for plant response parameters for Cultivar Y.....	146
Table 6.6. PLS regression results from plant growth and nectar parameter data of cultivars R, G and Y	152
Table 6.7. Summary of analysis of Variance (ANOVA) to assess the differences between the cultivars in terms of response to measured parameters.	153
Table 6.8. Interaction results for the parameters that showed a significant interaction between the cultivar (CV) and the soil type	154
Table 6.9. Calculated nectar potential (NP) and MSI value for cultivars R and G.	155
Table 8.1. HPLC data showing DHA and sugar component values for the cultivars normalised to 80 °BRIX.	193
Table 8.2. PLS models and Regression analysis results for nectar sugars in <i>L. scoparium</i> using ATR-FTIR and FT-Raman	197

List of Figures

Figure 2.1. <i>Leptospermum scoparium</i> (site=Mt Holdsworth Road, Wairarapa, New Zealand) A = photo showing flowers, B = seed capsules photo showing mature habit of <i>L. scoparium</i> C=ecological habit growing alongside a steam, D = Mānuka plant in peak flowering	10
Figure 2.2. Molecular structure of methylglyoxal	16
Figure 2.3. Concentrations of MGO (■) and sum of methoxylated phenolic components (□) in Mānuka honeys.	17
Figure 2.4. Proposed metabolic pathways for sugar metabolism in floral nectaries.....	19
Figure 2.5. Example of FTIR spectra of three carbohydrate components of the nectar of <i>L. scoparium</i> ;	25
Figure 2.6. Example of FT-Raman spectra of various carbohydrates in solution.	27
Figure 3.1. Floral Density habits of 10 cultivars BS, B, LG, MG, O, P, PU, R, Y and G	43
Figures 3.2-3.11. Photographs of cultivars used in the investigations in this thesis.	44-48
Figure 3.12. Picture showing in-line irrigation system to each individual pot	51
Figure 3.13. Glasshouse plant layout for experiment 1	51
Figure 3.14. Latin Square Design used in experiment 1 with two blocks of a 5 X 5 grid	53
Figure 3.15. In-line watering system for soil experiment and later photos showing plants and flowering in progress	56
Figure 3.16. Experiment 2 mid experiment showing plants growing on the ten different soils	56
Figure 3.17. Experiment 2 mid experiment showing plants in flower during the soils experiment	56
Figure 3.18. Glasshouse layout to investigate the influence of soil type on growth and flowering of <i>Leptospermum scoparium</i>	57
Figure 4.1. Cosmic spikes are visible as very narrow, high relative intensity peaks in spectral data	67

Figure 4.2. Illustration of high frequency noise in spectral data	67
Figure 4.3. Example of spectral data of an organic sample set showing some spectra with fluorescent signal overlaying in the background resulting in a broad band of increased height through the middle region of the spectrum with the FT-Raman peaks still showing above the background band.....	68
Figure 4.3a. Full FT-Raman spectral data set plot from a large range of samples. Note two obvious artefact outliers in the sample set	69
Figure 4.4a. Un-processed FT-Raman spectral data showing vertical shift in the spectral dataset	70
Figure 4.4b. FT-Raman spectral data set after linear baseline processing. Note alignment of most spectral data to the zero baseline	70
Figure 4.5. Camo software data analysis showing MSC scatter effects in spectral data, note the classic fan shape in the descriptive data. This would indicate that application of the MSC transformation may be more useful than SNV	72
Figure 4.6. Camo software data analysis showing data scatter after MSC transformation has been applied. Note that the data is now aligned	72
Figure 4.7. Line plot of spectral data after smoothing and alignment using SNV processing.....	73
Figure 4.8. Hotelling with a T^2 statistical analysis, outliers are displayed above the percentile line (Red line in graph) and can be marked as above for removal from the dataset before recalculating the PCA.....	74
Figure 4.9. Examples of the effect of 1 st and 2 nd derivatives on FTIR-ATR spectral data. A un-derivatised data, B 1 st derivation, C 2 nd derivation.....	75
Figure 4.10. Representation of data points in the variable space. The red line passes through the centre and direction of the largest variance and is PC1 (an eigenvector). T = the t-score value of each object point	78
Figure 4.11. Representation of data points in the variable space. The red line passes through the centre and direction of the largest variance and is PC1.....	79
Figure 4.12. Example of PCA on spectral data. Figure taken from G.P.S. Smith et. al. 2015 “Raman imaging of drug delivery systems”	80
Figure 4.13. Loadings plot of PC3 for the sub-region spectrum (1650–1150 cm^{-1}). The arrows show the wavenumbers with the most influence = the largest coefficient values (-ve or +ve) on PC3	82

Figure 4.14 Loadings plot of PC4 for the sub-region spectrum (1650–1150 cm⁻¹). Regions with the large magnitude coefficients (-ve or +ve) are shown in boxes..... 83

Figure 4.15. Correlation loadings plot. Wavenumbers in the outer ellipse have a higher influence on the corresponding model than wavenumbers in the inner ellipse. Outermost wavenumbers have the highest influence and the relative position in the X and Y axis indicate whether the correlation is positive or negative 84

Figure 4.16. Explained variance PC plot showing how much each PC contributes to the variation in the data set..... 85

Figure 4.17. HPLC data showing DHA values for the cultivars normalised to 80°BRIX. It should be noted that only five out of the seven initial cultivars were used in this model because only five cultivars from which nectar was collected flowered during this experiment 87

Figure 4.18. Explained variance plot for the PLS model 88

Figure 4.19. PLS component plot of factor 3 versus factor 1 separates the P and Y cultivars..... 89

Figure 4.20. Regression graph of PLS model, note the R^2 value of 0.78 and the inclusion of the first three factors for this model 90

Figure 5.1. A: Ternary diagram of the dominant sugars in *L. scoparium* nectar of ten different cultivars. B: PCA score plot of main sugar composition data from the same ten cultivars, separate colours represent different cultivars. ... 105

Figure 5.2. Regression analysis of nectar constituents: fructose versus glucose illustrating the negative correlation between fructose and glucose in *L. scoparium* nectar 108

Figure 5.3. Flowering period in potted plants a) the start of flowering in days numbered from the 1st of January in the main experimental year, b) the average total flowering period for each cultivar..... 112

Figure 6.1. PCA plots and related loading plots of the soil chemistry data for all ten soils. Individual soils are labelled and coloured separately to show soil groupings 138

Figure 6.2. Relative start day of flowering, total days flowering, DHA concentration normalised to 80 °BRIX, numbers of seed capsules, basal stem diameter, plant height, nectar yield and relative growth rates for each cultivar R, G and Y..... 140

Figure 7.1. <i>L. scoparium</i> genotypes used in the experiments. Genotypes are identified by unique codes.....	168
Figure 7.2. LCMS data showing DHA values for the cultivars normalised to 80°BRIX Y=Yellow; P=Pink; O=Orange; MG=Mint Green; B=Blue	170
Figure 7.3. The sub-region 1650–1150cm ⁻¹ of Raman spectra of leaf samples of <i>L. scoparium</i> after linear baseline correction. Y=Yellow; LG=Lime Green; P=Pink; O=Orange; BS=Blue Stripe; MG=Mint Green; B=Blue	173
Figure 7.4. Scores plot (PC3 and PC4) obtained using the sub-region spectrum (1650–1150cm ⁻¹), plots shows the initial separation between leaf spectra of the seven cultivars	173
Figure 7.5. Scores plot (PC2 and PC3) obtained using the same spectral region. Note the further separation of B from BS and MG from Y.	174
Figure 7.6. Loadings plots for PCs 2, 3 and 4. The main peak wavenumbers from these plots are used to indicate what compounds, based on their molecular structure or reference spectra, could be responsible for differentiating the cultivars.....	177
Figure 7.7. Regression graph of the PLS model, note the <i>R</i> ² value of 0.75 and the inclusion of the first three factors for this model	179
Figure 7.8. Graph of score plot from the PLS regression of FT-RAMAN spectra of leaves against DHA levels in the nectar of the flowers from these five cultivars.....	179
Figure 8.1. PCA of the main sugar composition (fructose, glucose and sucrose) of the nectar of the ten cultivars; B, BS, G, LG, O, P, PU, R and Y. Graph A= PCA score plot B = Loadings plot.....	194
Figure 8.2. Comparison of FTIR (A) and FT-RAMAN (B) spectra of 8% DHA, <i>L. scoparium</i> nectar and a composition of mock nectar made from the saccharide solutions fructose and glucose (ratio 2:1).....	195
Figure 8.3. DHA regression model plots from FTIR-ATR (A) and FT-Raman (B) spectral data. Also regression plots of predicted data against HPLC data for FTIR-ATR (C) and FT-Raman (D)	196
Figure 8.4. Prediction data plot from the FT-Raman regression model. Note samples with a larger standard deviation represented by the box around the mean are considered outliers in the prediction i.e. MG-4, O-6, MG-5 and P-5.....	198

Figure 9.1. Illustration representing the interplay of influences on plant response factors relevant to the UMF honey yield 209

Figure A.1. Illustrating the five stages of flower development in *Leptospermum scoparium*, nectar was collected at stage IV 219

Figure A.2. Photograph of a cross section through the middle of a *Leptospermum scoparium* flower. Dark red pigmented area is the hypanthium surface containing the nectaries. The moisture visible on the hypanthium surface is nectar..... 219

List of Abbreviations and Symbols

AAS	atomic absorption spectrophotometry
ACN	Acetonitrile
ANOVA	Analysis of Variance
ATR	Attenuated Total Reflectance (Attenuated Total Reflectance Fourier Transform
ATR-FTIR	(Infrared Spectroscopy
B	Boron
BS	Basal Stem
Ca	Calcium
CCD	Charged Couple Device
CCE	Calcium Carbonate Equivalent
CEC	Cation Exchange Capacity
Cl	Chloride
cm ⁻¹	Spectral Wavenumbers
Co	Cobalt
Cu	Copper
CV	Cultivar
DHA	Dihydroxyacetone
EDTA	Ethylene-diamine-tetra-acetic acid
Fe	Ferric/Iron
FP	Flowering Period
F	Fructose
FTIR	Fourier Transform Infrared Spectroscopy
FT-Raman	Fourier Transform Raman Spectroscopy
g	Grams
GC	Gas Chromatography
GC-FID	Gas Chromatography with Flame Ionization Detector
GLM	General Linear Model

G	Glucose
HA	Hydroxyacetone
HPLC	High Pressure Liquid Chromatography
HT	Plant Height
ICP-OES	Inductively Coupled Plasma Optical Emission Spectrometry
IR	Infrared
K	Potassium
kg	Kilograms
l	Litre
<i>L. scoparium</i>	<i>Leptospermum scoparium</i>
LCMS	Liquid Chromatography Mass Spectrometry
me	milli-equivalents of exchangeable base cation
Mg	Magnesium
mg	Milligrams
MGO	Methylglyoxal
Mj/m ²	Micro-Joules of light energy per meter squared
ml	Milli-litre
mM	Milli-Molar concentration
Mn	Manganese
MRPL	Mānuka Research Partnership Limited
MS	Mass Spectrometry
MSC	Multiplicative Scatter Correction
MSI	Mānuka Soil Index
mW	Milli-Watts
N	Nitrogen
Na	Sodium
NA	Not Applicable
NaOH	Sodium Hydroxide
NIPALS	Non-Linear Iterative Partial Least Squares

NIR	Near Infrared
NMR	Nuclear Magnetic Resonance
NP	Nectar Potential
NPA	Non-Peroxide Antibacterial Activity
P	Phosphate
PC	Principal Component
PCA	Principal Component Analysis
PFBHA	O-(2, 3, 4, 5, 6-pentafluorobenzyl) hydroxylamine
PGP	Primary Growth Partnership
pH	Acidity/Alkalinity level
PLS	Partial Least Squares
PLSR	Partial Least Squares Regression
R	Correlation Value
RGR	Relative Growth Rate
RI	Refractive Index
RMSE	Root Mean Square Error
RMSEC	Root Mean Square Error of Calibration
RMSECP	Root Mean Square Error of the Prediction model.
RMSECV	Root Mean Square Error of the Cross Validated model.
S	Sulphur/Sucrose
SC	Seed Capsule numbers
SD	Start Days = Start of Flowering Period in Days
SNV	Standard Normal Variate
SO ₄	Sulphate
µg	Micro-grams
µL	Micro-litres
UMF	Unique Mānuka Factor
UV/VIS	Ultra Violet/Visual Absorbance Spectra)
UV	Ultra Violet

Y, MG, G, B, BS, P, O, LG, PU and R Cultivar Codes

Yd Yield

Zn Zinc