



UNIVERSITI PUTRA MALAYSIA

***THERAPEUTIC EFFECTS OF *Cosmos caudatus Kunth* LEAF
EXTRACT IN THE PREVENTION AND TREATMENT OF OBESITY
IN SPRAGUE DAWLEY RATS***

HAFEEDZA BINTI ABDUL RAHMAN

FSTM 2015 25



THERAPEUTIC EFFECTS OF *Cosmos caudatus* Kunth LEAF EXTRACT IN THE PREVENTION AND TREATMENT OF OBESITY IN SPRAGUE DAWLEY RATS

By

HAFEEDZA BINTI ABDUL RAHMAN

Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirements for the Degree of Doctor of Philosophy

May 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia,
in fulfillment of the requirement for the degree of Doctor of Philosophy

**THERAPEUTIC EFFECTS OF *Cosmos caudatus* Kunth LEAF
EXTRACT IN THE PREVENTION AND TREATMENT OF OBESITY IN
SPRAGUE DAWLEY RATS**

By

HAFEEDZA BINTI ABDUL RAHMAN

May 2015

Chairman: Prof. Azizah Abdul Hamid, PhD
Faculty : Food Science and Technology

Obesity is a common nutritional disorder that has become one of the most important health issues of modern society around the world. Accumulating studies have shown that various herbs can be good sources of potent antioxidants. However, little information is available on the anti-obesity potential of these herbs. Preliminary study was carried out to investigate the anti-obesity and antioxidant activity of 8 common herbs, namely *Cosmos caudatus*, *Pluchea indica*, *Lawsonia inermis*, *Carica papaya*, *Piper betle*, *Andrographis paniculata*, *Pereskia bleo* and *Melicope lunu*. Anti-obesity activity was assessed using inhibition of pancreatic lipase (PL) and lipoprotein lipase (LPL) activity whereas antioxidant activity was measured using free radical DPPH (2,2-diphenyl-2-picrylhydrazyl) scavenging activity. Results of the preliminary study revealed that *C. caudatus* exhibited good anti-obesity activity as well as excellent antioxidant activity. Therefore, *C. caudatus* was selected for further study by extraction with different concentrations of ethanol (100%, 80%, 60%, 50% and 40%). Hundred percent ethanol extracts of *C. caudatus* showed highest activity in both anti-obesity ($21.8\pm1.5\%$ and $19.9\pm1.1\%$ in pancreatic and lipoprotein lipase assay) and antioxidant activity ($24.9\pm1.1\ \mu\text{g/mL}$) with total phenolic content of $865.8\pm5.0\ \text{mg GAE/g extract}$ and total flavonoid content of $398.8\pm34.8\ \text{mg RE/g extract}$ compared to that of other extracts. Strong positive correlation between antioxidant activity (DPPH) and both phenolic ($r = 0.708$) and flavonoid ($r = 0.766$) content was observed. The same trend existed between anti-obesity (PL and LPL) and that of phenolic ($r = 0.935$, 0.845) and flavonoid ($r = 0.945$, 0.835) compounds respectively. The extract consisted of various flavonoids (quercetin, quercitrin, catechin, epicatechin, kaempherol, rutin, and chlorogenic acid) as identified by nuclear magnetic resonance (NMR), liquid chromatography mass spectroscopy (LC-MS), and high performance liquid chromatography (HPLC). Based on the results obtained, 100% ethanol extract of *C. caudatus* was further assessed for the preventive and therapeutic effects of

obesity *in vivo*. In the preventive study, *C. caudatus* extract (175 mg/kg and 350 mg/kg body weight) was given to the lean *Sprague dawley* rats fed with high fat diet (HFD) for 11 weeks whereas in the treatment study, obese rats were treated with *C. caudatus* extract (200 mg/kg and 400 mg/kg body weight) for 9 weeks. In the preventive study, the extracts significantly suppressed the increase in body weight gain by 33.0 - 42.5% and percentage of abdominal fat by 33.0 - 42.0% when compared to that of control HFD group. However, it failed to reduce the weight and percentage of abdominal fat in HFD induced obese rats. Nevertheless, for both experiments the extract significantly suppressed the increase of plasma triglycerides, total cholesterol, LDL, insulin and leptin level. In addition, plasma ghrelin and adiponectin levels were increased. The extract also increased the fecal excretion of fat in rats (96.9 ± 10.5 - 114.6 ± 9.3 mg/g in preventive study and 87.7 ± 10.0 - 124.6 ± 20.2 mg/g in treatment study) when compared to that of control HFD groups (55.4 ± 6.6 - 58.2 ± 2.6 mg/g) respectively, suggesting that *C. caudatus* reduces the progression of obesity by inhibiting pancreatic lipase, leading to malabsorption of fat, validating the *in-vitro* results obtained in first part of the study. In the final part of the study, both urine and serum metabolites of rats were analyzed using (NMR) spectroscopy and multivariate data analysis (MVDA). Lean and obese rats were clearly discriminated from each other on Orthogonal partial least square (OPLS-DA) score plot proving the ability of the high fat diet used in inducing obesity. Metabolites associated with lipid, tricarboxylic acid cycle (TCA), glucose, amino acid, creatine and gut microbiota metabolism were found to be responsible for the discrimination observed. Interestingly, the therapeutic effects of *C. caudatus* extracts, specifically the low dose (200 mg/kg) were found to be better than that of Orlistat, based on the fact that the metabolic profiles of *C. Caudatus* treated groups were very similar to that of normal group. Betaine, succinate, 3-hydroxybutyrate, creatine, glycine, *N*-acetylglycine, pyruvate and glutamine were significantly increased/decreased towards the normal level. Finally, partial least square analysis (PLS-DA) showed that the obese group moved away from the position of lean group and after 9 weeks of treatment both treated groups were regulated back closer towards their healthy baseline levels, confirming the therapeutic effects of the extracts obtained from biochemical assays measured in the previous chapter. This study showed the anti-obesity effects of *C. caudatus* through inhibition of lipase activity as demonstrated by the increase in fecal fat content and also the positive effects on other obesity biomarkers measured. It also successfully demonstrated the ability of NMR based metabolomics in unraveling therapeutic effects of *C. caudatus* and further provides biochemical insights into the metabolic alterations associated with obesity. Results of the study suggest that *C. caudatus* has potential as a natural supplement or functional ingredient for the prevention and treatment of obesity.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**KESAN TERAPEUTIK EKSTRAK DAUN *Cosmos caudatus* Kunth
DALAM RAWATAN PENCEGAHAN DAN PEMULIHAN OBESITI
TERHADAP TIKUS SPRAGUE DAWLEY**

Oleh

HAFEEDZA BINTI ABDUL RAHMAN

Mei 2015

Pengerusi: Prof. Azizah Abdul Hamid, PhD
Fakulti : Sains dan Teknologi Makanan

Obesiti adalah gangguan pemakanan biasa yang telah menjadi salah satu isu kesihatan yang paling penting dalam masyarakat moden di seluruh dunia. Kajian terkumpul telah menunjukkan bahawa pelbagai herba boleh menjadi sumber antioksidan yang baik. Walau bagaimanapun, hanya sedikit maklumat didapati pada potensi anti-obesiti herba ini. Kajian awal telah dijalankan untuk menyiasat aktiviti anti-obesiti dan antioksidan 8 herba, iaitu *Cosmos caudatus*, *Pluchea indica*, *Lawsonia inermis*, *Carica papaya*, *Piper betle*, *Andrographis paniculata*, *Pereskia bleo* dan *Melicope lunu*. Keputusan kajian menunjukkan bahawa *C. Caudatus* menunjukkan aktiviti anti-obesiti serta antioksidan yang sangat baik. Aktiviti anti-obesiti dinilai menggunakan perencutan lipase pankreas (PL) dan lipase lipoprotein (LPL) aktiviti manakala aktiviti antioksidan dinilai menggunakan pemerangkapan radikal bebas DPPH (2,2-difenil-2-picrilhidrazil). *C. caudatus* telah dipilih untuk kajian lanjut oleh pengekstrakan etanol dengan kepekatan yang berbeza (100%, 80%, 60%, 50% dan 40%). Seratus peratus ekstrak etanol daripada *C. caudatus* menunjukkan aktiviti paling tinggi dalam kedua-dua aktiviti iaitu anti-obesiti ($21.8 \pm 1.5\%$ and $19.9 \pm 1.1\%$ dalam aktiviti pankreas dan lipoprotein lipase) dan antioksidan ($24.9 \pm 1.1\ \mu\text{g/mL}$) dengan jumlah kandungan fenolik sebanyak $865.8 \pm 5.0\ \text{mg GAE/g}$ ekstrak dan jumlah kandungan flavonoid sebanyak $398.8 \pm 34.8\ \text{mg RE/g}$ ekstrak berbanding dengan ekstrak lain. Korelasi positif yang kuat antara aktiviti antioksidan (DPPH) dan kedua-dua kandungan fenolik ($r = 0.708$) dan flavonoid ($r = 0.766$) diperhatikan. Trend yang sama wujud antara anti-obesiti (PL dan LPL) dan juga fenolik ($r = 0.935$, 0.845) dan flavonoid ($r = 0.945$, 0.835). Ekstrak ini terdiri daripada pelbagai flavonoids (kuersetin, kuersitrin, katekin, epikatekin, kaemferol, rutin dan asid klorogenik) dikenal pasti oleh resonans magnetik nuklear (NMR), cecair kromatografi spektroskopi jisim (LC-MS), dan kromatografi cecair berprestasi tinggi (HPLC). Berdasarkan keputusan yang diperolehi, 100% ekstrak etanol daripada *C. caudatus* terus dinilai untuk kesan pencegahan dan terapeutik

obesiti pada tikus *Sprague dawley* diberi makan diet lemak yang tinggi. Dalam kajian pencegahan, ekstrak *C. caudatus* (175 mg/kg dan 350 mg/kg berat badan) telah diberikan kepada tikus normal yang diberi makan dengan diet yang tinggi lemak selama 11 minggu manakala dalam kajian rawatan, tikus obes telah dirawat dengan ekstrak *C. caudatus* (200 mg/kg dan 400 mg/kg berat badan) untuk 9 minggu. Dalam kajian pencegahan, ekstrak berjaya menghalang peningkatan berat badan sebanyak 33.0 - 42.5% dan peratusan lemak sebanyak 33.0 - 42.0% di bahagian abdomen pada kumpulan yang dirawat berbanding kumpulan HFD kawalan. Walau bagaimanapun, ia gagal untuk mengurangkan berat badan dan peratusan lemak di bahagian abdomen tikus obes. Bagi kedua-dua kajian, ekstrak berjaya mengurangkan peningkatan trigliserida, jumlah kolesterol, LDL, insulin dan leptin. Di samping itu, paras grelin dan adiponektin juga meningkat. Ekstrak ini juga meningkatkan perkumuhan lemak dalam tinja tikus yang dirawat (96.9 ± 10.5 - 114.6 ± 9.3 mg/g dalam kajian pencegahan dan 87.7 ± 10.0 - 124.6 ± 20.2 mg/g dalam kajian rawatan) berbanding kumpulan HFD kawalan (55.4 ± 6.6 - 58.2 ± 2.6 mg/g) masing-masing, menunjukkan bahawa *C. caudatus* mengurangkan perkembangan obesiti dengan menghalang aktiviti lipase pankreas, yang membawa kepada kekurangan penyerapan lemak, mengesahkan keputusan in-vitro yang didapati di bahagian pertama kajian. Dalam bahagian akhir kajian ini, kedua-dua urin dan serum metabolit tikus telah dianalisis dengan menggunakan (NMR) spektroskopi dan analisis data multivariat (MVDA). Tikus normal dan obes jelas didiskriminasi antara satu sama lain pada ortogonal separa kurangnya persegi (OPLS-DA), membuktikan keupayaan diet lemak tinggi yang digunakan dalam mendorong obesiti. Metabolit yang berkaitan dengan lipid, kitaran asid trikarboxylic, glukos, asid amino, kreatin dan metabolisma microbiota usus didapati bertanggungjawab untuk diskriminasi yang diperhatikan. Menariknya, kesan terapeutik ekstrak *C. caudatus*, khususnya dos yang rendah (200 mg/kg) didapati lebih baik daripada Orlistat, berdasarkan fakta bahawa profil metabolik tikus yang dirawat *C. caudatus* adalah hampir sama dengan kumpulan normal. Betaine, sukinat, 3-hydroxybutyrate, kreatin, glysin, N-acetylglysin, piruvat dan glutamin telah meningkat/menurun dengan ketara ke arah tahap yang normal. Akhirnya, sebahagian analisis persegi kurangnya (PLS-DA) menunjukkan bahawa kumpulan yang obes beralih daripada kedudukan kumpulan normal dan selepas 9 minggu rawatan kedua-dua kumpulan dirawat bergerak lebih dekat ke arah tahap aras sihat mereka, mengesahkan kesan terapeutik ekstrak diperolehi daripada asai biokimia yang diukur dalam bab sebelumnya. Kajian ini menunjukkan kesan anti-obesiti *C. caudatus* melalui perencutan aktiviti lipase seperti yang ditunjukkan oleh peningkatan dalam kandungan lemak tinja dan juga kesan positif pada penanda biologi obesiti lain yang diukur. Ia juga berjaya memperbaiki keupayaan metabolik berdasarkan NMR untuk menunjukkan kesan terapeutik *C. caudatus* dan seterusnya memberikan pandangan biokimia ke dalam apa-apa perubahan metabolik yang berkaitan dengan obesiti. Keputusan kajian menunjukkan bahawa *C. caudatus* berpotensi sebagai makanan tambahan semula jadi atau bahan berfungsi untuk mencegah dan merawat obesiti.

ACKNOWLEDGEMENTS

First and foremost, I am grateful to Allah S.W.T for having made this project successful and for guiding me through the critical and difficult times.

I would like to express my special appreciation and thanks to my supervisor Professor Dr. Azizah Abdul Hamid, you have been a tremendous mentor for me. I would like to thank you for the encouragement, help and support throughout my research years. Your advice on both research as well as on my life have been priceless.

I am also truly grateful to my committee members, Professor Dr. Amin Ismail, Professor Dr. Nazamid Saari, and Assoc. Prof. Dr. Faridah Abas for serving as my committee members and for their help. I also want to thank all of you for your insightful suggestions and comments upon completing this thesis. I would especially like to thank all lecturers and staff at Faculty of Food Science and Technology.

My warmest thanks also goes to my family. Words cannot express how grateful I am to my parents, brothers and sister for all of the sacrifices that you've made on my behalf. Your prayers was what sustained me this far.

I would also like to thank all of my friends (Najla, Ahmed, Ani and Yana) who helped me in my data analysis and thesis writing, supported me to strive towards completing the thesis.

This thesis would have never been possible without my loving husband Ahmad Bazli. You were always around and helped me at times I thought that it is impossible to continue, thank you for your continued unfailing love, patience and support upon completing this thesis. I couldn't imagine doing my Ph.D without you.

Finally to my son Amir Hariz, the best gift from Allah SWT. I owe you lots and lots of fun hours. My bundle of joy, your smile gave me the courage to overcome the difficulties encountered in my pursuit of this Ph.D. Words cannot describe how grateful I am to both of you.

I certify that a Thesis Examination Committee has met on 11 May 2015 to conduct the final examination of Hafeedza binti Abdul Rahman on her thesis entitled "Therapeutic Effects of *Cosmos caudatus* Kunth Leaf Extract in the Prevention and Treatment of Obesity in Sprague Dawley Rats" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Abdulkarim Sabo Mohammed, PhD

Associate Professor

Faculty of Food Science and Technology

Universiti Putra Malaysia

(Chairman)

Tan Chin Ping, PhD

Professor

Faculty of Food Science and Technology

Universiti Putra Malaysia

(Internal Examiner)

Khozirah binti Shaari, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Internal Examiner)

Young Hae Choi, PhD

Senior Lecturer

Leiden University

Holland

(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 12 August 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Azizah Abdul Hamid, PhD

Professor

Faculty of Food Science and Technology

Universiti Putra Malaysia

(Chairman)

Amin Ismail, PhD

Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Member)

Nazamid Saari, PhD

Professor

Faculty of Food Science and Technology

Universiti Putra Malaysia

(Member)

Faridah Abas, PhD

Associate Professor

Faculty of Food Science and Technology

Universiti Putra Malaysia

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean,

School of Graduate Studies,

Universiti Putra Malaysia

Date:

Declaration by Graduate Student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations, have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institution;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules, 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules, 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia(Research) Rules, 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No: Hafeedza Binti Abdul Rahman (GS27327)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: _____
Name of Chairman of Supervisory Committee: Azizah Abdul Hamid

Signature: _____
Name of Member of Supervisory Committee: Amin Ismail

Signature: _____
Name of Member of Supervisory Committee: Nazamid Saari

Signature: _____
Name of Member of Supervisory Committee: Faridah Abas

TABLE OF CONTENTS

| | Page |
|--|-------|
| ABSTRACT | i |
| ABSTRAK | iii |
| ACKNOWLEDGEMENTS | v |
| APPROVAL | vi |
| DECLARATION | viii |
| LIST OF TABLES | xv |
| LIST OF FIGURES | xvii |
| LIST OF ABBREVIATIONS | xxi |
| LIST OF APPENDICES | xxiii |
| | |
| CHAPTER | |
| 1 GENERAL INTRODUCTION | 1 |
| | |
| 2 LITERATURE REVIEW | 3 |
| 2.1 <i>Cosmos caudatus</i> | 3 |
| 2.1.1 General description of <i>Cosmos caudatus</i> | 3 |
| 2.1.2 Nutritional composition of <i>Cosmos caudatus</i> | 3 |
| 2.1.3 Pharmacological Properties and bioactive compounds of <i>Cosmos caudatus</i> | 4 |
| 2.1.3.1 Antioxidant | 4 |
| 2.1.3.2 Antidiabetic | 5 |
| 2.1.3.3 Antihypertensive | 5 |
| 2.1.3.4 Antifungal, antimicrobial and antimutagen | 6 |
| 2.1.3.5 Antityrosinase | 6 |
| 2.2 Obesity | 6 |
| 2.2.1 Description and prevalence | 6 |
| 2.2.2 Causes of obesity | 7 |
| 2.2.3 Hormones implicated in obesity | 8 |
| 2.2.3.1 Leptin and Insulin | 8 |
| 2.2.3.2 Ghrelin | 9 |
| 2.2.3.3 Adiponectin | 10 |
| 2.2.4 Phytochemicals and obesity | 11 |
| 2.2.5 Obesity and its relation to oxidative stress | 12 |
| 2.3 Treatment for obesity | 13 |
| 2.3.1 Treatment of obesity using Drugs | 13 |
| 2.3.2 Treatment of obesity using plant-originated natural products | 14 |
| 2.3.2.1 Lipase inhibitory effect | 14 |
| 2.3.2.2 Appetite suppressive effect | 16 |
| 2.3.2.3 Stimulation of energy expenditure | 17 |
| 2.3.2.4 Inhibition of adipocyte differentiation | 17 |
| 2.3.2.5 Combination of multiple effects for obesity treatment | 18 |

| | | |
|----------|---|-----------|
| 2.4 | Lipases | 19 |
| 2.4.1 | Digestion and absorption of dietary fats | 19 |
| 2.5 | High fat diet induced obese rat models for obesity research | 21 |
| 2.6 | Metabolomics | 21 |
| 2.6.1 | Application of metabolomics in research on obesity | 22 |
| 2.6.2 | NMR-Based metabolomics | 23 |
| 2.6.3 | Identification of metabolites | 24 |
| 2.6.4 | Quantification of metabolites | 25 |
| 2.6.5 | Multivariate data analysis (MVDA) | 26 |
| 2.6.5.1 | Data processing | 26 |
| 2.6.5.2 | Statistical analysis | 26 |
| 3 | ANTI-OBESITY, ANTIOXIDANT ACTIVITY, PHENOLIC AND FLAVONOID CONTENT OF COSMOS CAUDATUS LEAF EXTRACTED WITH DIFFERENT CONCENTRATION OF ETHANOL | 28 |
| 3.1 | Introduction | 28 |
| 3.2 | Materials and Methods | 29 |
| 3.2.1 | Plant Materials | 29 |
| 3.2.2 | Preparation of extracts | 30 |
| 3.2.3 | Chemicals | 30 |
| 3.2.4 | <i>In-vitro</i> pancreatic lipase (PL) inhibitory assay | 30 |
| 3.2.5 | <i>In-vitro</i> lipoprotein lipase (LPL) inhibitory assay | 31 |
| 3.2.6 | Total phenolic content (TPC) | 32 |
| 3.2.7 | Total flavonoid content (TFC) | 32 |
| 3.2.8 | DPPH radical scavenging assay | 32 |
| 3.2.9 | Extraction and NMR measurement of <i>Cosmos caudatus</i> | 33 |
| 3.2.10 | Flavanoids determination using HPLC | 33 |
| 3.2.11 | Flavanoids determination using LCMS | 34 |
| 3.2.12 | Statistical analysis | 34 |
| 3.3 | Results and discussion | 35 |
| 3.3.1 | Pancreatic lipase (PL) inhibitory Activity | 35 |
| 3.3.2 | Lipoprotein lipase (LPL) Inhibitory Activity | 37 |
| 3.3.3 | Total phenolic content (TPC) | 39 |
| 3.3.4 | Total flavonoid content (TFC) | 41 |
| 3.3.5 | DPPH radical scavenging assay | 42 |
| 3.3.6 | Correlation of bioactive compounds with antioxidant and anti-obesity activity | 42 |
| 3.3.7 | Metabolite identification of <i>Cosmos caudatus</i> from ^1H -NMR spectra | 44 |
| 3.3.8 | Flavanoids determination by HPLC and LCMS | 49 |
| 3.4 | Conclusion | 53 |

| | | |
|----------|---|----|
| 4 | EFFECT OF <i>COSMOS CAUDATUS</i> EXTRACTS IN THE PREVENTION OF BODY WEIGHT GAIN IN LEAN RATS FED A HIGH FAT DIET | 54 |
| 4.1 | Introduction | 54 |
| 4.2 | Materials and Methods | 55 |
| 4.2.1 | Materials | 55 |
| 4.2.2 | Plant Materials | 55 |
| 4.2.3 | Sample preparation | 55 |
| 4.2.4 | Method of extraction | 56 |
| 4.2.5 | Experimental animals | 56 |
| 4.2.6 | Experimental design | 56 |
| 4.2.7 | Administration of the extract | 59 |
| 4.2.8 | Determination of body weight, food intake and energy intake | 59 |
| 4.2.9 | Collection of plasma, liver, lung, kidney, heart, testis, epididymal and perirenal fats, and feces | 59 |
| 4.2.10 | Determination of fecal fat content | 59 |
| 4.2.11 | Determination of plasma lipid parameters | 60 |
| 4.2.12 | Determination of plasma insulin, leptin, adiponectin and ghrelin levels | 60 |
| 4.2.13 | Determination of kidney and liver function test | 60 |
| 4.2.14 | Statistical analysis | 60 |
| 4.3 | Results and discussion | 61 |
| 4.3.1 | Effects of <i>Cosmos caudatus</i> extracts on body weight gain, abdominal fat mass, food and energy intake of rats | 61 |
| 4.3.2 | Effects of <i>Cosmos caudatus</i> extracts on organ weights of rats | 66 |
| 4.3.3 | Effects of <i>Cosmos caudatus</i> extracts on fecal fat content of rats | 68 |
| 4.3.4 | Effects of <i>Cosmos caudatus</i> extracts on plasma lipid profiles of rats | 69 |
| 4.3.5 | Effects of <i>Cosmos caudatus</i> extracts on plasma insulin, leptin, adiponectin and ghrelin levels of rats | 72 |
| 4.3.6 | Effects of <i>Cosmos caudatus</i> extracts on kidney and liver function test of rats | 77 |
| 4.4 | Conclusion | 80 |
| 5 | ANTIOBESITY EFFECTS OF <i>COSMOS CAUDATUS</i> EXTRACT ON HIGH FAT DIET INDUCED OBESE RATS | 81 |
| 5.1 | Introduction | 81 |
| 5.2 | Materials and Methods | 82 |
| 5.2.1 | Materials | 82 |
| 5.2.2 | Plant Materials | 82 |
| 5.2.3 | Sample preparation | 82 |
| 5.2.4 | Method of extraction | 82 |
| 5.2.5 | Experimental animals | 82 |
| 5.2.6 | Experimental design | 82 |

| | | |
|----------|---|-----|
| 5.2.7 | Administration of the extract | 84 |
| 5.2.8 | Determination of body weight, food intake and energy intake | 84 |
| 5.2.9 | Collection of plasma, liver, lung, kidney, heart, testis, epididymal and perirenal fats, and feces | 84 |
| 5.2.10 | Determination of fecal fat content | 84 |
| 5.2.11 | Determination of plasma insulin, leptin, adiponectin and ghrelin levels | 84 |
| 5.2.12 | Determination of plasma lipid profiles | 85 |
| 5.2.13 | Determination of kidney and liver function test | 85 |
| 5.2.14 | Statistical analysis | 85 |
| 5.3 | Results and discussion | 86 |
| 5.3.1 | Effects of <i>Cosmos caudatus</i> extracts on body weights, percent abdominal fat, food and caloric intake of obese rats | 86 |
| 5.3.2 | Effects of <i>Cosmos caudatus</i> extracts on fecal fat content of obese rats | 94 |
| 5.3.3 | Effects of <i>Cosmos caudatus</i> extracts on plasma insulin, leptin, adiponectin and ghrelin levels of obese rats | 95 |
| 5.3.4 | Effects of <i>Cosmos caudatus</i> extracts on lipid profiles of obese rats | 99 |
| 5.3.5 | Effects of <i>Cosmos caudatus</i> extracts on organ weights, liver and kidney function of obese rats | 102 |
| 5.4 | Conclusion | 107 |
| 6 | '¹H-NMR BASED METABOLOMIC ANALYSIS OF URINE AND SERUM FROM HIGH FAT DIET INDUCED OBESE AND COSMOS CAUDATUS TREATED RATS | 108 |
| 6.1 | Introduction | 108 |
| 6.2 | Materials and Methods | 109 |
| 6.2.1 | Materials | 109 |
| 6.2.2 | Methods | 109 |
| 6.2.2.1 | Collection of urine and serum of rats | 109 |
| 6.2.2.2 | ' ¹ H-NMR acquisition of urine and serum | 109 |
| 6.2.2.3 | ' ¹ H-NMR spectral data processing and multivariate data analysis | 110 |
| 6.3 | Results and discussion | 112 |
| 6.3.1 | Multivariate statistical analysis of urine and serum metabolites and identification of biomarker for obesity | 112 |
| 6.3.2 | Effect of <i>Cosmos caudatus</i> extracts on serum metabolites of treated rats | 125 |
| 6.4 | Conclusion | 149 |
| 7 | SUMMARY | 150 |

| | |
|-----------------------------|-----|
| REFERENCES | 153 |
| APPENDICES | 202 |
| BIODATA OF STUDENT | 224 |
| LIST OF PUBLICATIONS | 225 |



LIST OF TABLES

| Table | | Page |
|--------------|--|-------------|
| 2.1 | Scientific classification of <i>C. caudatus</i> | 3 |
| 3.1 | Parameters used in HPLC analysis of <i>C. caudatus</i> extract | 34 |
| 3.2 | Inhibitory effects (%) of different extracts of <i>C. caudatus</i> , Epicatechin and Orlistat on pancreatic lipase at different concentrations | 36 |
| 3.3 | Determination of DPPH, TPC and TFC in different extracts of <i>C. caudatus</i> | 40 |
| 3.4 | Pearson's correlation coefficients (<i>r</i>) between total phenolic content, total flavonoid content, antioxidant capacity, and lipases enzyme inhibitory activities of <i>C. caudatus</i> extracts | 43 |
| 3.5 | ¹ H-NMR chemical shifts (δ) and coupling constants (Hz) of metabolites identified in <i>C. caudatus</i> extract | 46 |
| 4.1 | Composition of normal and high fat diet used | 58 |
| 4.2 | Organ weights of rats treated with <i>C. caudatus</i> extracts for 11 weeks | 67 |
| 4.3 | The effects of <i>C. caudatus</i> extracts on plasma lipid profiles in rats after 11 weeks of treatment | 70 |
| 4.4 | The effects of <i>C. caudatus</i> extracts on plasma insulin, leptin, adiponectin and ghrelin level in obese rats after 11 weeks of treatment | 73 |
| 4.5 | The effects of <i>C. caudatus</i> extracts on liver and kidney function test in rats after 11 weeks of treatment | 78 |
| 5.1 | The effects of <i>C. caudatus</i> extracts on plasma insulin, leptin, adiponectin and ghrelin level in obese rats after 9 weeks of treatment | 96 |
| 5.2 | The effects of <i>C. caudatus</i> extracts on lipid profiles of obese rats after 9 weeks of treatment | 100 |
| 5.3 | The effects of <i>C. caudatus</i> extracts on organ weights of obese rats after 9 weeks of treatment | 103 |

| | | |
|-----|---|-----|
| 5.4 | The effects of <i>C. caudatus</i> extracts on liver and kidney function tests of obese rats after 9 weeks of treatment | 105 |
| 6.1 | ¹ H NMR assignment of the metabolites from <i>Sprague dawley</i> rat's urine and serum | 115 |
| 6.2 | Summary of parameters for assessment of the quality of OPLS-DA models | 117 |
| 6.3 | Summary of urine metabolites differentiating between obese and lean after 11 weeks induction of obesity | 123 |
| 6.4 | Summary of serum metabolites differentiating between obese and lean after 11 weeks induction of obesity | 124 |
| 6.5 | Significant metabolites in rat serum identified by ¹ H NMR and their variations between HFD obese group and ND lean group and treated groups (HFD + 200 mg/kg, HFD + 400 mg/kg and HFD + Orlistat) after 9 weeks of experimental period (n=6 for each group) | 134 |
| 6.6 | Summary of potential marker metabolites in rat serum identified by ¹ H NMR and their variations among groups after 9 weeks of treatment with <i>C. caudatus</i> extract | 137 |

LIST OF FIGURES

| Figure | | Page |
|---------------|---|-------------|
| 3.1 | Inhibitory effects (%) of different extracts of <i>C. caudatus</i> and Epicatechin on lipoprotein lipase (0.1 mg/mL) | 37 |
| 3.2 | Representative of 500 MHz ^1H -NMR spectra of <i>C. caudatus</i> extract from δ 0.50 to 10.0 | 44 |
| 3.3 | Expended 500 MHz ^1H -NMR spectra of <i>C. caudatus</i> extract from δ 6.1 to 7.1 | 45 |
| 3.4 | Expended 500 MHz ^1H -NMR spectra of <i>C. caudatus</i> extract from δ 7.1 to 8.1 | 45 |
| 3.5 | Structure of compounds found in <i>C. caudatus</i> leaf extract | 49 |
| 3.6 | Identification of rutin and quercetin 3-O-rhamnoside in <i>C. caudatus</i> extract by HPLC method | 50 |
| 3.7 | Identification of catechin in <i>C. caudatus</i> extract by HPLC method | 51 |
| 3.8 | Total ion chromatogram of <i>C. caudatus</i> extract | 52 |
| 4.1 | Experimental design for the determination of <i>C. caudatus</i> extract on prevention of obesity in lean rats fed a high fat diet | 57 |
| 4.2 | The effects of <i>C. caudatus</i> extracts on percent body weight gain in rats for 11 weeks of treatment | 62 |
| 4.3 | The effects of <i>C. caudatus</i> extracts on total percent body weight gain in rats | 63 |
| 4.4 | The effects of <i>C. caudatus</i> extracts on percentage of abdominal fats in rats for 11 weeks of treatment. | 64 |
| 4.5 | The effects of <i>C. caudatus</i> extracts on food and energy intake in rats for 11 weeks of treatment | 65 |
| 4.6 | The effects of <i>C. caudatus</i> extracts on fecal fat content in rats for 11 weeks of treatment | 68 |
| 5.1 | Experimental design for the determination on the treatment effects of <i>C. caudatus</i> extract in HFD induced | 83 |

obese rats

| | | |
|-----|--|-----|
| 5.2 | Effects of ND and HFD on body weights of rats during 11 weeks induction of obesity | 86 |
| 5.3 | Initial and final body weight of rats in each group after 9 weeks of treatment | 87 |
| 5.4 | The effects of <i>C. caudatus</i> extracts on body weights of obese rats during 9 weeks of treatment | 88 |
| 5.5 | The effects of <i>C. caudatus</i> extracts on daily food and energy intake in obese rats during 9 weeks of treatment | 89 |
| 5.6 | The effects of <i>C. caudatus</i> extracts on percentage of abdominal fats in obese rats after 9 weeks of treatment | 90 |
| 5.7 | The effects of <i>C. caudatus</i> extracts on fecal fat content in obese rats after 9 weeks of treatment | 94 |
| 6.1 | Representative of 500 MHz ^1H NMR spectra of rat urine from normal and obese groups | 113 |
| 6.2 | Representative of 500 MHz ^1H NMR spectra of rat serum from normal and obese groups | 114 |
| 6.3 | OPLS-DA scores plot derived from rat urine of lean ND (n=9) and obese HFD groups (n=24) after 11 weeks induction of obesity. | 118 |
| 6.4 | OPLS-DA scores plot derived from rat serum of lean ND (n=9) and obese HFD groups (n=17) after 11 weeks induction of obesity | 118 |
| 6.5 | OPLS-DA S-plot derived from rat urine of lean ND (n=9) and obese HFD groups (n=24) after 11 weeks induction of obesity | 120 |
| 6.6 | OPLS-DA S-plot derived from rat serum of lean ND (n=9) and obese HFD groups (n=17) after 11 weeks induction of obesity | 120 |
| 6.7 | OPLS-DA loading plot derived from rat urine of lean ND (n=9) and obese HFD groups (n=24) after 11 weeks induction of obesity | 121 |
| 6.8 | OPLS-DA loading plot derived from rat serum of lean ND (n=9) and obese HFD groups (n=17) after 11 weeks induction of obesity | 121 |

| | | |
|------|--|-----|
| 6.9 | OPLS-DA loading column plot derived from rat urine of lean ND (n=9) and obese HFD groups (n=24) after 11 weeks induction of obesity | 122 |
| 6.10 | OPLS-DA loading column plot derived from rat serum of lean ND (n=9) and obese HFD groups (n=17) after 11 weeks induction of obesity | 122 |
| 6.11 | The overlaid representative 500 MHz ^1H NMR spectra of rat serum from normal, <i>C. caudatus</i> , and Orlistat treated groups | 126 |
| 6.12 | OPLS-DA score plot of rat serum collected from obese HFD, lean ND, <i>C. caudatus</i> and Orlistat treated groups after 9 weeks of treatment | 128 |
| 6.13 | OPLS-DA scores plot derived from rat serum of lean ND (n=6) and obese HFD groups (n=6) after 9 weeks experimental period | 129 |
| 6.14 | OPLS-DA loading plot derived from rat serum of lean ND (n=6) and obese HFD groups (n=6) after 9 weeks experimental period | 129 |
| 6.15 | OPLS-DA scores plot derived from rat serum of obese HFD (n=6) and obese HFD200 groups (n=6) after 9 weeks of treatment with 200 mg/Kg of <i>C. caudatus</i> extract | 130 |
| 6.16 | OPLS-DA loading plot derived from rat serum of obese HFD (n=6) and obese HFD200 groups (n=6) after 9 weeks of treatment with 200 mg/Kg of <i>C. caudatus</i> extract | 130 |
| 6.17 | OPLS-DA scores plot derived from rat serum of obese HFD (n=6) and obese HFD400 groups (n=6) after 9 weeks of treatment with 400 mg/Kg of <i>C. caudatus</i> extract | 131 |
| 6.18 | OPLS-DA loading plot derived from rat serum of obese HFD (n=6) and obese HFD400 groups (n=6) after 9 weeks of treatment with 400 mg/Kg of <i>C. caudatus</i> extract | 131 |
| 6.19 | OPLS-DA scores plot derived from rat serum of obese HFD (n=6) and obese Orlistat groups (n=6) after 9 weeks of treatment with 30 mg/Kg of Orlistat | 132 |
| 6.20 | OPLS-DA loading plot derived from rat serum of obese | 132 |

| | | |
|------|---|-----|
| | HFD (n=6) and obese Orlistat groups (n=6) after 9 weeks of treatment with 30 mg/Kg of Orlistat | |
| 6.21 | PLS-DA score plots of 1H-NMR spectra of rat serum from low dose <i>C. caudatus</i> treated groups (HFD+200 mg/Kg extract) at different time points. Week 0 (Baseline); week 11 (Initial); week 20 (Final) | 138 |
| 6.22 | Permutation test of PLS-DA model from low dose <i>C. caudatus</i> treated groups (HFD + 200 mg/kg extract) at different time points. Week 0 (Baseline); week 11 (Initial); week 20 (Final) | 139 |
| 6.23 | PLS-DA score plots of 1H-NMR spectra of rat serum from high dose <i>C. caudatus</i> treated groups (HFD+400 mg/Kg extract) at different time points. Week 0 (Baseline); week 11 (Initial); week 20 (Final). | 139 |
| 6.24 | Permutation test of PLS-DA model from high dose <i>C. caudatus</i> treated groups (HFD+400 mg/Kg extract) at different time points. Week 0 (Baseline); week 11 (Initial); week 20 (Final). | 140 |

LIST OF ABBREVIATIONS

| | |
|----------------|---|
| ACC | Acetyl-CoA Carboxylase |
| ACE | Angiotensin Converting Enzyme |
| ACUC | Animal Care and Use Committee |
| ALP | Alkaline Phosphatase |
| ALT | Alanine Aminotransferase |
| AMPK | Monophosphate-Activated Protein Kinase |
| ANOVA | Analysis of variance |
| APOC-II | Apolipoprotein C II |
| AST | Aspartate Aminotransferase |
| BAT | Brown Adipose Tissue |
| BHA | Btylated hydroxyanisole |
| BHT | Butylated hydroxytoluene |
| BMI | Basal Metabolic Rate |
| BSA | Bovine Serum Albumin |
| CEBP- α | Cancer Enhancer Binding Protein - Alpha |
| cm | Centimetre |
| CNTF | Ciliary Neurotrophic Factor |
| d | Doublet |
| dd | Doublet of Doublets |
| DPPH | 1,1-diphenyl-2-picrylhydrazyl |
| ECG | Epicatechin Gallate |
| EGC | Epigallocatechin |
| EGCG | Epigallocatechin Gallate |
| EL | Endothelial Lipase |
| FDA | Food and Drug Administration |
| FFA | Free Fatty Acid |
| FTC | Ferric Thiocyanate |
| g | Gram |
| GA | Gallic Acid |
| GAE | Gallic Acid Equivalent |
| GGT | Ggamma-glutamyl Transferase |
| GhR | Ghrelin Receptor |
| GSE | Grape Seed Extract |
| HCA | Hydroxycitric Acid |
| HCL | Hydrochloric Acid |
| HFD | High Fat Diet |
| HL | Hepatic Lipase |
| 1H-NMR | Proton nuclear magnetic resonance |
| HPLC | High Performance Liquid Chromatography |
| Hz | Hertz |
| IC50 | Inhibition Concentration at 50 Percent |
| IDL | Intermediate Density Lipoprotein |
| J | Coupling constant in Hz |
| JAK/STAT | Janus Kinase/Signal Transducer |
| Kg | Kilogram |
| L | Litre |

| | |
|---------------|---|
| LCMS | Liquid Chromatography – Mass Spectroscopy |
| LDL | Low density lipoprotein |
| LPL | Lipoprotein Lipase |
| m | Multiplet |
| mg | Milligram |
| min | Minute |
| mL | Millilitre |
| mL | Micro litre |
| mm | Millimetre |
| mRNA | Messenger Ribose Nucleic Acid |
| MVDA | Multivariate Data Analysis |
| NADPH | Nicotinamide Adenine Dinucleotide Phosphate |
| NAOD | Sodium Deuterioxide |
| ND | Normal Diet |
| NHMS | National Health and Morbidity Survey |
| nm | Nanometer |
| NPY | Neuropeptide Y |
| OPLS-DA | Orthogonal Partial Least Square Data Analysis |
| PAI-1 | Plasminogen Activator Inhibitor Type 1 |
| PC | Principal Component |
| PL | Pancreatic Lipase |
| PLS-DA | Partial Least Square Data Analysis |
| PPAR | Peroxisome Proliferator Activated Receptor |
| PPM | Part Per Million |
| RE | Rutin Equivalent |
| ROS | Reactive oxygen species |
| rpm | Revolution per minute |
| s | Singlet |
| SOD | Superoxide dismutase |
| SPSS | Statistical Package for Social Science |
| TBA | Thiobarbituric Acid |
| TC | Total Cholesterol |
| TFC | Total flavanoid content |
| TG | Triglyceride |
| TNF- α | Tumor Necrosis Factor- α |
| TPC | Total phenolic content |
| TSP | Trimethylsilanepropionic Acid Sodium Salt |
| UV | Ultraviolet |
| VLDL | Very Low Density Lipoprotein |
| WAT | White Adipose Tissue |
| WHO | World Health Organization |
| δ | Chemical Shift in ppm |
| μg | Microgram |
| μm | Micro meter |
| $^{\circ}C$ | Degree Celsius |

LIST OF APPENDICES

| Appendix | | Page |
|----------|---|------|
| A | Standard curves | 202 |
| B | Determination of IC ₅₀ in DPPH assay for standard and samples | 207 |
| C | HPLC optimisation for <i>C. caudatus</i> extract | 209 |
| D | Total ion chromatogram for <i>C.caudatus</i> extract and flavonoids standards | 210 |
| E | Ethics approval | 215 |
| F | Voucher specimen of <i>C. caudatus</i> | 216 |
| G | Approval letter from Taman Pertanian, UPM | 217 |
| H | Picture of rats | 218 |
| I | Loading scatter plot | 221 |

CHAPTER 1

GENERAL INTRODUCTION

Obesity is a leading cause of death worldwide, affecting not only developed but also developing countries. Globally, it is estimated that over 1.4 billion adults are overweight with almost 500 million obese (WHO, 2013). The problem of obesity in Malaysia is at an alarming stage. Reports from the National Health and Morbidity Surveys (NHMS) in 1996, 2006 and 2011 respectively, showed that the prevalence of obesity among adults were increased by three-fold from 4.4% to 14% and 15.1% within 15-years period (NHMS 1996, NHMS 2006, NHMS 2011). Latest study showed that Malaysia is Southeast Asia's most obese country with 49% women and 44% men were either obese or overweight (Ng *et al.*, 2014). Looking at the statistics, scientists are searching for better approach to further understand the diseases and therefore can help in early detection, prevention, and offer solution for effective treatment to this ever-escalating problem.

Malaysia is a tropical country rich with medicinal plants and herbs. In Malaysia and other parts of the world, these medicinal herbs have long been used in the treatment of various ailments. As research on obesity and the use of pharmaceutical drugs in management of obesity is highly controversial and does not provide effective long-term solution, the role of medicinal herbs for treatment of obesity has gained much interest.

Cosmos caudatus, locally known as 'Ulam raja' is a well known herbs found mostly in all tropical regions including Mexico, Central and South America, Malaysia, Thailand and Indonesia (Samy *et al.*, 2005). In Malaysia, it is often consumed raw as salad with rice and food flavouring due to its unique taste and aroma (Shui *et al.*, 2005). Traditionally, it has been used to improve blood circulation, promote the formation of healthy bones, reduce body heat, promote fresh breath, treat infections associated with pathogenic microorganisms, lower high blood pressure and also useful in cleansing the blood (Burkill, 1966; Ismail, 2000; Shui *et al.*, 2005; Hassan 2006; Bodeker, 2009). This herb is also known to possess antioxidant, anti-diabetic, anti-hypertensive, anti-bacteria and anti-fungal properties (Rahalison *et al.*, 1991, Shui *et al.*, 2005, Rasdi *et al.*, 2010, Loh and Hadira, 2011).

Metabolomic approach has been shown to be able to distinguish the different phenotypes and discover potential biomarkers associated with certain phenotypes (Kim *et al.*, 2011). This is made possible with the help of high-throughput tool such as NMR. NMR-based metabolomics can provide a snapshot of metabolites and allow comprehensive metabolite profiling of body fluids. Moreover, it is nondestructive, compatible to both liquid and solid samples and requires no derivatisation with only little sample preparation steps required, making it the perfect tool for collecting a

wealth of data sets (Verpoorte *et al.*, 2007). With these benefits and abilities, metabolomics has become an invaluable tool in finding new biomarkers in obesity research and finally help in identifying mechanisms of obesity. This approach not only identifies, but also quantifies all metabolites in biological system in response to physiological or genetical modifications.

The metabolic effects of polyphenolic rich herbs are commonly assessed in animal studies and clinical trials by measuring plasma or serum concentrations of lipids, glucose, and other biochemical assays. However results are often inconsistent and do not reflect the overall effects of the herbs. Metabolomics provide better understanding on the effects of diet intervention on the metabolism with a holistic approach. This approach will allow for better understanding in terms of the mechanism of action and also help to identify biomarkers of effects that may not be possible with conventional methods (Scalbert *et al.*, 2005, Rezzi *et al.*, 2007). Therefore, to discover biomarkers associated with obesity and *C. caudatus* treatment, both conventional biochemical assays and ^1H -NMR based metabolomics approach was used to study the metabolic changes occurring in the urine and serum of obese rats fed a high fat diet and consequently treated with *C. caudatus* extracts. The combination of both traditional biochemical assay and ^1H -NMR based metabolomics will be able to give the more indepth picture in understanding the disease, evaluate the progression, determining the safety and efficacy of therapeutic interventions and also reveal the potential biomarkers altered with obesity and its treatment.

Hence, the objectives of this study were:

1. To determine the anti-obesity, antioxidant activity, phenolic and flavonoid content of *C. caudatus* extracted with different concentration of ethanol (100, 80, 60, 50 and 40%).
2. To determine the preventive effects of *C. caudatus* leaf extract on lean rats fed a high fat diet.
3. To determine to treatment effects of *C. caudatus* leaf extract on high fat diet induced obese rats.
4. To investigate the metabolic abnormalities in high fat diet induced obese rats and the metabolic alterations associated with the therapeutic effects of *C. caudatus* leaf extract using metabolomics approach.

REFERENCES

- Abas, F., Shaari, K., Lajis, N. H., Israf, D. A., and Umi Kalsom, Y. (2003). Antioxidative and radical scavenging properties of the constituents isolated from *Cosmos caudatus* Kunth. *Natural Product Science*, 9(4), 245-248.
- Abas, F., Nordin, H.L., Israf, D.A., Khozirah, S., and Umi Kalsom, Y. (2006). Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chemistry*, 95(4), 566-573.
- Abdollahi, M., and Afshar-Imani, B. (2003). A review on obesity and weight loss measures. *Middle East Pharmacy*, 11(5), 6-10.
- Addai, Z.R., Abdullah, A., and Abd. Mutualib, S. (2013). Effect of the extraction solvents on the phenolic content and antioxidant properties of two papaya cultivars. *Journal of medicinal plants research*, 7(47), 3354-3359.
- Afonso, M.S., De O Silva, A.M., Carvalho, E.B., Rivelli, D.P., Barros, S.B., Rogero, M.M., Lottenberg, A.M., Torres, R.P., and Mancini-Filho, J. (2013). Phenolic compounds from Rosemary (*Rosmarinus officinalis* L.) attenuate oxidative stress and reduce blood cholesterol concentrations in diet-induced hypercholesterolemic rats. *Nutrition and Metabolism (London)*, 10(1), 19.
- Ahn, J., Lee, H., Kim, S., Park, J., and Ha, T. (2008). The anti-obesity effect of quercetin is mediated by the AMPK and MAPK signaling pathways. *Biochemical and Biophysical Research Communications*, 373(4), 545-549.
- Akase, T., Shimada, T., Terabayashi, S., Ikeya, Y., Sanada, H., and Aburada, M. (2011). Antiobesity effects of *Kaempferia parviflora* in spontaneously obese type II diabetic mice. *Journal of Natural Medicines*, 65(1), 73-80.
- Al Awar, R., Obeid, O., Hwalla, N., and Azar, S. (2005). Postprandial acylated ghrelin status following fat and protein manipulation of meals in healthy young women. *Clinical Science (London)*, 109(4), 405- 411.
- Al-Waiz, M., Mikov, M., Mitchell, S.C., and Smith, R.L. (1992). The exogenous origin of trimethylamine in the mouse. *Metabolism*, 41(2), 135-136.
- Ala-Korpela, M. (2008). Critical evaluation of ^1H NMR metabonomics of serum as a methodology for disease risk assessment and diagnostics. *Clinical Chemistry and Laboratory Medicine*, 46(1), 27-42.

- American Psychiatric Publishing, (APA). (2000). Eating Disorders, In Diagnostic And Statistical Manual Of Mental Disorders. Fourth Edition, American Psychiatric Publishing, Arlington, VA.
- Ames, B.N. Shigenaga, M.K. Hagen, T.M. (1993). Oxidants, antioxidants and the degenerative diseases of aging. *Proceedings of the National Academic of Science of the United States of America*, 90(17), 7915-7922.
- Amin, K.A., and Nagy, M.A. (2009). Effect of Carnitine and herbal mixture extract on obesity induced by high fat diet in rats. *Diabetology and Metabolic Syndrome*, 1(1), 17.
- Andersen, R.E., Wadden, T.A., Bartlett, S.J., Vogt, R.A., and Weinstock, R.S. (1995). Relation of weight loss to changes in serum lipids and lipoproteins in obese women. *American Journal of Clinical Nutrition*, 62(2), 350-357.
- Aoki, F., Honda, S., Kishida, H., Kitano, M., Arai, N., Tanaka, H., Yokota, S., Nakagawa, K., Asakura, T., Nakai, Y., and Mae, T. (2007). Suppression by licorice flavonoids of abdominal fat accumulation and body weight gain in high-fat diet-induced obese C57BL/6J mice. *Bioscience Biotechnology and Biochemistry*, 71(1), 206-214.
- Arifin, N. (2005). Penyembuhan Semula Jadi Dengan Herba. PTS Litera Utama. 1st edition, 252 Pages.
- Arita, Y., Kihara, S., Ouchi, N., Takahashi, M., Maeda, K., Miyagawa, J., Hotta, K., Shimomura, I., Nakamura, T., Miyaoka, K., Kuriyama, H., Nishida, M., Yamashita, S., Okubo, K., Matsubara, K., Muraguchi, M., Ohmoto, Y., Funahashi, T., Matsuzawa, Y. (1999). Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochemical and Biophysical Research Communications*, 257(1), 79-83.
- Arthat, S.M., Lavie, C.J., Milani, R.V., and Ventura, H.O. (2008). The obesity paradox: impact of obesity on the prevalence and prognosis of cardiovascular diseases. *Postgraduate Medicine*, 120(2), 34–41.
- Arts, I.C.W., and Hollman, P.C.H. (2005). Polyphenols and disease risk in epidemiologic studies. *American Journal of Clinical Nutrition*, 81, 317-325.
- Asarian, L. and Geary, N. (2006). Modulation of appetite by gonadal steroid hormones. *Philos Trans R Soc Lond B Biol Sci*, 361(1471), 1251-1263.

- Attele, A.S., Zhou, Y.P., Xie, J.T., Wu, J.A., Zhang, L., Dey, L., Pugh, W., Rue, P.A., Polonsky, K.S., Yuan, C.S. (2002). Antidiabetic effects of Panax ginseng berry extract and the identification of an effective component. *Diabetes*, 51(6), 1851-1858.
- Azman, K.F., Amom, Z., Azlan, A., Esa, N.M., Ali, R.M., Shah, Z.M., and Kadir, K.K. (2011). Antiobesity effect of Tamarindus indica L. pulp aqueous extract in high-fat diet-induced obese rats. *Journal of Natural Medicines*, 66(2), 333-342.
- Backman, L., Hallberg, D., and Kallner, A. (1975). Amino acid pattern in plasma before and after jejuno ileal shunt operation for obesity. *Scandinavian Journal of Gastroenterology*, 10(8), 811-816.
- Bae, H., Jayaprakasha, G.K., Crosby, K., Jifon, J.L., and Patil, B.S. (2012). Influence of extraction solvents on antioxidant activity and the content of bioactive compounds in non-pungent peppers. *Plant Foods Human Nutrition*, 67(2), 120-128.
- Bamba, V., and Rader, D.J. (2007). Obesity and atherogenic dyslipidemia. *Gastroenterology*, 132(6), 2181-2190.
- Barness, L.A., Opitz, J.M., and Gilbert-Barness, E. (2007). Obesity: genetic, molecular, and environmental aspects. *American Journal of Medical Genetics*, 143(24), 3016-3034.
- Barr, J., Vazquez-Chantada, M., Alonso, C., Perez-Cormenzana, M., Mayo, R., Galan, A., Caballeria, J., Martin-Duce, A., Tran, A., Wagner, C., Luka, Z., Lu, S.C., Castro, A., Le Marchand Brustel, Y., Martinez Chanter, M.L., Veyrie, N., Clement, K., Tordjman, J., Gual., P and Mato, J.M. (2010) Liquid chromatography-mass spectrometry-based parallel metabolic profiling of human and mouse model serum reveals putative biomarkers associated with the progression of nonalcoholic fatty liver disease. *Journal of Proteome Research*, 9(9), 4501-4512.
- Barron, J.T., Barany, M., Gu, L., and Parrillo, J.E. (1998). Metabolic fate of glucose in vascular smooth muscle during contraction induced by noradrenaline. *Journal of Molecular and Cellular Cardiology*, 30(3), 709-719.
- Bednarek, M.A., Feighner, S.D., Pong, S.S., Mc Kee, K.K., Hreniuk, D.I., Silva M.V., Warren V.A., Howard A.D., Van Der Ploeg L.H., Heck J.V. (2000). Structure-function studies on the growth hormone-releasing peptide, ghrelin: minimal sequence of ghrelin necessary for activation of growth hormone secretagogue receptor. *Journal of Medicine and Chemistry*, 43(23), 4370-4376.

- Berg, A.H., Combs, T.P., Du, X., Brownlee, M., and Scherer, P.E. (2001). The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Natural Medicines*, 7(8), 947-953.
- Birari, R.B., and Bhutani, K.K. (2007). Pancreatic lipase inhibitors from natural sources: unexplored potential. *Drug Discovery Today*, 12(19-20), 879-889.
- Bjerrum, J.T., Nielsen, O.H., Wang, Y.L., and Olsen, J. (2008). Technology Insight: metabonomics in gastroenterology—basic principles and potential clinical applications. *Nature Clinical Practice Gastroenterology and Hepatology*, 5(6), 332-343.
- Bodeker, G. (2009). Health and Beauty from the Rainforest: Malaysian Traditions of Ramuan. Kuala Lumpur: Didier Millet. ISBN 978-981-4217-91-0.
- Bogdanov, M., Matson, W.R., Wang, L., Matson, T., Saunders Pullman, R., Bressman, S.S., and Beal, M. (2008). Metabolomic profiling to develop blood biomarkers for Parkinson's disease. *Brain*, 131(2), 389-396.
- Bose, M., Lambert, J., Ju, J., Reuhl, K., Shapes, S., and Yang, C. (2008). The major green tea polyphenol, (-)-epigallocatechin-3-gallate, inhibits obesity, metabolomic syndrome, and fatty liver disease in high fat fed mice. *Journal of Nutrition*, 138(9), 1677-1683.
- Bothwell, J.H., and Griffin, J.L. (2011). An introduction to biological nuclear magnetic resonance spectroscopy. *Biological Review of the Cambridge Philosophical Society*, 86(2), 493-510.
- Bracco, D., Ferrarra, J.M., Arnaud, M.J., Jequier, E., and Schutz, Y. (1995). Effects of caffeine on energy metabolism, heart rate, and methylxanthine metabolism in lean and obese women. *American Journal Physiology Endocrinology and Metabolism*, 269(4), 671-678.
- Brand-Williams, W., Bondent, V., and Bereset, C. (1995). Kinetics and mechanism of antioxidant activity using the DPPH. free radical methods. *Lebensmittel Wissenschaft und Technologiy*, 30(6), 609-615.
- Bray, G.A., and Popkin, B.M. (1998). Dietary fat intake does affect obesity! *American Journal of Clinical Nutrition*, 68(6), 1157-1173.
- Burkill, I.H. (1966). A dictionary of the economic products of the Malay Peninsula (Vols. 1 and 2). Kuala Lumpur, Malaysia: Ministry of Agriculture and Cooperative.

- Buss, J., Havel, P.J., Epel, E., Lin, J., Blackburn, E., Daubenmier, J. (2014). Associations of ghrelin with eating behaviors, stress, metabolic factors, and telomere length among overweight and obese women: preliminary evidence of attenuated ghrelin effects in obesity?. *Appetite*, 76, 84-94.
- Butterfield, D.A. (2002). Amyloid b-peptide (1–42)-induced Oxidative Stress and Neurotoxicity: Implications for Neurodegeneration in Alzheimer's Disease Brain. *Free Radical Research*, 36(12), 1307-1313.
- Cai, S., Wang, O., Wang, M., He, J., Wang, Y., Zhang, D., Zhou, F., and Ji, B. (2012). In vitro inhibitory effect on pancreatic lipase activity of subfractions from ethanol extracts of fermented Oats (*Avena sativa* L.) and synergistic effect of three phenolic acids. *Journal of Agriculture and Food Chemistry*, 60(29), 7245-7251.
- Cairns, E. (2005) Obesity: the fat lady sings? *Drug Discovery Today*, 10(5), 305-307.
- Calvani, R., Miccheli, A., Capuani, G., Tomassini Miccheli, A., Puccetti, C., Delfini, M., Iaconelli, A., Nanni, G., and Mingrone, G. (2010). Gut microbiome-derived metabolites characterize a peculiar obese urinary metabotype. *International Journal of Obesity*, 34(6), 1095-1098.
- Carey, M., Small, D., and Bliss, C. (1983). Lipid digestion and absorption. *Annual Review Physiology*, 45(1), 651-677.
- Carriere, F., Barrowman, J.A., Verger, R., and Laugier, R. (1993). Secretion and contribution to lipolysis of gastric and pancreatic lipases during a test meal in humans. *Gastroenterology*, 105(3), 876-888.
- Cevallos-Cevallos, J.M., De-Corcuera, J.R., Etxeberria, E., Danyluk M.D., and Rodrick, G.E. (2009). Metabolomic analysis in food science: a review. *Trends in Food Science and Technology*, 20(11), 557-566.
- Chagnac, A., Weinstein, T., Korzets, A., Ramadan, E., Hirsch, J., and Gafter, U. (2000). Glomerular hemodynamics in severe obesity. *American Journal of Physiology*, 278(5), 817-822.
- Chakrabarti, R. (2009). Pharmacotherapy of obesity: Emerging drugs and targets. *Expert Opinion on Therapeutic Targets*, 13, 195-207.
- Chandran, M., Phillips, S.A., Ciaraldi, T., and Henry, R.R. (2003). Adiponectin: more than just another fat cell hormone? *Diabetes Care*, 26(8), 2442-2450.

- Chantre, P., and Lairon, D., (2002). Recent findings of green tea extract A R25 (exolise) and its activity for the treatment of obesity. *Phytomedicine*, 9(1), 3-8.
- Chao, A.S., Jeon, S.M., Kim, M.J., Yeo, J., Seo, K., Choi, M.S., and Lee, M.K. (2010). Chlorogenic acid exhibits anti-obesity property and improves lipid metabolism in high fat diet induced obese mice. *Food and Chemical Toxicology*, 48(3), 937-943.
- Charles, R.A., Goh, S.Y. (2007). Point-of-care blood ketone testing: screening for diabetic ketoacidosis at the emergency department. *Singapore Medical Journal*, 48(11), 986-989.
- Chen, H.C., and Farese, R.V. (2002). Determination of adipocyte size by computer image analysis. *Journal of Lipid Research*, 43(6), 986-989.
- Chen, S.W., Chen, Y.X., Shi, J., Lin, Y., and Xie, W.F. (2006). The restorative effect of taurine on experimental nonalcoholic steatohepatitis. *Digestive Disease and Sciences*, 51(12), 2225–2234.
- Cho, A.S., Jeon, S.M., Kim, M.J., Yeo, J., Seo, K.I., Choi, M.S., and Lee, M.K. (2010). Chlorogenic acid exhibits anti-obesity property and improves lipid metabolism in high-fat diet-induced-obese mice. *Food Chemistry and Toxicology*, 48(3), 937-943.
- Choi, Y.H., Kim, H.K., Linthorst, H.J., Hollander, J.G., Lefeber, A.W., Erkelens, C., Nuzillard, J.M., and Verpoorte, R. (2006). NMR metabolomics to revisit the tobacco mosaic virus infection in Nicotiana tabacum leaves. *Journal of Natural Product*, 69(5), 742-748.
- Choi, H., Eo, H., Park, K., Jin, M., Park, E.J., Kim, S.H., Park, J.E., and Kim, S. (2007). A water-soluble extract from Cucurbita moschata shows anti-obesity effects by controlling lipid metabolism in a high fat diet- induced obesity mouse model. *Biochemical Biophysical Research Communications*, 359(3), 419-425.
- Choi, I., Kim, Y., Park, Y., Seog, H., Choi, H. (2007). Anti-obesity activities of fermented soygerm isoflavones by *Bifidobacterium breve*. *Biofactors*, 29(2-3), 105-112.
- Choo, J.J. (2003). Green tea reduces body fat accretion caused by high-fat diet in rats through beta-adrenoceptor activation of thermogenesis in brown adipose tissue. *Journal of Nutritional Biochemistry*, 14(11), 671-676.

- Chung, S., and Scanu, A.M. (1974). Continues pH stat titration method for the assay of lipoprotein lipase activity in vitro. *Analytical Biochemistry*, 62(1), 134-148.
- Coen, M., Holmes, E., Lindon, J.C., and Nicholson, J.K. (2009). NMR-based metabolic profiling and metabonomic approach to problems in molecular toxicology. *Chemical Research in Toxicology*, 21(1), 9-27.
- Conforti, F., Perri, V., Menichini,F., Marrelli,M., Uzunov,D., Statti, G.A., and Menichin, F. (2012). Wild Mediterranean Dietary Plants as Inhibitors of Pancreatic Lipase. *Phytotherapy Research*, 26(4), 600–604.
- Craig, S. A. (2004). Betaine in human nutrition American. *Journal of Clinical Nutrition*, 80(3), 539-549.
- Craig, A., Cloarec, O., Holmes, E., Nicholson, J.K., and Lindon, J.C. (2006). Scaling and normalization effects in NMR spectroscopic metabonomic data sets. *Analytical Chemistry*, 78(7), 2262-2267.
- Crozier, A., Jensen, E., Lean, M.E.J., and McDonald, M.S. (1997). Quantitative analysis of flavonoids by reversed phase high-performance liquid chromatography. *Journal of Chromatography A*, 761(1–2), 315–321.
- Cui, Q., Lewis, I.A., Hegeman, A.D., Anderson, M.E., Li, J., Schulte, C.F., Westler, W.M., Eghbalnia, H.R., Sussman M.R., and Markley, J.L., (2008). Metabolite identification via the Madison Metabolomics Consortium Database. *Nature Biotechnol*, 26(2), 162-164.
- Cummings, D.E., Purnell, J.Q., Frayo, R.S., Schmidova, K., Wisse, B.E., and Weigle, D.S. (2001). A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes*, 50(8), 1714-1719.
- Dai, J., and Mumper, R.J. (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.
- Dattilo A.M., and Kris-Etherton, P.M. (1992). Effects of weight reduction on blood lipids and lipoproteins: a meta analysis. *American Journal of Clinical Nutrition*, 56(2), 320-328.
- Daykin, C.A., Van Duynhoven, J.P., Groenewegen, A., Dachtler, M., Van Amelsvoort, J.M., and Mulder, T.P.J. (2005). Nuclear magnetic resonance spectroscopic based studies of the metabolism of black tea polyphenols in humans. *Journal of Agricultural and Food Chemistry*, 53(5), 1428-1434.

- De Groote, D., Van Belleghem, K., Devière, J., Van Brussel, W., Mukaneza, A., Amininejad, L. (2012). Effect of the intake of resveratrol, resveratrol phosphate and catechin rich grape seed extract on markers of oxidative stress and gene expression in adult obese subject. *Annals of Nutrition and Metabolism*, 61(1), 15-24.
- De la Garza, A.L., Milagro, F.I., Boque, N., Campion, J., and Martinez, J.A. (2011). Natural inhibitors of pancreatic lipase as new players in obesity treatment. *Planta Medica*, 77(8), 773-785.
- De Simone, G., and D'Addeo, G. (2008). Sibutramine: balancing weight loss benefit and possible cardiovascular risk. *Nutrition, Metabolism and Cardiovascular Diseases*, 18(5), 337-341.
- Décordé, K., Teissèdre, P.L., Sutra, T., Ventura, E., Cristol J.P., and Rouanet, J.M. (2009). Chardonnay grape seed procyanidin extract supplementation prevents high-fat diet-induced obesity in hamsters by improving adipokine imbalance and oxidative stress markers. *Molecular Nutrition and Food Research*, 53(5), 659-666.
- Dingemans, A.E., Bruna, M.J., and van Furth, E.F. (2002). Binge eating disorder: a review. *International Journal of Obesity and Related Metabolic Disorders*, 26(3), 299-307.
- Drew, B.S., Dixon, A.F., and Dixon, J.B. (2007). Obesity management: update on orlistat. *Vascular Health and Risk Management*, 3(6), 817-821.
- Duggan, G.E., Hittel, D.S., Hughey, C.C., Weljie, A., Vogel, H.J., and Shearer, J. (2011). Differentiating short- and long-term effects of diet in the obese mouse using ¹H-nuclear magnetic resonance metabolomics. *Diabetes, Obesity and Metabolism*, 13(9), 859–862.
- Dulloo, A.G., Duret, C., Rohrer, D. (1999). Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *American Journal of Clinical Nutrition*, 70(6), 1040–1045.
- Dumas, M.E., Barton, R.H., Toye, A., Cloarec, O., Blancher, C., Rothwell, A., Fearnside, J., Tatoud, R., Blanc, V., Lindon, J.C., Mitchell, S.C., Holmes, E., McCarthy, M.I., Scott, J., Gauguier, D., and Nicholson, J.K. (2006). Metabolic profiling reveals a contribution of gut microbiota to fatty liver phenotype in insulin-resistant mice. *Proceedings of the National Academic of Science of the United States of America*, 103(33), 12511-12516.
- Eriksson L, Johansson E, Kettaneh-Wold N, Trygg J, Wikstrom C, and Wold, S. (2006). Multi- and megavariate data analysis. Basic

Principles and Applications: Umetrics AB. 425 pages. ISBN-10: 91-973730-2-8.

- Ettinger, M.P., Littlejohn, T.W., Schwartz, S.L., Weiss, S.R., McIlwain, H.H., Heymsfield, S.B., Bray, G.A., Roberts, W.G., Heyman, E.R., Stambler, N., Heshka, S., Vicary, C., Guler, H.P. (2003). Recombinant variant of ciliary neurotrophic factor for weight loss in obese adults. *Journal of American Medical Association*, 289(14), 1826-1832.
- Eu, C.H., Lim, W.Y., Ton, S.H., Abdul Kadir, K. (2010). Glycyrrhizic acid improved lipoprotein lipase expression, insulin sensitivity, serum lipid and lipid deposition in high-fat diet-induced obese rats. *Lipids Health Disease*, 9(81).
- Farvid, M.S., Ng, T.W., Chan, D.C., Barrett, P.H., and Watts, G.F. (2005). Association of adiponectin and resistin with adipose tissue compartments, insulin resistance and dyslipidaemia. *Diabetes, Obesity and Metabolism*, 7, 406-413.
- Fasshauer, M., and Paschke, R. (2003). Regulation of adipocytokines and insulin resistance. *Diabetologia*, 46(12), 1594-603.
- Faujan, H.N., Noriham, A., Norrakiah, A.S., and Babji, A.S. (2009). Antioxidant activity of plants méthanolic extracts containing phenolic compounds. *African Journal of Biotechnology*, 8(3), 484-489.
- Fernández-Sánchez, A., Madrigal-Santillán, E., Bautista, M., Esquivel-Soto, J., Morales-González, A., Esquivel-Chirino, C., Durante-Montiel, I., Sánchez-Rivera, G., Valadez-Vega, C., and Morales-González, J.A. (2011). Inflammation, Oxidative Stress, and Obesity. *International Journal of Molecular Science*, 12, 3117-3132.
- Fielding, B.A., and Frayn, K.N. (1998). Lipoprotein lipase and the disposition of dietary fatty acids. *British Journal of Nutrition*, 80(6), 495-502.
- Food and Drug Administration (FDA). (2004). Calorie count: report of the working group on obesity, U.S. food and drug administrations, Washington D.C.
- Food and Drug Administration, (FDA). (2004).
- Fox, P.F., and Stepaniak, L. (1983). Isolation and some properties of extracellular heat-stable lipases from *Pseudomonas fluorescens* strain AFT 36. *Journal of Dairy Research*, 50(1), 77-89.

- Francis, G., Kerem, Z., Makkar, H.P., and Becker, K. (2002). The biological action of saponins in animal systems: a review. *British Journal of Nutrition*, 88(6), 587-605.
- Fried, M., Hainer, V., Basdevant, A., Buchwald, H., Dietel, M., Finer, N., Greve, J.W., Horber, F., Mathus-Vliegen, E., Scopinaro, N., Steffen, R., Tsigos, C., Weiner, R., and Widhalm, K. (2008) Interdisciplinary European guidelines on surgery for severe obesity. *Rozhl Chir*, 87(9), 468-476.
- Friedman, H., and Nylund, B. (1980). Intestinal fat digestion and absorption. *American Journal of Clinical Nutrition*, 33(5), 1108-1139.
- Friedman, J.M., and Halaas, J.L. (1998). Leptin and the regulation of body weight in mammals. *Nature*, 395(6704), 763–770.
- Friedman, J.M. (2000). Obesity in the new millennium. *Nature*, 404(6778), 632-634.
- Fukumoto, J., Iwai, W., and Tsujiska, Y. (1963). Studies on lipase purification and crystallization of a lipase secretion by *Aspergillus niger*. *Journal of General and Applied Microbiology*, 9, 353–361.
- Furukawa, S., Fujita, T., Shimabukuro, M., Iwaki, M., Yamada, Y., Nakajima, Y., Nakayama, O., Makishima, M., Matsuda, M., and Shimomura, I. (2004). Increased oxidative stress in obesity and its impact on metabolic syndrome. *Journal of Clinical Investigation*, 114(12), 1752–1761.
- Galhardo, J., Davis, N., Matthai, S., and Shield, J.P. (2010). European Medicines Agency withdrawal for sibutramine. *Archives of Disease in Childhood*, 95(10), 856.
- Gargouri, Y., Ransac, S., and Verger, R. (1997). Covalent inhibition of digestive lipases: an in vitro study. *Biochimica et Biophysica Acta*, 1344(1), 6–37.
- Geslain-Lane'elle, C. (2006). Conference on nutrition and health claims. Bologna: Noviembre.
- Gholamhoseinian, A., Shahouzehi, B., and Sharififar, F. (2010). Inhibitory effect of some plant extracts on pancreatic lipase. *International Journal of Pharmacology*, 6(1), 18-24.
- Gil-Campos, M., Cañete, R., and Gil, A. (2004). Hormones regulating lipid metabolism and plasma lipids in childhood obesity. *International Journal of Obesity and Related Metabolic Disorders*, 28(Suppl 3), 75-80.

- Glenny, A.M., O'Meara, S., Melville, A., Sheldon, T.A. and Wilson, C. (1997). The treatment and prevention of obesity: a systematic review of the literature. *International Journal of Obesity*, 21(9), 715–737.
- Golay, A., Swislocki, A.L., Chen, Y.D., Jaspan, J.B., and Reaven, G.M. (1986). Effect of obesity on ambient plasma glucose, free fatty acid, insulin, growth hormone, and glucagon concentrations. *Journal of Clinical Endocrinology and Metabolism*, 63(2), 481-484.
- Goldberg, I.J., and Merkel, M. (2001). Lipoprotein lipase: physiology, biochemistry, and molecular biology. *Frontiers in Bioscience*, 6, 388-405.
- Goldstein, D.J. (1992). Beneficial health effects of modest weight loss. *International Journal of Obesity*, 16(6), 397-415.
- Gordon, T., and Kannel, W.B. (1982). Multiple risk functions for predicting coronary heart disease: the concept, accuracy, and application. *American Heart Journal*, 103(6), 1031-1039.
- Gowda, G. A., Zhang, S., Gu, H., Asiago, V., Shanaiah, N., and Raftery, D. (2008). Metabolomics Based Methods for Early Disease Diagnostics: A Review. *Expert Review in Molecular Diagnosis*, 8(5), 617-633.
- Graf, B.A., Milbury, P.E., and Blumberg, J.B. (2005). Flavonols, flavonones, flavanones and human health: Epidemiological evidence. *Journal of Medicinal Food*, 8, 281-290.
- Grivet, J.P., and Delort, A.M. (2009). NMR for microbiology: In vivo and in situ applications. *Progress in Nuclear Magnetic Resonance Spectroscopy*, 549(1), 1–53.
- Gülçin, I., İrfan Kufrevioğlu, Ö., Oktay, M., Büyükkokkuroğlu, M.E. (2004). Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica L.*). *Journal of Ethnopharmacology*, 90(2-3), 205–215.
- Hagstrom, E., Arner, P., Ungerstedt, U., and Bolinder, J. (1990). Subcutaneous adipose tissue: a source of lactate production after glucose ingestion in humans. *American Journal of Physiology - Endocrinology and Metabolism*, 258(5 Part 1), 888-893.
- Hall, R.D., Brouwer, I.D., and Fitzgerald, M.A. (2008). Plant metabolomics and its potential application for human nutrition. *Physiologia Plantarum*, 132(2), 162-175.

- Hamad, E.M., Sato, M., Uzu, K., Yoshida, T., Higashi, S., Kawakami, H., Kadooka, Y., Matsuyama, H., Abd El-Gawad, I.A., and Imaizumi, K. (2009). Milk fermented by *Lactobacillus gasseri* SBT2055 influences adipocyte size via inhibition of dietary fat absorption in Zucker rats. *British Journal of Nutrition*, 101(5), 716-724.
- Hamosh, M. (1990). Lingual and gastric lipases. *Nutrition*, 6(6), 421-428.
- Han, L.K., Kimura, Y., and Okuda, H. (1999). Reduction in fat storage during chitin chitosan treatment in mice fed a high-fat diet. *International Journal of Obesity and Related Metabolic Disorders*, 23(2), 174-179.
- Han, L.K., Takaku, T., Li, J., Kimura, Y., and Okuda, H. (1999). Anti-obesity action of oolong tea. *International Journal of Obesity and Related Metabolic Disorders*, 23(1), 98-105.
- Han, L.K., Xu, B.J., Kimura, Y., Zheng, Y., and Okuda, H. (2000). *Platycodi radix* affects lipid metabolism in mice with high fat diet-induced obesity. *Journal of Nutrition*, 130(11), 2760-2764.
- Han, L.K., Kimura, Y., Kawashima, M., Takaku, T., Taniyama, T., Hayashi, T., Zheng, Y.N., and Okuda, H. (2001). Anti-obesity effects in rodents of dietary teasaponin, a lipase inhibitor. *International Journal of Obesity and Related Metabolic Disorders*, 25(10), 1459-1464.
- Han, L.K., Zheng, Y.N., Yoshikawa, M., Okuda, H., Kimura, Y. (2005). Anti-obesity effects of chikusetsusaponins isolated from *Panax japonicus* rhizomes. *BMC Complementary and Alternative Medicine*, 5, 9.
- Han, L.K., Nose, R., Li, W., Gong, X.J., Zheng, Y.N., Yoshikawa, M., Koike, K., Nikaido, T., Okuda, H., and Kimura, Y. (2006). Reduction of fat storage in mice fed a high-fat diet long term by treatment with saponins prepared from *Kochia scoparia* fruit. *Phytotherapy Research*, 20(10), 877-882.
- Han, X., Patters, A.B., Jones, D.P., Zelikovic, I., Chesney, R.W. (2006). The taurine transporter: mechanisms of regulation. *Acta physiologica (Oxford, England)*, 187(1-2), 61-73.
- Han, S.H., Sakuma, I., Shin, E.K., and Koh, K.K. (2009). Antiatherosclerotic and anti-insulin resistance effects of adiponectin: basic and clinical studies. *Progress in Cardiovascular Disease*, 52(2), 126-140.
- Hanefeld, M., and Sachse, G. (2002). The effects of orlistat on body weight and glycaemic control in overweight patients with type 2 diabetes: a

- randomized, placebo-controlled trial. *Diabetes Obesity Metabolism*, 4(6), 415-23.
- Harach, T., Aprikian, O., Monnard, I., Moulin, J., Membrez, M., Beolor, J.C., Raab, T., Mace, K., and Darimont, C. (2010). Rosemary (*Rosmarinus officinalis* L.) leaf extract limits weight gain and liver steatosis in mice fed a high-fat diet. *Planta Medica*, 76(6), 566-571.
- Harborne, J.B. (1998). Phytochemical methods: A guide to modern techniques on plant analysis. Kluwer Academic Publishers, United Kingdom. Third edition, p 299-312.
- Hardeman, W., Griffin, S., Johnston, M., Kinmonth, A.L., Wareham, N.J. (2000). Interventions to prevent weight gain: a systematic review of psychological models and behaviour change methods. *International Journal of Obesity and Related Metabolic Disorders*, 24(2), 131-143.
- Harte, R.A., Kirk, E.A., Rosenfeld, M.E., LeBoeuf, R.C. (1999). Initiation of hyperinsulinemia and hyperleptinemia is diet dependent in C57BL/6 mice. *Hormone and Metabolic Research*, 31(10), 570-575.
- Hassan, W.E., and Mahmood, M. (2006). Healing Herbs of Malaysia. Biotropics Malaysia Berhad. ISBN 978-983-99544-2-5.
- Havel, R. (1994). McCollum Award Lecture, 1993: Triglyceride-rich lipoproteins and atherosclerosis-new perspectives. *American Journal of Clinical Nutrition*, 59(4), 795-796.
- He, Q., Ren, P., Kong, X., Wu, Y., Wu, G., Li, P., Hao, F., Tang, H., Blachier, F., and Yin, Y. (2012). Comparison of serum metabolite compositions between obese and lean growing pigs using an NMR-based metabonomic approach. *Journal of Nutritional Biochemistry*, 23(2), 133-139.
- Higgins, S.C., Gueorguiev, M., and Korbonits, M. (2007). Ghrelin, the peripheral hunger hormone. *Annals of Medicine*, 39(2), 116-136.
- Hill, J.O., Wyatt, H.R., Reed, G.W., and Peters, J.C. (2003). Obesity and the environment: where do we go from here?. *Science*, 299(5608), 853-855.
- Hlais, S., Reslan, D.R., Sarieddine, H.K., Nasreddine, L., Taan, G., Azar, S., and Obeid, O.A. (2012) Effect of lysine, vitamin B(6), and carnitine supplementation on the lipid profile of male patients with hypertriglyceridemia: a 12-week, open-label, randomized, placebo-controlled trial. *Clinical Therapy*, 34(8), 1674-82.

- Ho, J.N., Jang, J.Y., Yoon, H.G., Kim, Y., Kim, S., Jun, W., and Lee, J. (2012). Anti-obesity effect of a standardised ethanol extract from Curcuma longa L. fermented with Aspergillus oryzae in ob/ob mice and primary mouse adipocytes. *Journal of Science and Food Agriculture*, 92(9), 1833-1840.
- Hoffler, U., Hobbie, K., Wilson, R., Bai, R., Rahman, A., Malarkey, D., Travlos, G., and Ghanayem, B.I. (2009). Diet-induced obesity is associated with hyperleptinemia, hyperinsulinemia, hepatic steatosis, and glomerulopathy in C57Bl/6J mice. *Endocrine*, 36(2), 311-325.
- Hofmann, A.F., and Borgstrom, B. (1964). The intraluminal phase of fat digestion in man: the lipid content of the micellar and oil phases of intestinal content obtained during fat digestion and absorption. *Journal of Clinical Investigations*, 43(2), 247-257.
- Holmes, E., Bonner, F., Gartland, K., and Nicholson, J. (1990). Proton NMR monitoring of the onset and recovery of experimental renal damage. *Journal of Pharmaceutical and Biomedical Analysis*, 8(8-12), 959-962.
- Holmes, E., Bonner, F.W., and Nicholson, J.K. (1997). ¹H NMR spectroscopic and histopathological studies on propyleneimine-induced renal papillary necrosis in the rat and the multimammate desert mouse (*Mastomys natalensis*). *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 116(2), 125-34.
- Holmes, E., Li, J.V., Athanasiou, T., Ashrafi, H., and Nicholson, J.K. (2011). Understanding the role of gut microbiome-host metabolic signal disruption in health and disease. *Trends in Microbiology*, 19(7), 349-359.
- Homans, A. L. and Fuchs, A. (1970). Direct bioautography on thin-layer chromatograms as a method for detecting fungitoxic substances. *Journal of Chromatography*, 51(2), 327-329.
- Hosoda, H., Kojima, M., and Matsuo, H. (2000). Purification and characterization of rat des-Gln14-Ghrelin, a second endogenous ligand for the growth hormone secretagogue receptor. *Journal of Biological Chemistry*, 275(29), 21995-22000.
- Hotamisligil, G.S., Shargill, N.S., and Spiegelman, B.M. (1993). Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science*, 259(5091), 87-91.

- Hotamisligil, G.S. (1999). The role of TNF- α and TNG receptors in obesity and insulin resistance. *Journal of Internal Medicine*, 245(6), 621-625.
- Hotta, K., Funahashi, T., Arita, Y., Takahashi, M., Matsuda, M., Okamoto, Y., Iwahashi, H., Kuriyama, H., Ouchi, N., Maeda, K., Nishida, M., Kihara, S., Sakai, N., Nakajima, T., Hasegawa, K., Muraguchi, M., Ohmoto, Y., Nakamura, T., Yamashita, S., Hanafusa, T., and Matsuzawa, Y. (2000). Plasma concentrations of an oval, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 20(6), 1595-1599.
- Hsu, C.Y., Chan, Y.P., and Chang, J. (2007). Antioxidant activity of extract from Polygonum cuspidatum. *Biological Research*, 40(1), 13-21.
- Hsu, C.L., and Yen, G.C. (2007). Effect of gallic acid on high fat diet-induced dyslipidaemia, hepatosteatosis and oxidative stress in rats. *British Journal of Nutrition*, 98(4), 727-735.
- Hsu, C.L., and Yen, G.C. (2007). Effects of flavonoids and phenolic acids on the inhibition of adipogenesis in 3T3-L1 adipocytes. *Journal of Agriculture and Food Chemistry*, 55(21), 8404-8410.
- Hsu, C.H., Tsai, T.H., Kao, Y.H., Hwang, K.C., Tseng, T.Y., Chou P. (2008). Effect of green tea extract on obese women: A randomized, double-blind, placebo-controlled clinical trial. *Clinical Nutrition*, 27(3), 363-370.
- Hsu, C.L., Wu, C.H., Huang, S.L., and Yen, G.C. (2009). Phenolic compounds rutin and o-coumaric acid ameliorate obesity induced by high fat diet. *Journal of Agricultural and Food Chemistry*, 57(2), 425-431.
- Hsu, C.H., Liao, Y.L., Lin, S.C., Tsai, T.H., Huang, C.J., and Chou, P. (2011). Does supplementation with green tea extract improve insulin resistance in obese type 2 diabetics? A randomized, double-blind, and placebo-controlled clinical trial. *Alternative Medicine Review*, 16(2), 157-163.
- Hu, E., Liang, P., and Spiegelman, B.M. (1996). AdipoQ is a novel adipose-specific gene dysregulated in obesity. *Journal of Biological Chemistry*, 271(18), 10697-10703.
- Huang, D., Ou, B. and Prior, R.L. (2005). The chemistry behind the antioxidant capacity assays. *Journal of Agriculture and Food Chemistry*, 53(6), 1841-1856.

- Hursel, R., Viechtbauer, W., Dulloo, A.G., Tremblay, A., Tappy, L., Rumpler, W., and Westerterp-Planteng, M.S. (2011). The effects of catechin rich teas and caffeine on energy expenditure and fat oxidation: a meta-analysis. *Obesity reviews*, 12(7), 573-581.
- Husain, G.M., Chatterjee, S.S., Singh, P.N., and Kumar, V. (2011). Hypolipidemic and Antiobesity-Like Activity of Standardised Extract of Hypericum perforatum L. in Rats. *International Scholarly Research Network ISRN Pharmacology*, 2011, Article ID 505247, 7 pages.
- Hussein, M.A. (2011). Anti-obesity, antiatherogenic, anti-diabetic and antioxidant activities of J. montana ethanolic formulation in obese diabetic rats fed high-fat diet. *Free Radicals and Antioxidants*, 1(1), 49-60.
- Ibarra, A., Cases, J., Roller, M., Chiralt-Boix, A., Coussaert, A., and Ripoll, C. (2011). Carnosic acid-rich rosemary (*Rosmarinus officinalis* L.) leaf extract limits weight gain and improves cholesterol levels and glycaemia in mice on a high-fat diet. *British Journal of Nutrition*, 106(8), 1182-1189.
- Ignjatovic, V., Ogru, E., Heffernan, M., Libinaki, R., Lim, Y., and Ng, F. (2000). Studies on the use of "slimax", a chinese herbal mixture, in the treatment of human obesity. *Pharmaceutical Biology*, 38(1), 30-35.
- Ikeda, I., Tsuda, K., Suzuki, M., Kobayashi, T., Unno, H., Tomoyori, H., Goto, H., Kawata, Y., Imaizumi, K., Nozawa, A., and Kakuda, T. (2005). Tea catechins with a galloyl moiety suppress postprandial hypertriacylglycerolemia by delaying lymphatic transport of dietary fat in rats. *Journal of Nutrition*, 135(2), 155-159.
- Ikhlas, B., Huda, N., and Ismail, N. (2012). Effect Of *Cosmos Caudatus*, *Polygonum Minus* And Bht On Physical Properties, Oxidative Process, And Microbiology Growth Of Quail Meatball During Refrigeration Storages. *Journal of Food Processing and Preservation*, 36(1), 55–66.
- Insull, W., Black, D., Dujovne, C., Hosking, J.D., Hunninghake, D., Keilson, L., Knopp, R., McKenney, J., Stein, E., and Troendle, A.J. (1994). Efficacy and safety of once-daily vs twice-daily dosing with fluvastatin, a synthetic reductase inhibitor, in primary hypercholesterolemia. *Archives of Internal Medicine*, 154(21), 2449-2455.
- Ismail, S. (2000). Sayuran tradisional ulam dan penyedap rasa. Universiti Kebangsaan Malaysia, Bangi. ISBN 967-942-459-6.

- Iwashita, S., Tanida, M., Terui, N., Ootsuka, Y., Shu, M., Kang, D., and Suzuki, M. (2002). Direct measurement of renal sympathetic nervous activity in high-fat diet-related hypertensive rats. *Life Science*, 71(5), 537–546.
- Jaeger, K.E., Ransac, S., Dijkstra, B.W., Colson, C., van Heuvel, M., and Misset, O. (1994). Bacterial lipases. *FEMS Microbiology Reviews*, 15(1), 29-63.
- Jäger, S., Trojan, H., Kopp, T., Laszczyk, M., and Scheffler, A. (2009). Pentacyclic triterpene distribution in various plants-rich sources for a new group of multi potent plant extracts. *Molecules*, 14(6), 2016-2031.
- Jang, D., Lee, G., Kim, J., Lee, Y., Kim, J., Kim, Y. (2008). A new pancreatic lipase inhibitor isolated from the roots of *Actinidia arguta*. *Archives of Pharmacal Research*, 31(5), 666-670.
- Jang, W.S. and Choung, S.Y. (2013). Antiobesity Effects of the Ethanol Extract of *Laminaria japonica* Areshoung in High-Fat-Diet-Induced Obese Rat. *Evidence Based Complementary Alternative Medicines*, 2013, 492807.
- Jansson, P.A., Larsson, A., Smith, U., and Lönnroth, P. (1994). Lactate release from the subcutaneous tissue in lean and obese men. *Journal of Clinical Investigations*, 93(1), 240–246.
- Jensen, R.G., Galluzzo, D.R., and Bush, V.J. (1990). Selectivity is an important characteristic of lipases (acylglycerol hydrolases). *Biocatalysis*, 3(4), 307-316.
- Jiang, C.Y., Yang, K.M., Yang, L., Miao, Z.X., Wang, Y.H., and Zhu, H.B. (2013). A ¹H NMR-Based Metabonomic Investigation of Time-Related Metabolic Trajectories of the Plasma, Urine and Liver Extracts of Hyperlipidemic Hamsters. *PLoS ONE*, 8(6), e66786.
- Johnston, J.I., and Franz, V. (1992). Reninangiotensin system: a dual tissue and hormonal system for cardiovascular control. *Journal of Hypertension*, 10(7), 13-26.
- Jones, D. (2011). Suspense Builds on Anti-Obesity Rollercoaster Ride. *Nature Reviews Drug Discovery*, 10(1), 5-6.
- Juhel, C., Armand, M., Pafumi, Y., Rosier, C., Vandermander, J., and Lairon, D. (2000). Green tea extract (AR25) inhibits lipolysis of triglycerides in gastric and duodenal medium in vitro. *Journal of Nutritional Biochemistry*, 11(1), 45-51.

- Jung, J.Y., Kim, I.Y., Kim, Y.N., Shin, J.H., Jang, Z.H., Lee, H.S., Hwang, G.S., and Seong, J.K. (2012). *1H NMR-based metabolite profiling of diet-induced obesity in a mouse model*. *Biochemistry and Molecular Biology Reports*, 45(7), 419-424.
- Kaddurah-Daouk, R., Kristal, B.S., and Weinshilboum, R.M. (2008). Metabolomics: a global biochemical approach to drug response and disease. *Annual Review of Pharmacology and Toxicology*, 48, 653-683.
- Kageyama, H., Hirano, T., Okada, K., Ebara, T., Kageyama, A., Murakami, T., Shioda, S., and Adachi, M. (2003). Lipoprotein lipase mRNA in white adipose tissue but not in skeletal muscle is increased by pioglitazone through PPAR-gamma. *Biochemical Biophysical Research Communications*, 305(1), 22-27.
- Kähkönen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K., Kujala, T.S., and Heinonen, M. (1999). Antioxidant Activity of Plant Extracts Containing Phenolic Compounds. *Journal of Agricultural and Food Chemistry*, 47(10), 3954-3962.
- Kahn, S.E., Hull, R.L., and Utzschneider, K.M. (2006). Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*, 444(7121), 840–846.
- Kalhan, S.C., Guo, L., Edmison, J., Dasarathy, S., McCullough, A.J., Hanson, R.W., and Milburn, M. (2011) Plasma metabolomic profile in nonalcoholic fatty liver disease. *Metabolism*, 60(3), 404-413.
- Kamisoyama, H., Honda, K., Tominaga, Y., Yokota, S., and Hasegawa, S. (2008). Investigation of the anti-obesity action of licorice flavonoid oil in diet-induced obese rats. *Bioscience, Biotechnology and Biochemistry*, 72(12), 3225-3231.
- Kand'ár R., and Záková P. (2008). Allantoin as a marker of oxidative stress in human erythrocytes. *Clinical Chemistry and Laboratory Medicine*, 46(9), 1270-1274.
- Kang, Y.J., Jung, U.J., Lee, M.K., Kim, H.J., Jeon, S.M., Park, Y.B., Chung, H.G., Baek, N.I., Lee, K.T., Jeong, T.S., Choi, M.S. (2008). Eupatilin, isolated from Artemisia princeps Pampanini, enhances hepatic glucose metabolism and pancreatic β -cell function in type 2 diabetic mice. *Diabetes Research and Clinical Practice*, 82(1), 25-32.
- Kao, Y.H., Hiipakka, R.A., and Liao, S. (2000). Modulation of endocrine systems and food intake by green tea epigallocatechin gallate. *Endocrinology*, 141(3), 980-987.

- Karamadoukis, L., Shivashankar, G.H., Ludeman, L., and Williams, A.J. (2009). An unusual complication of treatment with orlistat. *Clinical Nephrology*, 71(4), 430-432.
- Karmase, A., Birari, R., and Bhutani, K.K. (2013). Evaluation of anti-obesity effect of *Aegle marmelos* leaves. *Phytomedicine*, 20(10), 805-812.
- Kasparavičienė, G., Ramanauskienė, K., Svickas, A., Velziene, S., Kalveniene, Z., Kazlauskienė, D., Ragazinskiene, O., and Ivanauskas, K. (2013). Evaluation of total phenolic content and antioxidant activity of different *Rosmarinus officinalis* L. ethanolic extracts. *Biologija*, 59(1), 39-44.
- Kato, E., Nakagomi, R., Gunawan-Puteri, M., and Kawabata, J. (2013). Identification of hydroxychavicol and its dimers, the lipase inhibitors contained in the Indonesian spice, *Eugenia polyantha*. *Food Chemistry*, 136(3-4), 1239-1242.
- Kaur, G., and Kulkarni, S.K. (2001). Differential effect of polyherbal, antiobesity preparation, OB-200G in male and female mice and monosodium glutamate-treated rats. *Indian Journal of Experimental Biology*, 39(6), 551-557.
- Kazmi, I., Afzal, M., Rahman, S., Iqbal, M., Imam, F., and Anwar, F. (2012). Antiobesity potential of ursolic acid stearoyl glucoside by inhibiting pancreatic lipase. *European Journal of Pharmacology*, 709(1-3), 28-36.
- Keaney Jr, J.F., Larson, M.G., Vasan, R.S., Wilson, P.W.F., Lipinska, I., Corey, D., Massaro, J.M., Sutherland, P., Vita, J.A., Benjamin, E.J. (2003). Obesity and systemic oxidative stress clinical correlates of oxidative stress in the framingham study. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 23(3), 434-439.
- Keun, H.C. (2006). Metabonomic modeling of drug toxicity. *Pharmacology and Therapeutics*, 109(1-2), 92-106.
- Kim, M.S., Kim, J.K., Kim, H.J., Moon, S.R., Shin, B.Ch., Park, K.W., Yang, H.O., Kim, S.M., and Park, R. (2003). *Hibiscus* extract inhibits the lipid droplet accumulation and adipogenic transcription factors expression of 3T3-L1 preadipocytes. *Journal of Alternative and Complementary Medicine*, 9(4), 499-504.
- Kim, S.O., Yun, S.J., Jung, B., Lee, E.H., Hahm, D.H., Shim, I., Lee, H.J. (2004). Hypolipidemic effects of crude extract of adlay seed (*Coix lachrymajobi* var Mayuen) in obesity rat fed high fat diet: relations of TNF-a and leptin Mrna expressions and serum lipid levels. *Life Science*, 75(11), 1391-1404.

- Kim, H.Y., and Kang, M.H. (2005). Screening of Korean medicinal plants for lipase inhibitory activity. *Phytotherapy Research*, 19(4), 359-361.
- Kim, J.H., Hahm, D.H., Yang, D.C., Kim, J.H., Lee, H.J., Shim, I., (2005). Effect of crude saponin of Korean red ginseng on high-fat diet-induced obesity in the rat. *Journal of Pharmacological Sciences*, 97(1), 124-131.
- Kim, Y.J., Shin, Y.O., Ha, Y.W., Lee, S., Oh, J.K., and Kim, Y.S. (2006). Anti-obesity effect of *Pinellia ternata* extract in zucker rats. *Biological and pharmaceutical bulletin*, 29(6), 1278-1281.
- Kim, Y.J., and Park, T. (2008). Genes are differentially expressed in the epididymal fat of rats rendered obese by high fat diet. *Nutrition Research*, 28(6), 414-422.
- Kim, S.H., Yang, S.O., Kim, H.S., Kim, Y., Park, T., and Choi, H.K. (2009). ¹H-nuclear magnetic resonance spectroscopy-based metabolic assessment in a rat model of obesity induced by a high-fat diet. *Analytical and Bioanalytical Chemistry*, 395(4), 1117-1124.
- Kim, W.S., Lee, Y.S., Cha, S.H., Jeong, H.W., Choe, S.S., Lee, M.R., Oh, G.T., Park, H.S., Lee, K.U., Lane, M.D., and Kim, J.B. (2009). Berberine improves lipid dysregulation in obesity by controlling central and peripheral AMPK activity. *American Journal of Physiology Endocrinology and Metabolism*, 296(4), 812-819.
- Kim, J.Y., Park, J.Y., Kim, O.Y., Ham, B.M., Kim, H.J., Kwon, D.Y., Jang, Y., and Lee, J.H. (2010). Metabolic profiling of plasma in overweight/obese and lean men using ultra performance liquid chromatography and Q-TOF mass spectrometry (UPLC-Q-TOF MS). *Journal of Proteome Research*, 9(9), 4368-4375.
- Kim, M.J., Park, M.H., Jeong, M.K., JYeo, J.D., Cho, W., Chang, P.S., Chung, J.H., and Lee, J.H. (2010). Radical Scavenging Activity and Anti-obesity Effects in 3T3-L1 Preadipocyte Differentiation of Ssuk (*Artemisia princeps* Pamp.) Extract. *Food Science and Biotechnology*, 19(2), 535-540.
- Kim, H.K., Choi, Y.H., and Verpoorte, R. (2010a). NMR-based metabolomic analysis of plants. *Nature Protocols*, 5(3), 536-540.
- Kim, S.Y., Hyun, S.H., Yang, S.O., Choi, H.K., and Lee, B.Y. (2010b). ¹H-NMR-Based Discrimination of Thermal and Vinegar Treated Ginseng Roots. *Journal of Food Science*, 75(6), 577-581.
- Kim, I.Y., Jung, J., Jang, M., Ahn, Y.G., Shin, J.H., Choi, J.W., Sohn, M.R., Shin, S.M., Kang, D.G., Lee, H.S., Bae, Y.S., Ryu do, H., Seong,

- J.K., and Hwang, G.S. (2010) ^1H NMR-based metabolomic study on resistance to diet-induced obesity in AHNAK knock-out mice. *Biochemical and Biophysical Research Communications*, 403(3-4), 428-434.
- Kim, H.J., Kim, J.H., Noh, S., Hur, H.J., Sung, M.J., Hwang, J.T., Park, J.H., Yang, H.J., Kim, M.S., Kwon, D.Y., and Yoon, S.H. (2011). Metabolomic analysis of livers and serum from high-fat diet induced obese mice. *Journal of Proteome Research*, 10(2), 722-731.
- Kim, Y.S., Lee, Y., Kim, J., Sohn, E., Kim, C.S., Lee, Y.M., Jo, K., Shin, S., Song, Y., Kim, J.H., and Kim, J.S. (2012). Inhibitory Activities of Cudrania tricuspidata Leaves on Pancreatic Lipase In Vitro and Lipolysis In Vivo. *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 878365, 8 pages.
- Kim, M.J., Yang, H.J., Kim, J.H., Ahn, C.W., Lee, J.H., Kim, K.S., and Kwon, D.Y. (2013). Obesity-related metabolomic analysis of human subjects in black soybean peptide intervention study by ultraperformance liquid chromatography and quadrupole-time-of-flight mass spectrometry. *Journal of Obesity*, 2013, 874981.
- Kishino, E., Ito, T., Fujita, K., and Kiuchi, Y. (2006). A mixture of the Salacia reticulata (kotala himbutu) aqueous extract and cyclodextrin reduces the accumulation of visceral fat mass in mice and rats with high-fat diet-induced obesity. *Journal of Nutrition*, 136(2), 433-439.
- Klok ,M.D., Jakobsdottir, S., and Drent, M.L. (2007). The role of leptin and ghrelin in the regulation of food intake and body weight in humans: a review. *Obesity Review*, 8(1), 21-34.
- Knight, Z.A., Hannan, K.S., Greenberg, M.L., and Friedman, J.M. (2010). Hyperleptinemia Is Required for the Development of Leptin Resistance. *Plos ONE*, 5(6), e11376.
- Kobori, M., Masumoto, S., Akimoto, Y., and Oike, H. (2011). Chronic dietary intake of quercetin alleviates hepatic fat accumulation associated with consumption of a Western-style diet in C57/BL6J mice. *Molecular Nutrition and Food Research*, 55(4), 530-540.
- Koffi, E., Sea, T., Dodehe, Y., and Soro, S. (2010). Effect of solvent type on extraction of polyphenols from twenty three Ivorian plants. *Journal of Animal and Plant Sciences*, 5(3), 550-558.
- Kojima, M., Hosoda, H., Date, Y., Nakazato, M., Matsuo, H., and Kangawa, K. (1999). Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*, 402(6762), 656-660.

- Kolovou, G.D., Anagnostopoulou, K.K., and Cokkinos, D.V. (2005). Pathophysiology of dyslipidaemia in the metabolic syndrome. *Postgraduate Medical Journal*, 81(956), 358-366.
- Konishi, F. (1964). The relationship of urinary 17-hydroxycorticosteroids to creatinine in obesity *Metabolism*, 13(9), 847–851.
- Könner, A.C., Hess, S., Tovar, S., Mesaros, A., Sánchez-Lasheras, C., Evers, N., Verhagen, L.A., Brönnike, H.S., Kleinridders, A., Hampel, B., Kloppenburg, P., Brüning, J.C. (2011). Role for insulin signaling in catecholaminergic neurons in control of energy homeostasis. *Cell Metabolism*, 13(6), 720-728.
- Krauss, R.M., and Winston, M. (1998). Obesity impact on cardiovascular disease. *Circulation*, 98, 1472-1476.
- Krentz, A.J., and Bailey, C.J. (2005). Oral antidiabetic agents: current role in type 2 diabetes mellitus. *Drugs*, 65(3), 385-411.
- Kubota, N., Terauchi, Y., Miki, H., Tamemoto, H., Yamauchi, T., Komeda, K., Satoh, S., Nakano, R., Ishii, C., Sugiyama, T., Eto, K., Tsubamoto, Y., Okuno, A., Murakami, K., Sekihara, H., Hasegawa, G., Naito, M., Toyoshima, Y., Tanaka, S., Shiota, K., Kitamura, T., Fujita, T., Ezaki, O., Aizawa, S., and Kadokawa, T. (1999). PPAR gamma mediates high-fat diet-induced adipocyte hypertrophy and insulin resistance. *Molecular Cell*, 4(4), 597-609.
- Kulakowski, E.C., and Maturo, J. (1984). Hypoglycemic properties of taurine: not mediated by enhanced insulin release. *Biochemistry and Pharmacology*, 33(18), 2835–2838.
- Kuo, K.L., Weng, M.S., Chiang, C.T., Tsai, Y.J., Lin-Shiau, S.Y., and Lin, J.K. (2005). Comparative studies on the hypolipidemic and growth suppressive effects of oolong, black, pu-erh, and green tea leaves in rats. *Journal of Agriculture and Food Chemistry*, 53(2), 480-489.
- Kussmann, M., Raymond, F., and Affolter, M. (2006). OMICS-driven biomarker discovery in nutrition and health. *Journal of Biotechnology*, 124(4), 758–787.
- Labre`che, F.P., and Goldberg, M.S. (1997). Exposure to Organic Solvents and Breast Cancer in Women: A Hypothesis. *American Journal Of Industrial Medicine*, 32(1), 1-14.
- Lann, D., and LeRoith, D. (2007). Insulin Resistance as the Underlying Cause for the Metabolic Syndrome. *Medical Clinics of North America*, 91(6), 1063-1077.

- Lau, D.C., and Teoh, H. (2013). Benefits of modest weight loss on the management of type 2 diabetes mellitus. *Canadian Journal of Diabetes*, 37(2), 128-34.
- Lee, J.S., Lee, M.K., Ha, T.Y., Bok, S.H., Park, H.M., Jeong, K.S., Woo, M.N., Do, G.M., Yeo, J.Y., and Choi, M.S., (2006). Supplementation of whole persimmon leaf improves lipid profiles and suppresses body weight gain in rats fed high-fat diet. *Food and Chemical Toxicology*, 44(11), 1875-1883.
- Lee, R.A., and Balick, M.J. (2007). Indigenous use of *Hoodia gordonii* and appetite suppression. *Explore New York*, 3(4), 404-406.
- Lee, S.H., Bae, E.A., Park, E.K., Shin, Y.W., Baek, N.I., Han, E.J., Chung, H.G., and Kim, D.H. (2007). Inhibitory effect of eupatilin and jaceosidin isolated from *Artemisia princeps* in IgE-induced hypersensitivity. *International Immunopharmacology*, 7(13), 1678-1684.
- Levin, B.E., Dunn-Meynell, A.A., Balkan, B., Keesey, R.E. (1997). Selective breeding for diet-induced obesity and resistance in *Sprague dawley* rats. *American Journal of Physiology*, 273(2 Pt 2), 725-730.
- Levin, B.E., and Keesey, R.E. (1998). Defense of differing body weight set points in diet-induced obese and resistant rats. *American Journal of Physiology*, 274, 412–419.
- Lew, S.Q., and Bosch, J.P. (1991). Effect of diet on creatinine clearance and excretion in young and elderly healthy subjects and in patients with renal disease. *Journal of the American Society of Nephrology*, 2(4), 856–865.
- Li, B.H., Ma, X.F., Wu, X.D., and Tian, W.X. (2006). Inhibitory activity of chlorogenic acid on enzymes involved in the fatty acid synthesis in animals and bacteria. *IUBMB. Life*, 58(1), 39-46.
- Li, R.W., Theriault, A.G., Au, K., Douglas, T.D., Casaschi, A., Kurowska, E.M., and Mukherjee, R. (2006). Citrus polymethoxylated flavones improve lipid and glucose homeostasis and modulate adipocytokines in fructose-induced insulin resistant hamsters. *Life Science*, 79(4), 365–373.
- Li, F., Li, W., Fu, H., Zhang, Q., and Koike, K. (2007). Pancreatic lipase inhibiting triterpenoid saponins from fruits of *Acanthopanax senticosus*. *Chemical and Pharmaceutical Bulletin*, 55(7), 1087-1089.
- Li, H., Xie, Z., Lin, J., Song, H., Wang, Q., Wang, K., Su, M., Qiu, Y., Zhao, T., Song, K., Wang, X., Zhou, M., Liu, P., Zhao, G., Zhang, Q., and

- Jia, W. (2008). Transcriptomic and Metabonomic Profiling of Obesity-Prone and Obesity-Resistant Rats under High Fat Diet. *Journal of Proteome Research*, 7(11), 4775-4783.
- Li, S., Zhang, H.Y., Hu, C.C., Lawrence, F., Gallagher, K.E., Surapaneni, A. (2008). Assessment of diet-induced obese rats as an obesity model by comparative functional genomics. *Obesity*, 16, 811–818.
- Li, F., Xu, Q., Zheng, T., Huang, F., and Han, L. (2014). Metabonomic analysis of Allium macrostemon Bunge as a treatment for acute myocardial ischemia in rats. *Journal of Pharmaceutical and Biomedical Analysis*, 88, 225-34.
- Liao, P., Wei, L., Zhang, X., Li, X., Wu, H., Wu, Y., Ni, J., Pei, F. (2007). Metabolic profiling of serum from gadolinium chloride-treated rats by ^{1}H NMR spectroscopy. *Analytical Biochemistry*, 364(2), 112-21.
- Lin, S., Thomas, T.C., Storlien, L.H., and Huang, X.F. (2000). Development of high fat diet-induced obesity and leptin resistance in C57Bl/6J mice. *International journal of obesity and related metabolic disorders*, 24(5), 639-646.
- Lin, C.Y., Viant M.R., and Tjeerdenma, R.S. (2006). Metabolomics: Methodologies and Applications in the Environmental Sciences. *Journal of Pesticide Science*, 31(3), 245-251.
- Liu, Q., Raina, A.K., Smith, M.A., Sayre, L.M. and Perry, G. (2003). Hydroxynonenal, toxic carbonyls, and Alzheimer disease. *Molecular Aspects of Medicines*, 24(4-5), 305–313.
- Liu, R.H. (2003). Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals. *American Journal of Clinical Nutrition*, 78(3), 517-520.
- Liu, D.Z., Wang, F., Liao, T.G., Tang, J.G., Steglich, W., Zhu, H.J., and Liu, J.K. (2006). Vibralactone: a lipase inhibitor with an unusual fused beta-lactone produced by cultures of the basidiomycete *Boreostereum vibrans*. *Org. Letters*, 8(25), 5749-5752.
- Liu, J. Y., Chen, C.C., Wang, W.H., Hsu, J.D., Yang, M.Y., and Wang, C.J. (2006). The protective effects of Hibiscus sabdariffa extract on CCl₄-induced liver fibrosis in rats. *Food Chemistry and Toxicology*, 44(3), 336-343
- Liu, J.Y., Li, N., Yang, J., Li, N., Qui, H., Ai, D., Chiamvimonvat, N., Zhu, Y., and Hammock, B.D. (2010). Metabolic profiling of murine plasma reveals an unexpected biomarker in rofecoxib-mediated cardiovascular events. *Proceedings of the National Academy of Sciences of the United States of America*, 107(39), 17017-17022.

- Liu, I.M., Tzeng, T.F., Liou, S.S., and Chang, C.J. (2012). Regulation of obesity and lipid disorders by extracts from Angelica acutiloba root in high-fat diet-induced obese rats. *Phytotherapy Research*, 26(2), 223-230.
- Loh, S.P., and Hadira, O. (2011). In vitro inhibitory potential of selected Malaysian plants against key enzymes involved in hyperglycemia and hypertension. *Malaysian Journal of Nutrition*, 17(1), 77-86.
- Lopes, H.F., Martin, K.L., Nashar, K., Morrow, J.D., Goodfriend, T.L., and Egan, B.M. (2003). DASH diet lowers blood pressure and lipid-induced oxidative stress in obesity. *Hypertension*, 41(3), 422-30.
- Lucidi, P., Murdolo, G., Di Loreto, C., Parlanti, N., De Cicco, A., Ranchelli, A., Fatone, C., Taglioni, C., Fanelli, C., Santeusanio, F., and De Feo, P. (2004). Meal intake similarly reduces circulating concentrations of octanoyl and total ghrelin in humans. *Journal of Endocrinological Investigation*, 27(5), 12-15.
- Maestu, J., Jurima e, J., Valter, I., and Jurima e, T. (2008). Increases in ghrelin and decreases in leptin without altering adiponectin during extreme weight loss in male competitive bodybuilders. *Metabolism*, 57(2), 221-225.
- Maddux, B.A., See, W., Lawrence, J.C., Goldfine, A.I., Goldfine, I.D., and Evans, J.L. (2001). Protection against oxidative stress-induced insulin resistance in rat L6 muscle cells by micromolar concentrations of α-lipoic acid. *Diabetes*, 50(2), 404-410.
- Madsen, E.L., Rissanen, A., Bruun, J.M., Skogstrand, K., Tonstad, S., Hougaard, D.M., and Richelsen, B. (2008). Weight loss larger than 10% is needed for general improvement of levels of circulating adiponectin and markers of inflammation in obese subjects: a 3-year weight loss study. *European Journal of Endocrinology*, 158(2), 179-187.
- Madsen, A.N., Hansen, G., Paulsen, S.J., Lykkegaard, K., Tang-Christensen, M., Hansen, H.S. (2010). Long-term characterization of the diet-induced obese and diet-resistant rat model: a polygenic rat model mimicking the human obesity syndrome. *Journal of Endocrinology*, 206 287–296.
- Maeda, K., Okubo, K., Shimomura, I., Funahashi, T., Matsuzawa, Y., Matsubara, K. (1996). cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (adipose most abundant gene transcript 1). *Biochemical and Biophysical Research Communications*, 221(2), 286-289.

- Maeda, N., Shimomura, I., Kishida, K., Nishizawa, H., Matsuda, M., Nagaretani, H., Furuyama, N., Kondo, H., Takahashi, M., Arita, Y., Komuro, R., Ouchi, N., Kihara, S., Tochino, Y., Okutomi, K., Horie, M., Takeda, S., Aoyama, T., Funahashi, T., and Matsuzawa, Y. (2002). Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Natural Medicines*, 8(7), 731-737.
- Mahan, L.K., and Escott-Stump, S. (2008). Krause's food, nutrition, and diet therapy. 12th edition, Philadelphia: WB Saunders.
- Maher, A.D., Lindon, J.C., and Nicholson, J.K. (2009). HNMR-based metabolomics for investigating diabetes. *Future Medicinal Chemistry*, 1(4), 737-747.
- Mahmoud, R.H., and Elnour, W.A. (2013). Comparative evaluation of the efficacy of ginger and orlistat on obesity management, pancreatic lipase and liver peroxisomal catalase enzyme in male albino rats. *European Review for Medical and Pharmacological Sciences*, 17(1), 75-83.
- Mäkinen, V.P., Soininen, P., Forsblom, C., Parkkonen, M., Ingman, P., Kaski, K., Groop, P.H., and Ala-Korpela, M. (2008). ¹H NMR metabolomics approach to the disease continuum of diabetic complications and premature death. *Molecular Systems Biology*, 4(167), 1-6.
- Mamot, S. Traditional malaysian salads as a source of antioxidants. (2009). Prosiding Seminar Kimia Bersama UKM-ITB VII.
- Mark, A.L. (2006). Dietary therapy for obesity is a failure and pharmacotherapy is the future: a point of view. *Clinical and Experimental Pharmacology and Physiology*, 33(9), 857-862.
- Marloth, R. (1932). The Flora of South Africa with synopsis of the South African genera of Phanerogamous plant, Vol III, Wheldon and Wesley, London.
- Matsuzawa, Y., Funahashi, T., Kihara, S., and Shimomura, I. (2004). Adiponectin and metabolic syndrome. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 24(1), 29-33.
- McLaughlin, T., Abbasi, F., Kim, HS., Lamendola, C., Schaaf, P., Reaven, G.M. (2001). Relationship between insulin resistance, weight loss, and coronary heart disease risk in healthy, obese women. *Metabolism*, 50(7), 795-800.
- Mead, J.R., Irvine, S.A., Ramji, D.P. (2002). Lipoprotein lipase: structure, function, regulation, and role in disease. *Journal of Molecular Medicine (Berlin)*, 80(12), 753-69.

- Mediani, A., Abas, F., Khatib, A., Maulidiani, H., Shaari, K., Choi, Y.H., and Lajis, N.H. (2012). ¹H-NMR-based metabolomics approach to understanding the drying effects on the phytochemicals in *Cosmos caudatus*. *Food Research International*, 49(2), 763-770.
- Mediani, A., Abas, F., Ping, T.C., Khatib, A., and Lajis, N.H. (2012). Influence of growth stage and season on the antioxidant constituents of *Cosmos caudatus*. *Plant Foods Human Nutrition*, 67(4), 344-350.
- Mediani, A., Abas, F., Khatib, A., and Ping, T.C. (2013). *Cosmos Caudatus* as a Potential Source of Polyphenolic Compounds: Optimisation of Oven Drying Conditions and Characterisation of Its Functional Properties. *Molecules*, 18(9), 10452-10464.
- Mehran, A.E., Templeman, N.M., Brigidi, G.S., Lim, G.E., Chu, K.Y., Hu, X., Botezelli, J.D., Asadi, A., Hoffman, B.G., Kieffer, T.J., Bamji, S.X., Clee, S.M., Johnson, J.D. (2012). Hyperinsulinemia Drives Diet-Induced Obesity Independently of Brain Insulin Production. *Cell Metabolism*, 16(6), 691-692.
- Melis, G.C., ter Wengel, N., Boelens, P.G., and van Leeuwen, P.A. (2004). Glutamine: recent developments in research on the clinical significance of glutamine. *Current Opinion in Clinical Nutrition and Metabolic Care*, 7(1) 59-70.
- Melnikova, I., and Wages, D. (2006). Antiobesity therapies. *Nature Review Drug Discovery*, 5(5), 369–370.
- Michels, K.B., Giovannucci, E., Joshipura, K.J., Rosner, B.A., Stampfer, M.J., Fuchs, C.S., Colditz, G.A., Speizer, F.E., Willett, W.C. (2000). Prospective Study of Fruit and Vegetable Consumption and Incidence of Colon and Rectal Cancers. *Journal of the National Cancer Institute*, 92(21), 1740-1752.
- Mnafgui, K., Hamden, K., Salah, H.B., Kchaou, M., Nasri, M., Slama, S., Derbali, F., Allouche, N., and Elfeki, A. (2012). Inhibitory activities of *Zygophyllum album*: A natural weight lowering plant on key enzymes in high fat diet fed rats. *Evidence Based Complimentary and Alternative Medicine*, 2012, Article ID 620384, 9 pages.
- Modak, M., Dixit, P., Londhe, J., Ghaskadbi, S., and Devasagayam, T.P. (2007). Indian herbs and herbal drugs used for the treatment of diabetes. *Journal of Clinical Biochemistry and Nutrition*, 40(3), 163-173.
- Moon, H.S., Lee, H.G., Choi, Y.J., Kim, T.G., and Cho, C.S. (2007). Proposed mechanisms of (-)-epigallocatechin-3-gallate for anti-obesity. *Chemico-Biological Interactions*, 167(2), 85-98.

- Mooradian, A. D. (2009). Dyslipidemia in type 2 diabetes mellitus. *Nature Clinical Practice Endocrinology and Metabolism*, 5(3), 150–159.
- Moreno, D.A., Ilic, N., Poulev, A., Brasaemle, D.L., Fried, S.K., Raskin, I., (2003). Inhibitory effects of grape seed extract on lipases. *Nutrition*, 19(10), 876–879.
- Moreno, D.A., Ilic, N., Poulev, A., and Raskin, I. (2006). Effects of *Arachis hypogaea* nutshell extract on lipid metabolic enzymes and obesity parameters. *Life Science*, 78(24), 2797-2803.
- Moreno, D.A., Ripoll, C., Ilic, N., Poulev, A., Aubin, C., and Raskin, I. (2006). Inhibition of lipid metabolic enzymes using *Mangifera indica* extracts. *Journal of Food Agriculture and Environment*, 4(1), 21-26.
- Morrison, R.F., and Farmer, S.R. (1999). Insights into the transcriptional control of adipocyte differentiation. *Journal of Cellular Biochemistry*, 32-33, 59-67.
- Morrow, J.D. (2003). Is oxidative stress a connection between obesity and atherosclerosis? *Arteriosclerosis, Thrombosis, and Vascular Biology*, 23(3), 368-370.
- Mukherjee, M. (2003). Human digestive and metabolic lipases - a brief review. *Journal of Molecular Catalysis B: Enzymatic*, 22(5-6), 369–376.
- Mund, R.A., and Frishman, W.H. (2013). Brown adipose tissue thermogenesis: β 3-adrenoreceptors as a potential target for the treatment of obesity in humans. *Cardiology in Review*, 21(6), 265-269.
- Must, A., Spadano, J., Coakley, E.H., Field, A.E., Colditz, G., and Dietz, W.H. (1999). The disease burden associated with overweight and obesity. *Journal of the American Medical Association*, 282(16), 1523–1529.
- Mustafa, R. A., Hamid, A. A., Mohamed, S., and Bakar, F. A. (2010). Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*, 75(1), 28-35.
- Naik, G.H., Priyadarshini, K.I., Satav, J.G., Banavalikar, M.M., Sohoni, D.P., Biyani, M.K., and Mohan, H. (2003). Comparative antioxidant activity of individual herbal components used in Ayurvedic medicine. *Phytochemistry*, 63(1), 97-104.
- Nakagawa, K., Kishida, H., Arai, N., Nishiyama, T., and Mae, T. (2004). Licorice flavonoids suppress abdominal fat accumulation and

- increase in blood glucose level in obese diabetic KK-Ay mice. *Biological and Pharmaceutical Bulletin*, 27(11), 1775-1778.
- Nakai, M., Fukui, Y., Asami, S., Toyoda-Ono, Y., Iwashita, T., Shibata, H., Mitsunaga, T., Hashimoto, F., and Kiso, Y. (2005). Inhibitory effects of oolong tea polyphenols on pancreatic lipase in vitro. *Journal of Agriculture and Food Chemistry*, 53(11), 4593-4598.
- Nakazato, M., Murakami, N., Date, Y., Kojima, M., Matsuo, H., Kangawa, K., Matsukura, S. (2001). A role for ghrelin in the central regulation of feeding. *Nature*, 409(6817), 194-198.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E.C., Biryukov, S., Abbafati, C., Abera, S.F., Abraham, J.P., Abu-Rmeileh, N.M. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, 384(9945), 766-781.
- Nicholson, J. K., Timbrell, J. A., and Sadler, P. J. (1985). A high-resolution proton nuclear magnetic resonance approach to the study of hepatocyte and drug metabolism. Application to acetaminophen. *Molecular Pharmacology*, 27(6), 644-651.
- Nicholson, J.K., Connelly, J., Lindon, J.C., and Holmes, E. (2002). Metabonomics: a platform for studying drug toxicity and gene function. *Nature Reviews Drug Discovery*, 1(2), 153-161.
- Nicholson, J.K., and Lindon, J.C. (2008). Systems biology: Metabonomics," *Nature*, 455(7216), 1054-1056.
- Ninomiya, K., Matsuda, H., Shimoda, H., Nishida, N., Kasajima, N., Yoshino, T., Morikawa, T., and Yoshikawa, M. (2004). Carnosic acid, a new class of lipid absorption inhibitor from sage. *Bioorganic and Medicinal Chemistry Letters*, 14(8), 1943–1946.
- Norhanom, A. W., Ashril, Y., and Mustafa, A. M. (1999). Anti-tumour activity in local ulam. In Ali, A.M., Shaari, K., and Zakaria, Z. (Eds.), *Phytochemicals and biopharmaceutins from the Malaysian rainforest* (p 135-141). Kuala Lumpur, Malaysia: FRIM.
- Nukitrangsang, N., Okabe, T., Toda, T., Inafuku, M., Iwasaki, H., Yanagita, T., and Oku, H. (2011). Effect of Peucedanum japonicum Thunb on the expression of obesity-related genes in mice on a high-fat diet. *Journal of Oleo Science*, 60(10), 527-536.
- Oberbach, A., Blüher, M., Wirth, H., Till, H., Kovacs, P., Kullnick, Y., Schlichting, N., Tomm, J.M., Rolle-Kampczyk, U., Murugaiyan, J., Binder, H., Dietrich, A., and von Bergen, M. (2011). Combined

- proteomic and metabolomic profiling of serum reveals association of the complement system with obesity and identifies novel markers of body fat mass changes. *Journal of Proteome Research*, 10(10), 4769–4788.
- Ohara, Y., Peterson, T.E. and Harrison, D.G. (1933). Hypercholesterolemia increases endothelial superoxide anion production. *Journal of Clinical Investigation*, 91(6), 2546-2551.
- Ohashi, K., Ouchi, N., and Matsuzawa, Y. (2011). Adiponectin and Hypertension. *American Journal Of Hypertension*, 24(3), 263-269.
- Ohia, S.E., Opere, C.A., LeDay, A.M., Bagchi, M., Bagchi, D., and Stohs, S.J. (2002). Safety and mechanism of appetite suppression by a novel hydroxycitric acid extract (HCASX). *Molecular and Cellular Biochemistry*, 238(1-2), 89-103.
- Ohkoshi, E., Miyazaki, H., Shindo, K., Watanabe, H., Yoshida, A., and Yajima, H. (2007). Constituents from the leaves of *Nelumbo nucifera* stimulate lipolysis in the white adipose tissue of mice. *Planta Medica*, 73(12), 1255-1259.
- Oktay, M., Gülcin, I., Kufrevioglu, Ö.I. (2003). Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *Lebensmittel-Wissenschaft und Technologie*, 36(2), 263–271.
- Omole, O.O., Nappert, G., Naylor, J.M., Zello, G.A. (2001). Both L- and D-lactate contribute to metabolic acidosis in diarrheic calves. *Journal of Nutrition*, 131(8), 2128-2131.
- Ono, Y., Hattori, E., Fukaya, Y., Imai, S., and Ohizumi, Y. (2006). Anti-obesity effect of *Nelumbo nucifera* leaves extract in mice and rats. *Journal of Ethnopharmacology*, 106(2), 238-244.
- Osada, K., Suzuki, T., Kawakami, Y., Senda, M., Kasai, A., Sami, M., Ohta, Y., Kanda, T., and Ikeda, M. (2006). Dose-dependent hypocholesterolemic actions of dietary apple phenol in rats fed cholesterol. *Lipids*, 41(2), 133-139.
- Othman, A., Ismail, A., Abdul Ghani, N., and Adenan, I. (2007). Antioxidant capacity and phenolic content of cocoa beans. *Food Chemistry*, 100(4), 1523-1530.
- Ouchi, N., Kihara, S., Arita, Y., Maeda, K., Kuriyama, H., Okamoto, Y., Hotta, K., Nishida, M., Takahashi, M., Nakamura, T., Yamashita, S., Funahashi, T., and Matsuzawa, Y. (1999). Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation*, 100(25), 2473-2476.

- Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, Hotta K, Nishida M, Takahashi M, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y. (2000). Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- κ B signaling through a cAMP-dependent pathway. *Circulation*, 102(11), 1296-1301.
- Ouchi, N., Kihara, S., Funahashi, T., Matsuzawa, Y., and Walsh, K. (2003). Obesity, adiponectin and vascular inflammatory disease. *Current Opinion in Lipidology*, 14(6), 561-566.
- Oyejide, O.O., and Olushola, L. (2005). Hepatoprotective and antioxidant properties of extract of *Carmellia sinensis* (black tea) in rats. *African Journal of Biotechnology*, 4(12), 1432-1438.
- Ozata, M., Mergenb, M., Oktenlib, C., Aydinc, A., Sanisoglu, S.Y., Bolua, E., Yilmazb, M.I., Sayalc, A., Isimerc, A., and Ozdemira, C. (2002). Increased oxidative stress and hypozincemia in male obesity. *Clinical Biochemistry*, 35(8), 627-631.
- Padwal, R.S., and Majumdar, S.R. (2007). Drug treatments for obesity: orlistat, sibutramine, and rimonabant. *Lancet*, 369(9555), 71-77.
- Pagotto, U., Vanuzzo, D., Vicennati, V., and Pasquali, R. (2008). Pharmacological therapy of obesity. *Italian Journal of Cardiology*, 9(4 Suppl 1), 83-93.
- Pak Dek, M.S., Abdul Hamid, A., Osman, A., and Soh, C.S. (2008). Inhibitory effect of *Morinda citrifolia* L on lipoprotein lipase activity. *Journal of Food Science*, 73(8), 595-598.
- Pak Dek, M.S., Osman, A., Sahib, N.G., Saari, N., Markom, M., Abdul Hamid, A., and Anwar, F. (2011). Effects of extraction techniques on phenolic components and antioxidant activity of Mengkudu (*Morinda citrifolia* L.) leaf extracts. *Journal of Medicinal Plants Research*, 5(20), 5050-5057.
- Park, Y.S., Yoon, Y. and Ahn, H.S. (2007). *Platycodon grandiflorum* extract represses up-regulated adipocyte fatty acid binding protein triggered by a high fat feeding in obese rats. *World Journal of Gastroenterology*, 13(25), 3493-3499.
- Park, M.H., Kim, M.J., Cho, W.I., Chang, P.S., and Lee, J.H. (2009). Effects of treatments on the volatile distribution in *Artemisia princeps* Pampan. *Korean Journal of Food Science and Technology*, 41(5), 587-591.

- Permana, P.A., Del Parigi, A., and Tataranni, P.A. (2004). Microarray gene expression profiling in obesity and insulin resistance. *Nutrition*, 20(1), 134-138.
- Perry, B., Zhang, J., Sun, C., Saleh, T., and Wang, Y. (2012). Liuwei dihuang lowers body weight and improves insulin and leptin sensitivity in obese rats. *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 847167, 8 pages.
- Perumal, V., A. Hamid, A., Ismail, A., Saari, K., Abas, F., Ismail, I.S., Maulidiani, H. Lajis, N., Khatib, A. (2014). Effect Of Cosmos Caudatus Kunth Leaves On The Lipid Profile Of A Hyperlipidemia-Induced Animal Model. *Journal of Food Chemistry and Nutrition*, 2(1), 43-51.
- Pérusse, L., Després, J.P., Lemieux, S., Rice, T., Rao, D.C., Bouchard, C. (1996). Familial aggregation of abdominal visceral fat level: results from the Quebec family study. *Metabolism*, 45(3), 378-382.
- Pittler, M.H., Schmidt, K., and Ernst, E. (2005). Adverse events of herbal food supplements for body weight reduction: systematic review. *Obesity Review*, 6(2), 93-111.
- Podsędek, A., Majewska, I., Redzynia, M., Sosnowska, D., Koziołkiewicz, M. (2014). In vitro inhibitory effect on digestive enzymes and antioxidant potential of commonly consumed fruits. *Journal of Agriculture and Food Chemistry*, 62(20), 4610-4617.
- Pollare, T., Vessby, B., Lithell, H. (1991). Lipoprotein lipase activity in skeletal muscle is related to insulin sensitivity. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 11(5), 1192-1203.
- Posmyk, M.M., Kontek, R., and Janas, K.M. (2009). Antioxidant enzymes activity and phenolic compounds content in red cabbage seedlings exposed to copper stress. *Ecotoxicology and Environmetal safety*, 72(2), 596-602.
- Potenza, M.A., Marasciulo, F.L., Tarquinio, M., Tiravanti, E., Colantuono, G., Federici, A., Kim, J.A., Quon, M.J., and Montagnani, M. (2007). EGCG, a green tea polyphenol, improves endothelial function and insulin sensitivity, reduces blood pressure, and protects against myocardial I/R injury in SHR. *American Journal of Physiology - Endocrinology and Metabolism*, 292(5), 1378-1387.
- Poulos, S.P., Dodson, M.V., and Hausman, G.J. (2010). Cell line models for differentiation: preadipocytes and adipocytes. *Experimental Biology and Medicine (Maywood)*, 235(10), 1185-1193.

- Prentice, A.M. and Jebb, S.A. (2003). Fast foods, energy density and obesity: a possible mechanistic link. *Obesity Reviews*, 4(4), 187-194.
- Qader, S.W., Abdulla, M.A., Chua, L.S., Najim, N., Zain, M.M., and Hamdan, S. (2011). Antioxidant, total phenolic content and cytotoxicity evaluation of selected Malaysian plants. *Molecules*, 16(4), 3433-3443.
- Quettier-Deleu, C., Gressier, B., Vasseur, J., Dine, T., Brunet, C., Luyckx, M., Cazin, M., Cazin, J.C., Bailleul, F., and Trotin, F. (2000). Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. *Journal of Ethnopharmacology*, 72(1-2), 35-42.
- Qureshi, K., and Abrams, G.A. (2007). Metabolic liver disease of obesity and role of adipose tissue in the pathogenesis of nonalcoholic fatty liver disease. *World Journal of Gastroenterology*, 13(26), 3540-3553.
- Raftery, D., and G. A. Nagana Gowda, (2008). An Approaching New Wave of Multicomponent Biomarker Diagnostics?. *Journal of Urology*, 179(6), 2089-2090.
- Ragasa, C.Y., Nacpil, Z.D., Penalosa, B.A., Coll, J.C., and Rideout, J.A. (1997). Antimutagen and antifungal compounds from *Cosmos caudatus*. *Philippine journal of science*, 126(3), 199-206.
- Ragasa, C.Y., Nacpil, Z.D. Penalosa, B.A., Coll, J.C. and Rideout, J.A. (1999). Antimutagen and antifungal compounds from *Cosmos caudatus*. *Philippines Journal of Science*, 126(3), 199-206.
- Rahalison, L., Hamburger, M., Hostettmann, K., Monod, M., and Frenk, E. (1991). A bioautographic agar overlay method for the detection of antifungal compounds from higher plants. *Phytochemical Analysis*, 2(5), 199-203.
- Rahman, K. (2003). Garlic and aging: new insights into an old remedy. *Ageing Research Reviews*, 2(1), 39-56.
- Rains, T.M., Agarwal, S., Maki, K.C. (2011). Antiobesity effects of green tea catechins: a mechanistic review. *Journal of Nutritional Biochemistry*, 22(1), 1-7.
- Ramgopal, M., Attitala, I.H., Avinash, P., and Balaji, M. (2010). Evaluation of Antilipidemic and Anti Obesity Efficacy of *Bauhinia purpurea* Bark Extract on Rats Fed with High Fat Diet. *Academic Journal of Plant Sciences*, 3(3), 104-107.

- Rani, N., Sharma, S.K., and Vasudeva, N. (2012). Assessment of Antiobesity Potential of Achyranthes aspera Linn. Seed. *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 715912, 7 pages.
- Rani, N., Vasudeva, N., and Sharma, S.K. (2012). Quality assessment and anti-obesity activity of *Stellaria media* (Linn.) Vill. *BMC Complementary and Alternative Medicine*, 12(145).
- Rasdi, N.H., Samah, O.A., Sule, A., and Ahmed, Q.U. (2010). Antimicrobial studies of *Cosmos caudatus* Kunth. (Compositae). *Journal of Medicinal Plant Research*, 4(8), 669-673.
- Ratnawati, R., Indra, M.R., and Satuman, A. (2007). Epigallocatechin gallate of green tea inhibits proliferation, differentiation and TNF- α in the primary human visceral preadipocytes culture. *Majalah Ilmu Faal Indonesia*, 6(3), 160–168.
- Rayalam, S., Della-Fera, M.A., Baile, C.A. (2008). Phytochemicals and regulation of the adipocyte life cycle. *Journal of Nutritional Biochemistry*, 19(11), 717-726.
- Rebouche, C.J., and Seim, H. (1998). Carnitine metabolism and its regulation in microorganisms and mammals. *Annual Review of Nutrition*, 18, 39-61.
- Redinger, R.N. (2009). Fat storage and the biology of energy expenditure. *Translational Research*, 154(2), 52–60.
- Reinehr, T., Roth, C.L., Alexy, U., Kersting, M., Kiess, W., and Andler, W. (2005). Ghrelin levels before and after reduction of overweight due to a low-fat high-carbohydrate diet in obese children and adolescents. *International Journal of Obesity (London)*, 29(4), 362-368.
- Reitman, A., Friedrich, I., Ben Amotz, A., and Levy, Y. (2002). Low plasma antioxidants and normal plasma B vitamins and homocysteine in patients with severe obesity. *Israel Medical Association Journal*, 4(8), 590-593.
- Rezzi, S., Ramadan, Z., Fay, L.B., and Kochhar, S. (2007). Nutritional metabolomics: applications and perspectives. *Journal of Proteome Research*, 6(2), 513-525.
- Rivera, L., Morón, R., Sánchez, M., Zarzuelo, A., and Galisteo, M. (2008). Quercetin ameliorates metabolic syndrome and improves the inflammatory status in obese Zucker rats. *Obesity*, 16(9), 2081-2087.

- Roh, C., and Jung, Y. (2012). Screening of crude plant extracts with anti-obesity activity. *International Journal of Molecular Sciences*, 13(2), 1710–1719.
- Roongpisuthipong, C., Kantawan, R., Roongpisuthipong, W. (2007). Reduction of adipose tissue and body weight: effect of water soluble calcium hydroxycitrate in *Garcinia atroviridis* on the short term treatment of obese women in Thailand. *Asia Pacific Journal of Clinical Nutrition*, 16(1), 25-29.
- Rosenbaum, M., Nicolson, M., Hirsch, J., Heymsfield, S.B., Gallagher, D., Chu, F., and Leibel, R.L. (1996). Effects of gender, body composition, and menopause on plasma concentrations of leptin. *Journal of Clinical Endocrinology and Metabolism*, 81(9), 3424-3427.
- Ross, G.H., Hovde, L.B., Ibrahim, K.H., Ibrahim, Y.H., and Rotschafer, J.C. (2001). Comparison of once-daily versus twice-daily administration of cefdinir against typical bacterial respiratory tract pathogens. *Antimicrobial Agents and Chemotherapy*, 45(10), 2936-2938.
- Rudich, A., Tirosh, A., Potashnik, R., Hemi, R., Kanety, H., and Bashan, N. (1998). Prolonged oxidative stress impairs insulin-induced GLUT4 translocation in 3T3-L1 adipocytes. *Diabetes*, 47(10), 1562-1569.
- Ryan, A.S., Berman, D.M., Nicklas, B.J., Sinha, M., Gingerich, R.L., Meneilly, G.S., Egan, J.M., and Elahi, D. (2003). Plasma adiponectin and leptin levels, body composition, and glucose utilization in adult women with wide ranges of age and obesity. *Diabetes Care*, 26(8), 2383-2388.
- Ryo, M., Nakamura, T., Kihara, S., Kumada, M., Shibasaki, S., Takahashi, M., Nagai, M., Matsuzawa, Y., Funahashi, T. (2004). Adiponectin as a biomarker of the metabolic syndrome. *Circulation*, 68(11), 975-981. Saely, C.H., Geiger, K., Drexel, H. (2012). Brown versus white adipose tissue: a mini-review. *Gerontology*, 58(1), 15–23.
- Sahib, N.G., Abdul Hamid, A., Kitts, D., Purnama, M., Saari, N., and Abas, F. (2011). The effects of *morinda citrifolia*, *momordica charantia* and *centella asiatica* extracts on lipoprotein lipase and 3t3-l1 preadipocytes. *Journal of Food Biochemistry*, 35(4), 1186–1205.
- Sahib, N.G., Hamid, A.A., Saari, N., Abas, F., Pak Dek, M.S., Rahim, M. (2012). Anti-Pancreatic Lipase and Antioxidant Activity of Selected Tropical Herbs. *International Journal of Food Properties*, 15(3), 569-578.
- Salek, R.M., Maguire, M.L., Bentley, E., Rubtsov, D.V., Hough, T., Cheeseman, M., Nunez, D., Sweatman, B.C., Haselden, J.N., Cox,

- R.D., Connor, S.C., and Griffin, J.L. (2007). A metabolomic comparison of urinary changes in type 2 diabetes in mouse, rat, and human. *Physiological Genomics*, 29(2), 99-108.
- Samy, J., Sugumaran, M., Lee, K.L. (2005). Herbs of Malaysia: An Introduction to the Medicinal, Culinary, Aromatic and Cosmetic Use of Herbs. Times Editions. ISBN 983-300-179-3.
- Sanins, S.M., Nicholson, J.K., Elcombe, C., and Timbrell, J.A. (1990). Hepatotoxin-induced hypertaurinuria: a proton NMR study. *Archives of Toxicology*, 64(5), 407-411.
- Sato, M., Uzu, K., Yoshida, T., Hamad, E.M., Kawakami, H., Matsuyama, H., Abd El-Gawad, I.A., and Imaizumi, K. (2008). Effects of milk fermented by Lactobacillus gasseri SBT2055 on adipocyte size in rats. *British Journal of Nutrition*, 99(5), 1013-1017.
- Satoh, H. (2994). Cardioprotective actions of taurine against intracellular and extracellular calcium-induced effects. *Advances in Experimental Medicine and Biology*, 359, 181–196.
- Saxena, M., Saxena, J., Nema, R., Singh, D., and Gupta, A. (2013). Phytochemistry of Medicinal Plants. *Journal of Pharmacognosy and Phytochemistry*, 1(6), 168-182.
- Scalbert, A., Manach, C., Morand, C., and Remesy, C., and Jimenez, L. (2005). Dietary Polyphenols and the Prevention of Diseases. *Critical Reviews in Food Science and Nutrition*, 45(4), 287-306.
- Scheen, A.J., and Lefébvre, P.J. (1999). Pharmacological treatment of obesity: present status. *International Journal of Obesity and Related Metabolic Disorders*, 23(Suppl 1), 47-53.
- Schirra, H.J., Anderson, C.G., Wilson, W.J., Kerr, L., Craik, D.J., Waters, M.J., and Lichanska, A.M. (2008). Altered metabolism of growth hormone receptor mutant mice: a combined NMR metabonomics and microarray study. *PLoS One*, 3(7), Article ID e2764.
- Schramm, D.D., Karim, M., Schrader, H.R., Holt, R.R., Cardetti, M., Keen, C.L. (2003). *Journal of Agriculture and Food Chemistry*, 51(6), 1732-1735.
- Schwartz, M.W., Baskin, D.G., Bukowski, T.R., Kuijper, J.L., Foster, D., Lasser, G., Prunkard, D.E., Porte, D., Woods, S.C., Seeley, R.J., Weigle, D.S. (1996). Specificity of leptin action on elevated blood glucose levels and hypothalamic neuropeptide Y gene expression in ob/ob mice. *Diabetes*, 45(4), 531-535.

- Senior, J.R. (1964). Intestinal absorption of fats. *Journal of Lipid Research*, 5, 495-521.
- Serkova, N.J., Jackman, M., Brown, J.L., Liu, T., Hirose, R., Roberts, J.P., Maher, J.J., and Niemann, C.U. (2006). Metabolic profiling of livers and blood from obese Zucker rats. *Journal of Hepatology*, 44(5), 956–962.
- Shannon, J.R., Gottesdiener, K., Jordan, J., Chen, K., Flattery, S., Larson, P.J., Candelore, M.R., Gertz, B., Robertson, D., and Sun, M. (1999). Acute effect of ephedrine on 24-h energy balance. *Clinical Science (London)*, 96(5), 483-491.
- Shearer, J., Duggan, G., Weljie, A., Hittel, D.S., Wasserman, D.H., and Vogel, H.J. (2008). Metabolomic profiling of dietary-induced insulin resistance in the high fat-fed C57BL/6J mouse. *Diabetes, Obesity and Metabolism*, 10(10), 950-958.
- Shehzad, A., Iqbal, W., Shehzad, O., and Lee, Y.S. (2012). Adiponectin: Regulation of its production and its role in human diseases. *Hormones*, 11(1), 8-20.
- Shiiya, T., Nakazato, M., Mizuta, M., Date, Y., Mondal, M.S., Tanaka, M., Nozoe, S., Hosoda, H., Kangawa, K., and Matsukura, S. (2002). Plasma ghrelin levels in lean and obese humans and the effect of glucose on ghrelin secretion. *Journal of Clinical Endocrinology and Metabolism*, 87(1), 240-244.
- Shimada, M., Mochizuki, K., Sakurai, N., and Goda, T. (2007). Dietary supplementation with epigallocatechin gallate elevates levels of circulating adiponectin in non-obese Type 2 diabetic Gotoe Kakizaki rats. *Bioscience, Biotechnology and Biochemistry*, 71(8), 2079-2082.
- Shimoda, H., Seki, E., and Aitani, M. (2006). Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complementary and Alternative Medicine*, 6, 9.
- Shimomura, I., Funahashi, T., Takahashi, M., Maeda, K., Kotani, K., Nakamura, T., Yamashita, S., Miura, M., Fukuda, Y., Takemura, K., Tokunaga, K., and Matsuzawa, Y. (1996). Enhanced expression of PAI-1 in visceral fat: possible contributor to vascular disease in obesity. *Natural Medicines*, 2(7), 800-803.
- Shimura, S., Itoh, Y., Yamashita, A., Kitano, A., Hatano, T., Yoshida, T., Okuda, T. (1994). Inhibitory effects of flavonoids on lipase. *Nippon Shokuhin Kogyo Gakkaishi*, 41, 847-850.

- Shui, G., Leong, L.P., and Shih, P.W. (2005) Rapid screening and characterization of antioxidants of *Cosmos caudatus* using liquid chromatography coupled with mass spectrometry. *Journal of Chromatogr B*, 827(1), 127–138.
- Shuib, N.H., Shaari, K., Khatib, A., Maulidiani, Kneer, R., Zareen, S., Mohd. Raof, S., N., Lajis and Neto, V. (2011). Discrimination of young and mature leaves of *Melicope ptelefolia* using ^1H NMR and multivariate data analysis. *Food Chemistry*, 126(2), 640-645.
- Singhal, A., Jamieson, N., Fewtrell, M., Deanfield, J., Lucas, A., and Sattar, N. (2005). Adiponectin predicts insulin resistance but not endothelial function in young, healthy adolescents. *Journal of Clinical Endocrinology and Metabolism*, 90(8), 4615-4521.
- Sigleton, V.L., Orthofer, R., and Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin- Ciocalteu reagent. *Method Enzymology*, 299, 152-178.
- Singleton, V.L., and Rossi, J.A. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16(3), 144-158.
- Škottová, N., Kazdováb, L., Oliyarnyk, O., Večeřaa, R., Sobolová, L., and Ulrichovác, J. (2004). Phenolics-rich extracts from *Silybum marianum* and *Prunella vulgaris* reduce a high-sucrose diet induced oxidative stress in hereditary hypertriglyceridemic rats. *Pharmacological Research*, 50(2), 123-130.
- Slanc, P., Doljak, B., Kretf, S., Lunder, M., Janes, D., and Strukelj, B. (2009). Screening of selected food and medicinal plant extracts for pancreatic lipase inhibition. *Phytotherapy Research*, 23(6), 874-877.
- Slovacek, L., Pavlik, V., and Slovackova, B. (2008). The effect of sibutramine therapy on occurrence of depression symptoms among obese patients. *Nutrition, Metabolism and Cardiovascular Diseases*, 18(8), 43-44.
- Snow, V., Barry, P., Fitterman, N., Qaseem, A., Weiss, K. (2005). Clinical Efficacy Assessment Subcommittee of the American College of Physicians. Pharmacologic and surgical management of obesity in primary care: a clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*, 142(7), 525-531.

- Soares, J.B., Roncon-Albuquerque R., Leite-Moreira A. (2008). Ghrelin and ghrelin receptor inhibitors: agents in the treatment of obesity. *Expert Opinion Therapeutic Patents*, 12(9), 1177-1189.
- Sogawa, M., Seura, T., Kohno, S., Hirasaka, K., Yamaguchi, Y., Takagaki, R., Harada, A., Okumura, Y., Yamamoto, S., Kishi, K., and Nikawa, T. (2009). Awa (Tokushima) lactate-fermented tea as well as green tea enhance the effect of diet restriction on obesity in rats. *Journal of Medical Investigation*, 56(1-2), 42-48.
- Song, X., Wang, J., Wang, P., Tian, N., Yang, M., and Kong, L. (2013). ¹H NMR-based metabolomics approach to evaluate the effect of Xue-Fu-Zhu-Yu decoction on hyperlipidemia rats induced by high-fat diet. *Journal of Pharmaceutical and Biomedical Analysis*, 78-79, 202-210.
- Speakman, J.R. (2004). Obesity: The Integrated Roles of Environment and Genetics. *Journal of Nutrition*, 134(8), 2090-2105.
- Spratlin, J. L., Serkova, N.J., and Eckhardt, S.G. (2009). Clinical applications of metabolomics in oncology: a review. *Clinical Cancer Research*, 15(2), 431-440.
- Staiger, H., and Häring, H.U. (2005). Adipocytokines: fat-derived humoral mediators of metabolic homeostasis. *Experimental and Clinical Endocrinology and Diabetes*, 113(2), 67-79.
- Statnick, M.A., Beavers, L.S., Conner, L.J., Corominola, H., Johnson, D., Hammond, C.D., Rafaeloff-Phail, R., Seng, T., Sutter, T.M., Sluka, J.P., Ravussin, E., Gadski, R.A., and Caro, J.F. (2000). Decreased expression of apM1 in omental and subcutaneous adipose tissue of human with type 2 diabetes. *International Journal of Experimental Diabetes Research*, 1(2), 81-88.
- Stefan, N., and Stumvoll, M. (2002). Adiponectin – its role in metabolism and beyond. *Hormone and Metabolic Research*, 34(9), 469-474.
- Stein, C.J., and Colditz, G.A. (2004). The Epidemic of Obesity. *The Journal of Clinical Endocrinology and Metabolism*, 89(6), 2522-2525.
- Steppan, C.M., Bailey, S.T., Bhat, S., Brown, E.J., Banerjee, R.R., Wright, C.M., Patel, H.R., Ahima, R.S., and Lazar, M.A. (2001). The hormone resistin links obesity to diabetes. *Nature*, 409(6818), 307-312.
- Storey, M.L., Forshee, R.A., Weaver, A.R., and Sansalone, W.R. (2003). Demographic and lifestyle factors associated with body mass index

- among children and adolescents. *International Journal of Nutrition and Food Sciences*, 54(6), 491-503.
- Stunkard, A.J. (1996). Current views on obesity. *American Journal of Medicine*, 100(2), 230-236.
- Stunkard, A.J. (2002). Binge eating and the night eating disorder. In *Handbook of Obesity Treatment*. Wadden, T.A., and Stunkard, A.J. (Eds). New York, Guilford Press, 107–121.
- Stunkard, A.J. and Allison, K.C. (2003). Two forms of disordered eating in obesity: binge eating and night eating. *International Journal of Obesity and Related Metabolic Disorders*, 27(1), 1-12.
- Sugino, N. (1990). Evaluation of renal function in chronic renal disease with special reference to creatinine. *Japanese Journal of Nephrology*, 32(7), 745–750.
- Sugiyama, K., Akai, H., and Muramatsu, K. (1986). Effects of methionine and related compounds on plasma cholesterol level in rats fed a high cholesterol diet. *Journal of Nutritional Science and Vitaminology*, 32(5), 537-549.
- Sukrasno, S., Fidriany, I., Anggadiredja, K., Handayani, W. A., and Anam, K. (2011). Influence of drying method on flavonoid content of *Cosmos caudatus* (Kunth) leaves. *Research Journal of Medicinal Plant*, 5(2), 189-195.
- Sun, T. and Ho, C.T. (2005). Antioxidant activities of buckwheat extracts. *Food Chemistry*, 90(4), 743-749.
- Sun, Y.J., Wang, H.P., Liang, Y.J., Yang, L., Li, W., and Wu, Y.J. (2012) An NMR-based metabonomic investigation of the subacute effects of melamine in rats. *Journal of Proteome Research*, 11(4), 2544–2550.
- Sung, Y.Y., Yoon, T., Yang, W.K., Kim, S.J., Kim, D.S., and Kim, H.K. (2013). The Antioesity Effect of *Polygonum aviculare* L. Ethanol Extract in High-Fat Diet-Induced Obese Mice. *Evidence-Based Complementary and Alternative Medicine*, 2013, Article ID 626397, 11 pages.
- Svendsen A. (2000). Lipase protein engineering. *Biochimica Biophysica Acta*, 1543(2), 223-238.
- Tamura, A., Fukushima, M., Shimada, K.I., Han, K.H., Sekikawa, M., Watanabe, S., Nakano, N., Matsumoto, M., and Chiji, H. (2004). Cholesterol metabolism in rat is affected by protocatechuic acid. *Journal of Nutritional Science and Vitaminology*, 50(1), 13-18.

- Temple, N.J. and Gladwin, K.K. (2003). Fruit, vegetables, and the prevention of cancer: research challenges. *Nutrition*, 19(5), 467-470.
- Tentolouris, N., Kokkinos, A., Tsigos, C., Kyriaki, D., Doupis, J., Raptis, S.A., and Katsilambros, N. (2004). Differential effects of high-fat and high-carbohydrate content isoenergetic meals on plasma active ghrelin concentrations in lean and obese women. *Hormone and Metabolic Research*, 36(8), 559-563.
- Thielecke, F., Boschmann, M. (2009). The potential role of green tea catechins in the prevention of the metabolic syndrome – a review. *Phytochemistry*, 70(1), 11–24.
- Thurairajah, P.H., Syn, W.K., Neil, D.A., Stell, D., and Haydon, G. (2005). Orlistat (xenical) induced subacute liver failure. *European Journal of Gastroenterology and Hepatology*, 17(12), 1437-1438.
- Tian, W.X., Li, L.C., Wu, X.D. and Chen, C.C. (2004). Weight reduction by Chinese medicinal herbs may be related to inhibition of fatty acid synthase. *Life Science* 74(19), 2389-2399.
- Tian, N., Wang, J., Wang, P., Song, X., Yang, M., and Kong, L. (2013). NMR-based metabonomic study of Chinese medicine Gegen Qinlian Decoction as an effective treatment for type 2 diabetes in rats. *Metabolomics*, 9(6), 1228-1242.
- Tong, L.T., Katakura, Y., Kawamura, S., Baba, S., Tanaka, Y., Udon, M., Kondo, Y., Nakamura, K., Imaizumi, K., and Sato, M. (2010). Effects of Kurozu concentrated liquid on adipocyte size in rats. *Lipids in Health and Disease*, 9(134).
- Trivedi, D.K., and Iles, R.K. (2012). The Application of SIMCA P+ in Shotgun Metabolomics Analysis of ZIC®HILIC-MS Spectra of Human Urine - Experience with the Shimadzu IT-TOF and Profiling Solutions Data Extraction Software. *Journal of Chromatography Separation Technique*, 3(6).
- Tschöp, M., Weyer, C., Tataranni, P.A., Devanarayan, V., Ravussin, E., and Heiman, M.L. (2001). Circulating ghrelin levels are decreased in human obesity. *Diabetes*, 50(4), 707-709.
- Tsuda, T., Ueno, Y., Aoki, H., Koda, T., Horio, F., Takahashi, N., Kawada, T., and Osawa, T. (2004). Anthocyanin enhances adipocytokine secretion and adipocyte-specific gene expression in isolated rat adipocytes. *Biochemical and Biophysical Research Communications*, 316(1), 149-157.

- Tsujita, T., Takaichi, H., Takaku, T., Aoyama, S., and Hiraki, J., (2006). Antiobesity action of e-polylysine, a potent inhibitor of pancreatic lipase. *Journal of Lipid Research*, 47(8), 1852–1858.
- Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., and Gordon, J.I. (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*, 444(7122), 1027–1031.
- Turpin, P. (1985). A double blind study of the effectiveness of Beaufor betaine citrate ampules in the treatment of type IV hyperlipidemias. *Sem Hop*. 61, 2420-2434.
- Ulrich, E.L., Akutsu, H., Doreleijers, J.F., Harano, Y., Ioannidis, Y.E., Lin, J., Livny, M., Mading, S., Maziuk, D., Miller, Z., Nakatani, E., Schulte, C.F., Tolmie, D.E., Kent Wenger, R., Yao, H., and Markley, J.L. (2007). BioMagResBank. *Nucleic Acids Research*, 36(Database issue), 402-408.
- Utsunomiya, H., Yamakawa, T., Kamei, J., Kadonosono, K., and Tanaka, S. (2005). Anti-hyperglycemic effects of plum in a rat model of obesity and type 2 diabetes, Wistar fatty rat. *Biomedical Research*, 26(5), 193-200.
- Van Gaal, L., Mertens, I., Ballaux, D., and Verkade, H. (2004). Modern, new pharmacotherapy for obesity. A gastrointestinal approach. *Best Practice and Research Clinical Gastroenterology*, 18(6), 1049-1072.
- Van Guilder, G.P., Hoetzer, G.L., Greiner, J.J., Stauffer, B.L., and DeSouza, C.A. (2006). Influence of Metabolic Syndrome on Biomarkers of Oxidative Stress and Inflammation in Obese Adults. *Obesity (Silver Spring)*, 14(12), 2127-2131.
- Van Heerden, F.R., Vleggaar, R., Horak, R.M., Learmonth, R.A., Maharaj, V.J., and Whittal, R.D. (1998). Pharmaceutical compositions having appetite-suppressant activity. US patent US6376657 B1.
- Van Heerden, F.R. (2008). *Hoodia gordonii*: a natural appetite suppressant. *Journal of Ethnopharmacology*, 119(3), 434-437.
- Vaziri, N.D., Wang, X.Q., Oveis, F., and Rad, B. (2000). Induction of oxidative stress by glutathione depletion causes severe hypertension in normal rats. *Hypertension*, 36(1), 142-146.
- Verpoorte, R., Choi, Y.H., and Kim, H.K. (2007). NMR-based metabolomics at work in phytochemistry. *Phytochemistry Reviews*, 6(1), 3–14.
- Vetter, M.L., Faulconbridge, L.F., Webb, V.L., and Wadden, T.A. (2010).

- Behavioral and pharmacologic therapies for Obesity. *Nature Reviews Endocrinology*, 6(10), 578-588.
- Viant, M.R. (2008). Recent developments in environmental metabolomics. *Molecular BioSystems*, 4(10), 980-986.
- Vice, E., Privette, J.D., Hickner, R.C., and Barakat, H.A. (2005). Ketone body metabolism in lean and obese women. *Metabolism*, 54(11), 1542-1545.
- Vickers, S.P., Jackson, H.C., and Cheetham, S.C. (2011). The utility of animal models to evaluate anti-obesity agents. *British Journal of Pharmacology*, 164, 1248-1262.
- Vucenik, I., and Stains, J.P. (2012). Obesity and cancer risk: evidence, mechanisms, and recommendations. *Annals of the New York Academy of Sciences*, 1271(1), 37-43.
- Wadden, T.A., and Stunkard, A.J. (1985). Social and psychological consequences of obesity. *Annals of Internal Medicine*, 103(6 (Pt 2)), 1062-1067.
- Wadden, T.A. (1993). Treatment of obesity by moderate and severe caloric restriction. Results of clinical research trials. *Annals of Internal Medicine*. 119(7 Pt 2), 688-693.
- Walker, J.B. (1961). Metabolic control of creatine biosynthesis. II. Restoration of transamidinase activity following creatine repression. *Journal of Biological Chemistry*, 236, 493-498.
- Wang, Z., Klipfell, E., Bennett, B.J., Koeth, R., Levison, B.S., Dugar, B., Feldstein, A.E., Britt, E.B., Fu, X., Chung, Y.M., Wu, Y., Schauer, P., Smith, J.D., Allayee, H., Tang, W.H., DiDonato, J.A., Lusis, A.J., and Hazen, S.L. (2011). Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*, 472(7341), 57-63.
- Wang, X., Zhang, A., and Sun, H. (2012). Future perspectives of Chinese medical formulae: chinomedomics as an effector. *OMICS*, 16(7-8), 414-421.
- Wang, H., Bai, J., Chen, G., Li, W., Xiang, R., Su, G., and Pei, Y. (2013). A metabolic profiling analysis of the acute hepatotoxicity and nephrotoxicity of Zhusha Anshen Wan compared with cinnabar in rats using ^1H NMR spectroscopy. *Journal of Ethnopharmacology*, 146(2), 572-580.
- Wang, X., Wang, Q., Zhang, A., Zhang, F., Zhang, H., Sun, H., Cao, H., and Zhang, H. (2013). Metabolomics study of intervention effects of

- Wen-Xin-Formula using ultra high-performance liquid chromatography/mass spectrometry coupled with pattern recognition approach. *Journal of Pharmaceutical and Biomedical Analysis*, 74, 22-30.
- Wang, P.R., Wang, J.S., Yang, M.H., and Kong, L.Y. (2014). Neuroprotective effects of Huang-Lian-Jie-Du-Decoction on ischemic stroke rats revealed by ^1H NMR metabolomics approach. *Journal of Pharmaceutical and Biomedical Analysis*, 88, 106-116.
- Wei, L., Liao, P., Wu, H., Li, X., Pei, F., Li, W., and Wu, Y. (2008). Toxicological effects of cinnabar in rats by NMR-based metabolic profiling of urine and serum. *Toxicology and Applied Pharmacology*, 227(3), 417-429.
- Wei, H., Pasman, W., Rubingh, C., Wopereis, S., Tienstra, M., Schroen, J., Wang, M., Verheij, E., Van der Greef, J. (2012). Urine metabolomics combined with the personalized diagnosis guided by Chinese medicine reveals subtypes of pre-diabetes. *Molecular BioSystem*, 8(5), 1482-1491.
- Weigle, D.S. (2003). Pharmacological therapy of obesity: past, present, and future. *Journal of Clinical Endocrinology and Metabolism*, 88(6), 2462-2469.
- Weyer, C., Funahashi, T., Tanaka, S., Hotta, K., Matsuzawa, Y., Pratley, R.E., Tataranni, P.A. (2001). Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *Journal of Clinical Endocrinology and Metabolism*, 86(5), 1930-1935.
- White, R.T., Damm, D., Hancock, N., Rosen, B.S., Lowell, B.B., Usher, P., Flier, J.S., and Spiegelman, B.M. (1992). Human adipsin is identical to complement factor D and is expressed in high levels in adipose tissue. *Journal of Biological Chemistry*, 267(13), 9210-9213.
- Wickelgren I. (1998). Obesity: how big a problem?. *Science*, 280(5368), 1364-1367.
- Williams, H.R., Cox, I.J., Walker, D.G., North, B.V., Patel, V.M., Marshall, S.E., Jewell, D.P., Ghosh, S., Thomas, H.J., Teare, J.P., Jakobovits, S., Zeki, S., Welsh, K.I., Taylor Robinson, S.D., and Orchard, T.R. (2009) Characterization of inflammatory bowel disease with urinary metabolic profiling. *American Journal of Gastroenterology*, 104(6), 1435–1444.
- Wing, R.R., Lang, W., Wadden, T.A., Safford, M., Knowler, W.C., Bertoni, A.G., Hill, J.O., Brancati, F.L., Peters, A., and Wagenknecht, L.

- (2011). Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care*, 34(7), 1481-1486.
- Wishart, D.S., Tzur, D., Knox, C., Eisner, R., Guo, A.C., Young, N., Cheng, D., Jewell, K., Arndt, D., Sawhney, S., Fung, C., Nikolai, L., Lewis, M., Coutouly, M.A., Forsythe, I., Tang, P., Srivastava, S., Jeroncic, K., Stothard, P., Amegbey, G., Block, D., Hau, D.D., Wagner, J., Miniaci, J., Clements, M., Gebremedhin, M., Guo, N., Zhang, Y., Duggan, G.E., Macinnis, G.D., Weljie, A.M., Dowlatabadi, R., Bamforth, F., Clive, D., Greiner, R., Li, L., Marrie, T., Sykes, B.D., Vogel, H.J., Querengesser, L. (2007). HMDB: the human metabolome database. *Nucleic Acids Research*, 35(Database Issue), 521-526.
- Wishart, D.S. (2008). Metabolomics: applications to food science and nutrition research. *Trends in Food Science and Technology*, 19(9), 482-493.
- Wishart, D.S. (2008). Quantitative metabolomics using NMR. *Trends in Analytical Chemistry*, 27(3), 228-237.
- Wishart, D.S., Knox, C., Guo, A.C., Eisner, R., Young, N., Gautam, B., Hau, D.D., Psychogios, N., Dong, E., Bouatra, S., Mandal, R., Sinelnikov, I., Xia, J.G., Jia, L., Cruz, J.A., Lim, E., Sobsey, C.A., Srivastava, S., Huang, P., Liu, P., Fang, L., Peng, J., Fradette, R., Cheng, D., Tzur, D., Clements, M., Lewis, A., De Souza, A., Zuniga, A., Dawe, M., Xiong, Y.P., Clive, D., Greiner, R., Nazirova, A., Shaykhutdinov, R., Li, L., Vogel, H.J., Forsythe, I. (2009). HMDB: a knowledgebase for the human metabolome. *Nucleic Acids Research*, 37(Database Issue), 603-610.
- Wishart, D.S. (2011). Advances in metabolite identification. *Bioanalysis*, 3(15), 1769-1782.
- Wolfram, S., Wang, Y., and Thielecke, F. (2006). Anti-obesity effects of green tea: from bedside to bench. *Molecular Nutrition and Food Research*, 50(2), 176-187.
- Won, E.Y., Yoon, M.K., Kim, S.W., Jung, Y., Bae, H.W. (2013). Gender-Specific Metabolomic Profiling of Obesity in Leptin-Deficient ob/ob Mice by 1H NMR Spectroscopy. *PLoS ONE*, 8(10), e75998.
- Wong, H., and Schotz, M.C. (2002). The lipase gene family. *Journal of Lipid Research*, 43(7), 993-999.
- Woo, M.N., Bok, S.H., Lee, M.K., Kim, H.J., Jeon, S.M., Do, G.M., Shin, S.K., Ha, T.Y., and Choi, M.S. (2008). Anti-obesity and hypolipidemic effects of a proprietary herb and fiber combination

- (S&S PWL) in rats fed high-fat diets. *Journal of Medicinal Food*, 11(1), 169-178.
- Wood, P.D., Stefanick, M.L., Dreon, D.M., Frey Hewitt, B., Garay, S.C., Williams, P.T., Superko, H.R., Fortmann, S.P., Albers, J.J., Vranizan, K.M., Ellsworth, N.M., Terry, R.B., and Haskell, W.L. (1988). Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. *New England Journal of Medicine*, 319(18), 1173-1179.
- World Health Organisation (WHO), (1997). Obesity: preventing and managing the global epidemic. Report of a WHO consultation on obesity. Geneva: World Health Organisation.
- World Health Organization (WHO), (2003).
- World Health Organization (WHO), (2009). Obesity and overweight. Country and Regional Data. World Health Organization.
- World Health Organization (WHO), (2011).
- World Health Organization (WHO), (2014).
- Wu, G.Y., Gunasekara, A., Brunengraber, H., and Marliss, E.B. (1991). Effects of extracellular pH, CO₂, and HCO₃- on ketogenesis in perfused rat liver. *American Journal of Physiology*, 261(2 Pt 1), 221-226.
- Wu, B., Yan, S., Lin, Z., Wang, Q., Yang, Y., Yang, G., Shen, Z., and Zhang, W. (2008). Metabonomic study on ageing: NMR-based investigation into rat urinary metabolites and the effect of the total flavone of Epimedium. *Molecular Biosystem*, 4(8), 855-861.
- Xie, B., Waters, M.J., and Schirra, H.J. (2012). Investigating Potential Mechanisms of Obesity by Metabolomics. *Journal of Biomedicine and Biotechnology*, 2012, Article ID 805683, 10 pages.
- Xiong, Y., Shen, L., Liu, K.J., Tso, P., Xiong, Y., Wang, G., Woods, S.C., and Liu, M. (2010). Antiobesity and Antihyperglycemic Effects of Ginsenoside Rb1 in Rats. *Diabetes*, 59(10), 2505–2512.
- Xu, R.Y., Wan, Y.P., Tang, Q.Y., Wu, J., and Cai, W. (2008). The effects of high fat on central appetite genes in Wistar rats: a microarray analysis. *Clinica Chimica Acta*, 397(1-2), 96-100.
- Xu, E.Y., Schaefer, W.H., and Xu, Q. (2009). Metabolomics in pharmaceutical research and development: metabolites,

- mechanisms and pathways. *Current Opinion in Drug Discovery and Development*, 12(1), 40-52.
- Xu, J., Liu, C., Cai, S., Dong, J., Li, X., Feng, J., Chen, Z. (2013). Metabolomic Profilings of Urine and Serum from High Fat-Fed Rats via ^1H NMR Spectroscopy and Pattern Recognition. *Applied Biochemistry and Biotechnology*, 169(4), 1250-1261.
- Yamamoto, M., Shimura, S., Itoh, Y., Ohsaka, T., Egawa, M., and Inoue, S. (2000). Anti obesity effects of lipase inhibitor CT-II, an extract from edible herbs, Nomame Herba, on rats fed a high-fat diet. *International Journal of Obesity*, 24(6), 758-764.
- Yamauchi, T., Kamon, J., Waki, H., Terauchi, Y., Kubota, N., Hara, K., Mori, Y., Ide, T., Murakami, K., Tsuboyama-Kasaoka, N., Ezaki, O., Akanuma, Y., Gavrilova, O., Vinson, C., Reitman, M.L., Kagechika, H., Shudo, K., Yoda, M., Nakano, Y., Tobe, K., Nagai, R., Kimura, S., Tomita, M., Froguel, P., and Kadawaki, T. (2001). The fat derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nature Medicine*, 7(8), 941-946.
- Yang, C.S., Landau, J.M., Huang, M.T., and Newmark, H.L. (2001). Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annual Review of Nutrition*, 21, 381-406.
- Yang, M.H., Wang, C.H., and Chen, H.L. (2001). Green, oolong and black tea extracts module lipid metabolism in hyperlipidemia rats fed high sucrose diet. *Journal of Nutritional Biochemistry*, 12(1), 14-20.
- Yang, D.J., Chang, Y.Y., Hsu, C.L., Liu, C.W., Lin, Y.L., Lin, Y.H., Liu, K.C., Chen, Y.C. (2010). Antioesity and hypolipidemic effects of polyphenol-rich longan (*Dimocarpus longans* Lour.) flower water extract in hypercaloric dietary rats. *Journal of Agriculture and Food Chemistry*, 58(3), 2020-2027.
- Yang, M.Y., Peng, C.H., Chan, K.C., Yang, Y.S., Huang, C.N., and Wang, C.J. (2010). The hypolipidemic effect of *Hibiscus sabdariffa* polyphenols via inhibiting lipogenesis and promoting hepatic lipid clearance. *Journal of Agriculture and Food Chemistry*, 58(2), 850-859.
- Yao, L.H., Jiang, Y.M., Caffin N D'Arcy, B., Datta, N., Liu, X., Singanusong, R., and Xu, Y. (2006). Phenolic compounds in tea from Australian supermarkets. *Food Chemistry*, 96(4), 614-620.
- Ye, T., Mo, H., Shanaiah, N., Gowda, G.A., Zhang, S., and Raftery, D. (2009). Chemoselective ^{15}N tag for sensitive and high-resolution

- nuclear magnetic resonance profiling of the carboxyl-containing metabolome. *Analytical Chemistry*, 81(12), 4882-4888.
- Yoshikawa, M., Shimoda, H., Nishida, N., Takada, M., and Matsuda, H. (2002). Salacia reticulata and its polyphenolic constituents with lipaseinhibitory and lypophilic activities have mild antiobesity effects in rats. *Journal of Nutrition*, 132(7), 1819–1824.
- Yu, J.G., Javorschi, S., Hevener, A.L., Kruszynska, Y.T., Norman, R.A., Sinha, M., and Olefsky, J.M. (2002). The effect of thiazolidinediones on plasma adiponectin levels in normal, obese, and type 2 diabetic subjects. *Diabetes*, 51(10), 2968-2974.
- Yuan, G., Al-Shali, K.Z., and Hegele, R.A. (2007). Hypertriglyceridemia: its etiology, effects and treatment. *Canadian Medical Association Journal*, 176(8), 1113–1120.
- Yun, J.W. (2010). Possible anti-obesity therapeutics from nature – A review. *Phytochemistry*, 71(14-15), 1625-1641.
- Zai, H., and Maibach, H.I. (2006). Skin Antioxidants. In Antiaging: Physiology to Formulation. Carol Stream: Allured Publishing Corporation, 287-292.
- Zechner, R., Strauss, J., Frank, S., Wagner, E., Hofmann, W., Kratky, D. (2000). The role of lipoprotein lipase in adipose tissue development and metabolism. *International Journal of Obesity and Related Metabolic Disorders*, 24(Suppl 4), 53-56.
- Zeng, M., Liang, Y., Li, H., Wang, M., Wang, B., Chen, X., Zhou, N., Cao, D., and Wu, J. (2010). Plasma metabolic fingerprinting of childhood obesity by GC/MS in conjunction with multivariate statistical analysis. *Journal of Pharmaceutical and Biomedical Analysis*, 52(2), 265-272.
- Zhang, X., Wei, D., Yap, Y., Li, L., Guo, S., and Chen, F. (2007). Mass spectrometry based “omics” technologies in cancer diagnostics. *Mass Spectrometry Reviews*, 26(3), 403-431.
- Zhang, Y. F., Li, X. Y., Zou, D. J., Liu, W., Yang, J. L., Zhu, N., Huo, L., Wang, M., Hong, J., Wu, P., Ren, G., and Ning, G. (2008). Treatment of type 2 diabetes and dyslipidemia with the natural plant alkaloid berberine. *Journal of Clinical Endocrinology and Metabolism*, 93(7), 2559-2565.
- Zhang, J., Kang, M.J., Kim, M.J., Kim, M.E., Song, J.H., Lee, Y.M., and Kim, J.I. (2008). Pancreatic lipase inhibitory activity of taraxacum officinale in vitro and in vivo. *Nutrition Research and Practice*, 2(4), 200-203.

- Zhang, S., Nagana Gowda, G. A. Asiago, V., Shanaiah, N., Barbas, C., and Raftery, D. (2008). Correlative and quantitative ^1H NMR-based metabolomics reveals specific metabolic pathway disturbances in diabetic rats. *Analytical Biochemistry*, 383(1), 76-84.
- Zhang, S., Nagana Gowda, G.A., Ye, T., and Raftery, D. (2010). Advances in NMR-based biofluid analysis and metabolite profiling. *Analyst*, 135(7), 1490-1498.
- Zhang, A., Sun, H., Han, Y., Yuan, Y., Wang, P., Song, G., Yuan, X., Zhang, M., Xie, N., and Wang, X. (2012). Exploratory urinary metabolic biomarkers and pathways using UPLC-Q-TOF-HDMS coupled with pattern recognition approach. *Analyst*, 137(18), 4200-4208.
- Zhang, S., Myracle, A., Xiao, K., Yan, P., Ye, T., Ferruzzi, M., Janle, E., and Raftery, D. (2013). Metabolic Profiling of Green Tea Treatments in Zucker Diabetic Rats Using ^1H NMR. *International Journal of Food Sciences and Nutrition*, 3(6), 239-243.
- Zhao, L.C., Zhang, X.D., Liao, S.X., Gao, H.C., Wang, H.Y., and Lin, D.H. (2010). Ametabolic comparison of urinary changes in Zucker and GK rats. *Journal of Biomedicine and Biotechnology*, 2010, Article ID 431894, 6 pages.
- Zhao, L.C., Zhang, X.D., Wang, H.Y., and Lin, D.H. (2011). ^1H NMR-Based Metabonomic Analysis of Metabolic Changes of Serum and Liver in Zucker Obese Rats. *Analytical Letters*, 44(9), 1579–1590.
- Zheng, C.D., Duan, Y.Q., Gao, J.M., and Ruan, Z.G. (2010). Screening for anti-lipase properties of 37 traditional Chinese medicinal herbs. *Journal of Chinese Medicine Association*, 73(6), 319–324.
- Zhou, K., and Yu, L. (2004). Effects of extraction solvent on wheat bran antioxidant activity estimation. *LWT-Food Science and Technology*, 37(7), 717-721.