



## LJMU Research Online

**Buchanan, GM, Beresford, AE, Hebblewhite, M, Escobedo, FJ, De Klerk, HM, Donald, PF, Escribano, P, Koh, LP, Martinez-Lopez, J, Pettorelli, N, Skidmore, AK, Szantoi, Z, Tabor, K, Wegmann, M and Wich, SA**

**Free satellite data key to conservation**

<http://researchonline.ljmu.ac.uk/8963/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Buchanan, GM, Beresford, AE, Hebblewhite, M, Escobedo, FJ, De Klerk, HM, Donald, PF, Escribano, P, Koh, LP, Martinez-Lopez, J, Pettorelli, N, Skidmore, AK, Szantoi, Z, Tabor, K, Wegmann, M and Wich, SA (2018) Free satellite data key to conservation. *Science*. 361 (6398). pp. 139-140. ISSN**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

# Free satellite data crucial for research

Biodiversity is in crisis, with extinction rates orders of magnitude higher than background levels (1). Underfunded conservationists need to target their limited resources effectively. Over the past decade, satellite remote sensing has revolutionized our ability to monitor biodiversity globally, and is now used routinely, especially by nongovernmental organizations, to detect changes, set priorities, and target conservation action. The U.S. Geological Survey (USGS) unlocked high-resolution Landsat data in 2008 (2), making data now available online (3), and the Copernicus program from the European Commission subsequently made their data available too (4). These resources have been instrumental to biodiversity research. Assessments of environmental changes such as deforestation are now readily available. The current spatial and spectral resolution of Landsat data make them appropriate to many conservation applications, and although they are not always ideal, pragmatic researchers with limited resources use them regularly. Conservationists have already called for these data to remain free (5). Consequently, the news that USGS may charge for data (6) is deeply troubling.

A recent USGS survey on Landsat 9 design asks whether users would be prepared to pay \$50 to more than \$3000 for increased spectral and spatial resolution images (7). This pricing would put these images beyond the reach of conservationists. It would halt time-series analyses that have been useful in monitoring the effects of climate change, land-cover change, and ocean surfaces, likely hindering the achievement of the Sustainable Development Goals (8). We urge the USGS to reconsider their position and continue to provide data from the Landsat program freely to all users.

G. M. Buchanan,<sup>1\*</sup> A. E. Beresford,<sup>1</sup> M. Hebblewhite,<sup>2</sup> F. J. Escobedo,<sup>3</sup> H. M. De Klerk,<sup>4</sup> P. F. Donald,<sup>5</sup> P. Escribano,<sup>6</sup> L.P. Koh,<sup>7</sup> J. Martínez-López,<sup>8</sup> N. Pettorelli,<sup>9</sup> A. K. Skidmore,<sup>10</sup> Z. Szantoi,<sup>4</sup> K. Tabor,<sup>7</sup> M. Wegmann<sup>11</sup>, S. Wich<sup>12</sup>

<sup>1</sup>RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, Edinburgh, EH12 9DH, UK. <sup>2</sup>Wildlife Biology Program, Department of Ecosystem and Conservation Sciences, W.A. Franke College of Forestry and Conservation, University of Montana, Missoula, MT 59812, USA. <sup>3</sup>Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Bogotá D.C., 11122, Colombia. <sup>4</sup>Department of Geography and Environmental Studies, Stellenbosch University, Stellenbosch 7602, South Africa. <sup>5</sup>BirdLife International, David Attenborough Building, Pembroke Street, Cambridge, CB2 3QZ, UK. <sup>6</sup>CAESCG, University of Almería, Cañada de San Urbano s/n 04120 Almería, Spain. <sup>7</sup>Betty & Gordon Moore Center for Science, Conservation International, Arlington, VA 22202, USA. <sup>8</sup>BC3-Basque Centre for Climate Change, University of the Basque Country, 48940, Leioa, Spain. <sup>9</sup>Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY, UK. <sup>10</sup>University of Twente, Faculty of Geo-

Information Science and Earth Observation, 7500 AE Enschede, Netherlands. <sup>11</sup>Institute of Geography and Geology, 97074 Würzburg, Germany. <sup>12</sup>School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, L33AF.

\*Corresponding author. Email: graeme.buchanan@rspb.org.uk

## REFERENCES

1. J. DeVos *et al.* *Cons. Biol.* 29, 452 (2015).
2. C. Woodcock *et al.* *Science* 320, 1011 (2008).
3. <https://earthexplorer.usgs.gov/>
4. <http://copernicus.eu/>
5. W. Turner *et al.*, *Biol. Conserv.* 182, 173 (2015).
6. G. Popkin, *Nature* 556, 417 (2018).
7. USGS Landsat 9 user survey.
8. <https://sustainabledevelopment.un.org/?menu=1300>

10.1126/science.aau2650