



# Sterigmatocystin in foodstuffs and feed: aspects to consider

Carla Viegas, Janne Nurme, Elena Piecková & Susana Viegas

**To cite this article:** Carla Viegas, Janne Nurme, Elena Piecková & Susana Viegas (2018): Sterigmatocystin in foodstuffs and feed: aspects to consider, *Mycology*, DOI: [10.1080/21501203.2018.1492980](https://doi.org/10.1080/21501203.2018.1492980)

**To link to this article:** <https://doi.org/10.1080/21501203.2018.1492980>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 09 Jul 2018.



Submit your article to this journal



View Crossmark data



## Sterigmatocystin in foodstuffs and feed: aspects to consider

Carla Viegas<sup>a,b</sup>, Janne Nurme<sup>a</sup>, Elena Piecková<sup>c</sup> and Susana Viegas<sup>a,b</sup>

<sup>a</sup>H&TRC- Health & Technology Research Center, ESTeSL- Escola Superior de Tecnologia da Saúde, Instituto Politécnico de Lisboa, Lisbon, Portugal; <sup>b</sup>Centro de Investigação em Saúde Pública, Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal;

<sup>c</sup>Faculty of Medicine, Slovak Medical University, Bratislava, Slovakia

### ABSTRACT

Sterigmatocystin (STC) is a possible human carcinogen (2B) according to International Agency for Research on Cancer classification and has been associated with immunotoxic and immunomodulatory activity, together with mutagenic effects. It might be found in numerous substrates, from foods and feeds to chronically damp building materials and indoor dust. Although European Food Safety Authority concluded that the exposure to STC to be of low concern for public health, reinforces the need of data concerning exposure of European citizens. Climate change can represent an increased risk of exposure to STC since it is a crucial factor for agro-ecosystem powering fungal colonisation and mycotoxin production. This aspect can represent an increased risk for European countries with temperate climates and it was already reported by the scientific community.

### ARTICLE HISTORY

Received 2 March 2018

Accepted 21 June 2018

### KEYWORDS

Sterigmatocystin; foodstuffs and feed; climate change; increased risk; public health

## 1. Background

Food and feed can be contaminated with a mixture of mycotoxins, toxic secondary metabolites produced by fungal species. Fungal invasion and consequent mycotoxin production occurs both in the field and/or during the storage of crops (de Nijs et al. 2016). The level of contamination is dependent on many factors such as fungal interactions, type of crops and environmental factors (Battilani et al. 2012; Van der Fels-Klerx 2016). In food, other aspects can influence the profile and levels of the mycotoxins present, namely, industrial processes involved and, after being acquired, household preparation process (Nijs et al. 2016). In addition, we should consider that mycotoxins are extremely difficult to eliminate from food even after the cooking process because they are quite stable molecules. All this explains why mycotoxins can be present in the food or environment long after death and disintegration of the toxic fungus (Peraica et al. 1999; Halstensen 2008; Alborch et al. 2011; Viegas et al. 2015).

## 2. Aspergillus section Versicolores

The identification of aspergilli down to the species level was traditionally based on the morphological features (Raper and Fennell 1965). More recently, Houbraken et al.

(2014) and Hubka et al. (2014) proposed the system comprising four subgenera (*Aspergillus*, *Circumdati*, *Fumigati*, *Nidulantes*) with 20 sections based on phylogenetic approach. The official fungal DNA barcode applying ITS locus (Schoch et al. 2012) has seemed to be insufficient for accurate identification of aspergilli and their sexual morphs, so, additional marker analyses were developed: calmodulin gene(CaM), beta-tubuline (BenA) and RNA polymerase II second largest subunit (RPB2) sequencing, along with extrolite spectrum data as well (Samson et al. 2014; Frisvad 2015). Nowadays, 17 species are assigned to the *Aspergillus* section *Versicolores*: *A. amoenus*, *A. austroafricanus*, *A. creber*, *A. cvjetkovicii*, *A. fructus*, *A. griseoaurantiacus*, *A. hongkongensis*, *A. jensenii*, *A. pepii*, *A. protuberans*, *A. puulaauensis*, *A. subversicolor*, *A. sydowii*, *A. tabacinus*, *A. tennesseensis*, *A. venenatus* and *A. versicolor* (Jakšić Despot et al. 2017). According to the works by Chen et al. (2016) and Hubka et al. (2016), the section *Versicolores* was merged to the section *Nidulantes* as the *A. versicolor* clade, to maintain monophyly of *Aspergillus*. The newest combined phylogeny of aspergilli at the section *Versicolores* based on GenBank sequences was published by Jakšić Despot et al. (2017).

Along to the typical aspergillum – a biseriate head on a long stipe, several *A. versicolor* and *A. sydowii* isolates produce diminutive conidial heads resembling

**CONTACT** Carla Viegas  carla.viegas@estesl.ipl.pt

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

penicillate conidiophores. The recommended sexual name *Emericella* should be used for species in the section *Versicolores*. *Aspergillus versicolor* (Vuill.) Tirab. was historically the most commonly reported representative of the entire section. Its herbarium strain CBS 538.65 bears ITS barcode EF652442, *BenA* = EF652266, *CaM* = EF652354, *RPB2* = EF652178, followed by *A. sydowii* (Bainier and Sartory) Thom and Church, herb.: IMI 211,384, ITS barcode: EF652450, *BenA* = EF652302, *CaM* = EF652390, *RPB2* = EF652214 (Samson et al. 2014). So, that is why *A. versicolor* strains were reported growing worldwide, prospering in many habitats, including foodstuffs and feedstuffs as well as indoor environments, and being implicated in various human and animal health hazards, from mycoses to mycotoxicoses due to production of sterigmatocystin (STC). Of note, it has been already reported that their high prevalence in some specific occupational environments is a consequence of this evidence, such as swineries (Sabino et al. 2012; Viegas et al. 2013).

An extensive search was performed to identify scientific papers, available in different scientific databases and published after 2010, reporting *Aspergillus* section *Versicolores* presence in foodstuffs and feed samples. These species have been reported in several foodstuffs such as cocoa (Egbuta et al. 2014), maize (Makun et al. 2010), date fruits (Al-Bulushi et al. 2017), dry-cured meat products (Sonjak et al. 2011), coffee beans (Batista et al. 2003; Viegas et al. 2017), wheat (Tančinová and Labuda 2009; Al-Hazmi 2010), wheat and wheat products (Riba et al. 2008; Piotrowska 2013), rice (Lima et al. 2000; Aydin et al. 2011), cereals (Tabuc et al. 2010; Toffa et al. 2013), peanut (Toffa et al. 2013), ready-to-use vegetable salads (Kocić-Tanackov et al. 2010), vegetables (Accensi et al. 2004), honey (Kačániová et al. 2012), apples with rotting (Tančinová et al. 2013), puerh tea (Haas et al. 2013), frozen chicken (Darwish et al. 2016), dried raisins (Alghalibi and Shater 2004) and spice Kashmiri chilli mild (Hammami et al. 2014). Several different feeds were reported to be contaminated, namely, maize grain (Chemining'wa et al. 2009; Ayalew 2010), fish feed (Barbosa et al. 2013), feed mixture (Magnoli et al. 2002; Accensi et al. 2004; Labuda and Tančinová 2006), silage (El-Shanawany et al. 2005), soybean (Kačániov 2003), cereals (Accensi et al. 2004; Tabuc et al. 2009, 2011), raw materials like corn and barley rootlets

(Rosa et al. 2008), soybean meals (Al-Seen 2012) and sorghum (Silva et al. 2000) (Table 1).

### 3. STC production ability

Mycotoxin STC synthesis is restricted to species in four sections in *Aspergillus* (*Ochraceorosei*, *Versicolores*, *Nidulantes* and *Flavi*) (Rank et al. 2011). Most of the aspergillus species from the section *Versicolores* are able to produce STC, namely, *A. amoenus*, *A. creber*, *A. cvjetkovicii*, *A. fructus*, *A. griseoaurantiacus*, *A. hongkongensis*, *A. jensenii*, *A. pepii*, *A. protuberans*, *A. puulaauensis*, *A. subversicolor*, *A. tennesseensis*, *A. venenatus* and *A. versicolor* (Jurjievic et al. 2013; Visagie et al. 2014; Jakšić Despot et al. 2017). According to Frisvad (2015), the metabolic profile is unique for each fungal species entity with a high degree of chemoconsistency among different isolates of the particular species. But, so far, at the section *Versicolores*, there are chemical markers characterised just for *A. versicolor* and *A. sydowii* (Samson et al. 2010). Liquid Chromatography Mass Spectrometry (LC-MS) LC/MS-based methods proved to be accurate to identify chromatograms of fungal extracts in general. At the moment, some other extrolites of *A. versicolor* were found, like polyketides (Lee et al. 2010), stephacidin A and notoamide B (Greshock et al. 2008) or kipukasins (Jiao et al. 2007), respectively.

STC and dihydrosterigmatocystin are the penultimate precursors of aflatoxins – polyketide-derived furanocoumarins. It has been demonstrated that 25 identified genes clustered within a 70-kb DNA region in the chromosome are involved in their biosynthesis (Townsend 1997). The homologous genes and their corresponding enzymes acting in each bioconversion step in the biochemical pathway common to aflatoxins and STC were described later on as well (Yu et al. 2004). For example, in *A. nidulans*, the last in the row crucial gene seems to be *stcP* encoding O-methyltransferase B required for the conversion of dimethylsterigmatocystin to STC (Kelkar et al. 1996).

### 4. STC toxicity

STC is a possible human carcinogen (2B) according to IARC classification (McConnell and Garner 1994) and showed immunotoxic and immunomodulatory activity (Liu et al. 2014), together with mutagenic effects (Gao et al. 2015). It might be found in numerous

**Table 1.** *Aspergillus* section *Versicolor* presence in food and feed samples.

Setting	Food or feed sampled	Number of samples	Analytical methods	Results	Reference
Rough rice warehouse from Rio Grande do Sul Brazil	Polished rice Rice bran Rice hulls Sorghum	30 30 30 140	Plated onto Potato dextrose agar; plates were incubated for 7 days at 25°C Inoculated onto potato dextrose agar. Incubated at 25°C for 5 days and observed daily Inoculated onto DRBC agar; incubated in darkness at 28 for 7 days Plated directly onto filter paper moistened with sterile distilled water; 15 were plated without prior hypochlorite disinfection and 15 were disinfected with 1% hypochlorite for 2 min	Prevalence 6.6% 1.4% prevalence $1.0 \times 10^3$ – $5.0 \times 10^3$ (CFU g <sup>-1</sup> ) 3 isolates	Lima et al. (2000) Silva et al. (2000) Magnoli et al. (2002) Batista et al. (2003)
Freshly harvested and stored sorghum from the State of São Paulo, Brazil	Poultry feed	120	Inoculated onto Czapek-Dox agar with streptomycin; incubated at 25°C 10 days	Prevalence 9.5% Prevalence 14.5% Prevalence 8.3%	Accensi et al. (2004)
Mixed feed production plant from Cordoba, Argentina	Processed coffee beans ( <i>Coffea arabica</i> L.)	45	Inoculated to 1% glucose and 40% sucrose-Czapek's agar media; incubated at 28°C for 10 days	1 isolation from dried raisins	Alghalib and Shater (2004)
Eleven municipalities supermarkets from Brazil	Soybean	25	Inoculated onto Czapek-Dox agar with streptomycin; incubated at 25°C 10 days	2 isolates	Kacaniova (2003)
Agricultural companies in the years 2002 and 2003 from Slovak Republic	Mixed feeds Cereals Vegetables Dried raisins, figs and dates	147 117 36 20	Inoculated onto malt extract agar; incubated at 28°C for 7 days Inoculated onto abouraud's dextrose agar with chloramphenicol and cycloheximide; -incubated at 28°C for 1–2 weeks	Occurrence in 13 samples with 33 isolates	El-Shanawany et al. (2005)
Local factories	Silage	40	Inoculated onto Dichloran chloramphenicol peptone agar (DCPA); condial suspensions inoculated onto Czapek Yeast Extract Agar (CYA), Czapek Yeast Extract Agar CY20S and Malt extract agar (MEA) and incubated in the dark at 25°C for 7–14 days	Occurrence in 13 samples with 33 isolates	Labuda and Tančinová (2009)
Dried fruits from shops and markets in Yemen Republic	Feed mixture	108	Inoculated onto Dichloran Rose Bengal Chloramphenicol Agar (DRBC) Agar; incubated for 5–7 days at 28°C in the dark; colonies sub-cultured on malt extract agar; allowed to grow at 28°C for 7–10 days	Frequency ranged from 10% to 28%	Riba et al. (2008)
Farms, mills and retail markets at Assuit and Sohag governorates from Egypt	Wheat and wheat products	30 31 12 12	Inoculated onto DRBC; incubated at 28°C for 7 days	Corn prevalence 11.8% and barley rootlets prevalence 8.1%	Rosa et al. (2008)
Poultry farms from Slovakia	Tizi Ouzou region Flour mill Semolina mill and samples from Algeria region Cattle dairy feed from Rio de Janeiro State, Brazil	106	Inoculated onto Czapek dextrose agar; incubated at 25°C for 7–10 days	Positive results in 6 zones, percentage ranging from 0.6% to 6.6%	Cheminings'wa et al. (2009)
Setif region	Nine agro-ecological zones from Eastern Kenya	5 farmers per ecological zone			(Continued)



Table 1. (Continued).

Setting	Food or feed sampled	Number of samples	Analytical methods	Results	Reference
Cereal samples from Southeastern Romania	Corn Wheat Barley Wheat bran	54 35 21 56	Plated onto malt agar and malt salt agar; colonies were counted after 3, 5, and 7 days of culture at 25°C and 31°C Inoculated onto Dichloran chloramphenicol peptone agar (DCPA); 5–7 days of incubation at 25°C Inoculated onto potato dextrose agar (PDA); incubated at 25°C in dark for 7 days	Prevalence of 46.2% Prevalence of 22.8% Prevalence of 9.4% Occurrence in 5 samples with 8 isolates	Tabuc et al. (2009)
Mills from Slovakia					Tančinová and Labuda (2009)
Wheat from different production areas and sold Jeddah market, Saudi Arabia	Wheat grain	30		2 × 10 <sup>2</sup> colony/g 3 × 10 <sup>3</sup> colony/g	Al-Hazmi (2010)
2004/2005 harvest in Ethiopia	Maize grain	17	Inoculated onto Dichloran Glycerol (DG)18 agar; Incubated at room temperature for 1–2 weeks Inoculated to DRBC agar; Incubated at 25°C for 5–7 days	Surface prevalence 23.5%	Ayalew (2010)
Fresh ready-to-use salads made of different types of vegetables in Serbia	Salads	17	Inoculated onto PDA; incubated 5–7 days at 28°C	Prevalence of 1.78%	Kocić-Tanackov et al. (2010)
Maize field, market and storage facilities from three States in Nigeria	Five different food commodities	343	Applied onto malt and agar with 6% NaCl; incubated for 5–7 days at 25°C	10.7% prevalence	Makun et al. (2010)
Cereal samples from different locations in Banat region, Romania	Cereal grain (corn, wheat, barley, oats)	56	Inoculated onto DRBC agar; incubated at 25°C for 5–7 days; Colonies sub-cultured on malt extract agar incubated at 25°C for 5–7 days		Tabuc et al. (2010)
Two rice-growing areas Uzunkopru and Ipsala in the Thrace region from Turkey	Rice	100	Inoculated to MEA and to MEA with 5% NaCl; isolates inoculated onto MEA and grown for 7 days at 25°C in the dark	2 positive samples	Aydin et al. (2011)
Slovenian meat-processing plant	Dry-cured meat products	75 items of different dry-cured meat products		2 isolates	Sonjak et al. (2011)
Cereals produced in the South East part of Romania between 2008 and 2010	Feed cereals (maize, wheat, barley, oats, rye, soya, sunflower, colza, rice, triticale)	86	Plated onto malt agar medium and malt salt agar medium; incubated 7 days at 25°C and 31°C	2008 ( <i>n</i> = 42) 38.0% 2009 ( <i>n</i> = 32) 31.2% 2010 ( <i>n</i> = 12) 58.3%	Tabuc et al. (2011)
Soybean meal collected from fodder markets at Jeddah, Saudi Arabia	Soybean meals	20	Inoculated onto Czapek-Dox agar with 1% streptomycin; incubated at 25°C 10 days	1.4 × 10 <sup>2</sup> count/g	Al-Seeni (2012)
Bee-keepers (apairy honey) from Slovakia and other countries	Honey	53	The cultures grew under specific conditions on Czapek-Dox agar and Malt agar	3 positive samples with frequency 6.82%	Kačániová et al. (2012)
Two tilapia farms in the Rio de Janeiro State, Brazil	Fish feed	60	Inoculated onto DRBC and DG18; Incubated at 25°C for 5–7 days	16% relative density	Barbosa et al. (2013)
Different stores and markets in south-western Nigeria	Processed cocoa	22	Inoculated onto PDA, OAES Aohio agricultural experimental station agar, MEA and CYA; incubated 4–7 days at 25°C	2.0 × 10 <sup>4</sup> (CFU/g) 4.6%	Egbuta et al. (2013)

(Continued)

**Table 1.** (Continued).

Setting	Food or feed sampled	Number of samples	Analytical methods	Results	Reference
Tea shops from the province of Yunnan in south-western China	Pu-erh tea	36	Plated onto MEA, Sabouraud Glucose Agar and DG18; Incubated at 25°C 7–14 days	2.8% prevalence	Haas et al. (2013)
Cereal breakfast products from Lodz (Poland) markets	Cereal snacks with cinnamon, cornflakes, cornflakes with nuts and honey, multi-cereal products, cereal products with chocolate and muesli with dried fruit, nuts, cereal and coconut flakes	15	Inoculated onto DRBC agar and DG18; incubated at 25°C for 7 days	Cereal snacks with cinnamon – 37%; cornflakes with nuts and honey – 24%; multi-cereal products – 15%; muesli – 20%; cereal products with chocolate – 50% and 10%	Piotrowska (2013)
Apples with the characters of rotting bought in the commercial network from Slovakia	Rotting apples	30	Producers of rotting were inserted directly onto MEA Cultivated 5–7 days in the dark at 25 ± 1 °C. <i>Aspergillus</i> sp. isolated to C/A, MEA, CY20S, CREA creatine sucrose agar and YES; cultivation proceeded for 5–7 days in the dark at 25 ± 1 °C	1 isolate	Tančinová et al. (2013)
Cereals and peanut from local markets from Republic of Niger	Cereals (rice, maize, sorghum, millet) and peanut	81	Plated onto PDA, CYA, MEA; incubated MEA 30°C, PDA 27°C, CYA 25°C for 7 days	4 strains	Toffa et al. (2013)
Powdered spices from Qatar local markets	Spices: chili, Kashmiri chili hot, Kashmiri chili mild, basil, oregano, ginger, curry, cumin, turmeric, tandoori masala, garam masala, black pepper, garlic and coriander	14	Inoculated onto DRBC agar; Incubated at 28°C 5 days	In Kashmiri chili mild 3 isolates	Hammami et al. (2014)
Frozen chicken from different localities in Zagazig city, Egypt	Breast Thigh Gizzards Livers	20 20 20 20	Culturing on malt extract agar media and Czapeck-Dox agar with 5% Nacl; incubation in dark at 25°C for 5–7 days	10% 10% 14–15% 4–5%	Darwish et al. (2016)
Several date palm from Oman region	Date palm (Tamar stored dates)	3 date fruits from each cultivar.	Illumina MiSeq Sequencing Analysis	Frequency of occurrence in Khenizi mesocarp 0.32 and skin 0.065	Al-Bulushi et al. (2017)
Green coffee beans from <i>Coffea arabica</i> (Arabica) and <i>Coffea canephora</i> var. <i>robusta</i> (Robusta coffee) to be roasted in Portugal	Samples from different countries of origin	28	The washed supernatant (100 µl) was plated in MEA and DG18; incubated at 27°C for 5 to 7 days.	6.1 prevalence	Viegas et al. (2017)

substrates, from foods and feeds to chronically damp building materials and indoor dust.

In 2013, the European Food Safety Authority (EFSA) was asked by the European Commission to deliver a scientific opinion on STC in food and feed. The Panel on Contaminants in the Food Chain (CONTAM) from EFSA was responsible for this opinion. However, due to the absence of exposure data for the European population, the margin of exposure approach for substances that are genotoxic and carcinogenic could not be applied for STC, and therefore, the risk of STC for human health was not characterised.

Despite this, it was possible to collect all the available information related to STC toxicokinetics, toxicity, mode of action and dose-response assessment by comparing with aflatoxin B1. The following information was available in the EFSA Journal and, more precisely, in the scientific opinion on the risk for public and animal health related to the presence of STC in food and feed (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

This report concludes the exposure to STC to be of low concern for public health based on the relative carcinogenic potency of STC and AFB1 and exposure data. However, the need of data concerning exposure of European citizens was also mentioned (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

#### **4.1. STC toxicokinetics**

There is limited information available related to STC toxicokinetics. However, the accessible data suggests that absorption of STC is limited following oral exposure.

In the same way, data on the biotransformation of STC is also insufficient. Few studies published to date indicate that phase I metabolism of STC comprises cytochrome P450 (CYP450)-mediated formation of a reactive epoxide as well as monohydroxylation and dihydroxylation reactions. In a more detailed manner, STC is metabolised in the liver and lung by various CYP450 enzymes into different hydroxymetabolites and its reactive exo-epoxide that readily forms DNA adducts (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013; Walkow et al. 1985).

As phase II metabolites, a glucuronide of STC and of monohydroxy-STC has been observed and

reported, together with a sulphate conjugate of monohydroxy-STC and a glutathione adduct of a monooxygenated STC. Excretion of both conjugated parent STC and its hydroxylated metabolites occurs via bile and urine. Nevertheless, the structure of most of these metabolites is not completely known and more research is necessary to allow the availability of more detailed information (Walkow et al. 1985; EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

#### **4.2. Toxicity of STC**

Due to the structural similarities, aflatoxins and STC share relevant toxic effects, including genotoxicity and carcinogenicity (Miller and Trenholm 1994; EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013). However, in contrast to aflatoxins, only limited information on occurrence and toxicity of STC is available.

Liver and kidneys are the target organs of acute toxicity. However, the acute oral toxicity is relatively low (range between 120 and 166 mg/kg body weight). STC is hepatotoxic in rat, mouse, monkey and guinea pig. The incidence of hepatocellular necrosis and haemorrhages increases with dose and duration of exposure. In the kidney, hyaline degeneration, tubular necrosis and haemorrhages were described in rats and/or monkeys exposed to STC (Purchase and Van Der Watt 1969; EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

Results from *in vivo* and *in vitro* studies suggest that STC may have also immunomodulatory activity, but strong conclusions cannot be drawn (Huang et al. (2002), Xing et al. (2005), and Zhang et al. (2012) cited in EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013)).

STC is mutagenic in both bacterial and mammalian cell assays after metabolic activation. Subsequently, STC induces chromosomal damage both *in vitro* and *in vivo* in experimental animals (Curry et al. 1984; Ueda et al. 1984; Mori et al. 1986; Crofton-Sleigh et al. 1993; Abdel-Wahhab et al. 2005).

Various studies aimed to compare the genotoxicity of STC and AFB1. However, the uncertainty regarding their actual concentration in the test system, the efficiency of the activation/detoxification metabolic routes and the repair rate of induced lesions does not allow a direct comparison of the

relative mutagenic potency of these mycotoxins (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

In previous studies, tumourigenicity of STC was observed after oral, intraperitoneal, subcutaneous and/or dermal administration in the animal species tested (rat, mouse, Mongolian gerbils, monkey and fish). After oral exposure, premalignant and malignant lesions such as hepatocellular carcinomas (HCC), haemangiosarcomas in the liver, angiosarcomas in the brown fat, lung adenomas and incidental findings in other organs were reported (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

Base on the available information, the CONTAM Panel of EFSA concluded that STC is genotoxic and carcinogenic (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

Additionally, a study developed by Miller et al. (2010), which considered exposure of STC via the indoor environment, concluded that following inhalation (intratracheal installation) of STC, a non-specific but severe inflammatory response of the lung tissue was observed. Similarly, severe cytotoxic and inflammatory damage of lung tissue as well as breaking down of self-cleaning mechanism of airways in rats *in vivo* were observed after intratracheal instillation of STC containing complex extrolites of an *A. versicolor* strain of indoor origin in the studies by Piecková et al. (2011, 2015).

#### **4.3. STC mode of action**

The mode of action of STC can be described as follows. Phase I metabolism results in metabolic activation that promotes the formation of N7-guanyl DNA adducts. These adducts are likely to be responsible for the STE mutagenic effects (Essigmann et al. 1979, 1980). A dose-dependent formation of DNA adducts of STC was found in the concentration range between 1 and 3 mg STC per liver (Reddy et al. 1985; cited in EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

STC induces cytotoxicity, inhibition of cell cycle and mitosis, as well as an increased formation of reactive oxygen species (ROS) and lipid peroxidation *in vivo* (Kawai et al. 1984; Ueno et al. 1995; Xie et al. 2000; Sivakumar et al. 2001; Bünger et al. 2004).

The conclusion made by EFSA Panel on Contaminants in the Food Chain (CONTAM) (2013) was that the genotoxicity of STC is based on the

formation of DNA adducts that, if unrepaired, increase the likelihood of mutation fixation. Moreover, when comparing with AFB1, most *in vitro* studies with purified DNA indicate that the level of induced N7-guanyl adducts is higher after AFB1 than STC exposure, supporting the view that AFB1 is a more potent liver carcinogen than STC. Various *in vitro* and *in vivo* investigations have demonstrated that STC exerts cytotoxicity, inhibition of cell cycle and mitosis, as well as an increased ROS formation and lipid peroxidation *in vivo*. However, most of the *in vitro* assays have been conducted with rather high STC concentrations, not representing the real human exposure scenario that should be a chronic exposure. Therefore, the observed effects of those studies have to be interpreted with caution not allowing to make conclusions regarding the potential adverse effects of (low dose) dietary exposure to STC (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

More recently, a study developed by Wang et al. (2015) tried to confirm that STC exposure is a risk factor for oesophageal cancer and that STC may induce DNA damage and G2 phase arrest in immortalised human oesophageal epithelial cells (Het-1A). Indeed, the study developed allowed to conclude that STC can induce different cell cycle arrest in primary human oesophageal epithelial cells and immortalised human oesophageal epithelial cells *in vitro* (Wang et al. 2015).

In 2017, Jiang and co-authors aimed to investigate whether checkpoint adaptation occurs in GES-1 Cellosaurus cell line (GES-1) cells treated with STC. The results suggested that STC induces an initial G2 arrest that is subsequently followed by G2 phase checkpoint adaptation, which may potentially promote genomic instability and result in tumorigenesis (Jiang et al. 2017).

Additionally, a study developed by Huang et al. (2014) in human pulmonary cells *in vitro* observed that STC induced DNA damage and affected key proteins involved in cell cycle regulation to trigger genomic instability, which may be a potential mechanism underlying the developmental basis of lung carcinogenesis.

#### **4.4. Dose-response modelling**

Despite the evidence on genotoxicity and carcinogenicity, only a limited tumourigenicity database was

available for dose-response assessment since most of the studies published have several limitations that do not allow to be used for dose-response modelling (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

Being aware of this limitation, the CONTAM Panel of EFSA compared the carcinogenic potency of STC and AFB1 in the BMD10 values. After the comparison of the BMD10 of STC for the occurrence of haemangiosarcomas and that of AFB1 for the occurrence of HCC, the CONTAM Panel concluded the carcinogenic potency of STC is approximately three orders of magnitude lower than that of AFB1 (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2013).

## 5. Food and feed contamination already reported

The natural occurrence of STC in foodstuffs and feed has been reported in a limited number of surveys (Table 2). Reports from year 2000 to 2017 have presented STC contamination in several foodstuffs such as rice (Sawane and Sawane 2014; Mo et al. 2015; Rofiat et al. 2015; Bertuzzi et al. 2017), bread (Veršilovskis and Bartkevičs 2012), wheat bran (Tančinová and Labuda 2009), grain (Veršilovskis et al. 2008a; Mo et al. 2015; ), maize (Wirth et al. 2012; Mo et al. 2015; ), groundnuts (Wirth et al. 2012), peanut seed (Youssef et al. 2008), coffee bean (Bokhari and Aly 2009; Culliao and Barcelo 2015), beer (Veršilovskis et al. 2008b), cheese (Veršilovskis et al. 2009), cereal grains, cereal products (Mo et al. 2015) and almond seed (Yassin et al. 2013). Additionally, some reports in animal feed found contamination in grass (Nichea et al. 2015), feed mixture (Labuda and Tančinová 2006; Wirth et al. 2012), wheat, corn, barley, soybean, sunflower cake (Kovalenko et al. 2011), silage (El-Shanawany et al. 2005), straw and hay (Mol et al. 2014).

Liquid chromatography-tandem mass spectrometry was used for STC detection in nine studies with limits of detection (LOD) ranging from 0.03 to 2.0 µg/kg and five out of nine had also limits of quantification (LOQ) ranging from 0.1 to 0.5 µg/kg (Wirth et al. 2012; Veršilovskis et al. 2008a; b, 2009; Veršilovskis and Bartkevičs 2012; Mol et al. 2014; Rofiat et al. 2015; Nichea et al. 2015; Bertuzzi et al. 2017). Also thin-layer chromatography (El-Shanawany et al. 2005; Labuda

and Tančinová 2006; Youssef et al. 2008; Bokhari and Aly 2009; Tančinová and Labuda 2009; Sawane and Sawane 2014; Culliao and Barcelo 2015), high-performance liquid chromatography (Veršilovskis et al. 2008a; b; Yassin et al. 2013; Culliao and Barcelo 2015) and enzyme-linked immunoassay (Kovalenko et al. 2011) analytical methods were used to determine STC in samples. However, in some of these studies, LOD or LOQ values were not available.

## 6. Climate change influence on STC production

Climate change has been occurring since the earth existed, and global temperatures normally show that 7 of the top 10 warmest years on record have occurred since the 1990s. The decade of 2000–2009 was the warmest period worldwide (EPA 2010). By 2100, the atmospheric concentration of CO<sub>2</sub> is predicted to rise up to the range of 540 and 970 ppm above the current concentration. Together with other greenhouse gases such as CH<sub>4</sub>, this will lead to a predicted global temperature increase of 1.1–6.4°C, depending on different models used and global region (In et al. 2007; Battilani et al. 2012). There is increased risk of European countries with temperate climates to have the higher exposure to fungi and mycotoxins due to climate change, which has already been identified by some authors (Paterson and Lima 2011; Battilani et al. 2012). The climate of these countries will probably become warmer reaching temperatures of 33°C, which is, for instance, a temperature very close to the optimal temperature for *Aspergillus* section *Versicolores* growth (30°C) (Atalla et al. 2003) and STC production (optimal temperature between 23°C and 29°C) (Atalla et al. 2003). However, effects of climate change on fungal species distribution and activity are difficult to predict because they are influenced in many different ways such as fungal characteristics, host features and availability, and competitive interactions between microbiota. In addition, environmental variables such as temperature, water availability and atmospheric CO<sub>2</sub> and the interaction of these variables make it difficult to predict their influence on fungal distribution (Boddy 1984) and, consequently, mycotoxin presence in food and feed even higher (In et al. 2007). Mycotoxins are profoundly dependent on climate, plant and storage-associated problems, and also influenced by non-infectious factors (e.g.

**Table 2.** Sterigmatocystin prevalence in foodstuffs and feed samples.

Setting	Food and feed sampled	Number of samples	Extraction method	Analytical method	Results (quantitative and qualitative)	Reference
Assuit and Sohag farms, mills and retail markets	Silage	40	Chloroform extract	TLC	2 samples	El-Shanaway et al. (2005)
Poultry farms	Feed mixture	108 samples, 5132 isolates	chloroform:methanol (2:1, v/v)	TLC	10 isolates	Labuda and Tančinová (2006)
Different parts of Latvia	Grain samples	1. In 2006 95 samples. 2. In 2007 total 120 samples	84% acetonitrile in water	-LC-MS/MS analysis -An electrospray ionization (ESI)-	1. 13.7% (13 samples) from 2006 (0.7 to 2. 35% (42 samples) from 2007 (1 to 47 µg/kg)	Veršilovskis et al. (2008a)
Latvia local supermarkets	Beer (9 dark, 17 light)	26	Acetonitrile:water (10:90, v/v)	HPLC-UV analysis	In 2 samples: 4.0 µg/l (light beer), 7.8 µg/l (dark beer)	Veršilovskis et al. (2008b)
Peanut seed	Fruit=20 Roasted=20 Roasted with salt=20	60	Cyclohexane for 10 h using Soxlet type extractor	TLC	Roasted seed: 3 samples 16.8 µg/kg, 12.2 µg/kg. 14.8 µg/kg Roasted with salt seed: in 1 sample 12.2 µg/kg	Youssef et al. (2008)
Saudi city of Jeddah markets	Coffee bean seed	30	Chloroform and purified by phenyl-bond solid phase	TLC	In 3 samples: 13 ng/g, 11 ng/g, 5 ng/g	Bokhari and Aly (2009)
Mills	Wheat bran	8 isolates from 5 samples	Chloroform:methanol 2:1 v/v	TLC	All 7 isolates	Tančinová and Labuda (2009)
Latvia and Belgium local supermarkets	Cheese	Latvia = 8 Belgium = 13	Acetonitrile-water (90 : 10, v/v) and n-hexane	LC-MS/MS system	Latvia – in 4 samples 2 × 0.04, 0.07, 0.03 (µg/kg).	Veršilovskis et al. (2009)
Pig farms 2006–2009	Several samples from different materials	Wheat – 93 Corn – 111 Barley – 146 Soybean – 15 Sunflower cake – 53 Mixed feed for piglets of group 0–2 months/2–4 months, sows and fattening – 120/94	—	ELISA method	Belgium – 3 samples 1.23, 0.03, 0.52 (µg/kg). 2008: 8% of corn samples; 2006: 7–17% of wheat, soybean, pea, and mixed feeds for piglets of age groups 0–2 and 2–4 months; in 2007, 9–22% of samples of wheat, corn, barley, pea, bran, sunflower cake, and mixed feeds for piglets of age group 0–2 months and sows; in 2009, 8–50% of samples of all feed types, except mixed feeds for piglets of age group 2–4 months and sows	Kovalenko et al. (2011)
Foods and feedstuffs from Burkina Faso	Maize = 26, feed = 4 and others = 30	Total 122	Acetonitrile/water/acetic acid (79:20:1, v/v/v)	LC-MS/MS analysis	Maize: 2 samples, median 2.3 µg/kg Feed: 3, median 6.5 µg/kg Others 2 median 6.7 µg/kg	Wirth et al. (2012)
Foods and feedstuffs from Mozambique	Maize = 13, groundnuts = 23, feed = 10 and others = 7				Maize: 1, median 2.7 µg/kg. Groundnuts: 1, median 9.7 µg/kg. Feed: 1, median 11 µg/kg. Others: 2, median 26.1 µg/kg	Veršilovskis et al. (2012)
Bread samples from Riga local supermarkets	Plain rye bread Mixed rye-wheat bread Plain wheat bread	6 9 14	16% water in acetonitrile	LC-MS/MS and ESI	In 1 sample = 2.4 µg/kg In 1 sample = 7.1 µg/kg In 3 samples = 4.4 µg/kg; 3.2 µg/kg; 5.6 µg/kg	
Different locations in Riyadh city, Kingdom of Saudi Arabia	Almond seed	20	Acetone, chloroform	HPLC	300–440 ppb	Yassin et al. (2013)

(Continued)

**Table 2.** (Continued).

Setting	Food and feed sampled	Number of samples	Extraction method	Analytical method	Results (quantitative and qualitative)	Reference
India rice	Rice	36	Chloroform/methanol (2:1, v/v), mycelia	TLC	5 isolates out of 12	Sawane A and Sawane M (2014)
Animal Feed	Straw	149	Acetonitrile/water (84/16) containing 1% formic acid	LC-MS/MS analysis	2 samples 0.001–0.038 mg/kg	Mol et al. (2014)
2 beef cattle farms	Hay Grass	43	106 samples in 2011 and 69 samples in 2014	Acetonitrile/water/acetic acid (79:20:1, v/v/v).	22 samples 0.0012–0.17 mg/kg 90% in 2011 and 60% in 2014	Nichea et al. (2015)
143 rice processors	Milled rice	38	Acetonitrile/water/acetic acid (79:20:1,v/v/v)	LC-MS/MS with ESI source and HPLC System	50% of rice samples, mean value 19 µg/kg, maximum value 125 µg/kg	Rofiat et al. (2015)
Domestic and imported cereals and cereal products from different countries in different regions within the European Union	Cereal grains: Wheat Rye Maize Rice Barley Oats	221 35 33 28 59 51	Acetonitrile-water 80+20 v/v	LC-MS/MS analysis	Wheat: 12 samples (9 = LOD-0.5 µg/kg, 3 = 0.5–1.5 µg/kg); rye: 2 samples (1 = LOD-0.5 µg/kg, 1 = 0.5–1.5 µg/kg); maize: 2 samples (0.5–1.5 µg/kg); Rice: 27 samples (8 = LOD-0.5 µg/kg, 8 = 0.5–1.5 µg/kg, 10 = 1.5–5 µg/kg, 1 = 5 µg/kg); barley: 1 sample (1.5–5 µg/kg); oats: 11 samples (4 = LOD-0.5 µg/kg, 5 = 0.5–1.5 µg/kg, 1 = 1.5–5 µg/kg, 1 = 5 µg/kg)	Mo et al. (2015)
	Cereal products: Grain milling products, Rice, Pasta, Bread and rolls, Breakfast cereals, Fine bakery ware, Cereal-based infant food	125 89 115 143 97 90 54	Cereal products: Grain milling products, Rice, Pasta, Bread and rolls, Breakfast cereals, Fine bakery ware, Cereal-based infant food		Grain milling products: 6 samples (5 = LOD-0.5 µg/kg, 1 = 0.5–1.5 µg/kg); rice: 19 samples (12 = LOD-0.5 µg/kg, 5 = 0.5–1.5 µg/kg, 2 = 1.5–5 µg/kg); pasta: 6 samples (OD-0.5 µg/kg; bread/rolls: 10 samples (6 = LOD-0.5 µg/kg, 2 = 0.5–1.5 µg/kg, 2 = 1.5–5 µg/kg); breakfast cereals: 18 samples (11 = LOD-0.5 µg/kg, 6 = 0.5–1.5 µg/kg, 1 = 1.5–5 µg/kg); fine bakery ware: 6 samples (5 = LOD-0.5 µg/kg, 1 = 0.5–1.5 µg/kg); cereal-based infant food: 4 samples (2 = LOD-0.5 µg/kg, 2 = 0.5–1.5 µg/kg);	
Philippines coffee beans from production Farms and storage facilities	Coffee beans Paddy rice	42 49	Chloroform	TLC and HPLC	22 samples 0.0012–0.17 mg/kg Mean concentrations 55.8, 16.1 and 193.7 µg/kg	Culliao and Barcelo (2015)
Retail and wholesale resources	Processed rice	83	Acetonitrile-water 80/20v/v	LC-MS/MS system.	All samples in the range of 0.29–15.85 µg/kg	Bertuzzi et al. (2017)
					In 21% of samples ( <i>n</i> = 19) 0.12 and 1.32 µg/kg (brown rice)	

bioavailability of (micro)nutrients, insect damage, and other pests attack) that are also driven by climatic conditions. Therefore, climate represents the crucial factor for agro-ecosystem powering fungal colonisation and mycotoxin production (Magan et al. 2003).

## 7. Future perspectives regarding food and feed contamination

The report by the CONTAM Panel from EFSA mentioned the need for more occurrence data on STC in food and feed across European countries to allow an accurate assessment of dietary exposure. Furthermore, the prediction of climate change and how it can influence fungal contamination and mycotoxin production should be considered. Therefore, besides not knowing in detail what is the actual exposure to this mycotoxin in Europe, the new scenario of climate change brings new challenges due to a probable new exposure trends, particularly in countries with temperate climate.

## Acknowledgments

The authors are grateful for the support given by the Environment and Health Research Group from Escola Superior de Tecnologia da Saúde de Lisboa and Associação Para o Desenvolvimento de Conhecimento e Inovação (POLITEC & ID).

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

The authors are grateful for the support given by the Environment and Health Research Group from Escola Superior de Tecnologia da Saúde de Lisboa and Associação Para o Desenvolvimento de Conhecimento e Inovação (POLITEC & ID)

## References

- Abdel-Wahhab MA, Hasan AM, Aly SE, Mahrous KF. 2005. Adsorption of sterigmatocystin by montmorillonite and inhibition of its genotoxicity in the Nile tilapia fish (*Orechromis niloticus*). *Mutat Res.* 582:20–27.
- Accensi F, Abarca ML, Cabañas FJ. 2004. Occurrence of *Aspergillus* species in mixed feeds and component raw materials and their ability to produce ochratoxin A. *Food Microbiol.* 21(5):623–627.
- Alborch L, Bragulat MR, Abarca ML, Cabañas FJ. 2011. Effect of water activity, temperature and incubation time on growth and ochratoxin A production by *Aspergillus niger* and *Aspergillus carbonarius* on maize kernels. *Int J Food Microbiol.* 147:53–57.
- Al-Bulushi IM, Bani-Uraba MS, Guizani NS, Al-Khusaibi MK, Al-Sadi AM. 2017. Illumina MiSeq sequencing analysis of fungal diversity in stored dates. *BMC Microbiol.* 17:72.
- Alghalibi SMS, Shater ARM. 2004. Mycoflora and mycotoxin contamination of some dried fruits in Yemen Republic. *Ass Univ Bull Environ Res.* 7:2.
- Al-Hazmi NA. 2010. Determination of zearalenone (ZEA) in wheat samples collected from jeddah market, Saudi Arabia. *Afr J Microbiol Res.* 4(23):2513–2519.
- Al-Seeni MN. 2012. Natural occurrence of heavy metal, fungi and mycotoxins in soybean meal samples used in animal feeding in Saudi Arabia. *Afr J Biotechnol.* 11(38):9288–9294.
- Atalla MM, Hassanein NM, El-Beih AA, Youssef YA. 2003. Mycotoxin production in wheat grains by different *Aspergilli* in relation to different relative humidities and storage periods. *Nahrung.* 47:6–10.
- Ayalew A. 2010. Mycotoxins and surface and internal fungi of maize from Ethiopia. African. Vol. 10. *Journal of Food Agriculture Nutrition and Development*; p. 9.
- Aydin A, Aksu H, Gunzen U. 2011. Mycotoxin levels and incidence of mould in Turkish rice. *Environ Monit Assess.* 178:271–280.
- Barbosa TS, Pereyra CM, Soleiro CA, Dias E, Oliveira A, Keller KM, Silva P, Cavagliere LR, Rosa C. 2013. Mycobiota and mycotoxins present in finished fish feeds from farms in the Rio de Janeiro State, Brazil. *Int Aquatic Res.* 5:3.
- Batista LR, Chalfoun SM, Prado G, Freitas Schwan R, E. Wheals A. 2003. Toxigenic fungi associated with processed (green) coffee beans (*Coffea arabica L.*). *Int J Food Microbiol.* 85 (3):293–300.
- Battilani P, Rossi V, Giorni P, Pietri A, Gualla A, Van der Fels-Klerx HJ, Booij CJH, Moretti A, Logrieco A, Toscano P, et al. 2012. Modelling, predicting and mapping the emergence of aflatoxins in cereals in the EU due to climate change. Scientific report submitted to EFSA [Internet]. [place unknown]. Available from: <http://www.efsa.europa.eu/en/supporting/pub/en-223>
- Bertuzzi T, Romani M, Rastelli S, Mulazzi A, Pietri A. 2017. Sterigmatocystin occurrence in paddy and processed rice produced in Italy in the years 2014–2015 and distribution in milled rice fractions. *Toxins.* 9(3):86.
- Boddy L. 1984. The micro-environment of basidiomycete mycelia in temperate deciduous woodlands. In: Jennings DH, Rayner ADM, Eds. *Ecology and physiology of the fungal mycelium*. Cambridge: Cambridge University Press; p. 261–289.
- Bokhari FM, Aly M. 2009. Evolution of traditional means of roasting and mycotoxins contaminated coffee beans in Saudi Arabia. *Adv Biol Res (Rennes).* 3(3–4):71–78.
- Bünger J, Westphal G, Monnich A, Hinnendahl B, Hallier E, Müller M. 2004. Cytotoxicity of occupationally and environmentally relevant mycotoxins. *Toxicology.* 202:199–211.
- Cheminig'wa GN, Gathumbi JK, Njenga LN, Muthomi JW. 2009. The occurrence of aflatoxins in maize and distribution

- of mycotoxin-producing fungi in Eastern Kenya. *Plant Pathol*. J. 8(3):113–119.
- Chen AJ, Frisvad JC, Sun BD, Varga J, Kocsube S, Dijksterhuis J, Kim DH, Hong SB, Houbraken J, Samson RA. **2016**. *Aspergillus* section *Nidulantes* (formerly *Emericella*): polyphasic taxonomy, chemistry and biology. *Stud Mycol*. 84:1–118.
- Crofton-Sleigh C, Doherty A, Ellard S, Parry EM, Venitt S. **1993**. Micronucleus assays using cytochalasin-blocked MCL-5 cells, a proprietary human cell line expressing five human cytochromes P-450 and microsomal epoxide hydrolase. *Mutagenesis*. 8:363–372.
- Culliao AGL, Barcelo JM. **2015**. Fungal and mycotoxin contamination of coffee beans in Benguet province, Philippines. *Food Additives & Contaminants: Part A*. 32(2):250–260.
- Curry PT, Reed RN, Martino RM, Kitchin RM. **1984**. Induction of sister-chromatid exchanges in vivo in mice by the mycotoxins sterigmatocystin and griseofulvin. *Mutat Res*. 137:111–115.
- Darwish WS, Bayomi RM, El-Moaty AM, Gad TM. **2016**. Mould contamination and aflatoxin residues in frozen chicken meat-cuts and giblets. *Jpn J Vet Res*. 64(2):167–171.
- De Nijs M, Mengelers MJB, Boon PE, Heyndrickx E, Hoogenboom LAP, Lopez P, Mol HGJ. **2016**. Strategies for estimating human exposure to mycotoxins via food. *World Mycotoxin Journal Special issue: Mycotoxins in a changing world*. 9(5):831–845.
- EFSA Panel on Contaminants in the Food Chain (CONTAM). **2013**. Scientific opinion on the risk for public and animal health related to the presence of sterigmatocystin in food and feed. *EFSA J*. 11(6):3254.
- Egbuta M, Mulunda F, Njobeh PB, Phoku JZ, Chilaka CA, Dutton MF, Dutton F. **2014**. Isolation of filamentous fungi species contaminating some nigerian food commodities. *J Food Res*. 4(1):38–50.
- El-Shanawany AA, Eman Mostafa M, Barakat A. **2005**. Fungal populations and mycotoxins in silage in Assiut and Sohag governorates in Egypt, with a special reference to characteristic *Aspergilli* toxins. *Mycopathologia*. 159:281–289.
- Essigmann JM, Barker LJ, Fowler KW, Francisco MA, Reinhold VN, Wogan GN. **1979**. Sterigmatocystin-DNA interactions: identification of a major adduct formed after metabolic activation in vitro. *Proc Natl Acad Sci U S A*. 76:179–183.
- Essigmann JM, Donahue PR, Story DL, Wogan GN, Brunengraber H. **1980**. Use of the isolated perfused rat liver to study carcinogen-DNA adduct formation from aflatoxin B1 and sterigmatocystin. *Cancer Res*. 40:4085–4091.
- European Food Safety Authority. **2010**. Management of left-censored data in dietary exposure assessment of chemical substances. *Eur Food Saf Auth J*. 8:1–96.
- Frisvad JC. **2015**. Taxonomy, chemodiversity, and chemoconsistency of *Aspergillus*, *Penicillium*, and *Talaromyces* species. *Front Microbiol*. 5:773–783.
- Gao W, Jiang L, Ge L, Chen M, Geng C, Yang G, Li Q, Ji F, Yan Q, Zou Y, et al. **2015**. Sterigmatocystin-induced oxidative DNA damage in human liver-derived cell line through lysosomal damage. *Toxicol Vitro*. 29:1–7.
- Greshock T, Grubbs AW, Jiao P, Wicklow DT, Gloer JB, Williams RM. **2008**. Isolation, structure elucidation, and biomimetic total synthesis of versicolamide B, and the isolation of antipodal (-)-stephacidin A and (+)-notoamide B from *Aspergillus versicolor* NRRL 35600. *Angewandte Chemie*. 47:3573–3577.
- Haas D, Pfeifer B, Reiterich C, Partenheimer R, Reck B, Buzina W. **2013**. Identification and quantification of fungi and mycotoxins from Pu-erh tea. *Int J Food Microbiology*. 166(2):316–322.
- Halstensen AS. **2008**. Species-specific fungal DNA in airborne dust as surrogate for occupational mycotoxin exposure? *Int J Mol Sci*. 9:2543–2558.
- Hammami W, Fiori S, Al Thani R, Ali Kali N, Balmas V, Miglieli Q, Jaoua S. **2014**. Fungal and aflatoxin contamination of marketed spices. *Food Control*. 37:177–181.
- Houbraken J, De Vries RP, Samson RA. **2014**. Modern taxonomy of biotechnologically important *Aspergillus* and *Penicillium* species. *Adv Appl Microbiol*. 86:199–249.
- Huang S, Wang J, Xing L, Shen H, Yan X, Wang J, Zhang X. **2014**. Impairment of cell cycle progression by sterigmatocystin in human pulmonary cells in vitro. *Food Chem Toxicol*. 66:89–95.
- Huang X, Zhang X, Yan X, Yin G. **2002**. Effects of sterigmatocystin on interleukin-2 secretion of human peripheral blood mononuclear cells in vitro. *J Hyg Res*. 31:112–114.
- Hubka V, Nováková A, Kolarik A, Jurjevic Z, Peterson SW. **2014**. Revision of *Aspergillus* section *flavipes*: seven new species and proposal of section *Jani* sect. *Nov. Mycologia*. 107(1):169–208.
- Hubka V, Nováková A, Peterson SW, Stephen W, Frisvad JC, Sklenar F, Matsuzawa T, Kubátová A, Kolarik M. **2016**. A reappraisal of *Aspergillus* section *nidulantes* with descriptions of two new sterigmatocystin-producing species. *Plant Systematics and Evolution*. 302:1267–1299.
- In: IPCC, Pachauri RK, Reisinger A, ed. **2007**. Climate change 2007: synthesis report. Contribution of working groups I, II and III to the fourth assessment report of the intergovernmental panel on climate change. Geneva: IPCC.
- Jakšić Despot D, Kocsbá S, Bencsik O, Kecskeméti A, Szekeres A, Vágvölgyi C, Varga J, Šegvić Klarić M. **2017**. New sterigmatocystin-producing species of *Aspergillus* section *Versicolores* from indoor air in Croatia. *Mycological Prog*. 16(1):63–72.
- Jiang X, Wang J, Xing L, Shen H, Lian W, Yi L, Zhang D, Yang H, Liu J, Zhang X. **2017**. Sterigmatocystin-induced checkpoint adaptation depends on Chk1 in immortalized human gastric epithelial cells in vitro. *Arch Toxicol*. 1(91):259–270.
- Jiao P, Mudur SV, Gloer JB, Wicklow DT. **2007**. Kipukasins, nucleoside derivatives from *Aspergillus versicolor*. *J Nat Prod*. 70:1308–1311.
- Jurjevic Z, Peterson SW, Solfrizzo M, Peraica M. **2013**. Sterigmatocystin production by nine newly described *Aspergillus* species in section *Versicolores* grown on two different media. *Mycol Res*. 29:141–145.
- Kačániov M. **2003**. Feeding soybean colonization by microscopic fungi. *Trakya Univ J Sci*. 4(2):165–168.
- Kačániová M, Krňazovická V, Felšciová S, Rovná K. **2012**. Microscopic fungi recovered from honey and their toxigenicity. *J Environ Sci Health*. 47(11):1659–1664.

- Kawai K, Nakamaru T, Nozawa Y, Maebayashi Y, Yamazaki M, Natori S. 1984. Inhibitory effect of sterigmatocystin and 5,6-dimethoxysterigmatocystin on ATP synthesis in mitochondria. *Appl Environ Microbiol.* 48:1001–1003.
- Kelkar HS, Keller NP, Adams TH. 1996. *Aspergillus nidulans* stcP encodes an O-methyltransferase that is required for sterigmatocystin biosynthesis. *Appl Environ Microbiol.* 62:4296–4298.
- Kocić-Tanackov SD, Dimić GR, Lević JT, Pejin DJ, Pejin JD, Jajić IM. 2010. Occurrence of potentially toxic mould species in fresh salads of different kinds of ready-for-use vegetables. *APTEFF.* 41:1–203.
- Kovalenko AV, Soldatenko NA, Fetisov LN, Strel'tsov NV. 2011. More accurate determination of the minimum allowable level of sterigmatocystin in piglet feed. *Russian Agric Sci.* 37(6):504–507.
- Labuda R, Tančinová D. 2006. Fungi recovered from Slovakian poultry feed mixtures and their toxinogenicity. *Ann Agric Environ Med.* 13:193–200.
- Lee YM, Li H, Hong J, Cho HJ, Bae KS, Kim MA, Kim D-K, Jung JH. 2010. bioactive metabolites from the sponge-derived fungus *aspergillus versicolor*. *Arch Pharmacological Res.* 33:231–235.
- Lima CAP, Orsi R, Dilkin P, Correa B. 2000. Mycoflora and aflatoxigenic species in derivatives of milled rice. *Ciênc Tecnol Aliment.* 20(1):37–39.
- Liu Y, Du M, Zhang G. 2014. Proapoptotic activity of aflatoxin B1 and sterigmatocystin in HepG2 cells. *Toxicol Rep.* 1:1076–1086.
- Magan N, Hope R, Cairns V, Aldred D. 2003. Post-harvest fungal ecology: impact of fungal growth and mycotoxin accumulation in stored grain. *Eur J Plant Pathol.* 109 (7):723–730.
- Magnoli C, Chiacchiera SM, Miazzo R, Palacio G, Angeletti A, Hallak C, Dalcero A. 2002. The mycoflora and toxicity of feedstuffs from a production plant in Cordoba, Argentina. *Mycotoxin Res.* 18(1):7–22.
- Makun HA, Anjorin ST, Moronfoye B, Adejo FO, Afolabi OA, Fagbayibo G, Balogun BO, Surajudeen AA. 2010. Fungal and aflatoxin contamination of some human food commodities in Nigeria. *Afr J Food Sci.* 4(4):127–135.
- McConnell IR, Garner RC. 1994. DNA adducts of aflatoxins, sterigmatocystin and other mycotoxins. In: IARC scientific publications. (125):49–55.
- Miller JD, Sun M, Gilyan A, Roy J, Rand TG. 2010. Inflammation-associated gene transcription and expression in mouse lungs induced by low molecular weight compounds from fungi from the built environment. *Chem Biol Interact.* 183:113–124.
- Miller JD, Trenholm L, eds. 1994. Mycotoxins in grain: compounds other than aflatoxin: ACCI Press.
- Mo HGJ, Pietri A, MacDonald SJ, Anagnostopoulos C, Spanjer M. 2015. on sterigmatocystin in food. EFSA Supporting Publication. 12(3):774E.
- Mol JGJ, De Rijk TC, Van Egmond H, De Jong J. 2014. Occurrence of mycotoxins and pesticides in straw and hay used as animal feed. Wageningen RIKILT Wageningen UR (RIKILT report 2014.006)
- Mori H, Sugie S, Yoshimi N, Kitamura J, Niwa M, Hamasaki T, Kawai K. 1986. Genotoxic effects of a variety of sterigmatocystin-related compounds in the hepatocyte/DNA-repair test and the *Salmonella* microsome assay. *Mutat Res.* 173:217–222.
- Nichea MJ, Palacios SA, Chiacchiera SM, Sulyok M, Krška R, Chulze SN, Torres AM, Ramirez ML. 2015. Presence of multiple mycotoxins and other fungal metabolites in native grasses from a wetland ecosystem in Argentina intended for grazing cattle. *Toxins.* 7(8):3309–3329.
- Peraica M, Radic B, Lucic A, Pavlovic M. 1999. Toxic effects of mycotoxins in humans. *Who.* 77:754–766.
- Paterson RRM, Lima N. 2011. Further mycotoxin effects from climate change. *Food Res Int* [Internet]. 44:2555–2566. Available from: <https://doi.org/10.1016/j.foodres.2011.05.038>
- Piecková E, Hurbánková M, Černá S, Majorošová M, Kováčiková Z, Pangallo D. 2011. Respiratory toxicity of *Aspergillus versicolor*: the most common indoor mould in Slovakia. In: Brebbia CA, Egli M, Knets I, Of R M, Popov V, Eds. Environmental health and biomedicine 15. Southampton: Wit Press; p. 135–145.
- Piecková E. 2015. Domestic environment – indoor mycobiota as a public health risk factor. In: Viegas C, Pinheiro C, Sabino R, Viegas S, Branda J, Veríssimo C, Eds. Environmental mycology in public health. Fungi and mycotoxins risk assessment and management. Elsevier – Academic Press; p. 129–146. <https://www.elsevier.com/books/environmental-mycology-in-public-health/viegas/978-0-12-411471-5>
- Piotrowska M. 2013. Contamination of breakfast cereal products by fungi and mycotoxins – a potential risk for consumer's health. *Biotechnol Food Sci.* 77(1):3–10.
- Purchase IF, Van Der Watt JJ. 1969. Acute toxicity of sterigmatocystin to rats. *Food Cosmet Toxicol.* 7:135–139.
- Rank C, Nielsen KF, Larsen TO, Varga J, Samson RA, Frisvad JC. 2011. Distribution of sterigmatocystin in filamentous fungi. *Fungal Biol.* 115:406–420.
- Raper KB, Fennell DL. 1965. The genus *Aspergillus*: Williams and Wilkins Company, Baltimore. 686.
- Reddy MV, Irvin TR, Randerath K. 1985. Formation and persistence of sterigmatocystin DNA adducts in rat-liver determined via P-32-postlabeling analysis. *Mutat Res.* 152:85–96.
- Riba A, Mokrane S, Mathieu F, Lebrihi A, Sabaou N. 2008. Mycoflora and ochratoxin A producing strains of *Aspergillus* in Algerian wheat. *Int J Food Microbiol.* 122(1–2):85–92.
- Rofiat AS, Fanelli F, Atanda O, Sulyok M, Cozzi G, Bavaro S, Krška R, Logrieco AF, Ezekiel CN. 2015. Fungal and bacterial metabolites associated with natural contamination of locally processed rice (*Oryza sativa* L.) in Nigeria. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 32(6):950–959.
- Rosa CAR, Cavagliere LR, Ribeiro JMM, Keller KM, Alonso VA, Chiacchiera SM, Dalcero AM, Lopes CWG. 2008. Mycobiota and naturally-occurring ochratoxin A in dairy cattle feed from Rio de Janeiro State, Brazil. *World Mycotoxin Journal.* 1(2):195–201.
- Sabino R, Faísca VM, Carolino E, Veríssimo C, Viegas C. 2012. Occupational exposure to aspergillus by swine and poultry farm workers in Portugal. *J Toxicol Environ Health.* 75:1381–1391.

- Samson RA, Houbraken J, Thrane U, Frisvad JC, Andersen B. **2010**. Food and indoor fungi: CBS-KNAW fungal biodiversity centre. Utrecht. 390.
- Samson RA, Visagie CM, Houbraken J, et al. **2014**. Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology*. 78:141–173.
- Sawane A, Sawane M. **2014**. Mycotoxicogenicity of *Aspergillus*, *Penicillium* spp. isolated stored rice. *Int J Curr Microbiol Appl Sciences*. 3(11):116–121.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CV, Chen W, Consortium FB. **2012**. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. In: Janzen DH, Ed. *Proceedings of the national academy of sciences of the United States of America*, Philadelphia; p. 6241–6246.
- Silva JB, Pozzi CR, Mallozzi MAB, Ortega EM, Corre B. **2000**. Mycoflora and occurrence of aflatoxin B1 and fumonisin B1 during storage of Brazilian sorghum. *J Agric Food Chem*. 48:4352–4356.
- Sivakumar V, Thanislass J, Niranjali S, Devaraj H. **2001**. Lipid peroxidation as a possible secondary mechanism of sterigmatocystin toxicity. *Hum Exp Toxicol*. 2:398–403.
- Sonjak S, Ličena M, Frisvad JC, Gunde-Cimerman N. **2011**. The mycobiota of three dry-cured meat products from Slovenia. *Food Microbiol*. 28(3):373–376.
- Tabuc C, Marin D, Guerre P, Sesan T, Bailly JD. **2009**. Molds and mycotoxin content of cereals in Southeastern Romania. *J Food Prot*. 72(3):662–665.
- Tabuc C, Stroia C, Neacsu A. **2010**. Incidence of *Aspergillus* strains and of aflatoxin B1 in cereals in South-Western Romania. *Res J Agric Sci*. 42(2):42–47.
- Tabuc C, Tarantu I, Calin L. **2011**. Survey of moulds and mycotoxin contamination of cereals in South-Eastern Romania in 2008–2010. *Archiva Zootechnica*. 14(4):25–38.
- Tančinová D, Barboráková Z, Kačinová J, Mašková Z, Volčková M. **2013**. The occurrence of micromycetes in apples and their potential ability to produce mycotoxins. *J Microbiology, Biotechnol Food Sci*. 2(Special issue 1):1800–1807.
- Tančinová D, Labuda R. **2009**. Fungi on wheat bran and their toxinogeneity. *Ann Agric Environ Med*. 16:325–331.
- Toffa DD, Mahnine N, Ouaffak L, El Abidi A, El Alaoui Faris FZ, Zinedine A. **2013**. First survey on the presence of ochratoxin A and fungi in raw cereals and peanut available in the Republic of Niger. *Food Control*. 32:558–562.
- Townsend CA. **1997**. Progress towards a biosynthetic rationale of the aflatoxin pathway. *Pure Appl Chem*. 58:227–238.
- Ueda N, Fujie K, Gotoh-Mimura K, Chattopadhyay SC, Sugiyama T. **1984**. Acute cytogenetic effect of sterigmatocystin on rat bone-marrow cells in vivo. *Mutat Res*. 139:203–206.
- Ueno Y, Umemori K, Niimi E, Tanuma S, Nagata S, Sugamata M, Ihara T, Sekijima M, Kawai K, Ueno I, et al. **1995**. Induction of apoptosis by T-2 toxin and other natural toxins in HL-60 human promyelotic leukemia cells. *Nat Toxins*. 3:129–137.
- Van der Fels-Klerk HJ, Liu C, Battilani P. **2016**. Modelling climate change impacts on mycotoxin contamination. *World Mycotoxin J [Internet]*. 9:1–10. Available from:<http://www.wageningenacademic.com/doi/10.3920/WMJ2016.2066>
- Veršilovskis A, Bartkevičs V. **2012**. Stability of sterigmatocystin during the bread making process and its occurrence in bread from the Latvian market. *Mycotoxin Res*. 28:123.
- Veršilovskis A, Bartkevičs V, Mikelsone V. **2008a**. Sterigmatocystin presence in typical Latvian grains. *Food Chemistry* 109(1):243–248.
- Veršilovskis A, De Saeger S, Mikelsone V. **2008b**. Determination of sterigmatocystin in beer by high performance liquid chromatography with ultraviolet detection. *World Mycotoxin Journal*. 1(2):161–166.
- Veršilovskis A, Van Peteghem C, De Saeger S. **2009**. Determination of sterigmatocystin in cheese by high-performance liquid chromatography-tandem mass spectrometry. *Food Additives & Contaminants: Part A*. 26(1):127–133.
- Viegas C, Pacífico CFaria T. **2017**. Fungal contamination in green coffee beans samples: a public health concern. *Journal Of Toxicology and Environmental Health, Part a*. doi:10.1080/15287394.2017.1286927
- Viegas C, Sabino R, Viegas S, Veríssimo C. **2013**. Occupational exposure to toxicigenic *Aspergillus versicolor* in Portuguese swine. *Int Symp Occup Saf Hyg*. 3: 433–434.
- Viegas S, Veiga L, Figueiredo P, Almeida A, Carolino E, Viegas C. **2015**. Assessment of workers' exposure to aflatoxin B1 in a Portuguese waste industry. *Ann Occup Hyg*. 59(2):173–181.
- Visagie CM, Hirooka Y, Tanney JB, Whitfield E, Mwange K, Meijer M, Amend AS, Seifert KA, Samson RA. **2014**. *Aspergillus*, *Penicillium* and *Talaromyces* isolated from house dust samples collected around the world. *Stud Mycol*. 78:63–139.
- Walkow J, Sullivan G, Maness D, Yakatan GJ. **1985**. Sex and age differences in the distribution of 14C-sterigmatocystin in immature and mature rats: a multiple dose study. *J Am Coll Toxicol*. 4:45–51.
- Wang J, Huang S, Xing L, Cui J, Tian Z, Shen H, Jiang X, Yan X, Wang J, Zhang X. **2015**. Sterigmatocystin induces G1 arrest in primary human esophageal epithelial cells but induces G2 arrest in immortalized cells: key mechanistic differences in these two models. *Arch Toxicol*. 89(11):2015–2025.
- Warth B, Parich A, Atehnkeng J, Bandyopadhyay R, Schuhmacher R, Sulyok M, R. K. **2012**. Quantitation of mycotoxins in food and feed from burkina faso and Mozambique using a modern LC-MS/MS multitoxin method. *J. Agric. Food Chem*. 60(36):9352–9363.
- Xie T-X, Misumi J, Aoki K, Zhao W-Y, Liu S-Y. **2000**. Absence of p53-mediated G1 arrest with induction of MDM2 in sterigmatocystin-treated cells. *Int J Oncol*. 17:737–742.
- Xing LX, Zhang XH, Li YH, Yan X, Wang J, Wang F. **2005**. Effects of sterigmatocystin on HLA- I expression of human peripheral blood mononuclear cells in vitro. *J Hyg Res*. 34:454–456.
- Yassin MA, Ama E-S, Moslem MA, El-Naggar MA. **2013**. Mycobiota of almond seeds and the toxigenicity of some involved genera. *Life Sci J*. 10(4):1088–1093.
- Youssef MS, Omo E-M, Ibrahim YM. **2008**. Mycobiota and mycotoxins of Egyptian peanut (*Arachis Hypogaea* L.) seeds. *Int J Bot*. 4(4):349–360.

Yu J, Chang P-K, Ehrlich KC, Cary JW, Bhatnagar D, Cleveland TE, Payne GA, Linz JE, Woloshuk CP, Bennett JW. **2004**. Clustered pathway genes in aflatoxin biosynthesis. *Appl Environ Microbiol.* 70:1253–1262.

Zhang Y, Yao ZG, Wang J, Xing LX, Xia Y, Zhang XH. **2012**. Effects of sterigmatocystin on TNF-alpha, IL-6 and IL-12 expression in murine peripheral blood mononuclear cells and peritoneal macrophages *in vivo*. *Mol Med Rep.* 5:1318–1322.