

Haemoparasites of the pied flycatcher: inter-population variation in the prevalence and community composition

Anna Dubiec¹, Edyta Podmokła², Iga Harnist¹ and Tomasz D. Mazgajski¹

¹Museum and Institute of Zoology, Polish Academy of Sciences, Wilcza 64, 00-679 Warszawa, Poland and ²Institute of Environmental Sciences, Jagiellonian University, Gronostajowa 7, 30-387 Kraków, Poland

Research Article

Cite this article: Dubiec A, Podmokła E, Harnist I, Mazgajski TD. Haemoparasites of the pied flycatcher: inter-population variation in the prevalence and community composition. *Parasitology* <https://doi.org/10.1017/S0031182017001913>

Received: 17 February 2017
Revised: 14 September 2017
Accepted: 21 September 2017

Key words:

Co-infection; diversity index; haemosporidians; phylogenetic relationships; trypanosomes; wildlife diseases

Author for correspondence:

Edyta Podmokła, E-mail: edytapodmokla@gmail.com

Abstract

The prevalence and community composition of haemoparasites can substantially differ among avian host populations, which may lead to different selection pressures. Therefore, information about these parameters is crucial for understanding, e.g. the inter-population variation in host life history traits. Here, we molecularly screened a population of a long-distance migrant, the pied flycatcher *Ficedula hypoleuca*, from central Poland for the presence of three genera of blood parasites: *Haemoproteus*, *Plasmodium* and *Trypanosoma*. The infection rate in this population was the highest for haemosporidians (86.8%) and one of the highest for trypanosomes (39.7%) among the thus far screened breeding populations of this species. The haemosporidian community was composed of six *Haemoproteus/Plasmodium* lineages, and the trypanosome community – 4 species and a parasite assigned to genus level. Trypanosomes were dominated by *T. culicavium*, a recently described species, corroborating the prediction that insectivorous songbirds are vertebrate hosts of this parasite. Host sex and age did not explain variation in infection incidence except for the higher trypanosome infection rates in males. A comparison of the study population with three other breeding populations previously screened molecularly for haemosporidians showed some geographic differences. This study confirms the importance of examining local parasite communities across a host distribution range.

Introduction

Parasites are one of the most important factors exerting selection pressures in animal populations (Haldane, 1949). Generally, each species may act as a competent host for a very versatile and large community of parasites. However, only rarely do these communities show high similarity in terms of prevalence and diversity across the distribution range of the host (Freeman-Gallant *et al.* 2001; Poulin, 2003). Differences in parasite communities should translate into diverse selection pressures acting upon the host, and this in turn could result in variations of life-history traits among host populations. Consequently, in order to understand the role of parasites in shaping inter-population differences in fitness-associated traits, it is necessary to examine local parasite communities and investigate the patterns of variation in infection/infestation rates.

In birds, among the commonest parasites are vector-transmitted protozoans, which complete some stages of the life cycle in the blood of the vertebrate host (therefore, they are generally referred to as blood parasites or haemoparasites). This group includes, among others, parasites from the genera *Plasmodium*, *Haemoproteus* and *Trypanosoma* (Clark *et al.* 2009). Avian haemosporidians and trypanosomes are transmitted by blood-sucking arthropods. Specifically, *Plasmodium* is transmitted by mosquitoes (Culicidae), *Haemoproteus* by hippoboscids (Hippoboscidae) and biting midges (Ceratopogonidae) and *Trypanosoma* by black flies (Simuliidae), hippoboscids (Hippoboscidae), biting midges (Ceratopogonidae), *Culex* spp. mosquitoes (Culicidae) and dermanyssid mites (Dermanyssidae) (Baker, 1976; Molyneux, 1977; Votýpka *et al.* 2012). Despite being transmitted by the same type of vectors (blood-sucking arthropods) haemosporidians and trypanosomes differ in their transmission modes: haemosporidians enter the host organism through the bite of the vector, and trypanosomes – through ingestion of the vector or contamination of abraded skin or conjunctiva with material containing infective parasite stages (Desser *et al.* 1975; Molyneux, 1977; Votýpka and Svobodová, 2004; Votýpka *et al.* 2012). Parasites from all three genera can negatively affect components of fitness in birds (Merino *et al.* 2000; Knowles *et al.* 2010; Martínez-de la Puente *et al.* 2010), although a number of studies failed to find such associations (Dufva, 1996; Siikamäki *et al.* 1997). The introduction at the beginning of the XXI century of molecular techniques for detection and identification of these parasites not only improved the detectability of infection (Hellgren *et al.* 2004; Waldenström *et al.* 2004), but also uncovered their extreme genetic diversity (Bensch *et al.* 2009). Importantly, by using a molecular approach, it was possible to show that genetic lineages identified as the same morphospecies may, in fact, represent separate biological entities (e.g. Bensch *et al.* 2004; Nilsson *et al.* 2016).

Examination of different populations of the same host species in the context of infection rates and diversity of blood parasites has shown that both parameters can vary geographically (Bensch and Åkesson, 2003; Pagenkopp *et al.* 2008; Szöllösi *et al.* 2011; Spurgin *et al.* 2012), even within a short distance (Sol *et al.* 2000; Ferrer *et al.* 2012). However, the geographic variation of these parameters is not a universal phenomenon. In some host species, it has not been found or was restricted to a subgroup of individuals (Swanson *et al.* 2014; Svoboda *et al.* 2015). For example, in the breeding populations of the migratory lark sparrow (*Chondestes grammacus*) located across the USA, geographic differences in the overall prevalence were present only in juvenile birds, but not in the adults (Swanson *et al.* 2014). Similarly, adult bluethroats (*Luscinia svecica*) sampled in seven populations ranging in location from central to northern Europe did not differ in overall infection rates (Svoboda *et al.* 2015).

In the present study, we molecularly examine the prevalence and community composition of parasites from the *Haemoproteus*, *Plasmodium* and *Trypanosoma* genera in a population of the pied flycatcher (*Ficedula hypoleuca*) located in central Poland. The pied flycatcher is a long-distance migrant which breeds in most of Europe and western Asia and winters in West Africa (Cramp and Perrins, 1993; Ouweland *et al.* 2016). It is a model species in the evolutionary ecology of cavity-nesting long-distance migrants with several closely monitored populations located across its breeding range (e.g. Laaksonen *et al.* 2015; Ouweland *et al.* 2016). To date, several studies looked into the community of blood parasites in this species, but most of them were based on blood smear screening (Bennett *et al.* 1994; Dale *et al.* 1996; Siikamäki *et al.* 1997; Sanz *et al.* 2001a) and consequently may have underestimated infection rates and the diversity of these parasites (Fallon *et al.* 2003; Garamszegi, 2010; Nilsson *et al.* 2016). Importantly, this is one of the few studies molecularly examining avian trypanosomes. In contrast to haemosporidian parasites, infections with *Trypanosoma* have been largely understudied in birds, although such infections may have serious consequences for a host's fitness, including the study species (Rätti *et al.* 1993; Sanz *et al.* 2001a; Dyrz *et al.* 2005, also see Siikamäki *et al.* 1997; Sanz *et al.* 2001b), and are known to widely occur in birds (Sehgal *et al.* 2001; Oakgrove *et al.* 2014). Additionally, we: (i) studied the relationship between infection incidence and two host traits commonly considered in the analyses of the presence of infection, namely sex and age (McCurdy *et al.* 1998; Deviche *et al.* 2001; van Oers *et al.* 2010), and (ii) investigated geographic variation in the prevalence and diversity of haemosporidian parasites (*Haemoproteus* and *Plasmodium*) among four molecularly screened breeding populations of the pied flycatcher using data from the current and previously published studies (Wiersch *et al.* 2007; Kulma *et al.* 2013).

Materials and methods

Data were collected from a small population (up to 17 pairs per year) of nest box-breeding pied flycatchers in 2011–2014. The study site is located in Sękocin Stary, 10 km south-west of Warsaw's city borders, Poland (52°05'N, 20°52'E). All study plots except for one are located in a pine (*Pinus sylvestris*) forest with an admixture of deciduous trees [birch (*Betula* spp.), sessile (*Quercus petraea*) and pedunculate (*Quercus robur*) oaks and common hornbeam (*Carpinus betulus*)] with a moderate to dense understory. Nest boxes (for their dimensions and distribution see Dubiec and Mazgajski, 2013) are primarily occupied by great tits (*Parus major*), blue tits (*Cyanistes caeruleus*) and pied flycatchers, and occasionally by coal tits (*Periparus ater*). The number of nest boxes varied among breeding seasons, from 156 in 2011 to 259 in 2014. Pied flycatchers bred only in pine-

dominated plots. We additionally monitored over 35 nest boxes, differing in size and the height of attachment, located in close vicinity to the study plots. Each year at least one pied flycatcher breeding attempt was recorded in these boxes.

Before each breeding season nest boxes were cleaned of remaining nest material and feces of roosting birds. Regular controls of boxes, in order to record phenological data and breeding parameters, started in the beginning of April. Adult birds were caught when the nestlings were 13 days old (hatching day = day 1) or older, either by mist-netting or using traps installed inside the box. Birds were ringed, sexed, aged, measured (data not shown) and sampled for blood. Sexing and ageing (as yearlings or at least 2 years old) were based on plumage characteristics following Svensson (1994). In the case of re-trapped birds, the assigned age was later verified based on ringing records from previous seasons. Blood, obtained by puncturing the brachial vein, was collected into non-heparinized capillaries placed in 96% ethanol. Until the molecular analyses were performed, blood samples were stored at ambient temperature. Because of the brood mortality at the early post-hatching stage or unsuccessful catching trials (mostly because the birds were not present in the vicinity of the box during a catching trial), a few individuals were not caught.

DNA was extracted using the standard ammonium acetate precipitation method (Bruford *et al.* 1998). DNA isolates were checked for concentration and purity (absorbance ratios: A260/A280 and A260/A230) using a NanoDrop spectrophotometer. Samples were screened for the presence of blood parasites: (1) from the *Haemoproteus* and *Plasmodium* genera by amplifying a 478 bp fragment of the mitochondrial *cyt b* gene, using the nested polymerase chain reaction (Waldenström *et al.* 2004; Cosgrove *et al.* 2008); (2) from the *Trypanosoma* genus by amplifying a 326 bp fragment of the small subunit ribosomal RNA (SSU rRNA) gene as described in Sehgal *et al.* (2001). Four μ L of final PCR products were run on 2% agarose gels stained with GelRed (Biotium, Hayward, CA, USA) and visualized under ultraviolet light. Each plate contained a positive (DNA from individuals with confirmed infection based on blood smear screening) and negative (ddH₂O) control, to monitor for possible contamination or failures during PCRs. The quality of the DNA isolate of all samples yielding no PCR product was checked with sex-specific primers (Griffiths *et al.* 1998). PCR products of positive samples were purified with FastAP (Fermentas) and then sequenced directly (mostly unidirectionally, except for a subset of trypanosome PCR products, which were sequenced bidirectionally) with an automated ABI 3130 DNA analyser (Applied Biosystems) using BigDye terminator ver. 3.1 (Applied Biosystems). The obtained sequences were aligned using BioEdit software (Hall, 1999) and compared with the MalAvi database (Bensch *et al.* 2009) in the case of *Plasmodium* and *Haemoproteus* and with the GenBank database in the case of *Trypanosoma*.

In total, we screened 97 samples from 68 individuals. Twenty-one individuals were sampled more than once. To avoid the issue of pseudoreplication, only one record per multiply sampled individuals was selected for further analyses. The record was selected either randomly (most cases) or non-randomly when information was lacking (age of the individual) in the datasets collected in different seasons. In such cases, the record with the full set of data was chosen for statistical analyses. The final data set included 43 females and 25 males.

The association between occurrences of parasites representing different genera within a host was analysed with Fisher's exact test. To assess the association between the probability of infection (separately for each parasite genera) and sex and age of the host generalized linear mixed models (GLMM) with a binomial error distribution and a logit link function were used. The individual

age and sex were defined as fixed categorical variables, a year as a random variable and infection status (0 for absence and 1 for presence) as a response variable. The interaction between sex and age was introduced into the model, but removed if not significant. The model was run using a dataset reduced by four females, because these individuals were not aged (final data set: females – 18 yearlings and 21 older birds, males – 12 yearlings and 13 older birds). Because *Plasmodium* infections occurred at a low rate (see 'Results' section), the analyses of haemoparasite co-occurrence and age- and sex-related variation were limited to *Haemoproteus* and *Trypanosoma* infections. GLMM models were run using the glmer function in the package 'lme4' (Bates *et al.* 2015) in R (R Development Core Team, 2014) with the Laplace approximation of maximum likelihood (Bolker *et al.* 2009).

Geographic variation in the prevalence and ecological diversity of haemosporidian parasites among the breeding populations of the pied flycatcher was estimated for three previously molecularly screened populations – Lingen and Bahrddorf in Germany (sampling period: 1993 for Bahrddorf and 1996 for Lingen; Wiersch *et al.* 2007) and the island of Öland in Sweden (sampling period: 2004–2010, except 2008; Kulma *et al.* 2013) – and the population examined in the present study (Fig. 1). In the case of four mitochondrial sequences from German populations (one out of six sequences in Lingen and three out of four sequences in

Bahrddorf), it was not possible to assign them to specific lineages from the MalAvi database because only information about the GenBank entry most closely matching each of these sequences (the similarity was over 98% in all cases) was provided. These entries were used in the current study as a reference to assign the sequence to a parasite genus. Two ecological diversity indices were used: the species richness (S – the total number of detected lineages) and the Shannon index (H ; combining information on species richness and relative abundance). The diversity indices were calculated in the 'vegan' package (Oksanen *et al.* 2016) in R, while the 95% confidence intervals (CIs) of the Shannon index and differences in those indexes between populations were calculated in the 'boot' package (Canty and Ripley, 2015) according to the recommendations of Gardener (2014). CIs were estimated based on 1000 bootstrap resamples (Custom R Command: 'H_boot' from Gardener, 2014). The statistical significance of the difference in diversity index between populations was estimated using a randomization method based on 2000 bootstrap resamples (Custom R Command: 'H_bss' from Gardener, 2014).

The phylogenetic analysis includes sequences of *Haemoproteus* and *Plasmodium* lineages from the present study and two other European breeding populations of pied flycatchers [Lingen (five sequences that could be matched with specific lineages from the MalAvi database) and Öland] as well as additional lineages found previously in this species (birds caught on breeding




Fig. 1. Distribution and prevalence of *Plasmodium* and *Haemoproteus* parasites across four breeding populations of the pied flycatcher molecularly screened for the presence of parasites: Bahrddorf and Lingen in Germany (Wiersch *et al.* 2007), the island of Öland in Sweden (Kulma *et al.* 2013) and Sękocin Stary in Poland (present study). Pie charts show the parasite genus frequencies of each population. Sample sizes: $N_{\text{Bahrddorf}} = 72$; $N_{\text{Lingen}} = 147$; $N_{\text{Öland}} = 353$; $N_{\text{Sękocin Stary}} = 68$. Note that *Haemoproteus* infection rates recorded in Bahrddorf and Lingen populations may be underestimated as a consequence of the applied protocol (see 'Discussion' for details).

grounds either during or outside of the breeding season), which we extracted from the MalAvi database (version: Malavi8120153_04102016; Bensch *et al.* 2009). Bayesian reconstructions were carried out in MrBayes v. 3.1.2 (Ronquist and Huelsenbeck, 2003), following a General Time Reversible model including invariable sites and variation among sites (GTR + I + G) suggested by the software jModelTest 2.1.10 (Darriba *et al.* 2012). The *Leucocytozoon* spp. lineage SISKIN2 was used as an outgroup and two separate runs of 10 million generations were conducted with a sampling frequency of every 100th generation. The burn-in fraction was set to 0.25.

Results

Infection rates of the study population of pied flycatchers varied from very frequent for *Haemoproteus* (79.4%), through moderate for *Trypanosoma* (39.7%) to rare for *Plasmodium* (8.8%). Parasites from the *Haemoproteus* and *Plasmodium* genera were represented by two and four cytochrome *b* lineages, respectively (Table 1). The most common lineage – PFC1, representing the morphospecies *H. pallidus* – was found in 96.3% of *Haemoproteus*-infected individuals. Other lineages occurred at a frequency of <5%, including two lineages detected only in a single individual each (Table 1B). Only two birds (2.9%) carried mixed haemosporidian infections: one individual with two *Haemoproteus* lineages and the other with *Haemoproteus* and *Plasmodium* lineages. In the case of *Trypanosoma* parasites, five types of parasite SSU rRNA sequences were detected: four assigned to a species level and one assigned to a genus level (Table 1C). The most common species – *T. culicaviium*, occurred in 59.2% of *Trypanosoma*-infected individuals. Only one bird (1.5%) carried a mixed (double) infection.

Among all screened individuals, 35.3% carried simultaneous infections of *Haemoproteus* and *Trypanosoma* parasites. Because of differences between males and females in *Trypanosoma* infection rates (see below), the analyses of the association between the occurrences of these two parasite genera within an individual host were performed separately for each sex. The occurrence of *Haemoproteus* was not associated with the occurrence of

Trypanosoma either in males or females (Fisher's exact test, males: $P = 0.565$, females: $P = 0.290$).

The probability of carrying a *Haemoproteus* infection was not associated with either the sex or age of the host (sex: males – 88.0%, females – 69.2%, age: yearlings – 83.3%, older birds – 70.6%; Table 2). In the case of *Trypanosoma*, the probability of infection varied in relation to the host's sex, with males being more frequently infected than females (sex: males – 56.0%, females – 30.8%, age: yearlings – 40.0%, older birds – 41.2%; Table 2).

The prevalence of haemosporidian infections varied among four breeding populations of the pied flycatcher, with the lowest incidence of infection in the two German populations, intermediate in Öland and the highest in the population from central Poland ($\chi^2 = 213.81$, $df = 3$, $P < 0.001$, Table 3). Also, the proportion of infections caused by each parasite genus differed among populations (Fisher's exact test, $P < 0.001$). In general, infections in the two German populations were dominated by *Plasmodium*, while in the two other populations, by *Haemoproteus* (Fig. 1, Table 3). Lineage richness was the highest in the Öland population with over twice as many lineages as in each of the three other populations (Table 3), while the Shannon index was over two times lower in Sękocin Stary than in the three other populations (Table 3). However, only the difference between the Sękocin Stary and Öland populations was statistically significant (Öland vs Sękocin Stary: $z = 5.79$, $P < 0.001$; Lingen vs Sękocin Stary: $z = 1.92$, $P = 0.08$; Lingen vs Öland: $z = 0.82$, $P = 0.42$; Lingen vs Bährdorf: $z = 0.66$, $P = 0.53$; Öland vs Bährdorf: $z = 1.53$, $P = 0.14$; Sękocin Stary vs Bährdorf: $z = 0.51$, $P = 0.62$). Among 18 lineages detected in Lingen, Öland and Sękocin Stary, only one (pSGS1) was common for all populations, while 66.7% of the lineages occurred only in a single population (Fig. 2).

Lineages included in the phylogenetic analysis (18 lineages from Lingen, Öland, Sękocin Stary and five additional lineages detected in pied flycatchers on breeding grounds, Fig. 2) form two major clades of haemosporidian parasites: *Haemoproteus* and *Plasmodium*. The lineages found in all three breeding populations are dispersed in the clades and do not form monophyletic groups.

Table 1. The community of blood parasites [genera *Haemoproteus* (A), *Plasmodium* (B) and *Trypanosoma* (C)] in the pied flycatcher population from central Poland (Sękocin Stary) in 2011–2014 ($n = 68$)

Parasite taxon	Lineage	GenBank accession no.	$N_{\text{occurring}}$	Prevalence, %
(A)				
<i>Haemoproteus pallidus</i>	PFC1	JX026906	52	76.5
<i>Haemoproteus majoris</i>	PHSIB1	AF495565	2	2.9
(B)				
<i>Plasmodium ashfordi</i>	GRW02	AF254962	1	1.5
<i>Plasmodium</i> sp.	GRW09	FJ404701	3	4.4
<i>Plasmodium</i> sp.	PLOPRIO1	FJ404702	1	1.5
<i>Plasmodium relictum</i>	SGS1	AF495571	2	2.9
(C)				
<i>Trypanosoma avium</i>		AY099320	2	2.9
<i>Trypanosoma bennetti</i>		JF778738	5	7.4
<i>Trypanosoma corvi</i>		JN006854	4	5.9
<i>Trypanosoma culicaviium</i>		HQ107967	16	23.5
<i>Trypanosoma</i> sp.		JN006850	1	1.5

The dataset includes only one record per individual for birds sampled more than once. The prevalence of each parasite taxon is presented as the percentage of individuals harbouring infection (either as a single or as a multiple infection) with a given taxon.

Table 2. The association between host's sex and age and the probability of infection with parasites from the *Haemoproteus* and *Trypanosoma* genera

	Predictor	Coefficient	s.e.	z	P
<i>Haemoproteus</i>	Sex	1.19	0.71	1.67	0.094
	Age	-0.75	0.63	-1.19	0.236
<i>Trypanosoma</i>	Sex	1.12	0.57	1.98	0.048
	Age	0.07	0.54	0.13	0.897

Data were analysed with generalized linear mixed models (GLMM) with sex and age as categorical explanatory variables and study year as a random variable (effect not shown). Non-significant interactions were removed from the model. Significant terms ($P < 0.05$) are shown in bold.

Table 3. Prevalence and diversity of haemosporidian parasites in four molecularly screened breeding populations of the pied flycatcher: Bahrdorf and Lingen in Germany (Wiersch *et al.* 2007), the island of Öland in Sweden (Kulma *et al.* 2013) and Sękocin Stary in Poland (present study)

Population	N	N _{inf}	Prevalence		Species richness (<i>Plasmodium</i> , <i>Haemoproteus</i>)	Shannon index (asymptotic 95% CIs)
			<i>Plasmodium</i>	<i>Haemoproteus</i>		
Bahrdorf	72	5	5 (6.9%)	0	4 (4, 0)	1.332 (0.50–1.33)
Lingen	147	10	8 (5.4%)	2 (1.4%)	6 (5, 1)	1.696 (0.94–1.75)
Öland	353 ^a	216 ^b	21 (5.9%)	191 (54.1%)	14 (10, 4)	1.473 (1.29–1.58)
Sękocin Stary	68	59 ^c	7 (10.3%)	53 (77.9%)	6 (4, 2)	0.643 (0.31–0.92)

N and N_{inf} refer to the number of screened samples/individuals and the number of samples/individuals positive for the presence of a parasite.

^aIncludes multiple samples from the same individuals, the total number of screened individuals: 319.

^bIncludes four individuals with unidentified infection.

^cIncludes two individuals with mixed infections: hPFC1/pSGS1 and hPFC1/hPHSIB1.

Discussion

The incidence of infection with haemosporidians and trypanosomes in a population of the pied flycatcher located in central Poland is one of the highest reported so far for this species. Specifically, we show that the prevalence of infection with *Haemoproteus* and *Plasmodium* parasites reaches 86.8%, while infection with *Trypanosoma* – 39.7%. Previous studies in populations of this species screened either molecularly or using blood smears showed the prevalence range between 6.9 and 61.2% in the case of haemosporidians (Bennett *et al.* 1994; Dale *et al.*

1996; Siikamäki *et al.* 1997; Sanz *et al.* 2001a; Wiersch *et al.* 2007; Kulma *et al.* 2013) and between 1.1 and 40.6% in the case of trypanosomes (Bennett *et al.* 1994; Dale *et al.* 1996; Siikamäki *et al.* 1997; Sanz *et al.* 2001a). The detected haemosporidian prevalence is also much higher than the prevalence (estimated based on molecular screening) reported in two closely related species – the collared (*Ficedula albicollis*) and the semi-collared flycatchers (*Ficedula semitorquata*) (Szöllösi *et al.* 2008; Kulma *et al.* 2013; Peev *et al.* 2016).

The comparison of four pied flycatcher populations, which were screened molecularly for the presence of haemosporidian




Fig. 2. Phylogenetic relationships of 23 parasite lineages detected in pied flycatchers (8 of *Haemoproteus* spp. and 15 of *Plasmodium* spp.) from the three analysed breeding populations [Lingen (Germany), Öland (Sweden) and Sękocin Stary (Poland)]; indicated with colour boxes) and previously found in this species (birds caught on breeding grounds either during or outside of the breeding season) extracted from the MalAvi database. In the case of the Lingen population one lineage was excluded from the analysis because, based on available information, it was not possible to match it with the lineage in the MalAvi database. The phylogenetic tree was constructed from partial cytochrome *b* sequences (479 bp) and rooted with *Leucocytozoon* spp. lineage SISKIN2. Nodal support values near branches indicate Bayesian posterior probabilities.

infections, showed some geographic differences both in the infection rate and the diversity of the haemosporidian community. Based on the overall prevalence and frequency of infection with *Plasmodium* and *Haemoproteus* parasites, four populations can be clustered into two groups: one containing both German populations (Wiersch *et al.* 2007), and the other containing the populations from Öland, Sweden (Kulma *et al.* 2013) and Sękocin Stary, Poland (present study). Populations from Bährdorf and Lingen were characterized by a low prevalence (6.9%) and the dominance of *Plasmodium* infections, while populations from Öland and central Poland by a high prevalence (61.2–86.8%) and the dominance of *Haemoproteus* lineages. It has to be noted, however, that *Haemoproteus* infection rates recorded in two German populations, may be underestimated as a consequence of the applied protocol. In contrast to Kulma *et al.* (2013) and our study, which assessed infection status based solely on Waldenström *et al.* (2004) protocol, Wiersch *et al.* (2007) additionally carried out an initial screening and chose for the subsequent analysis only positive samples. Since the primer set used in this step, targeting 18 SSU rRNA gene fragment, presumably detects mostly *Plasmodium* specimens (Hulier *et al.* 1996), it may, in general, fail at amplification of the majority of *Haemoproteus* parasites. Because this specific primer set was only very occasionally used in avian haemosporidian studies (Ribeiro *et al.* 2005; Wiersch *et al.* 2007), it is very difficult to reliably assess the success rate of *Haemoproteus* detection with its use. Although Öland and Sękocin Stary populations are similar in some aspects, they differ in some indices of parasite diversity. For example, the Öland population is characterized by higher species richness (the total number of detected lineages) and ecological diversity (as described by the Shannon index). In the case of the *Haemoproteus* community a noticeable difference between these two populations is a fairly common occurrence of two lineages on Öland (PFC1 – 42.0%, PHSIB1 – 34.4%) and an almost complete dominance of one lineage in Sękocin Stary (PFC1 – 96.3%). The higher lineage richness reported for Öland may be at least partly associated with the much larger sample size, since the number of detected lineages increases with the number of screened individuals (Neto *et al.* 2015). In general, the geographic variation of parasite infection rates and diversity indicates that pied flycatchers from different breeding populations may be exposed to different parasite-mediated selection pressures.

Observed differences in the composition of haemosporidian communities as well as in the incidence of infection in geographically distant populations of the same host species might result from various ecological, climatic and population factors, which affect haemosporidian distribution directly or indirectly by shaping the abundance and diversity of vectors (e.g. Lachish *et al.* 2011; Krama *et al.* 2015). It is currently not possible to assess to what extent vectors drive the observed patterns because the distribution of vectors, both on wintering and breeding grounds, is poorly understood. That is especially true for *Haemoproteus* vectors.

The current study is one of the few molecularly characterizing the incidence and richness of trypanosomes in populations of a specific avian host, as opposed to avian guilds (e.g. Soares *et al.* 2016; Valkiūnas *et al.* 2016). The infection rate with trypanosomes in the population from central Poland is over three times higher than the estimates in most pied flycatcher populations screened up to date. Specifically, the incidence of infection varied between 1.1% in south-eastern Sweden, 12–16.2% in central Finland, 15% in south-eastern Norway, 16% in central Spain and 40.6% in northern Finland (Bennett *et al.* 1994; Dale *et al.* 1996; Siikamäki *et al.* 1997; Sanz *et al.* 2001a). Because the aforementioned prevalences were derived from blood smear screening, they may be underestimated in comparison with PCR-based

diagnostics, which have been shown to be generally more sensitive in detecting *Trypanosoma* (Sehgal *et al.* 2001; Nagata, 2006, but see Valkiūnas *et al.* 2016). Moreover, actual prevalences, regardless of the technique used (smear- or PCR-based diagnostics using blood), may be underestimated because in infected birds *Trypanosoma* parasites are abundant in the bone marrow, while they usually occur at very low densities or may be absent in the peripheral blood (Apanius, 1991).

Trypanosome infections in the study population were dominated by *T. culicavium* – a recently described novel species of avian trypanosomes, which is transmitted by mosquitoes (subgenera *Culex* and *Barraudius*) (Votýpka *et al.* 2012). Votýpka *et al.* (2012), based on experimentally identified transmission mode (ingestion of infected vectors) and detection of the parasite in two individuals of the collared flycatcher, suggested that insectivorous songbirds are natural vertebrate hosts of this species. By demonstrating the detection of this trypanosome in another insectivorous bird species, our study confirms this statement. Information about the diet of adult pied flycatchers is very limited, but available data indicate that they ingest Diptera, although at a much lower frequency than other prey (mainly Hymenoptera and Coleoptera) (Bibby and Green, 1980; Silverin and Andersson, 1984). So far, *T. culicavium* has only been detected in European mosquitoes (Votýpka *et al.* 2012; Svobodová *et al.* 2015a). Among the three other detected trypanosome species, *T. corvi* [vector: hippoboscids flies (Mungomba *et al.* 1989; Votýpka *et al.* 2004)] has been shown to be transmitted in Europe (Zidková *et al.* 2012), *T. avium* [vector: blackflies (Bennett, 1961; Votýpka and Svobodová, 2004)] – both in Europe and Africa (Svobodová *et al.* 2015b; Valkiūnas *et al.* 2016), while *T. bennetti* (vectors currently unknown) – in Europe (Zidková *et al.* 2012). However, since information about the vectors and occurrence of trypanosome species in birds is currently very limited, the areas of transmission of these parasites are not definite.

The incidence of infection with *Haemoproteus* and *Trypanosoma* in the study population was unrelated. In birds, the correlated occurrence of these parasites has been found in some [a positive association in white-winged crossbills (*Loxia leucoptera*)] (Deviche *et al.* 2010), but not in other studies (Oakgrove *et al.* 2014; Svobodová *et al.* 2015a). In general, a non-independent incidence of infection may occur if, e.g. infection with one parasite causes immunosuppression allowing the other parasite to successfully invade and develop in the host (positive association) or if infection with one parasite prevents the development of the other parasite *via* cross-immunity (negative association) (Pedersen and Fenton, 2007).

We also looked at each parasite genus separately to search for any relationship between infection incidence and two host traits: age and sex. Despite the fact that susceptibility to parasitic infection is known to increase with age as new infections accumulate during life (Deviche *et al.* 2001; Podmokła *et al.* 2014), we did not find such associations. In the case of the host sex, we found that males were more prone to be infected with *Trypanosoma* than females, while there was no difference between sexes in the probability of infection with *Haemoproteus*. Sex differences in parasite prevalence are generally being attributed to an unequal exposure of the sexes to vectors (van Oers *et al.* 2010; Egizi *et al.* 2014) and differences between males and females in the immune-endocrine system (Schuurs and Verheul, 1990).

In conclusion, by using the molecular approach we showed that the pied flycatcher population from central Poland exhibits one of the highest infection rates with haemoparasites across the breeding range of this species. The comparison of molecularly screened breeding populations revealed some geographic differences in haemosporidian communities, which indicates that pied flycatchers from different populations may experience

diverse parasite-mediated selection pressures. However, to be able to confidently make such a conclusion, more thorough study is needed, including of populations with balanced and synchronized sampling, conditions which were not met by the current study. The latter condition means that samples should be collected in the same years as well as at the same point in a breeding period, since both factors are known to be associated with changes in the prevalence and/or composition of the parasite community (annual variation: Bensch and Åkesson, 2003; Bensch *et al.* 2007; Knowles *et al.* 2011, but see Pagenkopp *et al.* 2008; variation within a season: Szöllösi *et al.* 2016, Dubiec *et al.* 2017).

Acknowledgments. We thank Bartosz Matuszczak for help in the field, Przemysław Chylarecki for statistical advice and Barbara Przybylska for improving the manuscript's English. The study conforms to the legal requirements of Poland.

Financial support. This work was supported by the Polish Ministry of Science and Higher Education/National Science Centre to TDM [grant number N304 345139].

References

- Apanius V (1991) Avian trypanosomes as models of hemoflagellate evolution. *Parasitology Today* **7**, 87–90.
- Baker JR (1976) Biology of the trypanosomes of birds. In Lumsden WHR and Evans DA (eds). *Biology of the Kinetoplastida*, London: Academic Press, pp. 131–174.
- Bates D, Mächler M, Bolker B and Walker S (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1–48.
- Bennett GF (1961) On the specificity and transmission of some avian trypanosomes. *Canadian Journal of Zoology* **39**, 17–33.
- Bennett GF, Siikamäki P, Rätti O, Allander K, Gustafsson L and Squires-Parsons D (1994) Trypanosomes of some Fennoscandian birds. *Memórias do Instituto Oswaldo Cruz* **89**, 531–537.
- Bensch S and Åkesson S (2003) Temporal and spatial variation of hematozoans in Scandinavian willow warblers. *Journal of Parasitology* **89**, 388–391.
- Bensch S, Pérez-Tris J, Waldenström J and Hellgren O (2004) Linkage between nuclear and mitochondrial DNA sequences in avian malaria parasites: multiple cases of cryptic speciation? *Evolution* **58**, 1617–1621.
- Bensch S, Waldenström J, Jonzén N, Westerdahl H, Hansson B, Sejberg D and Hasselquist D (2007) Temporal dynamics and diversity of avian malaria parasites in a single host species. *Journal of Animal Ecology* **76**, 112–122.
- Bensch S, Hellgren O and Pérez-Tris J (2009) Malavi: a public database of malaria parasites and related haemosporidians in avian hosts based on mitochondrial cytochrome b lineages. *Molecular Ecology Resources* **9**, 1353–1358.
- Bibby CJ and Green RE (1980) Foraging behaviour of migrant pied flycatchers, *Ficedula hypoleuca*, on temporary territories. *The Journal of Animal Ecology* **49**, 507–521.
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH and White J-SS (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution* **24**, 127–135.
- Bruford M, Hanotte O, Brookfield J and Burke T (1998) Multilocus and single-locus DNA fingerprinting. In Hoelzel A (ed.). *Molecular Genetic Analysis of Populations: A Practical Approach*, Oxford, UK: IRL Press, pp. 287–336.
- Canty A and Ripley B (2015) Boot: Bootstrap R (S-Plus) Functions. *R package version 1.3–15*.
- Clark P, Boardman W and Raidal S (2009) *Atlas of Clinical Avian Hematology*. Oxford: Wiley-Blackwell.
- Cosgrove CL, Wood MJ, Day KP and Sheldon BC (2008) Seasonal variation in Plasmodium prevalence in a population of blue tits *Cyanistes caeruleus*. *Journal of Animal Ecology* **77**, 540–548.
- Cramp S and Perrins CM (1993) *Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Volume VII Flycatchers to Shrikes*. Oxford, UK: Oxford University Press.
- Dale S, Kruszewicz A and Slagsvold T (1996) Effects of blood parasites on sexual and natural selection in the pied flycatcher. *Journal of Zoology* **238**, 373–393.
- Darriba D, Taboada GL, Doallo R and Posada D (2012) Jmodeltest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**, 772–772.
- Desser SS, McIver SB and Jez D (1975) Observations on the role of simuliids and culicids in the transmission of avian and anuran trypanosomes. *International Journal for Parasitology* **5**, 507–509.
- Deviche P, Greiner EC and Manteca X (2001) Seasonal and age-related changes in blood parasite prevalence in dark-eyed juncos (*Junco hyemalis*, Aves, Passeriformes). *Journal of Experimental Zoology* **289**, 456–466.
- Deviche P, Fokidis HB, Lerbour B and Greiner E (2010) Blood parasitaemia in a high latitude flexible breeder, the white-winged crossbill, *Loxia leucop-tera*: contribution of seasonal relapse versus new inoculations. *Parasitology* **137**, 261–273.
- Dubiec A and Mazgajski TD (2013) Nest mass variation over the nesting cycle in the pied flycatcher (*Ficedula hypoleuca*). *Avian Biology Research* **6**, 127–132.
- Dubiec A, Podmokła E and Gustafsson L (2017) Intra-individual changes in haemosporidian infections over the nesting period in great tit females. *Parasitology Research* **116**, 2385–2392.
- Dufva R (1996) Blood parasites, health, reproductive success, and egg volume in female great tits *Parus major*. *Journal of Avian Biology* **27**, 83–87.
- Dyrce A, Wink M, Kruszewicz A and Leisler B (2005) Male reproductive success is correlated with blood parasite levels and body condition in the promiscuous aquatic warbler (*Acrocephalus paludicola*). *The Auk* **122**, 558–565.
- Egizi AM, Farajollahi A and Fonseca DM (2014) Diverse host feeding on nesting birds may limit early-season West Nile virus amplification. *Vector-Borne and Zoonotic Diseases* **14**, 447–453.
- Fallon SM, Ricklefs RE, Swanson BL and Bermingham E (2003) Detecting avian malaria: an improved polymerase chain reaction diagnostic. *Journal of Parasitology* **89**, 1044–1047.
- Ferrer ES, García-Navas V, Sanz JJ and Ortego J (2012) Molecular characterization of avian malaria parasites in three Mediterranean blue tit (*Cyanistes caeruleus*) populations. *Parasitology Research* **111**, 2137–2142.
- Freeman-Gallant CR, O'Connor KD and Breuer ME (2001) Sexual selection and the geography of *Plasmodium* infection in Savannah sparrows (*Passerculus sandwichensis*). *Oecologia* **127**, 517–521.
- Garamszegi LZ (2010) The sensitivity of microscopy and PCR-based detection methods affecting estimates of prevalence of blood parasites in birds. *Journal of Parasitology* **96**, 1197–1203.
- Gardener M (2014) *Community Ecology: Analytical Methods Using R and Excel*. Exeter: Pelagic Publishing.
- Griffiths R, Double MC, Orr K and Dawson RJG (1998) A DNA test to sex most birds. *Molecular Ecology* **7**, 1071–1075.
- Haldane JBS (1949) Disease and evolution. *La Ricerca Scientifica Supplemento* **19**, 1–11.
- Hall TA (1999) Bioedit: a user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucleic Acids Symposium Series* **41**, 95–98.
- Hellgren O, Waldenström J and Bensch S (2004) A new PCR assay for simultaneous studies of Leucocytozoon, Plasmodium, and Haemoproteus from avian blood. *Journal of Parasitology* **90**, 797–802.
- Hulier E, Pétrou P, Snounou G, Nivez M-P, Miltgen F, Mazier D and Rénia L (1996) A method for the quantitative assessment of malaria parasite development in organs of the mammalian host. *Molecular and Biochemical Parasitology* **77**, 127–135.
- Knowles SCL, Palinauskas V and Sheldon BC (2010) Chronic malaria infections increase family inequalities and reduce parental fitness: experimental evidence from a wild bird population. *Journal of Evolutionary Biology* **23**, 557–569.
- Knowles SCL, Wood MJ, Alves R, Wilkin TA, Bensch S and Sheldon BC (2011) Molecular epidemiology of malaria prevalence and parasitaemia in a wild bird population. *Molecular Ecology* **20**, 1062–1076.
- Krama T, Krams R, Cirule D, Moore FR, Rantala MJ and Krams IA (2015) Intensity of haemosporidian infection of parids positively correlates with proximity to water bodies, but negatively with host survival. *Journal of Ornithology* **156**, 1075–1084.
- Kulma K, Low M, Bensch S and Qvarnström A (2013) Malaria infections reinforce competitive asymmetry between two Ficedula flycatchers in a recent contact zone. *Molecular Ecology* **22**, 4591–4601.
- Laaksonen T, Sirkkiä PM, Calhim S, Brommer JE, Leskinen PK, Primmer CR, Adamík P, Artemyev AV, Belskii E, Both C, Bureš S, Burgess MD, Doligez B, Forsman JT, Grinkov V, Hoffmann U, Ivankina E, Král M, Krams I, Lampe HM, Moreno J, Mägi M,

- Nord A, Potti J, Ravussin P-A and Sokolov L (2015) Sympatric divergence and clinal variation in multiple coloration traits of *Ficedula* flycatchers. *Journal of Evolutionary Biology* **28**, 779–790.
- Lachish S, Knowles SCL, Alves R, Wood MJ and Sheldon BC (2011) Fitness effects of endemic malaria infections in a wild bird population: the importance of ecological structure. *Journal of Animal Ecology* **80**, 1196–1206.
- Martínez-de la Puente J, Merino S, Tomas G, Moreno J, Morales J, Lobato E, García-Fraile S and Belda EJ (2010) The blood parasite *Haemoproteus* reduces survival in a wild bird: a medication experiment. *Biology Letters* **6**, 663–665.
- McCurdy DG, Shutler D, Mullie A and Forbes MR (1998) Sex-biased parasitism of avian hosts: relations to blood parasite taxon and mating system. *Oikos* **82**, 303–312.
- Merino S, Moreno J, Jose Sanz J and Arriero E (2000) Are avian blood parasites pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). *Proceedings of the Royal Society B: Biological Sciences* **267**, 2507–2510.
- Molyneux DH (1977) Vector relationships in the Trypanosomatidae. *Advances in Parasitology* **15**, 1–82.
- Mungomba LM, Molyneux DH and Wallbanks KR (1989) Host-parasite relationship of *Trypanosoma corvi* in *Ornithomyia avicularia*. *Parasitology Research* **75**, 167–174.
- Nagata H (2006) Reevaluation of the prevalence of blood parasites in Japanese passerines by using PCR based molecular diagnostics. *Ornithological Science* **5**, 105–112.
- Neto JM, Pérez-Rodríguez A, Haase M, Flade M and Bensch S (2015) Prevalence and diversity of plasmodium and haemoproteus parasites in the globally-threatened aquatic warbler *Acrocephalus paludicola*. *Parasitology* **142**, 1183–1189.
- Nilsson E, Taubert H, Hellgren O, Huang X, Palinauskas V, Markovets MY, Valkiūnas G and Bensch S (2016) Multiple cryptic species of sympatric generalists within the avian blood parasite *Haemoproteus majoris*. *Journal of Evolutionary Biology* **29**, 1812–1826.
- Oakgrove KS, Harrigan RJ, Loiseau C, Guers S, Seppi B and Sehgal RNM (2014) Distribution, diversity and drivers of blood-borne parasite co-infections in Alaskan bird populations. *International Journal for Parasitology* **44**, 717–727.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH and Wagner H (2016) *Vegan: Community Ecology Package. R package version 2.3–5*.
- Ouwehand J, Ahola MP, Aulsems ANMA, Bridge ES, Burgess M, Hahn S, Hewson CM, Klaassen RHG, Laaksonen T, Lampe HM, Velmala W and Both C (2016) Light-level geolocators reveal migratory connectivity in European populations of pied flycatchers *Ficedula hypoleuca*. *Journal of Avian Biology* **47**, 69–83.
- Pagenkopp KM, Klicka J, Durrant KL, Garvin JC and Fleischer RC (2008) Geographic variation in malarial parasite lineages in the common yellowthroat (*Geothlypis trichas*). *Conservation Genetics* **9**, 1577–1588.
- Pedersen AB and Fenton A (2007) Emphasizing the ecology in parasite community ecology. *Trends in Ecology & Evolution* **22**, 133–139.
- Peev S, Zehindjiev P, Ilieva M, Träff J, Briedis M and Adamík P (2016) Haemosporidian blood parasite diversity and prevalence in the semi-collared flycatcher (*Ficedula semitorquata*) from the eastern Balkans. *Parasitology International* **65**, 613–617.
- Podmokla E, Dubiec A, Drobnik SM, Arct A, Gustafsson L and Cichoń M (2014) Determinants of prevalence and intensity of infection with malaria parasites in the blue tit. *Journal of Ornithology* **155**, 721–727.
- Poulin R (2003) The decay of similarity with geographical distance in parasite communities of vertebrate hosts. *Journal of Biogeography* **30**, 1609–1615.
- R Development Core Team (2014) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rätti O, Dufva R and Alatalo RV (1993) Blood parasites and male fitness in the pied flycatcher. *Oecologia* **96**, 410–414.
- Ribeiro SE, Sebaio F, Branquinho FCS, Marini MÂ, Vago AR and Braga ÉM (2005) Avian malaria in Brazilian passerine birds: parasitism detected by nested PCR using DNA from stained blood smears. *Parasitology* **130**, 261–267.
- Ronquist F and Huelsenbeck JP (2003) MrBayes 3: bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Sanz JJ, Arriero E, Moreno J and Merino S (2001a) Female hematozoan infection reduces hatching success but not fledging success in pied flycatchers *Ficedula hypoleuca*. *The Auk* **118**, 750–755.
- Sanz JJ, Arriero E, Moreno J and Merino S (2001b) Interactions between hemoparasite status and female age in the primary reproductive output of pied flycatchers. *Oecologia* **126**, 339–344.
- Schuurs AHWM and Verheul HAM (1990) Effects of gender and sex steroids on the immune response. *Journal of Steroid Biochemistry* **35**, 157–172.
- Sehgal RNM, Jones HI and Smith TB (2001) Host specificity and incidence of *Trypanosoma* in some African rainforest birds: a molecular approach. *Molecular Ecology* **10**, 2319–2327.
- Siikamäki P, Rätti O, Hovi M and Bennett GF (1997) Association between haematozoan infections and reproduction in the pied flycatcher. *Functional Ecology* **11**, 176–183.
- Silverin B and Andersson G (1984) Food composition of adult and nestling pied flycatchers, *Ficedula hypoleuca*, during the breeding period. *Var Fagelvarld* **43**, 517–524.
- Soares L, Ellis VA and Ricklefs RE (2016) Co-infections of haemosporidian and trypanosome parasites in a North American songbird. *Parasitology* **143**, 1930–1938.
- Sol D, Jovani R and Torres J (2000) Geographical variation in blood parasites in feral pigeons: the role of vectors. *Ecography* **23**, 307–314.
- Spurgin LG, Illera JC, Padilla DP and Richardson DS (2012) Biogeographical patterns and co-occurrence of pathogenic infection across island populations of Berthelot's pipit (*Anthus berthelotii*). *Oecologia* **168**, 691–701.
- Svensson L (1994) *Identification Guide to European Passerines*. Stockholm: Svensson.
- Svoboda A, Marthinsen G, Pavel V, Chutný B, Turčoková L, Lifjeld JT and Johnsen A (2015) Blood parasite prevalence in the Bluethroat is associated with subspecies and breeding habitat. *Journal of Ornithology* **156**, 371–380.
- Svobodová M, Weidinger K, Peške L, Volf P, Votýpka J and Voříšek P (2015a) Trypanosomes and haemosporidia in the buzzard (*Buteo buteo*) and sparrowhawk (*Accipiter nisus*): factors affecting the prevalence of parasites. *Parasitology Research* **114**, 551–560.
- Svobodová M, Volf P and Votýpka J (2015b) Trypanosomatids in ornithophilic bloodsucking Diptera. *Medical and Veterinary Entomology* **29**, 444–447.
- Swanson BL, Lyons AC and Bouzat JL (2014) Distribution, prevalence and host specificity of avian malaria parasites across the breeding range of the migratory lark sparrow (*Chondestes grammacus*). *Genetica* **142**, 235–249.
- Szöllösi E, Hellgren O and Hasselquist D (2008) A cautionary note on the use of nested PCR for parasite screening – An example from avian blood parasites. *Journal of Parasitology* **94**, 562–564.
- Szöllösi E, Cichoń M, Eens M, Hasselquist D, Kempnaers B, Merino S, Nilsson J-Å, Rosivall B, Rytönen S, Török J, Wood MJ and Gáramszegi LZ (2011) Determinants of distribution and prevalence of avian malaria in blue tit populations across Europe: separating host and parasite effects. *Journal of Evolutionary Biology* **24**, 2014–2024.
- Szöllösi E, Gáramszegi LZ, Hegyi G, Laci M, Rosivall B and Török J (2016) Haemoproteus infection status of collared flycatcher males changes within a breeding season. *Parasitology Research* **115**, 4663–4672.
- Valkiūnas G, Jezhova TA and Sehgal RNM (2016) Deforestation does not affect the prevalence of a common trypanosome in African birds. *Acta Tropica* **162**, 222–228.
- van Oers K, Richardson DS, Sæther SA and Komdeur J (2010) Reduced blood parasite prevalence with age in the Seychelles warbler: selective mortality or suppression of infection? *Journal of Ornithology* **151**, 69–77.
- Votýpka J, Szabová J, Rádrová J, Zídková L and Svobodová M (2012) *Trypanosoma culicavium* sp. nov., an avian trypanosome transmitted by *Culex* mosquitoes. *International Journal of Systematic and Evolutionary Microbiology* **62**, 745–754.
- Votýpka J and Svobodová M (2004) *Trypanosoma avium*: experimental transmission from black flies to canaries. *Parasitology Research* **92**, 147–151.
- Votýpka J, Lukeš J and Oborník M (2004) Phylogenetic relationship of *Trypanosoma corvi* with other avian Trypanosomes. *Acta Protozoologica* **43**, 225–231.
- Waldenström J, Bensch S, Hasselquist D and Östman Ö (2004) A new nested polymerase chain reaction method very efficient in detecting Plasmodium and Haemoproteus infections from avian blood. *Journal of Parasitology* **90**, 191–194.
- Wiersch SC, Lubjuhn T, Maier WA and Kampen H (2007) Haemosporidian infection in passerine birds from lower Saxony. *Journal of Ornithology* **148**, 17–24.
- Zídková L, Cepicka I, Szabová J and Svobodová M (2012) Biodiversity of avian trypanosomes. *Infection, Genetics and Evolution* **12**, 102–112.