

**Transition-Metal-Substituted Diphosphenes. 29.<sup>1</sup> Cycloaddition Reactions of the Diphosphenyl Complex  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe-P=P-Mes}^*$  ( $\text{Mes}^* = 2,4,6\text{-}t\text{Bu}_3\text{C}_6\text{H}_2$ ) with Hexafluoroacetone. X-ray Structure Analyses of  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})\text{Fe-P(=PMes}^*)\text{OC}(\text{CF}_3)_2\text{C}(\text{O})$  and  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe-P-P}(\text{Mes}^*)\text{OC}(\text{CF}_3)_2$**

Lothar Weber,<sup>\*,†</sup> Susanne Buchwald,<sup>†</sup> Dieter Lentz,<sup>‡</sup> Dagmar Preugschat,<sup>‡</sup>  
Hans-Georg Stammer,<sup>†</sup> and Beate Neumann<sup>†</sup>


Fakultät für Chemie, Universität Bielefeld, Universitätsstrasse 25, 4800 Bielefeld 1, Germany,  
and Institut für Anorganische und Allgemeine Chemie der Freien Universität Berlin,  
Fabeckstrasse 34-38, 1000 Berlin 33, Germany

Received April 1, 1992

**Summary:** The diphosphenyl complex  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe-P=P-Mes}^*$  ( $\text{Mes}^* = 2,4,6\text{-}t\text{Bu}_3\text{C}_6\text{H}_2$ ) undergoes a [3 + 2] dipolar cycloaddition with hexafluoroacetone to give the metalla heterocycle  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})\text{Fe-P(=PMes}^*)\text{OC}(\text{CF}_3)_2\text{C}(\text{O})$  with a remarkably short Fe-P bond (2.084 (4) Å) and an exocyclic P=P bond. When stored in solution at -40 °C, this complex partly rearranges to the metalated 1-oxa-2,3-diphosphetane  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe-P-P}(\text{Mes}^*)\text{OC}(\text{CF}_3)_2$ . The molecular structures of both isomers were elucidated by single-crystal X-ray analyses.

Electron-releasing organometallic complex fragments as substituents markedly enhance the nucleophilicity of a diphosphene. In molecules such as  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})(\text{L})\text{M-P=P-R}$  ( $\text{M} = \text{Fe}$ ,  $\text{L} = \text{CO}$ ,  $\text{R} = 2,4,6\text{-}t\text{Bu}_3\text{C}_6\text{H}_2$  ( $\text{Mes}^*$ ),  $2,4,6\text{-}(\text{CF}_3)_3\text{C}_6\text{H}_2$ ,  $2,6\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3$ ,  $\text{C}(\text{SiMe}_3)_3$ ;  $\text{M} = \text{Ru}$ ,  $\text{Os}$ ,  $\text{L} = \text{CO}$ ,  $\text{R} = \text{Mes}^*$ ;  $\text{M} = \text{Mn}$ ,  $\text{Re}$ ,  $\text{L} = \text{NO}$ ,  $\text{R} = \text{Mes}^*$ ) the HOMO is mainly represented by the lone pair at the metalated phosphorus atom. This is documented by a number of reactions of  $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe-P=P-Mes}^*$  (1) with organic and organometallic electrophiles. Compound 1 is converted into oxaphosphenolene I by treatment with acrolein, methacrolein, and methyl vinyl ether.<sup>2</sup> Azodicarboxylates and azodiamides give rise to the formation of oxadiazaphosphenolene II,<sup>3</sup> whereas the reaction of 1 with 1,2,4-triazoline-3,5-diones furnished the first 1,2-diaza-3,4-diphosphetidines III.<sup>4</sup> [2 + 2] cycloadditions to 1,2-diphosphetanes IV are observed with fumarodinitrile, maleodinitrile dimethyl fumarate, dimethyl maleate,<sup>5</sup> and maleimides.<sup>6</sup> In all these processes we assume that the ring formation is initiated by the attack of the lone pair of the metalated P atom at the LUMO of the electrophile.


The step from electron-poor alkenes and azo compounds to other electrophiles with heteroatomic double bonds is obvious. Here we report on the chemical behavior of 1 toward anhydrous hexafluoroacetone, which is known to undergo oxidative additions to trivalent phosphorus compounds.<sup>7</sup> Low-coordinated phosphorus systems such as



$\text{R}^1, \text{R}^2 = \text{H}, \text{Me}$ ;  $\text{R}^3 = \text{OEt}, \text{OtBu}, \text{OCH}_2\text{Ph}, \text{NC}_5\text{H}_{10}$ ;  $\text{E} = \text{CN}, \text{CO}_2\text{Me}$

$\text{Aryl} = \text{Ph}, 4\text{-EtOC}_6\text{H}_4$ ;  $[\text{Fe}] = (\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Fe}$

iminophosphanes are converted by hexafluoroacetone to  $\lambda^5$ -oxaphosphiranes V (for  $\text{R} = \text{Me}_3\text{Si}$ )<sup>8</sup> or to  $\lambda^3$ -1-oxa-3-aza-2-phosphetanes VI (for  $\text{R} = \text{alkyl}$ ).<sup>9</sup> A different mode of reaction, however, is encountered with 1 and the ketone.



$\text{R}^1 = \text{Me}_3\text{Si}$  or  $t\text{Bu}$        $\text{R}^1 = t\text{Bu}, \text{R}^2 = i\text{Pr}$   
 $\text{R}^2 = \text{Me}_3\text{Si}$

Condensing an excess of gaseous hexafluoroacetone into

(8) Rösenthaller, G. V.; Sauerbrey, K.; Schmutzler, R. *Chem. Ber.* 1978, 11, 3105.

(9) Niecke, E.; Gudat, D.; Schoeller, W. W.; Rademacher, P. *J. Chem. Soc., Chem. Commun.* 1985, 15, 1050.

<sup>†</sup>Universität Bielefeld.

<sup>‡</sup>Freie Universität Berlin.

(1) Part 28: Weber, L.; Bastian, H.; Boese, R.; Stammer, H.-G.; Neumann, B. *Chem. Ber.*, in press.

(2) Weber, L.; Frebel, M.; Boese, R. *Organometallics* 1989, 8, 1718.

(3) Weber, L.; Bastian, H.; Müller, A.; Bögge, H. *Organometallics* 1991, 10, 2; *Z. Naturforsch., B* 1992, 47, 231.

(4) Weber, L.; Bastian, H.; Boese, R.; Stammer, H.-G. *J. Chem. Soc., Chem. Commun.* 1991, 24, 1778.

(5) Weber, L.; Frebel, M.; Boese, R. *Chem. Ber.* 1990, 123, 733.

(6) Weber, L.; Frebel, M.; Müller, A.; Bögge, H. *Organometallics* 1991, 10, 1130.

(7) Witt, M.; Dhathathreyan, K. S.; Roesky, H. W. *Adv. Inorg. Chem. Radiochem.* 1986, 30, 223 (review).





Figure 1. Molecular structure of 3.

Table I. Selected Bond Lengths (Å) and Angles (deg) for 3

Fe-P(1)	2.084 (4)	P(2)-C(16)	1.860 (13)
Fe-C(11)	1.795 (16)	O(1)-C(11)	1.128 (19)
Fe-C(13)	1.937 (14)	O(2)-C(12)	1.427 (15)
P(1)-P(2)	2.014 (5)	O(3)-C(13)	1.191 (16)
P(1)-O(2)	1.647 (9)	C(12)-C(13)	1.640 (19)
Fe-P(1)-P(2)	140.3 (2)	P(1)-O(2)-C(12)	115.6 (7)
Fe-P(1)-O(2)	109.5 (3)	O(2)-C(12)-C(13)	111.8 (10)
P(2)-P(1)-O(2)	109.9 (3)	Fe-C(13)-C(12)	116.3 (9)
P(1)-P(2)-C(16)	99.0 (4)	P(1)-Fe-C(13)	86.4 (4)

a pentane solution of 1 at  $-196^\circ\text{C}$  and warming to  $20^\circ\text{C}$  afforded the cycloadduct 3 as an orange crystalline solid. No other product could be observed in the reaction mixture by means of  $^{31}\text{P}$  NMR spectroscopy, and no intermediate was detected during the course of the reaction.



The structure of 3 was assigned on the basis of spectral evidence<sup>10</sup> and confirmed by the single-crystal X-ray diffraction study.<sup>11</sup> The  $^{31}\text{P}$  NMR spectrum exhibits two doublets at  $\delta$  483.49 and 180.68 with the large coupling

(10) 3:  $^1\text{H}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  1.33 (s, *p*-*t*Bu), 1.49 [s,  $\text{C}_6(\text{CH}_3)_5$ ], 1.65 (s, *o*-*t*Bu), 1.68 (s, *o*-*t*Bu), 7.56 (m, *m* aryl H);  $^{19}\text{F}$  NMR (84.2 MHz,  $n\text{-C}_4\text{H}_9$ ,  $\text{CFCl}_3$  standard)  $\delta$  -70.55 (q,  $^4J_{\text{FF}} = 7.3$  Hz,  $\text{CF}_3$ ), -71.35 (q,  $^4J_{\text{FF}} = 7.3$  Hz,  $\text{CF}_3$ );  $^{31}\text{P}$  NMR (40 MHz,  $n\text{-C}_4\text{H}_9$ , 85%  $\text{H}_3\text{PO}_4$  standard)  $\delta$  483.49 (d,  $^1J_{\text{PP}} = 633$  Hz, Fe-P), 180.68 (d,  $^1J_{\text{PP}} = 633$  Hz, P-Mes\*).

(11) Crystal data for complex 3: space group  $P2_1/c$ ,  $a = 21.809$  (4) Å,  $b = 10.066$  (2) Å,  $c = 16.250$  (3) Å,  $\beta = 95.87$  (1) $^\circ$ ,  $V = 3560$  (2) Å<sup>3</sup>,  $Z = 4$ ,  $\rho_{\text{calc}} = 1.344$  g cm<sup>-3</sup>, Mo  $K\alpha$  (graphite monochromator,  $\lambda = 0.71073$  Å),  $\omega$ -scan data collection at 183 K ( $3.0 \leq 2\theta \leq 50.0^\circ$ ), 6228 unique reflections, 2383 unique observed reflections ( $F > 4.0\sigma(F)$ ), Siemens  $P2_1$  four-circle diffractometer, structure solved by direct methods and refinement by full-matrix least squares, with use of the Siemens SHELTLX PLUS software on a Micro VAXII computer. All non-hydrogen atoms were refined anisotropically with 232 parameters (hydrogen atoms in calculated positions riding on the corresponding C atoms),  $U(\text{H}) = 0.08$  Å<sup>2</sup>,  $R = 0.106$ ,  $R_w = 0.072$ ,  $w^{-1} = \sigma^2(F)$ , and maximum rest electron density 0.88 e/Å<sup>3</sup>.





Figure 2. Molecular structure of 4.

Table II. Selected Bond Lengths (Å) and Angles (deg) for 4

Fe-P(1)	2.327 (4)	P(1)-C(13)	1.927 (13)
Fe-C(11)	1.779 (15)	P(2)-O(3)	1.722 (8)
Fe-C(12)	1.744 (15)	O(3)-C(13)	1.409 (13)
P(1)-P(2)	2.249 (5)	O(1)-C(11)	1.139 (18)
P(2)-C(16)	1.865 (11)	O(2)-C(12)	1.144 (18)
Fe-P(1)-P(2)	110.7 (2)	O(3)-P(2)-C(16)	96.6 (5)
Fe-P(1)-C(13)	118.5 (4)	P(1)-Fe-C(11)	103.0 (4)
P(1)-P(2)-C(16)	105.9 (4)	P(1)-Fe-C(12)	92.0 (5)
P(1)-C(13)-O(3)	102.2 (8)	C(11)-Fe-C(12)	90.4 (7)
P(2)-O(3)-C(13)	104.2 (7)		

constant  $^1J_{\text{PP}} = 633$  Hz, suggesting the presence of a P=O double bond in 3. In the  $^{19}\text{F}$  NMR spectrum two quartets at  $\delta$  -70.55 and -71.35 ( $^4J_{\text{FF}} = 7.3$  Hz) agree with two magnetically nonequivalent  $\text{CF}_3$  groups with no PF coupling and infer the absence of any direct PC( $\text{CF}_3$ )<sub>2</sub> linkage. The appearance of two discrete singlets for the *o*-*tert*-butyl substituents of the Mes\* ring in the  $^1\text{H}$  NMR spectrum at  $\delta$  1.65 and 1.68 is due to the chiral Fe center in 3. The IR spectrum (Nujol mull) displays only one intense band at  $\nu = 1960$  cm<sup>-1</sup> for the stretching mode of one terminal CO ligand. The acyclic carbonyl group gives rise to a band at 1650 cm<sup>-1</sup> of medium intensity.

The most interesting feature of the molecular structure of 3 (Figure 1 and Table I) is the geometry of the nearly planar five-membered metalla heterocycle (mean deviation from the plane 0.024 Å). The Fe-P bond of 2.084 (4) Å is remarkably short and strongly suggests multiple-bond contributions. Similarly short Fe-P bond distances were reported for complexes 5 (2.117 (2) Å)<sup>12</sup> and 6 (2.112 (1) and 2.202 (1) Å).<sup>13</sup> In 1 the Fe-P contact was determined



(Aryl = 2,6-*t*Bu<sub>2</sub>-4-MeC<sub>6</sub>H<sub>2</sub>)

to be 2.260 (1) Å.<sup>14</sup> The exocyclic P-P bond length (2.014 (5) Å) is comparable to the P-P bond in 1 (2.027 (3) Å). The bond length P(1)-O(2) (1.647 (9) Å) is shorter than the sum of the covalent single-bond radii for P (1.10 Å) and O (0.66 Å).<sup>15</sup> The atoms Fe, P(1), P(2), and O(2) are

(12) Weber, L.; Frebel, M.; Boese, R. *New J. Chem.* 1989, 13, 303.  
(13) Bartlett, R. A.; Dias, H. V. R.; Flynn, K. M.; Olmstead, M. M.; Power, P. P. *J. Am. Chem. Soc.* 1987, 109, 5699.

(14) Weber, L.; Reizig, K.; Bungardt, D.; Boese, R. *Organometallics* 1987, 6, 110.

(15) Pauling, L. *The Nature of the Chemical Bond*; Cornell University Press: Ithaca, NY, 1960; p 224.

located in the same plane (mean deviation from plane 0.023 Å).

Crystals of **3** were grown from pentane solutions at -40 °C in several experiments. In one sample after 4 weeks of crystallization compound **3** decomposed to some extent. The <sup>31</sup>P NMR spectrum displayed a doublet at δ 194.68 (<sup>1</sup>J<sub>PP</sub> = 94 Hz) and a doublet of quartets at δ 136.00 (<sup>1</sup>J<sub>PP</sub> = 94, <sup>3</sup>J<sub>PF</sub> = 26 Hz) in addition to the resonances of **3**. Both compounds were present in the ratio 3:4 = 3:1. The PP coupling constant is consistent with a PP single bond, whereas the size of the PF coupling suggests the presence of a PC(CF<sub>3</sub>)<sub>2</sub> group. Both complexes could not be separated on a preparative scale.

A red crystal of **4** was picked out of the crop and submitted to an X-ray diffraction study (Figure 2 and Table II). The analysis shows the presence of an essentially planar 1-oxa-2,3-diphosphetane (mean deviation from plane 0.01 Å) which is linked to the (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)(CO)<sub>2</sub>Fe fragment via an Fe-P single bond (2.327 (4) Å). Compound **4** is obviously the result of a formal [2 + 2] cycloaddition between **1** and hexafluoroacetone. The P-P bond of 2.249 (5) Å reveals a bond order of unity. In the four-membered ring the oxygen atom of the ketone is added to the arylated P atom (P(2)-O(3) = 1.722 (8) Å). In keeping with this, the (CF<sub>3</sub>)<sub>2</sub>C fragment in **4** is connected

to the metalated phosphorus via a long P-C single bond (P(1)-C(13) = 1.927 (13) Å), which is quite common for diphosphetane derivatives.<sup>5,6</sup>

Due to the longer distance P(1)-P(2) the bond angles at the phosphorus atoms are markedly more acute (C(13)-P(1)-P(2) = 72.3 (4)°, P(1)-P(2)-O(3) = 81.3 (3)°) as compared to the angles at carbon and oxygen (102.2 (9) and 104.2 (7)°, respectively). The *E* configuration of **1** has been maintained throughout the cycloaddition.

At room temperature solutions of **3** and **4** in benzene decompose within 1 week to give **1** and unidentified species. We suggest that the formation of **3** and **4** is initiated by a [2 + 1] cycloaddition to the transient adduct **2**. Subsequent scission of the P-C linkage in **2** and attack of the carbanion at a positively polarized carbon atom of a terminal carbonyl ligand affords metallaheterocycle **3** (path a), whereas P-O bond rupture in **2** and attack of the oxygen at the arylated P atom yields **4**. The dissociation of **3** into **1** and free (CF<sub>3</sub>)<sub>2</sub>CO and a subsequent [2 + 2] cycloaddition to compound **4** cannot be excluded at the moment. The nature of the transformation of **3** into **4** is presently under investigation.

**Acknowledgment.** Our work was generously supported by the Deutsche Forschungsgemeinschaft, Bonn, Germany, the Fonds der Chemischen Industrie, Frankfurt, Germany, and BASF AG, Ludwigshafen, Germany. This is gratefully acknowledged.

**Supplementary Material Available:** Listings giving details of the crystal data, data collection, and structure solution and refinement and tables of atomic coordinates, bond lengths and angles, and thermal parameters for **3** and **4** (17 pages). Ordering information is given on any current masthead page.

OM9201831

(16) Crystal data for complex **4**: space group *C2/c*, *a* = 46.97 (3) Å, *b* = 9.316 (6) Å, *c* = 16.611 (5) Å, β = 101.52 (3)°, *V* = 7121 (6) Å<sup>3</sup>, *Z* = 8, ρ<sub>calcd</sub> = 1.344 g cm<sup>-3</sup>, Mo Kα (graphite monochromator), λ = 0.71073 Å, ω-scan data collection at 179 K (3.0 ≤ 2θ ≤ 45.0°), 4661 unique reflections, 2143 unique observed reflections (*F* > 4.0σ(*F*)), Siemens P2<sub>1</sub> four-circle diffractometer, structure solved by direct methods and refinement by full-matrix least squares, with use of the Siemens SHELXTL PLUS software on a Micro VAXII computer. All non-hydrogen atoms were refined anisotropically with 254 parameters (hydrogen atoms in calculated positions, riding on the corresponding C atoms), *U*(H) = 0.08 Å<sup>2</sup>, *R* = 0.080, *R*<sub>w</sub> = 0.058, *w*<sup>-1</sup> = σ<sup>2</sup>(*F*), maximum rest electron density 0.50 e/Å<sup>3</sup>.

## Platinum-Complex-Catalyzed 1,4-Disilylation of 1,3-Dienes Using Organodisilanes: Remarkable Effect of a Phenyl Functionality on a Silicon Atom

Yasushi Tsuji,\* Rochel M. Lago, Shingo Tomohiro, and Hiroyuki Tsuneishi

Department of Chemistry, Faculty of Engineering, Gifu University, Gifu 501-11, Japan

Received March 20, 1992

**Summary:** Organodisilanes having a phenyl functionality on the silicon atom smoothly react with 1,3-dienes at 130 °C to afford the corresponding 1,4-disilylation adducts in high yields in the presence of platinum catalyst. However, other substituents such as Me, *n*-Bu, *t*-Bu, CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, CH<sub>2</sub>CH=CH<sub>2</sub>, and CH=CH<sub>2</sub> on the silicon atom are totally ineffective.

Activation of a silicon-silicon bond by a transition-metal catalyst<sup>1</sup> and subsequent insertions of 1,3-dienes,<sup>2</sup> acety-

lenes,<sup>3</sup> and olefins<sup>4a,b</sup> into silicon-silicon linkages are of current interest. 1,4-Disilylation of 1,3-dienes is especially important, since it affords adducts<sup>5</sup> having two versatile allylic silane<sup>6</sup> moieties. Transition-metal-catalyzed 1,4-

(3) (a) Ito, Y.; Suginome, M.; Murakami, M. *J. Org. Chem.* 1991, 56, 1948. (b) Yamashita, H.; Catellani, M.; Tanaka, M. *Chem. Lett.* 1991, 241. (c) Watanabe, H.; Kobayashi, M.; Higuchi, K.; Nagai, Y. *J. Organomet. Chem.* 1980, 186, 51. (d) Matsumoto, H.; Matsubara, I.; Kato, T.; Shono, K.; Watanabe, H.; Nagai, Y. *J. Organomet. Chem.* 1980, 199, 43. (e) Watanabe, H.; Kobayashi, M.; Saito, M.; Nagai, Y. *J. Organomet. Chem.* 1981, 216, 149. (f) Tamao, K.; Hayashi, T.; Kumada, M. *J. Organomet. Chem.* 1976, 114, C19. (g) Sakurai, H.; Kamiyama, Y.; Nakadaira, Y. *J. Am. Chem. Soc.* 1975, 97, 931.

(4) (a) Murakami, M.; Anderson, P. G.; Suginome, M.; Ito, Y. *J. Am. Chem. Soc.* 1991, 113, 3987. (b) Hayashi, T.; Kobayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, M. *Organometallics* 1990, 9, 280. (c) Hayashi, T.; Kawamoto, A. M.; Kobayashi, T.; Tanaka, M. *J. Chem. Soc., Chem. Commun.* 1990, 563.

(5) (a) Hiyama, T.; Obayashi, M.; Mori, I.; Nozaki, H. *J. Org. Chem.* 1983, 48, 912. (b) Richter, W. J.; Neugebauer, B. *Synthesis* 1985, 1059. (c) Dunogues, J.; Arreguy, B.; Biran, C.; Calas, R.; Piscioti, F. *J. Organomet. Chem.* 1973, 63, 119.

(6) (a) Bassindale, A. R.; Taylor, P. G. In *The Chemistry of Organic Silicon Compounds*; Patai, S.; Rappoport, Z., Eds.; Wiley: Chichester, U.K., 1989; Part 2, Chapter 14, pp 893-963. (b) Colvin, E. W. *Silicon Reagents in Organic Synthesis*; Academic: London, 1988; pp 25-37.

(1) (a) Yamamoto, K.; Hayashi, A.; Suzuki, S.; Tsuji, J. *Organometallics* 1987, 6, 974. (b) Rich, J. D. *Organometallics* 1989, 8, 2609. (c) Rich, J. D. *J. Am. Chem. Soc.* 1989, 111, 5886. (d) Ito, Y.; Matsuura, T.; Murakami, M. *J. Am. Chem. Soc.* 1988, 110, 3692. (e) Kraft, T. E.; Rich, J. D.; McDermott, P. J. *J. Org. Chem.* 1990, 55, 5430.

(2) (a) Tamao, K.; Okazaki, S.; Kumada, M. *J. Organomet. Chem.* 1978, 146, 87. (b) Matsumoto, H.; Shono, K.; Wada, A.; Matsubara, I.; Watanabe, H.; Nagai, Y. *J. Organomet. Chem.* 1980, 199, 185. (c) Okinoshima, H.; Yamamoto, K.; Kumada, M. *J. Am. Chem. Soc.* 1972, 94, 9263. (d) Carlson, C. W.; West, R. *Organometallics* 1983, 2, 1801. (e) Sakurai, H.; Kamiyama, Y.; Nakadaira, Y. *Chem. Lett.* 1975, 887. (f) Sakurai, H.; Eriyama, Y.; Kamiyama, Y.; Nakadaira, Y. *J. Organomet. Chem.* 1984, 264, 229.