

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Solomon, AW; Willis, R; Pavluck, AL; Alemayehu, W; Bakhtiari, A; Bovill, S; Chu, BK; Courtright, P; Dejene, M; Downs, P; Flueckiger, RM; Haddad, D; Hooper, PJ; Kalua, K; Kebede, B; Kello, AB; Macleod, CK; McCullagh, S; Millar, T; Mpyet, C; Ngondi, J; Nwobi, B; Olobio, N; Onyebuchi, U; Rotondo, LA; Sarr, B; Shafi, O; Sokana, O; West, SK; Foster, A; For The Global Trachoma Mapping Project, (2018) Quality Assurance and Quality Control in the Global Trachoma Mapping Project. *The American journal of tropical medicine and hygiene*. ISSN 0002-9637 DOI: <https://doi.org/10.4269/ajtmh.18-0082>

Downloaded from: <http://researchonline.lshtm.ac.uk/4648678/>

DOI: [10.4269/ajtmh.18-0082](https://doi.org/10.4269/ajtmh.18-0082)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

In order to provide our readers with timely access to new content, papers accepted by the American Journal of Tropical Medicine and Hygiene are posted online ahead of print publication. Papers that have been accepted for publication are peer-reviewed and copy edited but do not incorporate all corrections or constitute the final versions that will appear in the Journal. Final, corrected papers will be published online concurrent with the release of the print issue.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

SOLOMON AND OTHERS

GLOBAL TRACHOMA MAPPING PROJECT

Quality Assurance and Quality Control in the Global Trachoma Mapping Project

Anthony W. Solomon,^{1,2*} Rebecca Willis,³ Alexandre L. Pavluck,³ Wondu Alemayehu,^{4,5} Ana Bakhtiari,³ Sarah Bovill,⁶ Brian K. Chu,³ Paul Courtright,⁷ Michael Dejene,⁸ Philip Downs,⁶ Rebecca M. Flueckiger,³ Danny Haddad,^{9,10} P. J. Hooper,³ Khumbo Kalua,¹¹ Biruck Kebede,¹² Amir Bedri Kello,^{7,13} Colin K. Macleod,⁶ Siobhain McCullagh,⁶ Tom Millar,⁶ Caleb Mpyet,^{7,14,15} Jeremiah Ngondi,^{16,17} Benjamin Nwobi,¹⁸ Nicholas Olobio,¹⁸ Uwazoeke Onyebuchi,¹⁸ Lisa A. Rotondo,¹⁷ Boubacar Sarr,¹⁹ Oumer Shafi,¹² Oliver Sokana,²⁰ Sheila K. West,²¹ and Allen Foster¹ for the Global Trachoma Mapping Project

¹*Clinical Research Department, London School of Hygiene & Tropical Medicine, London, United Kingdom;* ²*London Centre for Neglected Tropical Disease Research, London, United Kingdom;* ³*Task Force for Global Health, Decatur, Georgia;* ⁴*The Fred Hollows Foundation Ethiopia, Addis Ababa, Ethiopia;* ⁵*Berhan Public Health and Eye Care Consultancy, Addis Adaba, Ethiopia;* ⁶*Sightsavers, Haywards Heath, West Sussex, United Kingdom;* ⁷*Division of Ophthalmology, Kilimanjaro Centre for Community Ophthalmology, University of Cape Town, Cape Town, South Africa;* ⁸*Michael Dejene Public Health Consultancy Services, Addis Ababa, Ethiopia;* ⁹*Emory Eye Center, Atlanta, Georgia;* ¹⁰*Orbis International, New York, New York;* ¹¹*Blantyre Institute for Community Outreach, Blantyre, Malawi;* ¹²*Federal Ministry of Health, Addis Ababa, Ethiopia;* ¹³*Light for the World, Addis Ababa, Ethiopia;* ¹⁴*Sightsavers Nigeria, Kaduna, Nigeria;* ¹⁵*Department of Ophthalmology, Jos University, Jos, Nigeria;* ¹⁶*The Carter Center, Atlanta, Georgia;* ¹⁷*RTI International, Washington, District of Columbia;* ¹⁸*National Trachoma Control Program, Department of Public Health, Federal Ministry of Health, Abuja, Nigeria;* ¹⁹*Ministère de la Santé et de la Prévention Médicale, Dakar, Senegal;* ²⁰*Eyecare Department, Ministry of Health, Honiara, Solomon Islands;* ²¹*Dana Center for Preventive Ophthalmology, The Wilmer Institute, Johns Hopkins School of Medicine, Baltimore, Maryland*

* Address correspondence to Anthony W. Solomon, Department of Control of Neglected Tropical Diseases, World Health Organization, Ave. Appia 20, Geneva 1211, Switzerland. E-mail: solomona@who.int

Abstract.

In collaboration with the health ministries that we serve and other partners, we set out to complete the multiple-country Global Trachoma Mapping Project. To maximize the accuracy and reliability of its outputs, we needed in-built, practical mechanisms for quality assurance and quality control. This article describes how those mechanisms were created and deployed. Using expert opinion, computer simulation, working groups, field trials, progressively accumulated in-project experience, and external evaluations, we developed 1) criteria for where and where not to undertake population-based prevalence surveys for trachoma; 2) three iterations of a standardized training and certification system for field teams; 3) a customized Android phone-based data collection app; 4) comprehensive support systems; and 5) a secure end-to-end pipeline for data upload, storage, cleaning by objective data managers, analysis, health ministry review and approval, and online display. We are now supporting peer-reviewed publication. Our experience shows that it is possible to quality control and quality assure prevalence surveys in such a way as to maximize comparability of prevalence estimates between countries and permit high-speed, high-fidelity data processing and storage, while protecting the interests of health ministries.

INTRODUCTION

Trachoma is the leading infectious cause of blindness.¹ To help direct global elimination of trachoma as a public health problem by 2020,² the Global Trachoma Mapping Project (GTMP) aimed to complete the baseline trachoma map worldwide.³ Technical, scientific, and financial oversight to the GTMP was provided through a complex network of partners with complementary mandates, skills, and capacities, including national governments, academic institutions, and nongovernmental organizations. A true international collaboration,⁴ the GTMP, delivered high-quality⁵ population-based prevalence data on trachoma at unprecedented speed and scale.

Although the singular form of the word “project” is used in its title, the GTMP was actually a series of 55 trachoma mapping projects, each of which mapped between one^{6–8} and 91⁹ evaluation units (EUs) for trachoma. A project covered the trachoma mapping needs of a whole country, or of a regional state (Ethiopia) or state (Nigeria). In some projects, a phased approach was used, initially mapping a small number of EUs in which the likelihood of trachoma being a public health problem was felt to be the greatest, on the basis that mapping might be extended if prevalence was found to be high and not extended if it was not. Individual projects were owned and operated by health ministries or the local equivalent.^{10,11} Each EU was mapped using a population-based prevalence survey powered to be 95% confident of detecting an expected 10% prevalence of the sign “trachomatous inflammation–follicular”¹² in 1- to 9-year olds, with absolute precision of 3% and a design effect of 2.65.¹⁰

The template methodology has been described in detail elsewhere.¹⁰ The present article documents the steps that were taken in each constituent project, and at global level, to adhere to the tenets of that template and to try to maximize the accuracy and application of the output. In the spirit of full disclosure, it also lists quality assurance and quality control measures that we did not take, either because doing so would have been too expensive or impractical or because the prompt to do so came with experience. Some measures in the latter category have been introduced for baseline, impact, and surveillance trachoma prevalence surveys supported by Tropical Data (www.tropicaldata.org),^{13,14} following the end of the GTMP.

METHODS

Expert opinion, distilled through a series of teleconferences of the GTMP’s Methodologies and Prioritization Working Groups,¹⁰ was used to develop criteria for where to map and where not to map. We used computer simulation to confirm that population-based prevalence surveys were needed for mapping,¹⁵ rather than a quicker and epidemiologically dirtier approach. We held meetings and teleconferences of each of the four Working Groups (Methodologies, Prioritization, Tools, and Training), and convened the GTMP Advisory Committee to oversee development of pilot systems that were then trialed in the field in Oromia, Ethiopia, in October 2012.¹⁰ The training system, electronic data collection app and field methodologies were all subsequently refined and enhanced as a result of this experience.

The GTMP was formally launched on December 17, 2012, and supported trachoma prevalence survey fieldwork until January 19, 2016, operating in a total of 29 countries. It acquired and processed data on 625,541 households and 2,667,457 examined people. Improvements were progressively introduced during rollout, with identification of issues that warranted improvement facilitated by weekly teleconferences of the core GTMP team, 10 periodic meetings of the Advisory Committee, and formal midterm and end-of-project evaluations by (different) external consultants.

RESULTS

The criteria used for determining where to map and where not to map are given in the Panel. The original field team training manual¹⁶ was superseded by two revisions: version 2 from May 24, 2013,¹⁷ and version 3 from August 15, 2014,¹⁸ with projects beginning after those dates using the updated versions. Quality assurance and quality control points used in the GTMP's systems and methodologies are presented in a series of tables, covering issues relating to scope of mapping (Table 1); survey methodology (Supplemental Table 1), planning, budgeting, and logistics (Supplemental Table 2); training (Supplemental Table 3); survey implementation and field support (Supplemental Table 4); data entry (Supplemental Table 5); data management (Supplemental Table 6); data storage (Supplemental Table 7); and interpretation, reporting, and application of results (Supplemental Table 8). Common to all of these system details was a sequence of development through expertise and experience, consensus building, process design, operationalization, feedback, and follow-up. Measures were put in place through a combination of first-principles thinking (e.g., Supplemental Table 3, row 1), critical review of our own and others' previous work (e.g., Supplemental Table 1, row 4), and progressively accumulated experience (e.g., Supplemental Table 2, row 2)

PANEL

Criteria for where to map and where not to map used by the GTMP, 2012–2016.

Where to map:
• where, on the basis of historical data on trachoma in that district, current data on trachoma in adjacent districts, socioeconomic conditions, and access to water and sanitation, the population is very likely to be trachoma endemic; or
• where trichiasis surgery is being performed by local health-care providers; or
• where individuals with trichiasis are presenting to local health-care providers; or
• where individuals with trichiasis are being identified as part of community outreach campaigns
Where not to map:
• where there is no justification to believe trachoma might be endemic, based on the previously discussed data; or
• where epidemiologically valid prevalence data collected within the last 10 years are already available; or
• where undertaking mapping might put field teams at a security risk; or
• where the responsible authorities, following in-depth discussions, do not prioritize elimination of trachoma as a public health problem

DISCUSSION

“An expert,” Niels Bohr is reported to have said, “is a person who has found out by his own painful experience all the mistakes that one can make in a very narrow field.”³² In that sense, we regard ourselves as approaching expert status in the conduct of population-based prevalence surveys in developing countries. The “painful experience” part of our journeys to this point means that this article was not written to give its authors an opportunity to claim epidemiological superiority over those who have designed, supervised, participated in, or paid for population-based trachoma surveys conducted outside the GTMP. During the course of our careers, we have scattered survey design flaws over the trachoma-endemic globe; we have tried to document those mistakes here. Within the GTMP, we still did not achieve perfection, having had to balance our desire to achieve it with the knowledge that doing so would have reduced efficiency. As a particular example, we are aware that GTMP field teams often failed to enumerate residents who were eligible to be examined but did not participate,³³ despite the fact that our system facilitated it and our field team training system specified doing so. We think that the team members were anxious that they would face supervisors' criticism if they achieved much less than 100% enrolment, which is a training and communication issue that we tried (and continue to try) to address; we believed that pausing

fieldwork to alter enumeration habits would not have been productive. It should also be noted that a proportion of the potential problems that we list in the table as things that we attempted to avoid or correct in the GTMP are actually ghosts-of-problems-future that we have not necessarily yet encountered in real life. However, recognizing and confronting both previous failures and near-misses is important, and this article is an attempt to comprehensively catalog both the errors that we prevented or detected using the systems and methodologies of the GTMP and those that we continue to make and will try to eliminate, where possible, in the next phase^{13,14} of population-based data collection to guide trachoma elimination. If our experience can be used to help others strengthen the design and execution of their community-based surveys at the same time, it will be a double win.

Field-based surveys are complex undertakings, with many moving parts. We set out to generate a whole-of-process system with as few visible joins as possible, supporting survey implementation from the point of determining whether a survey was justified, through to interpreting health ministry-approved data and applying those data for the purposes of improving public health. In such a system, an error in the design or execution of one part of the process can have catastrophic effects on the project as a whole. Before the launch of the GTMP, therefore, we attempted to ensure that all phases of the implementation process had been planned to the fullest possible extent, with decisions made for one phase complementary to decisions made for the others; this article in part demonstrates the fruit of those efforts. As a high-profile endeavor within the neglected tropical diseases sector, within which there are many competing priorities, failure of the GTMP's systems to work as promised might have had reputational consequences for progress against trachoma internationally.

In that context, implementation of a purely electronic data pathway from collection through to display and application carried some risk, both in terms of risk of failure of a system built specifically to serve the GTMP, and in terms of the challenge of convincing scores of health ministries and other partner organizations to simultaneously jump with us from paper to silicon. An occasional objection raised was that without paper forms, we would not have the “original record” and would, therefore, be unable to investigate apparent problems in the data; this objection ignores the fact that irremediable errors are also made when recording data on paper, including many (such as skipped fields, out-of-range values, and illegible handwriting) that our app prevented by design. We believe that our recorders' error rate (estimated on the basis of the data on trichiasis in children—all reports of which were verified [Supplemental Table 4, row 3]—at 1.4 errors per 10,000 keystrokes) compares favorably with previously published data on error rates of data entry operators. Rabbitt found that when individuals were asked to electronically record answers to a question with two possible responses (an analogy from our surveys would be, “Is there trichiasis in the right eye?”), the observed error rate was six per 1,000.³⁴ An outstanding question is whether estimates of trichomatous trichiasis prevalence in adults should be automatically corrected downward to account for the inevitability of these occasional errors, on the basis that when recording the presence or absence of a rare event, erroneous entries are considerably more likely to bias prevalence estimates upward than downward.

The aforementioned question may leave the impression that we felt that we engaged in a high-stakes game by setting up to complete the GTMP and choosing electronic data capture. We would, therefore, be remiss if we failed to acknowledge that (other than in terms of scale and standardization) the GTMP was the setting for an evolution rather than a revolution in trachoma surveys. Our collective efforts outlined here owe much to others.^{24,35–40} We think we have built on that previous work by making electronic data capture the emerging standard for neglected tropical disease epidemiology, by highlighting the need for certification of

clinical examination accuracy in field surveys, by emphasizing data quality, and by the measures that we have undertaken to ensure local ownership.^{5,11}

Supporting health ministries to fulfill their mandate to lead and encouraging appropriate contributions and buy-in from all relevant stakeholders are extremely important issues in their own right.¹¹ They are also an important step to quality assure future prevalence surveys (which will be required to assess the impact of interventions on progress toward elimination⁴¹) because increasing local capacity creates more equal partnerships that will be primed to work together on robust survey designs in the next round.

We are open to constructive criticism from and future collaboration with others and look forward to continuing to adapt and improve as we work toward a world in which surveys to estimate the prevalence of trachoma eventually become unnecessary.⁴²

Received January 30, 2018.

Accepted for publication April 27, 2018.

Note: Supplemental tables appear at www.ajtmh.org.

Acknowledgments:

The Global Trachoma Mapping Project was supported by a grant from the United Kingdom's Department for International Development (DFID; ARIES: 203145) to Sightsavers, which led a consortium of nongovernmental organizations and academic institutions to support ministries of health to complete baseline trachoma mapping worldwide; the United States Agency for International Development (USAID), through the ENVISION project implemented by RTI International under cooperative agreement number AID-OAA-A-11-00048; and the END in Asia project implemented by FHI360 under cooperative agreement number OAA-A-10-00051.

Financial support: A committee established in March 2012 to examine issues surrounding completion of global trachoma mapping was initially funded by a grant from Pfizer to the International Trachoma Initiative. A. W. S. was a Wellcome Trust Intermediate Clinical Fellow (098521) at the London School of Hygiene & Tropical Medicine. L. A. R. reports grants from USAID, during the conduct of the study. O. S. reports grants from United Kingdom Department for International Development (DFID; ARIES: 203145) during the conduct of the study. A. W. S. reports grants from United Kingdom Department for International Development (DFID; ARIES: 203145), grants from United States Agency for International Development (AID-OAA-A-11-00048), grants from United States Agency for International Development (OAA-A-10-00051), and grants from Wellcome Trust (098521), during the conduct of the study.

Disclaimer: The content of this manuscript is solely the responsibility of the named authors and does not necessarily represent the decisions, policy, or views of the institutions with which they are affiliated or of the agencies that funded the work. None of the funders had any role in project design, in project implementation or analysis, or interpretation of data, in the decisions on where, how, or when to publish in the peer-reviewed press, or in preparation of the manuscript.

The Global Trachoma Mapping Project Investigators are Agatha Aboe, Liknaw Adamu, Wondu Alemayehu, Menbere Alemu, Neal D. E. Alexander, Ana Bakhtiari, Berhanu Bero, Sarah Bovill, Simon J. Brooker, Simon Bush, Brian K. Chu, Paul Courtright, Michael Dejene, Paul M. Emerson, Rebecca M. Flueckiger, Allen Foster, Solomon Gadisa, Katherine Gass, Teshome Gebre, Zelalem Habtamu, Danny Haddad, Erik Harvey, Dominic Haslam, Khumbo Kalua, Amir B. Kello, Jonathan D. King, Richard Le Mesurier, Susan Lewallen, Thomas M. Lietman, Chad MacArthur, Colin Macleod, Silvio P. Mariotti, Anna Massey, Els Mathieu, Siobhain McCullagh, Addis Mekasha, Tom Millar, Caleb Mpyet, Beatriz Muñoz, Jeremiah Ngondi, Stephanie Ogden, Alex Pavluck, Joseph Pearce, Serge Resnikoff, Virginia Sarah, Boubacar Sarr, Alemayehu Sisay, Jennifer L. Smith, Anthony W. Solomon, Jo Thomson, Sheila K. West, and Rebecca Willis. Authors' addresses: Anthony W. Solomon, Control of Neglected Tropical Diseases, World Health Organization, Geneva, Switzerland, E-mail: solomona@who.int. Rebecca Willis, Alexandre L. Pavluck, Ana Bakhtiari, Brian K. Chu, and P. J. Hooper, Task Force for Global Health, Decatur, GA, E-mails: rwillis@taskforce.org, apavluck@gmail.com, abakhtiari@taskforce.org, bchu@taskforce.org, and phooper@taskforce.org. Wondu Alemayehu, The Fred Hollows Foundation, Addis Ababa, Ethiopia, and Berhan Public Health and Eye Care Consultancy, Addis Ababa, Ethiopia, E-mail: walemayehu@yahoo.com. Sarah Bovill, Philip Downs, Colin K. Macleod, Siobhain McCullagh, and Tom Millar, Sightsavers, Haywards Heath, West Sussex, United Kingdom, E-mails: sbovill@sightsavers.org, pdowns@sightsavers.org, cmacleod@sightsavers.org, smccullagh@sightsavers.org, and tmillar@sightsavers.org. Paul Courtright, Division of Ophthalmology, Kilimanjaro Centre for Community Ophthalmology, University of Cape Town, Cape Town, South Africa, E-mail: pcourtright@kcco.net. Michael Dejene, Michael Dejene Public Health Consultancy Services, Addis Ababa, Ethiopia, E-mail: michael.dejene@gmail.com. Rebecca M. Flueckiger, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom, and Task Force for Global Health, Decatur, GA, E-mail:

r.m.flueckiger@gmail.com. Danny Haddad, Emory Eye Center, Emory University, Atlanta, GA, E-mail: dhaddad@emory.edu. Khumbo Kalua, Blantyre Institute for Community Outreach, Blantyre, Malawi, E-mail: director@bicomalawi.org. Biruck Kebede, Federal Ministry of Health, Addis Ababa, Ethiopia, E-mail: biruck1@gmail.com. Amir Bedri Kello, Light for the World, Addis Ababa, Ethiopia, and Division of Ophthalmology, Kilimanjaro Centre for Community Ophthalmology, University of Cape Town, Cape Town, South Africa, E-mail: amirbedrikello@gmail.com. Caleb Mpyet, Sightsavers Nigeria, Kaduna, Nigeria, Department of Ophthalmology, Jos University, Jos, Nigeria, and Division of Ophthalmology, Kilimanjaro Centre for Community Ophthalmology, University of Cape Town, Cape Town, South Africa, E-mail: mpyetc@yahoo.com. Jeremiah Ngondi, The Carter Center, Atlanta, GA, and RTI International, Washington, DC, E-mail: jn250@cam.ac.uk. Benjamin Nwobi, Nicholas Olobio, and Uwazoeke Onyebuchi, National Trachoma Control Program, Department of Public Health, Federal Ministry of Health, Abuja, Nigeria, E-mails: emekanwobi@gmail.com, olobio@yahoo.com, and happywaez@yahoo.com. Lisa A. Rotondo, RTI International, Washington, DC, E-mail: lrotondo@rti.org. Boubacar Sarr, Ministère de la Santé et de la Prévention Médicale, Dakar, Senegal, E-mail: bouksarr@yahoo.fr. Oumer Shafi, Federal Ministry of Health, Disease Control, Addis Ababa, Ethiopia, E-mail: oumerjaji@gmail.com. Oliver Sokana, Eyecare Department, Ministry of Health, Honiara, Solomon Islands, E-mail: osakana@moh.gov.sb. Sheila K. West, Dana Center for Preventive Ophthalmology, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, MD, E-mail: shwest@jhmi.edu. Allen Foster, Clinical Research Department, London School of Hygiene & Tropical Medicine, London, United Kingdom, E-mail: allen.foster@Lshmt.ac.uk.

REFERENCES

- <jrn>1. Bourne RR et al.; Vision Loss Expert Group, 2013. Causes of vision loss worldwide, 1990–2010: a systematic analysis. *Lancet Glob Health* 1: e339–e349.</jrn>
- <bok>2. World Health Organization, 2012. *Accelerating Work to Overcome the Global Impact of Neglected Tropical Diseases: A Roadmap for Intervention*. Geneva, Switzerland: WHO.</bok>
- <jrn>3. Solomon AW, Kurylo E, 2014. The global trachoma mapping project. *Community Eye Health* 27: 18.</jrn>
- <bok>4. Rosenberg ML, Hayes ES, McIntyre MH, Neil N, 2010. *Real Collaboration: What it Takes for Global Health to Succeed*. Berkeley, CA: University of California Press.</bok>
- <jrn>5. Engels D, 2016. The global trachoma mapping project: a catalyst for progress against neglected tropical diseases. *Ophthalmic Epidemiol* 23: 1–2.</jrn>
- <jrn>6. Taleo F et al.; Global Trachoma Mapping Project, 2017. Integrated mapping of yaws and trachoma in the five northern-most Provinces of Vanuatu. *PLoS Negl Trop Dis* 11: e0005267.</jrn>
- <jrn>7. Cama A et al.; Global Trachoma Mapping Project, 2017. Prevalence of signs of trachoma, ocular *Chlamydia trachomatis* infection and antibodies to Pgp3 in residents of Kiritimati Island, Kiribati. *PLoS Negl Trop Dis* 11: e0005863.</jrn>
- <jrn>8. Macleod CK et al., 2016. Low prevalence of ocular *Chlamydia trachomatis* infection and active trachoma in the western division of Fiji. *PLoS Negl Trop Dis* 10: e0004798.</jrn>
- <jrn>9. Abdala M et al., 2017. The epidemiology of trachoma in Mozambique: results of 96 population-based prevalence surveys. *Ophthalmic Epidemiol* ■■■: 1–10; Epub ahead of print.</jrn>
- <jrn>10. Solomon AW et al., 2015. The global trachoma mapping project: methodology of a 34-country population-based study. *Ophthalmic Epidemiol* 22: 214–225.</jrn>
- <jrn>11. Heggen AE, Solomon AW, Courtright P, 2016. Perspectives of national coordinators and partners on the work of the global trachoma mapping project. *Ophthalmic Epidemiol* 23: 366–372.</jrn>

- <jrn>12. Thylefors B, Dawson CR, Jones BR, West SK, Taylor HR, 1987. A simple system for the assessment of trachoma and its complications. *Bull World Health Organ* 65: 477–483.</jrn>
- <jrn>13. Hooper PJ, Millar T, Rotondo LA, Solomon AW, 2016. Tropical data: a new service for generating high quality epidemiological data. *Community Eye Health J* 29: 38.</jrn>
- <unknown>14. World Health Organization, 2016. *Tropical Data: A WHO-Led Initiative to Help National Programmes Collect and Do More with Their Data*. Available at: http://www.who.int/trachoma/news/News_Trachoma_Tropical_Data_launch/en/. Accessed July 21, 2016.</unknown>
- <jrn>15. Smith JL, Sturrock HJ, Olives C, Solomon AW, Brooker SJ, 2013. Comparing the performance of cluster random sampling and integrated threshold mapping for targeting trachoma control, using computer simulation. *PLoS Negl Trop Dis* 7: e2389.</jrn>
- <bok>16. Courtright P, Gass K, Lewallen S, MacArthur C, Pavluck A, Solomon A, West S, 2012. *Global Trachoma Mapping Project: Training for Mapping of Trachoma*. London, United Kingdom: International Coalition for Trachoma Control. Available at: <http://www.trachomacoalition.org/node/358>.</bok>
- <bok>17. Courtright P, Gass K, Lewallen S, MacArthur C, Pavluck A, Solomon A, West S, 2013. *Global Trachoma Mapping Project: Training for Mapping of Trachoma (Version 2)*. London, United Kingdom: International Coalition for Trachoma Control. Available at: <http://www.trachomacoalition.org/node/357>.</bok>
- <bok>18. Courtright P, Gass K, Lewallen S, MacArthur C, Pavluck A, Solomon A, West S, 2014. *Global Trachoma Mapping Project: Training for Mapping of Trachoma (Version 3)*. London, United Kingdom: International Coalition for Trachoma Control. Available at: <http://www.trachomacoalition.org/node/122>.</bok>
- <jrn>19. Ukety OT, 1993. Trachoma in northeastern Zaire: a study of 22 cases. *Ann Soc Belg Med Trop* 73: 61–66.</jrn>
- <jrn>20. Kilangalanga J et al.; Global Trachoma Mapping Project, 2017. Trachoma in the Democratic Republic of the Congo: results of 46 baseline prevalence surveys conducted with the global trachoma mapping project. *Ophthalmic Epidemiol* ■■■: 1–9; Epub ahead of print.</jrn>
- <jrn>21. Phiri I et al.; Global Trachoma Mapping Project, 2017. The burden of and risk factors for trachoma in selected districts of Zimbabwe: results of 16 population-based prevalence surveys. *Ophthalmic Epidemiol* ■■■: 1–11; Epub ahead of print.</jrn>
- <jrn>22. Mwingira UJ et al., 2016. Progress of trachoma mapping in Mainland Tanzania: results of baseline surveys from 2012 to 2014. *Ophthalmic Epidemiol* 23: 373–380.</jrn>
- <jrn>23. Ko R, Macleod C, Pahau D, Sokana O, Keys D, Burnett A, Willis R, Wabulembo G, Garap J, Solomon AW, 2016. Population-based trachoma mapping in six evaluation units of Papua New Guinea. *Ophthalmic Epidemiol* 23: 22–31.</jrn>
- <bok>24. Negrel AD, Taylor HR, West S, 2001. *Guidelines for Rapid Assessment for Blinding Trachoma (WHO/PBD/GET/00.8)*. Geneva, Switzerland: World Health Organization.</bok>
- <jrn>25. Miller H, Gallego G, Rodriguez G, 2010. Clinical evidence of trachoma in Colombian Amerindians of the Vaupes Province. *Biomedica* 30: 432–439.</jrn>

- <bok>26. Solomon AW, 2013. *Report on a Field Visit: Trachoma in Colombia*. London, United Kingdom: London School of Hygiene & Tropical Medicine/Global Trachoma Mapping Project.</bok>
- <bok>27. Berhane Y, Worku A, Bejiga A, 2006. *National Survey on Blindness, Low Vision and Trachoma in Ethiopia*. Addis Ababa, Ethiopia: Federal Ministry of Health of Ethiopia.</bok>
- <bok>28. Sub-Saharan African Medical and Research Foundation, 2010. *Prevalence of Trachoma in Five Zones of Oromia Region, Ethiopia*. Addis Ababa, Ethiopia: Oromia Regional Health Bureau.</bok>
- <jrn>29. Bero B et al.; Global Trachoma Mapping Project, 2016. Prevalence of and risk factors for trachoma in Oromia regional state of Ethiopia: results of 79 population-based prevalence surveys conducted with the global trachoma mapping project. *Ophthalmic Epidemiol* 23: 392–405.</jrn>
- <jrn>30. Marks M, Vahi V, Sokana O, Puiahi E, Pavluck A, Zhang Z, Dalipanda T, Bottomley C, Mabey DC, Solomon AW, 2014. Mapping the epidemiology of yaws in the Solomon Islands: a cluster randomized survey. *Am J Trop Med Hyg* 92: 129–133.</jrn>
- <jrn>31. Sokana O, Macleod C, Jack K, Butcher R, Marks M, Willis R, Chu BK, Posala C, Solomon AW, 2016. Mapping trachoma in the Solomon Islands: results of three baseline population-based prevalence surveys conducted with the global trachoma mapping project. *Ophthalmic Epidemiol* 23: 15–21.</jrn>
- <jrn>32. Coughlan R, 1954. Dr Edward Teller's magnificent obsession. *Life* ■■■: 61–74.</jrn>
- <jrn>33. Kalua K, Phiri M, Kumwenda I, Masika M, Pavluck AL, Willis R, Mpyet C, Lewallen S, Courtright P, Solomon AW; Global Trachoma Mapping Project, 2015. Baseline trachoma mapping in Malawi with the global trachoma mapping project (GTMP). *Ophthalmic Epidemiol* 22: 176–183.</jrn>
- <jrn>34. Rabbitt P, 1990. Age, IQ and awareness, and recall of errors. *Ergonomics* 33: 1291–1305.</jrn>
- <conf>35. World Health Organization, 2010. *Report of the 3rd Global Scientific Meeting on Trachoma*. Johns Hopkins University, Baltimore, MA, July 19–20, 2010 (WHO/PBD/2.10). Geneva, Switzerland: WHO.</conf>
- <jrn>36. Pavluck A, Chu B, Mann Flueckiger R, Ottesen E, 2014. Electronic data capture tools for global health programs: evolution of LINKS, an Android-, web-based system. *PLoS Negl Trop Dis* 8: e2654.</jrn>
- <bok>37. Smith PG, Morrow RH, 1996. *Field Trials of Health Interventions in Developing Countries: A Toolbox*, 2nd edition. London, United Kingdom: Macmillan.</bok>
- <bok>38. World Health Organization, 1993. *Primary Health Care Level Management of Trachoma (WHO/PBL/93.33)*. Geneva, Switzerland: WHO.</bok>
- <bok>39. Solomon AW, Zondervan M, Kuper H, Buchan JC, Mabey DCW, Foster A, 2006. *Trachoma Control: A Guide for Programme Managers*. Geneva, Switzerland: WHO.</bok>
- <jrn>40. Ngondi J, Reacher M, Matthews F, Brayne C, Emerson P, 2009. Trachoma survey methods: a literature review. *Bull World Health Organ* 87: 143–151.</jrn>

<bok>41. World Health Organization Strategic and Technical Advisory Group on Neglected Tropical Diseases, 2015. *Technical Consultation on Trachoma Surveillance. September 11–12, 2014, Task Force for Global Health, Decatur, USA (WHO/HTM/NTD/2015.02)*. Geneva, Switzerland: WHO.</bok>

<bok>42. World Health Assembly, 1998. *Global Elimination of Blinding Trachoma*. 51st World Health Assembly, Geneva, 16 May 1998, Resolution WHA51.11. Geneva, Switzerland: World Health Organization.</bok>

TABLE 1

Preemptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues relating to the scope of mapping

No.	The GTMP...	...To reduce the impact of, or avoid...	...Which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	Systematically discussed countries (and administrative divisions within countries) with individuals who had local knowledge, in an effort to uncover available evidence for possible trachoma endemicity, with documentation of evidence, and action where needed	Lack of expressed need to map in areas where mapping is needed	Delay in identification of endemic areas, delay in elimination program initiation, and failure to achieve GET2020	The GTMP systematically discussed the need for trachoma surveys in the Democratic Republic of the Congo with key informants ¹⁹ and the Ministère de la Santé ²⁰
		Lack of expressed need to map in areas where trachoma was historically found but has now disappeared	Continuing uncertainty and repeated reexamination of the same evidence over the need or otherwise to conduct mapping	In 1982, a study of prevalence and causes of blindness and low vision was conducted in eight provinces of Indonesia; trachoma was one of the top 10 causes; by 2013, trachoma had disappeared (unpublished Indonesia Ministry of Health data)
2	(Where evidence to justify mapping was of low quality) undertook mapping using a phased approach	Failure to take into account prevalence estimates in adjacent areas, as they accrued, in decision-making on whether there was a need to map	Excessive use of resources to document the absence of trachoma at baseline, or delay in identification of endemic areas, delay in elimination program initiation, and failure to achieve GET2020	The GTMP phased survey rollout in the Democratic Republic of the Congo, ²⁰ Yemen (manuscript in preparation) and Zimbabwe ²¹
3	(Where evidence to justify mapping was completely absent but suspicion of trachoma existed) provided technical and financial support to undertake preliminary survey work to determine whether baseline population-based prevalence surveys were	Expressed need to map in areas where mapping was not needed	Excessive use of resources to document the absence of trachoma at baseline	The GTMP undertook preliminary survey work in Tanzania to rule out areas unlikely to have trachoma as a public health problem ²²
		Lack of expressed need to map in areas	Delay in identification of endemic areas,	The GTMP undertook preliminary survey

	needed	where mapping is needed	delay in elimination program initiation, and failure to achieve GET2020	work in Papua New Guinea to provide evidence to justify population-based prevalence surveys ²³
4	Used a positive trachoma rapid assessment ²⁴ or other positive preliminary data on the presence of trachoma to initiate support for a population-based prevalence survey in the corresponding administrative area, as soon as possible	Assumption that data from a trachoma rapid assessment provide prevalence estimates	Maximally biased estimate of prevalence potentially used for programmatic decision-making	The GTMP did this for the duration of its operation
5	Initiated contact with health ministries of countries that may have been trachoma endemic (and responded to us on learning about the GTMP), then engaged in discussions to determine whether mapping was needed	Countries being isolated from the international trachoma community	Delay in identification of endemic areas, delay in elimination program initiation, and failure to achieve GET2020	Colombia identified trachoma in communities in the Amazon rainforest, near to the border with Brazil, between 2003 and 2006, ²⁵ but limited international engagement occurred until the GTMP visited in 2013 ²⁶
6	Undertook detailed discussions with health ministries over the benefits and risks associated with using the standardized systems and approaches of the GTMP for trachoma mapping, as opposed to completing trachoma mapping via other means	Incomplete uptake of standardized systems and approaches developed by the GTMP, and/or the incomplete use of funds allocated to the GTMP	Heterogeneity of approaches and/or failure to meet donors' expectations	The GTMP did this for the duration of its operation
7	Channeled financial resources donated by bilateral organizations to undertake baseline trachoma mapping in any country where baseline mapping was justified	Domestic funds available to map insufficient to meet clear needs	Delay in identification of endemic areas, delay in elimination program initiation, and failure to achieve GET2020	A national survey of blindness, low vision, and trachoma in Ethiopia in 2005–2006 ²⁷ showed that trachoma was highly and widely endemic in Oromia, the largest regional state. But by 2012, survey work had been undertaken in only 10 of Oromia's then-current 265 rural districts. ²⁸ The GTMP supported mapping of the rest of the regional state ²⁹
8	Encouraged health ministries to piggyback collection of data on other diseases of local importance, advocated to funders to secure permission to do so, and	Co-endemic diseases with data needs not mapped with baseline trachoma surveys	Lost opportunity for achieving efficiencies in the use of human and financial resources	In two EUs of the Solomon Islands and one EU of Vanuatu, the GTMP collected population-based data on the prevalence of yaws and trachoma at

	provided technical support to adjust fieldwork protocols and data collection tools as needed			the same time ^{6,30,31}
9	Supplemented hour-by-hour communication with weekly formal teleconferences of the core project group, to review progress and plan activities, country by country	Centralization of information and decision-making in the hands of one individual or one partner organization	Lost opportunities to benefit from complimentary experiences and to hear dissenting voices	The GTMP held weekly formal teleconferences for the duration of its operation

GET20202 = global elimination of trachoma as a public health problem by 2020; GTMP = Global Trachoma Mapping Project; EU = evaluation unit.

Quality Assurance and Quality Control in the Global Trachoma Mapping Project

SUPPLEMENTAL TABLE 1

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to survey methodology

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	asked countries to prepare a first draft of the survey protocol	local adoption of a methodology without local understanding of why each of its elements was important	failure to build local capacity.	The GTMP did this in each of its constituent projects.
		imposition of locally inappropriate survey elements if local partners feel inhibited about challenging a template	difficulties in survey implementation.	
2	ensured that the draft survey protocol was consistent with WHO recommendations, working with the health ministry and local partners to refine the draft as needed	international inconsistency	prevalence estimates that could not be compared between settings.	The GTMP did this in each of its constituent projects.
3	assisted countries to frame EUs of appropriate sizes (generally at the level of the local administrative unit for health care management and ideally containing	framing of inappropriately large EUs	potential to miss significant pockets of disease.	In Yobe State, Nigeria, a previous population-based trachoma prevalence survey covered a population of > 2 million people in

	populations of 100,000–250,000 persons ³⁵⁾			a single EU. ⁴³
		framing of inappropriately small EUs	excessive use of resources for mapping, or extrapolation of results from a small EU to provide prevalence estimates for a larger population. ⁴⁴	In the Solomon Islands, the regions of Rennell-Bellona (estimated population 3041) and Temotu (estimated population 21 362) had sufficiently similar socio-economic and environmental characteristics to be combined to form a single EU. ³¹
4	assisted countries to design epidemiologically valid cluster selection methods within EUs	selection of too many clusters	excessive use of resources for mapping.	Due to a misunderstanding, twice the required number of clusters were selected and visited in one GTMP-Mozambique EU that had been formed by combining 2 adjacent districts. We subsequently included explicit discussion of the implications of EU formation on cluster selection in our conversations with health ministries.
		selection of too few clusters	potential to miss significant pockets of disease.	In some trachoma prevalence surveys conducted prior to the GTMP, 7–14 clusters were selected for inclusion. ^{45–47}
		selection of clusters using a biased	generation of inaccurate	All GTMP-supported surveys applied epidemiologically-

		methodology	prevalence estimates.	appropriate cluster selection methods.
5	assisted countries to design epidemiologically valid household selection methods within clusters	selection of too many households	excessive use of resources for mapping, either through enrolment of an excessive total number of households per EU, or through making the workload required in one cluster too great to be reliably completed by one team in a single day.	In 1999, in order to select an appropriate sub-village for a longitudinal study, AWS and his team examined 5527 of 5703 residents (some or all residents of 1099 of 1103 households) of a single trachoma-endemic village in Tanzania, ⁴⁸ for reasons that he is now unable to fully explain. It took him and one other experienced grader nearly 3 months of full-time work.
		selection of too few households	failure to achieve an appropriate sample size, or inefficiency through having field teams unproductive for long periods of each day.	All GTMP-supported surveys involved selection of 25–45 households per selected cluster.
		selection of households using a biased methodology	generation of inaccurate prevalence estimates.	For the first few clusters enrolled in the Democratic Republic of the Congo, teams only visited households that had children; this was noted by our data manager and corrected via a telephone call.
6	selected a fixed number of households per cluster, rather than a fixed number of	perceived pressure to enrol individuals causing coercion of	abrogation of ethical	The GTMP did this in each of its constituent projects, except in Viet

	individuals	cluster residents or biased sampling, particularly towards the end of the day in the field	responsibilities, or selection bias.	Nam. ⁴⁹
7	developed a standard smartphone app for data collection and supplied it for use in all countries, allowing any changes in practice to be quickly incorporated through the app	lag in practice behind policy, which lags behind learning	continued implementation of known flaws.	The GTMP's purpose was to complete baseline trachoma mapping, but it soon received requests to support impact and surveillance surveys ⁴¹ for trachoma, too. At these phases of programme evolution, the prevalence of trichiasis "unknown to the health system" ³⁵ is important. This led rapidly to the incorporation into the standard survey of questions about previous management in eyes recorded as having trichiasis. ¹⁸
8	designed surveys to include collection of global positioning system data from each household enrolled	failure to include independently verifiable geolocation of selected households and clusters	inability to ensure that households and clusters have been appropriately enrolled.	In one constituent project of the GTMP, one team logged all households in 2 clusters at a single location in the national capital, several hundred kilometres from the EU ostensibly being surveyed. Following further investigation on the ground by the health ministry, those data were rejected and the

^a This excludes cases that have already been operated on, for which operations have been refused, or which are already scheduled to receive operations.

				team members redeployed away from the GTMP.
9	enrolled, as standard, all residents aged ≥ 1 year in each household selected for inclusion	failure to examine adults (or examination of adults only if they lived in the same house as 1–9-year-olds ⁵⁰)	inability to estimate a meaningful prevalence (or generation of a potentially biased estimate of the prevalence ⁵⁰) of trachomatous trichiasis.	Within the GTMP, 3 EUs in Chad and 3 EUs in Egypt had to be re-surveyed because field teams only examined 1–9-year-olds; in Cambodia, only households in which 1–9-year-olds lived were enrolled ⁵⁰ ; in Viet Nam, only 1–9-year-olds and ≥ 50 -year-olds were enrolled. ⁴⁹
		failure to examine children ⁴⁷	inability to estimate a meaningful prevalence of trachomatous inflammation—follicular.	The GTMP did this in each of its constituent projects.
		creation of an incentive for household residents very keen to be examined, or very keen not to be examined, to misrepresent their age	bias in prevalence estimates.	In recent trials of a trachomatous trichiasis-only survey methodology, when only those aged ≥ 40 years were examined, unexpectedly large numbers of individuals claiming to be aged 40–45 years were enrolled. ⁵¹
10	supported health ministries to obtain local ethical clearance before surveys started	neglect of locally important ethical considerations in survey design	failure to “take into consideration the laws and regulations of the country or countries in which the research is to be performed as well as applicable international norms	The GTMP did this in each of its constituent projects.

			and standards”, as required by the Declaration of Helsinki. ⁵²	
--	--	--	---	--

SUPPLEMENTAL TABLE 2

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to planning, budgeting and logistics

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	closely reviewed budgets against the agreed methodology and a standard budget template and ensured that the methodology was consistently reflected in the budgeting assumptions ⁵³	essential activities omitted, or excessive resources requested to undertake mapping	mapping activities not aligned with agreed methodology, inefficient use of resources, or shortfall in funding with consequent failure of one or more of the GTMP’s constituent project.	Resources to support field team supervisors added to budgets in several projects. In one country, excessive requests were trimmed, resulting in a budget reduction of 28%.
2	provided Android smartphones for survey teams, with (if possible) survey forms already pre-loaded	possible local purchase of phones with outdated versions of the Android operating system and/or difficulties in ensuring correct software installation	delays in commencing surveys, or reversion to the use of paper-based data collection.	The GTMP tried to do this in each of its constituent projects. Where it did not (because, for example, it was difficult to import phones, or lead times were too short), phone cost was often higher, and internet bandwidth occasionally made it challenging to download survey software.

3	provided binocular, 2.5× magnifying loupes for graders to use	lack of provision of loupes by programmes; or provision of loupes that were uncomfortable to wear, prone to breakage, or of the wrong magnification	failure to use loupes, or use of loupes with the wrong magnification, leading to reduction in diagnostic accuracy.	The GTMP did this in each of its constituent projects.
---	---	---	--	--

SUPPLEMENTAL TABLE 3

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to training

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	provided a standard training system for graders and recorders, ¹⁸ and required individuals to be certified in its use before using it to train others ¹⁰	different training systems, or different interpretation of the same training system, delivered to teams in different projects; and duplication of effort	variable quality training.	Over the course of its 3-year operation, the GTMP used 3 successive versions of a standardized training system. ¹⁰
2	provided all elements of the standard training system in English, French, Arabic, Spanish and Portuguese, ¹⁸ having had the accuracy of translation in each case checked by ophthalmologists with that language as a mother tongue	differential attention to quality control and quality assurance in different settings, depending on language preference	variable quality training.	All 3 iterations of the GTMP's standardized training system were made available in 5 languages. ¹⁰
3	trained local trachoma grader trainers, or (where there were insufficient human resources to meet the local training need) identified and funded GTMP-certified grader trainers from elsewhere to participate in training	human resources available to train local staff insufficient to meet the agreed need	delay in identification of endemic areas, delay in elimination programme initiation, and failure to achieve GET2020.	The GTMP funded a GTMP-certified grader trainer from the Solomon Islands to visit Vanuatu to train and certify local personnel as trachoma graders ⁶ ; there were many other instances of similar collaboration within the GTMP.

4	assisted newly GTMP-certified trainers undertaking their first GTMP training week by providing a fellow trainer with previous experience of delivering GTMP training	trepidation or uncertainty	Inconsistency in field team training between sites.	The GTMP tried to do this in each of its constituent projects; it was not possible in some instances because of travel restrictions.
5	(where there was an absolute shortage of medical and paramedical personnel to train as graders) brokered secondments of GTMP-certified graders from elsewhere to assist with mapping	understaffed field teams	scarce personnel deployed to the field for long periods to complete surveys, or mapping not attempted at all.	The GTMP did this on several occasions; the details are somewhat politically sensitive.
6	insisted on strict adherence to the definitions of signs in the WHO simplified trachoma grading scheme ¹²	often: over-diagnosis, because graders do not want to ignore obvious pathology; this stems from confusion between the role of a grader contributing to a prevalence survey (where definitions must be clear cut), and the role of a clinician caring for an individual (where knowledge of the natural history of disease and factors other than the patient's clinical signs contribute to formulating the management plan)	overestimation of the prevalence of trachoma, potentially leading to inappropriate allocation of resources for trachoma elimination.	In Togo in 2009, graders classified individuals with fewer than 5 follicles as having TF. ⁵⁴
7	undertook all grader training in known trachoma-endemic areas, and incorporated examination in the field as	deployment of graders who may not have previously had an opportunity to examine real subjects with the signs they are	uncertainty about grader competence.	The GTMP supported grader trainees from Lao People's Democratic Republic ⁵⁹ and Cambodia ⁵⁰ to be trained in

	part of the training process	then asked to identify ⁵⁵⁻⁵⁸		Ethiopia, at the invitation of the Oromia Regional Health Bureau.
8	only deployed graders who had demonstrated their trachoma grading competency through formal inter-grader agreement exercises on real subjects, using the assessments of a GTMP-certified grader trainer ¹⁰ as the gold standard	deployment of graders whose competency had been assessed only through grading of slides or photographs of trachoma ^{55-58,60}	uncertainty about grader competence.	The GTMP did this in each constituent project, other than Viet Nam, ⁴⁹ where travel of grader trainees to a more highly endemic country could not be undertaken. Though not allowing previously-experienced graders who did not pass the test to continue was controversial at the beginning, it was subsequently seen as an important demonstration of how important quality was to the GTMP. In some contexts, however, managing disappointed grader trainees became an issue in its own right.
9	set the standard for passing the inter-grader agreement exercise as a kappa of ≥ 0.70 for the presence or absence of the sign “trachomatous inflammation—follicular” in children aged 1–9 years	use of percentage agreement with the grader trainer, ^{43,46,61-63,64} or an unspecified measurement, ^{45,65} as the criterion for diagnostic accuracy	uncertainty about grader competence.	The GTMP did this in each of its constituent projects, other than Viet Nam (see above).
10	made the training system available for countries engaged in impact surveys for trachoma, even if they were conducted	different training systems delivered to teams in different settings; and duplication of effort to create	variable quality of training.	The GTMP training system was made available to India, Mali and Nepal for use in trachoma prevalence surveys that health

	without other involvement of the GTMP	training systems		ministries in those countries had planned to implement independently.
--	---------------------------------------	------------------	--	---

SUPPLEMENTAL TABLE 4

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to survey implementation and field support

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	deployed in-service supervisors, each of whom were required to first pass the formal inter-grader agreement exercises on real subjects, using as the gold standard the assessments of a GTMP-certified grader trainer	lack of appropriate supervision ^b	drift in accuracy of grading over time; errors in application of fieldwork protocol; or unreported social, economic, health or supply issues that could adversely affect field team performance.	The GTMP did this in each of its constituent projects.
2	rapidly reviewed raw data	over-estimation of the mean number of residents per household in available census data	failure to examine a number of individuals in each EU that would permit calculation of prevalence estimates with acceptable precision. ^c	In Southern Nations, Nationalities and Peoples' Region of Ethiopia, ⁶⁶ we requested that 4 clusters be added to an EU because there were too few 1–9-year-olds examined.

^b We intend to further improve the standard and consistency of supervision in trachoma impact and surveillance surveys, through the use of a dedicated training package for supervisors

^c We did not keep a sufficiently close eye on this issue early in project implementation. (In Oromia, for example, there were 2 EUs in which 651 and 653 1–9-year-olds were initially examined, but because of delays in data upload, teams had moved to other zones before this came to light.) Pre-GTMP surveys which stipulated a given number of subjects to be examined per cluster did not run this risk, but instead risked biased selection.

		under-estimation of the mean number of residents per household in available census data	examination of more individuals in each EU than necessary to permit calculation of prevalence estimates with acceptable precision, leading to inefficient use of resources. ^d	In Guinea, where 23 clusters were included per EU, the range of 1–9-year-olds examined per EU was 1113–3137.
3	telephoned field supervisors as soon as a record of trachomatous trichiasis in a child was identified	erroneous recording of the presence of trachomatous trichiasis in a child	potential mobilisation of a paediatric ophthalmologist or oculoplastic surgeon to provide service; or if undetected, tacit encouragement of a lack of concentration in the field.	In raw data from 55 projects, 519 cases of trichiasis were reported amongst 1 146 644 1–9-year-olds; 249 of those cases were confirmed when checked with field teams.
4	discussed and resolved fieldwork problems as they arose	uncertainty, confusion, inconsistency between teams	reductions in the accuracy and/or repeatability of prevalence estimates.	The GTMP did this in each of its constituent projects.

^d We did not do this well enough. Pre-GTMP surveys which stipulated a given number of subjects to be examined per cluster did not run this risk, but instead risked biased selection.

SUPPLEMENTAL TABLE 5

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to data entry

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	undertook all primary data recording in electronic format, using a purpose-built Android smartphone app, LINKS-GTMP ^{10,36}	failure to ensure high-fidelity transfer of data from paper to electronic format for the purposes of data analysis, and/or long delays while paper-based surveys are managed (photocopied, double entered, etc.)	data used to generate prevalence estimates not reflecting the findings observed in the field, and/or long intervals between survey completion and programmatic decision-making.	The GTMP did this in each of its constituent projects.
2	ensured that the LINKS-GTMP app did not permit fields to be skipped ^e	failure to collect or record available data	missing data, and uncertainty in analyses.	The GTMP did this in each of its constituent projects.
3	included “don’t know” and/or “other” options in all multiple choice questions	data recorders being forced to stop, enter junk data or use a parallel reporting system	junk data, loss of system integrity or inability to proceed.	The GTMP did this in each of its constituent projects.
4	used check screens requesting recorders to verify data just entered	lack of flagging of entry of rare outcomes, such as trachomatous	missed opportunities to correct erroneous keystrokes at source.	The GTMP did this in each of its constituent projects.

^e The dangers inherent in allowing skip fields are illustrated by the GPS data collected (for each household) in the GTMP. Because GPS signals are sometimes difficult or impossible to access, LINKS-GTMP allows recorders to proceed to the next question without collecting GPS coordinates, which teams were instructed to press only after the Android had tried to triangulate its location for at least 60s without success. In one project, a group of recorders rapidly developed a habit of always skipping the GPS field. In our next-generation app developed for Tropical Data, skipping GPS data collection is only possible after system-driven timeout.

		trichiasis		
5	restricted responses to sensible ranges – age, for example, could only be recorded as 1–100 years; for those reporting an age at last birthday of > 100 years, 100 years was recorded	errors	potentially uninterpretable data.	The GTMP did this in each of its constituent projects.

SUPPLEMENTAL TABLE 6

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to data management

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	undertook data cleaning centrally (by BKC from 12/2012–06/2013; by RW from 07/2013–07/2015; by RW and AB from 07/2015–01/2016; identical algorithms were used throughout)	inconsistency and/or lack of objectivity in data cleaning	random error or bias.	The GTMP did this in each of its constituent projects.
2	checked that clusters lay within the boundaries of the EU by comparing the mean GPS coordinates of the households enrolled in a cluster with the EU shape-file ^f	inclusion of data from clusters that were inadvertently selected from outside the boundaries of the EU	potentially inaccurate estimation of trachoma prevalence for the EU.	The GTMP did this in each of its constituent projects.

^f Accurate shape-files were often unavailable. In such cases, discussion with local staff was used to resolve uncertainties.

3	adjusted trachomatous inflammation—follicular prevalence data by age of subjects examined ¹⁰	implied failure to recognise that those examined in a house-to-house survey may not necessarily be representative of the underlying population, because of absence, refusal, or illness	bias in estimates of prevalence.	The GTMP did this in each of its constituent projects.
4	adjusted trachomatous trichiasis data by age and gender of subjects examined ¹⁰	implied failure to recognise that those examined in a house-to-house survey may not necessarily be representative of the underlying population, because of absence, refusal, or illness	bias in estimates of prevalence of trachomatous trichiasis, usually leading to overestimation.	The GTMP did this in each of its constituent projects.
5	equally weighted cluster-level outcome proportions when considering overall prevalence	over-emphasis of results on clusters with larger numbers or examined individuals	sampling or participation bias.	The GTMP did this in each of its constituent projects.
6	in projects commencing after 15 August 2014, classified trichiasis as trachomatous trichiasis if and only if the eyelid demonstrated easily visible conjunctival scarring ¹² , or could not be everted	implied failure to recognise that not all trichiasis is due to trachoma	misclassification bias in estimates of prevalence of trachomatous trichiasis, leading to overestimation.	Since adding collection of data on conjunctival scarring, 69% of all trichiasis recorded in GTMP surveys has been trachomatous trichiasis.
7	developed standard data analysis algorithms to be run in R	human error	erroneous output.	The GTMP did this in each of its constituent projects.
8	double-checked all output by eye	human error, or unintended consequences of the use of standard algorithms	erroneous outputs.	The GTMP did this in each of its constituent projects.

9	made the R code used freely available to any interested party (including as Supplementary information to this paper)	higher barrier to replication of analyses	scepticism.	For the R code, please see Supplementary information.
10	used a 2-stage health ministry data approval process and insisted on health ministry ownership of data ¹⁰	lack of acknowledgement of national ownership of data	delay or failure in application of outputs to disease control.	The GTMP did this in each of its constituent projects. (The lack of an analogous process for routinely engaging, at the planning stage, government officials responsible for water, sanitation and hygiene (WASH) occasionally led to tensions over household-level WASH data, which were collected in nearly all constituent projects of the GTMP. ^{67,68})

SUPPLEMENTAL TABLE 7

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to data storage

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	programmed LINKS-GTMP to automatically upload data from micro-SD cards to a Cloud-located database, as soon as a data-enabled mobile phone network or WiFi network was detected	electronic data maintained in only one place, or in an insufficient number of places	need for repeat survey in the event of loss, damage or corruption of storage media.	The GTMP did this in each of its constituent projects.
2	applied 128-bit encryption at the transport layer, ¹⁰ and carried out subsequent data review, cleaning and approval only through a secure website with transport layer security, IP-restricted firewall, and site authentication and authorization, to which access could be gained only through password-protected login.	paper-based or electronic data containing personally identifiable information stored without adequate security	potential inadvertent disclosure of personally identifiable information on survey subjects, violating standard ethical principles for investigators.	The GTMP did this in each of its constituent projects; for Tropical Data, ^{13, 14} 256-bit encryption is being used, including encryption of data at rest.
3	went to extreme lengths to recover data stored on micro-SD cards that were, in the rare case, damaged in the field	loss of data and potential loss of faith in the reliability of data storage on Android smartphones	possible need to repeat surveys, and possible need to design a new system.	The GTMP lost data from 1 cluster (due to a lost Android), of a total of 12,631 clusters visited. Where Androids were damaged or corrupted, all stored data were

				recovered.
--	--	--	--	------------

SUPPLEMENTAL TABLE 8

Pre-emptive and corrective measures put in place by the GTMP to avoid pitfalls inherent in trachoma mapping: issues related to interpretation, reporting and application of results

	The GTMP...	...to reduce the impact of, or avoid...	...which otherwise might have led to...	Examples of instances where this measure helped (or might have helped)
1	built capacity amongst collaborating partners in an effort to ensure that disaggregated data, if examined, were interpreted correctly	assumption that cluster-level data can be used to provide prevalence estimates for the cluster, ⁶⁴ or for a subdivision of the EU ⁴³	decisions made using data not powered to provide the estimates generated.	The GTMP did this in each of its constituent projects.
2	calculated 95% confidence intervals for prevalence estimates by bootstrapping adjusted cluster-level proportions, with replacement, over 10,000 replications	confidence intervals around prevalence estimates not calculated, not calculated using an appropriate methodology, or not reported	assessment of precision of prevalence estimates made difficult.	The GTMP did this in each of its constituent projects.
3	provided epidemiologist support to local authors, as needed, to draft project-specific manuscripts once each project was complete; and brokered agreement to publish the resulting papers, if accepted under the journal's normal criteria, in a series of supplements to a peer-reviewed journal, with financial support from the project to make those	data not published in peer-reviewed journals	lower visibility of data and of local efforts.	There has been considerable interest in the published output of the GTMP, with a number of special issues of <i>Ophthalmic Epidemiology</i> ^{5,81} produced.

	papers open-access ^{8,9,22,23,29,31,50,59,66,69-80}			
4	automatically channelled prevalence category data to the open access website of the Global Atlas of Trachoma, ^{82,83} with explicit health ministry agreement, as soon as they were fully approved by the health ministry	prevalence category data not made widely available	data not accessible for public health decision-making.	The GTMP did this in each of its constituent projects.
5	(if the health ministry explicitly agreed), electronically transferred prevalence data to the International Trachoma Initiative, so that they could be used, where indicated, to justify applications for donated azithromycin for trachoma elimination purposes	manual transfer of data	increased effort, delays, errors.	GTMP data have leveraged a probable donation of 283 283 514 doses of azithromycin from Pfizer (including doses already donated, those approved for donation, and doses projected for future donation based on standard numbers of treatment rounds).

SUPPLEMENTAL REFERENCES

43. Mpyet C, Ogoshi C, Goyol M, 2008. Prevalence of trachoma in Yobe State, north-eastern Nigeria. *Ophthalmic Epidemiol* 15: 303-7.
44. Courtright P, Sheppard J, Schachter J, Said ME, Dawson CR, 1989. Trachoma and blindness in the Nile Delta: current patterns and projections for the future in the rural Egyptian population. *Br J Ophthalmol*. 73: 536-40.
45. Ndayishimiye O et al., 2011. Population-based survey of active trachoma in 11 districts of Burundi. *Ophthalmic Epidemiol* 18: 146-9.
46. Ngondi J, Onsarigo A, Adamu L, Matende I, Baba S, Reacher M, Emerson P, Zingesser J, 2005. The epidemiology of trachoma in Eastern Equatoria and Upper Nile States, southern Sudan. *Bull World Health Organ*. 83: 904-12.
47. Regassa K, Teshome T, 2004. Trachoma among adults in Damot Gale District, South Ethiopia. *Ophthalmic Epidemiol* 11: 9-16.
48. Solomon AW, 2003. Rational use of azithromycin in the control of trachoma: using quantitative PCR to assess distribution of infection and impact of treatment. Department of Infectious and Tropical Diseases. London: University of London, 259.
49. Hiep NX et al., 2017. Trachoma in Viet Nam: results of 11 surveillance surveys conducted with the Global Trachoma Mapping Project. [submitted].
50. Meng N, Seiha D, Thorn P, Willis R, Flueckiger RM, Dejene M, Lewallen S, Courtright P, Solomon AW, Global Trachoma Mapping Project, 2016. Assessment of Trachoma in Cambodia: Trachoma Is Not a Public Health Problem. *Ophthalmic Epidemiol*: 1-5.
51. World Health Organization Strategic and Technical Advisory Group on Neglected Tropical Diseases, 2018. *Design and validation of a trachomatous trichiasis-only survey (WHO/HTM/NTD/PCT/2017.08)*. Geneva: World Health Organization.
52. World Medical Association, 2013. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 310: 2191-4.
53. Trotignon G et al., 2017. The cost of mapping trachoma: Data from the Global Trachoma Mapping Project. *PLoS Negl Trop Dis* 11: e0006023.
54. Centers for Disease Control and Prevention, 2011. *Prevalence assessment of trachoma, Togo*. Atlanta: Centers for Disease Control and Prevention.
55. Noa Noatina B, Kagmeni G, Mengouo MN, MOUNGUI HC, Tarini A, Zhang Y, Bella AL, 2013. Prevalence of trachoma in the Far North region of Cameroon: results of a survey in 27 Health Districts. *PLoS Negl Trop Dis* 7: e2240.
56. Noa Noatina B, Kagmeni G, Souleymanou Y, MOUNGUI HC, Tarini Hien A, Akame J, Zhang Y, Bella AL, 2014. Prevalence of trachoma in the north region of Cameroon: results of a survey in 15 health districts. *PLoS Negl Trop Dis* 8: e2932.
57. Yayemain D, King JD, Debrah O, Emerson PM, Aboe A, Ahorsu F, Wanye S, Ansah MO, Gyapong JO, Hagan M, 2009. Achieving trachoma control in Ghana after implementing the SAFE strategy. *Trans R Soc Trop Med Hyg* 103: 993-1000.
58. Dorkenoo AM, Bronzan RN, Ayena KD, Anthony G, Agbo YM, Sognikin KS, Dogbe KS, Amza A, Sodahlon Y, Mathieu E, 2012. Nationwide integrated mapping of three neglected tropical diseases in Togo: countrywide implementation of a novel approach. *Trop Med Int Health* 17: 896-903.
59. Southisombath K, Sisalermsak S, Chansan P, Akkhavong K, Phommala S, Lewallen S, Courtright P, Solomon AW, Global Trachoma Mapping Project, 2016. National Trachoma Assessment in the Lao People's Democratic Republic in 2013-2014. *Ophthalmic Epidemiol*: 1-7.
60. Kalua K, Chirwa T, Kalilani L, Abbenyi S, Mukaka M, Bailey R, 2010. Prevalence and risk factors for trachoma in central and southern Malawi. *PLoS One* 5: e9067.

61. Jip NF, King JD, Diallo MO, Miri ES, Hamza AT, Ngondi J, Emerson PM, 2008. Blinding trachoma in Katsina state, Nigeria: population-based prevalence survey in ten local government areas. *Ophthalmic Epidemiol* 15: 294-302.
62. Hassan A et al., 2011. The prevalence of blinding trachoma in northern states of Sudan. *PLoS Negl Trop Dis* 5: e1027.
63. Berhane Y et al., 2007. Prevalence of trachoma in Ethiopia. *Ethiop J Health Dev* 21: 211-5.
64. Kur LW et al., 2009. Trachoma in Western Equatoria State, Southern Sudan: implications for national control. *PLoS Negl Trop Dis* 3: e492.
65. Dolin PJ, Faal H, Johnson GJ, Ajewole J, Mohamed AA, Lee PS, 1998. Trachoma in The Gambia. *Br J Ophthalmol*. 82: 930-3.
66. Adera TH et al., 2016. Prevalence of and Risk Factors for Trachoma in Southern Nations, Nationalities, and Peoples' Region, Ethiopia: Results of 40 Population-Based Prevalence Surveys Carried Out with the Global Trachoma Mapping Project. *Ophthalmic Epidemiol* 23: 84-93.
67. Garn JV et al., 2018. Sanitation and water supply coverage thresholds associated with active trachoma: modeling cross-sectional data from 13 countries. *PLoS Negl Trop Dis* 12: e0006110.
68. Boisson S, Engels D, Gordon BA, Medlicott KO, Neira MP, Montresor A, Solomon AW, Velleman Y, 2016. Water, sanitation and hygiene for accelerating and sustaining progress on neglected tropical diseases: a new Global Strategy 2015-20. *Int Health*. 8 Suppl 1: i19-i21.
69. Mpyet C et al., 2016. Trachoma Mapping in Gombe State, Nigeria: Results of 11 Local Government Area Surveys. *Ophthalmic Epidemiol* 23: 406-411.
70. Kalua K, Chisambi A, Chinyanya D, Kamwendo Z, Masika M, Willis R, Flueckiger RM, Pavluck AL, Solomon AW, Global Trachoma Mapping Project, 2016. Completion of Baseline Trachoma Mapping in Malawi: Results of Eight Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic Epidemiol*: 1-7.
71. Mpyet C et al., 2016. Prevalence of Trachoma in Katsina State, Nigeria: Results of 34 District-Level Surveys. *Ophthalmic Epidemiol* 23: 55-62.
72. Omar FJ et al., 2016. Baseline Trachoma Surveys in Kaskazini A and Micheweni Districts of Zanzibar: Results of Two Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic Epidemiol*: 1-6.
73. Mpyet C et al., 2016. Prevalence of Trachoma in Bauchi State, Nigeria: Results of 20 Local Government Area-Level Surveys. *Ophthalmic Epidemiol* 23: 39-45.
74. Elshafie BE et al., 2016. The Epidemiology of Trachoma in Darfur States and Khartoum State, Sudan: Results of 32 Population-Based Prevalence Surveys. *Ophthalmic Epidemiol* 23: 381-391.
75. Adamu Y et al., 2016. Prevalence of Trachoma in Benishangul Gumuz Region, Ethiopia: Results of Seven Population-Based Surveys from the Global Trachoma Mapping Project. *Ophthalmic Epidemiol* 23: 70-76.
76. Sherief ST et al., 2016. The Prevalence of Trachoma in Tigray Region, Northern Ethiopia: Results of 11 Population-Based Prevalence Surveys Completed as Part of the Global Trachoma Mapping Project. *Ophthalmic Epidemiol* 23: 94-99.
77. Muhammad N et al., 2016. Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys. *Ophthalmic Epidemiol* 23: 46-54.
78. Adamu MD et al., 2016. Prevalence of Trachoma in Niger State, North Central Nigeria: Results of 25 Population-Based Prevalence Surveys Carried Out with the Global Trachoma Mapping Project. *Ophthalmic Epidemiol* 23: 63-69.
79. Abashawl A et al., 2016. Prevalence of Trachoma in Gambella Region, Ethiopia: Results of Three Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic Epidemiol* 23: 77-83.

80. Bio AA et al., 2017. Prevalence of trachoma in northern Benin: results from 11 population-based prevalence surveys covering 26 districts. *Ophthalmic Epidemiol* 24: 265-73.
81. Haddad D, Nwobi B, Schmidt E, Courtright P, 2016. Elimination of Trachoma-Knowing Where to Intervene. *Ophthalmic Epidemiol* 23: 345-346.
82. Smith J, Mann R, Haddad D, Polack S, Kurylo E, Brooker S, 2016. *Global Atlas of Trachoma: an open-access resource on the geographical distribution of trachoma* (www.trachomaatlas.org). Atlanta: International Trachoma Initiative.
83. Smith JL, Haddad D, Polack S, Harding-Esch EM, Hooper PJ, Mabey DC, Solomon AW, Brooker S, 2011. Mapping the global distribution of trachoma: why an updated atlas is needed. *PLoS Negl Trop Dis* 5: e973.