

PAPER • OPEN ACCESS

Analysis and comparison of tropospheric scintillation prediction models at Covenant University

To cite this article: S. A. Akinwumi *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **173** 012015

View the [article online](#) for updates and enhancements.

Related content

- [Impact of meteorological parameters over Covenant University, Ota, Nigeria](#)
E O Falayi, M R Usikalu, T V Omotosho *et al.*
- [Dust pollution and its potential health risk exposure to inhabitants of Covenant University and canaanland, Ogun state, Nigeria](#)
Omeje Maxwell, O. Adewoyin Olusegun, S. Joel Emmanuel *et al.*
- [2nd International Conference on Science and Sustainable Development \(ICSSD 2018\)](#)



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Analysis and comparison of tropospheric scintillation prediction models at Covenant University

S. A. Akinwumi¹, T. V. Omotosho¹, M. R. Usikalu¹, T. A. Adagunodo¹, M. O. Adewusi², O. O. Ometan²

¹Department of Physics, College of Science and Technology Covenant University
PMB 1023, Ota, Ogun state, Nigeria

²Department of Physics, Lagos state University, Ojo, Lagos state, Nigeria

oluwasayo.akinwumi@covenantuniversity.edu.ng

Abstract. Knowledge of tropospheric scintillation is an important phenomenon in the design of satellite communication system. One year (January 2015-December 2015) scintillation data extracted from Astra 2E/2F/2G Satellite link measurement installed at Covenant University, Ota (Lat: 6.7 °N, Long: 3.23 °E) southwest Nigeria, at an elevation angle of 59.9° and a frequency of 12.245 GHz was used in this study. The analysis and the result were compared with some reputable scintillation prediction models so as to obtain best performance model for Ota region. From the result, it was discovered that the Karasawa model gives the lowest percentage error rate for both fade and enhancement of about 0.57% at 0.1 percentage of time and 6.93% at 0.01 percentage of time respectively and therefore was best found fit for the prediction of propagation impairment for the region. However, the model should be tested further using higher frequency band such as Ka and V bands to confirm the accuracy of the model. The information provided in this study will help in fade margin for antenna sizing and performance needed for satellite communication link in the region.

Keywords: Tropospheric scintillation, Attenuation prediction, Satellite communication, Ku band, Performance evaluation

1. INTRODUCTION

Tropospheric scintillation occurrences transpires to be one of the key signal impairments that affect earth-satellite path [1-6] in modern earth-space communication systems, most importantly at higher frequency bands (both uplink and downlink). The influence of scintillation on radio wave signal transmission cannot be over highlighted due to its constant variation in phase and amplitude which affect signal power [7-8]. At short microwave or millimeter-wave bands, scintillation intensity increase with decrease in elevation angle, antenna size and with increase in frequencies [9]. Yet, the tropospheric scintillation is an intricate occurrence on earth-space transmission path, which includes the presence of gasses [10], and margin level up to altitudes of about 20 km within the tropical troposphere [11].

These disparities in turn transform the amplitude and phase of the received electric field. In actual fact scintillation intensity is a variable that can be influenced by atmospheric conditions [8], and the subsequent variability in tropospheric scintillation strength has a significant impact on the statistics of the scintillation process [12]. Also, tropospheric scintillation is acknowledged to show a robust correlation with some climatic parameters such as temperature, pressure and humidity [7].

2. METHODS AND DATA ANALYSIS

The tropospheric data site is located at the Covenant University, Ota from Astra 2E/2F/2G beacon satellite 12.245 GHz (Lat: 6.7 °N, Long: 3.23 °E, Elev. Angle: 59.9°) at a sample rate of 1 second. The data for this study were measured from January 2015 to December 2015. However, the non-rainy days were separated from rainy days for the analysis by using Davis automatic weather station and



spectrum analyser at rain rate 0 mm/h observed for non-rainy events, meanwhile the rain rate above 0 mm/h were removed from the corresponding days and time data within the period of observation because the rain rate above 0 mm/h indicate the presence of rain. A monthly average data were used as reference data signal level and was subtracted from daily measured received data signal level in other to obtained the non-rainy attenuation on every one minutes for each clear-sky day [1]. Subsequent filtering procedure result in data which consists of positive (enhancement) and negative (fade) scintillation amplitude that is above or below the mean level.

Also, the ground tropospheric scintillation measured data were compared with some of the existing scintillation prediction models. In this study, the considered models that predict the variance of signal log-amplitude are: ITU-R model [13], Karasawa model [14], Otung model [15], Van de Kamp model [16]. Finally, performance evaluation of each of the tropospheric scintillation was tested based on the fractional percentage error as presented in Eq. (1).

$$\text{Error } (\varepsilon) = \frac{\text{predicted} - \text{measured}}{\text{measured}} \times 100\% \quad (1)$$

Table 1: Percentage error of the models

% of time	ITU-R	Fade			Enhancement		
		Kasarawa	Otung	Van de Kamp	Kasarawa	Otung	Van de Kamp
0.01	55.77558	28.02516	190.9347	-39.1306	6.92731	71.48973	-31.7537
0.1	22.36777	0.570562	45.80179	-54.7107	-10.2135	1.607013	-50.2639
1	-2.97606	-20.2577	-25.8888	-66.6068	-24.1086	-51.4064	-64.6062
10	-33.2918	-51.3273	-56.6163	-78.5266	-44.9347	-98.2782	-77.5358

3. RESULT AND DISCUSSION

Table 1 shows the tropospheric scintillation prediction models with statistical comparison between fade (negative) and enhancement (positive). It can be observed that Karasawa model gave the lowest percentage error for fade of about 0.57% at 0.1% of time and 28.03% at 0.01% of time respectively. The next model to that of Karawasa is ITU-R, which is of about 22.37% at 0.1%. Same was observed for enhancement at 0.01% of time followed by ITU-R. However, at 1 and 10% of time, ITU-R is observed to have the least percentage error. Figure 1 demonstrates the assessment of the ground data in Ota and several models prediction for both fade and enhancement, in order to understand the limits of each predicted model and the degree of validity in Ota region.

For fade signal scintillation amplitude, Karasawa model (0.1%) and ITU-R model (1%) showed a very close agreement with the ground data values in Ota almost for all the percentage of time predicted. This was followed by Otung model, which slightly deviated from other models at 0.01%. The Van de Kamp model variance from other models may be attributed to the scintillation measurement during the presence of heavy clouds and rainfall. However, enhancement signal scintillation amplitude also shows a close relationship between measure data and Karasawa predicted model at all level of percentage of time (most especially at 0.1%). This closeness may be because the model was developed during non-rainy period with strong contribution from water vapour obtain from surface temperature and relative humidity [6]. No result for ITU-R model in enhancement because the model was only designed to yield result for signal fade.




Fig. 1: Comparison of Ota data with Four existing models for fade and enhancement

4. CONCLUSION

Evaluation of four existing clear-sky scintillation models namely: ITU-R, Karasawa, Otung and Van de Kamp models have been presented in this study. This existing models were compared with the ground data in Ota obtained from Astra 2E/2F/2G satellite beacon at 12.245 GHz located at Covenant University. The ground measurements from Ota have confirmed that Karasawa model gave the best prediction for tropospheric scintillation intensity for Ota and its environment.

ACKNOWLEDGMENT

The authors are grateful to Covenant University for sponsoring the publication of this research

REFERENCES

- [1] Omotosho, T. V., Akinwumi, S. A., Usikalu, M. R., Ometan, O. O., Adewusi, M. O and Abdullah, M. (2017). Analysis of non-rainy attenuation on earth-space path in Ota, Southwest, Nigeria. IOP Conf. Series: Journal of Physics: Conf. Series 852 (2017) 012039 doi:10.1088/1742-6596/852/1/012039.
- [2] Akinwumi, S. A., Omotosho, T. V., Usikalu, M. R., Adewusi, M. O and Ometan, O. O. (2016). Atmospheric gases attenuation in West Africa. 2016 IEEE Radio and Antenna Days of the Indian Ocean, RADIO 2016; Hotel Le Recif Saint-Gilles Les Bains; Reunion.
- [3] Afolabi, O. O., Adagunodo, T. A., Ayorinde, T. T., Rabi, A. B., Akinwumi, S. A., Usikalu, M. R. (2017). Impact of Magnetic Activity on Occurrence of Ionospheric Amplitude Scintillation over Lagos, Nigeria. IEEE RADIO 2017 at Protea Hotel by Marriott Sea Point, Cape Town, South Africa. DOI: 10.23919/RADIO.2017.8242229. <http://ieeexplore.ieee.org/document/8242229>

- [4] Alagbe G.A., Afolabi O.O., Adagunodo T.A., Rabiou A.B., Akinwumi S.A., Omotosho T.V. (2017). Study of Ionospheric Amplitude Scintillation during Geomagnetic Activities of 2012 at Low Latitude Region. *Journal of Informatics and Mathematical Sciences*, 9(2): 251 – 256.
- [5] Cheng, Y. C. and Mandeep, J. S. (2014). Comparison of tropospheric scintillation prediction models of the Indonesian climate. *Earth, Planets and Space* 2014, 66:64.
- [6] Mandeep, J. S., Syed, S. H., Kiyoshi, I., Kenji, T., Mitsuyoshi, I. (2006). Analysis of tropospheric scintillation intensity on earth to space in Malaysia. *Amer J App Sci* 3(9):2029–2032.
- [7] Omotosho, T. V., Akinwumi, S. A., Usikalu, M. R., Ometan, O. O., and Adewusi, M. O. (2016). Tropospheric Scintillation and its Impact on Earth-Space Satellite Communication in Nigeria. 2016 IEEE Radio and Antenna Days of the Indian Ocean, RADIO 2016; Hotel Le Recif Saint-Gilles Les Bains; Reunion.
- [8] Adagunodo, T.A., Akinwumi, S.A., Omotosho, T.V., Akinyemi, M.L. 2017. Estimation of Specific Attenuation of Radio Signal in Southwest Nigeria. IEEE RADIO 2017 at Protea Hotel by Marriott Sea Point, Cape Town, South Africa.
DOI:10.23919/RADIO.2017.8242225.<http://ieeexplore.ieee.org/document/8242225>
- [9] Harris, R. A. (2002). COST Action 255, Radiowaves propagation modeling for SATCOM services at Ku-Band and above, European Space Agency publication Division Noordwijk, Netherlands.
- [10] Akinwumi, S.A., Omotosho, T.V., Usikalu, M.R., Ometan, O.O., Adewusi, M.O., Adagunodo, T.A. (2018). Study of Oxygen and Water Vapour Attenuation in West Africa. *Interciencia Journal*, 43(1): 180 – 191.
- [11] Mandeep, J.S. and Hassan, S. I. (2004). Comparison of 1-minute rainfall rate distribution for tropical and equatorial climates. *Space Commun* 19:193–198.
- [12] Gibson, J. D. (2002). *The Communication Handbook*, second edition, CRC Press, LCC, 59-1 to 59-16, 2002.
- [13] ITU-R; Recommendation P.618, 2012. Propagation data and prediction methods required for the design of Earth-space telecommunication system.
- [14] Karasawa Y., Yamada M., Allnut J. E. (1988). A new prediction method for tropospheric scintillation on Earth-space paths. *IEEE Trans Ant Prop* 36(11): 1608-1614.
- [15] Otung I. E., (1996). Prediction of tropospheric amplitude scintillation on a satellite link. *IEEE Trans Ant Prop* 44(12): 1600-1608.
- [16] Van De Kamp M. M. J. L., Tervonen J. K., Salonen E. T., Poirares B. J. P. V. (1999). Improved models for long-term prediction of tropospheric scintillation on slath paths. *IEEE Trans Ant Prop* 47(2): 249-260.