

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Molecular and Cellular Endocrinology

journal homepage: www.elsevier.com/locate/mce

Thyroid hormone receptors and ligands, tissue distribution and sexual behavior

Eleonora Carosa ^a, Andrea Lenzi ^b, Emmanuele A. Jannini ^{c,*}^a Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy^b Chair of Endocrinology, Department of Experimental Medicine, University of Rome Sapienza, Rome, Italy^c Chair of Endocrinology & Medical Sexology (ENDOSEX), Department of Systems Medicine, University of Rome Tor Vergata, Rome, Italy

ARTICLE INFO

Article history:

Received 7 September 2017

Received in revised form

7 November 2017

Accepted 8 November 2017

Available online 23 November 2017

Keywords:

Thyroid hormone receptors

Thyroid hormones

Testis

Ovaries

Genitals

Sexual function

Sexual behavior

Sexual dysfunction

Thyroid dysfunction

Contents

1. Introduction	50
2. Thyroid hormones and thyroid hormone receptors	50
3. Thyroid hormone receptors expression in sexual organs	51
3.1. Male	51
3.1.1. Thyroid hormone receptors in testis	51
3.1.2. Thyroid hormone receptors in epididymis	51
3.1.3. Thyroid hormone receptors in corpora cavernosa	51
3.2. Female	51
3.2.1. Thyroid hormone receptors in ovary	51
3.2.2. Thyroid hormone receptors in oviduct	52
3.2.3. Thyroid hormone receptors in uterus	52
3.2.4. Thyroid hormone receptors in vagina	52
3.3. Brain	53
4. Thyroid and sexual function in male	53
4.1. Thyroid and ejaculation	54
4.2. Thyroid and erectile dysfunction	54

* Corresponding author. Chair of Endocrinology and Medical Sexology (ENDOSEX), Department of Systems Medicine, University of Rome Tor Vergata, Via Montpellier, 1, 00133 Rome, Italy.

E-mail address: eajannini@gmail.com (E.A. Jannini).

<https://doi.org/10.1016/j.mce.2017.11.006>

0303-7207/© 2017 Elsevier B.V. All rights reserved.

4.3. Thyroid and hypoactive sexual desire	54
5. Thyroid and sexual function in female	54
6. Thyroid and sexual behavior	56
7. Thyroid and sexual orientation	56
8. Conclusion	57
References	57

1. Introduction

The thyroid hormones (THs) triiodothyronine (T_3) and tetraiodothyronine, or thyroxine (T_4), not only dramatically impact on development and differentiation, but also on the sexual and reproductive function. There is large body of literature, in fact, on the effects of THs on the reproductive function in both humans (Poppe and Velkeniers, 2004; Wajner et al., 2009) and animals (Hapon et al., 2010; Nelson et al., 2011).

For a long time the gonads were thought to be unresponsive to THs, but TH receptors (TR) were discovered in rat (Jannini et al., 1990; Palmero et al., 1988) and then in human testis (Jannini et al., 2000). In women, the association of menstrual disturbance with thyroid disease was described as early as 1840 by von Basedow, but the discovery of TRs in the ovary was carried out at the end of last century (Wakim et al., 1994b). Therefore, the link between thyroid and reproductive function was well established. Since then, research has shown that thyroid dysfunction is associated with an adverse effect on fertility, both in men (Wagner et al., 2009) and women (Dittrich et al., 2011). There is also evidence that THs can affect the sex steroid hormone axis (Bagamasbad and Denver, 2011), consequently sexual hormones and the pituitary gland can mediate the action of THs on the reproductive physiology.

While the effects of THs on fertility have been widely studied, little is known about their influence on sexual function. In the last few years, an increasing number of evidences have shown the influence of THs on male sexual function, particularly on ejaculation control as well on desire and erectile function (Carani et al., 2005; Corona et al., 2012b; Di Sante et al., 2016). The female sexual function and the relationship with thyroid function is still less studied. Furthermore, studies conducted on animals have shown the presence of TRs in the male (Carosa et al., 2010) and female genitalia (Rodriguez-Castelan et al., 2017). Moreover, knockout mice for TRs showed alterations in sexual behavior (Dellovade et al., 2000).

The purpose of this review is to summarize and discuss the available data on the influence of THs on male and female sexual function to understand the molecular mechanisms of the influence of the thyroid gland on sexual behavior and function.

2. Thyroid hormones and thyroid hormone receptors

TH action is mainly dependent on the binding of the cognate with specific receptors, the most important being at nuclear level (Mendoza and Hollenberg, 2017). Thyroid hormone receptors (TRs) belong to the steroid/thyroid receptors superfamily that evolved from a single ancestor gene (Carosa et al., 1998; Kostrouch et al., 1998). TR_α and TR_β are the product of two distinct genes that are further differentially spliced into $TR_{\alpha 1}$, $TR_{\alpha 2}$ (Jannini et al., 1992; Sap et al., 1986; Weinberger et al., 1986), and $TR_{\beta 1}$ and $TR_{\beta 2}$ (Lazar et al., 1988; Mitsushashi et al., 1988). $TR_{\alpha 1}$ and $TR_{\beta 1}$ are widely expressed and act as thyroid hormone-dependent transcription factors,

inducing or repressing gene expression in response to T_3 . The differentially spliced product $TR_{\alpha 2}$ does not bind the hormone and exerts a dominant negative effect on the action of other TRs (Katz et al., 1995; Koenig et al., 1989). Although two different genes encode the two TR isoforms, they share a very high degree of structural homology and the more conserved part of protein is the DNA binding domain (Fig. 1). On the DNA, TRs recognize the sequence AGGTCA, to which they can bind as a monomer. In the thyroid responsive element (TRE) of the promoters of responsive genes this sequence is repeated with a space of 4 bp to permit the DNA bind of the heterodimer TR/retinoid X receptor (RXR). The presence of RXR increases the binding affinity (Lazar, 2003).

Thyroid hormones are hydrophobic and for a long time it was thought they entered the cytoplasm by passive diffusion. In recent years, it has become clear that this is not the case and several TH transport proteins have been identified (Friesema et al., 2003; Schweizer et al., 2014). To date, the most clinically relevant THs transporter is the monocarboxylate transporter 8 (MCT8), which is associated with a THs insensitivity syndrome (Schwartz et al., 2005). It is now clear that TH transporters can modulate the real effect of THs because they regulate, at the tissue level, the availability of THs. Deeping on THs transporters is beyond the purpose of this review, but the future will show if their alterations may involve sexual function. Here we can highlight that MCT8 is expressed both in the testis and in the ovary (Friesema et al., 2003; Romano et al., 2017).

Inside cells, THs bind TRs to regulate target gene expression. The interaction of T_3 with the ligand domain induces a conformation change of the hormone-binding pocket. The change of conformation appears to be critical for the release of corepressor and the recruitment of coactivators on the regulated gene's promoters. In the classic view (Fig. 2), T_3 action is based on its positive effect on target genes. In the absence of T_3 , the TRs recruit a multiprotein complex that includes nuclear corepressors, which in turn recruit histone deacetylase 3 (HDAC3) and other proteins to mediate transcriptional repression via histone deacetylation (Sun et al., 2013; You et al., 2013). The presence of T_3 leads to dismissal of




Fig. 1. Schematic representation of human TR isoforms and their domain functions.




Fig. 2. Classical view of THs action. The interaction of T_3 with TH induces the release of corepressor and the recruitment of coactivators on the regulated gene's promoters. In the absence of T_3 the TRs recruit a multiprotein complex that includes nuclear corepressors, which in turn recruit HDAC3 and other proteins to mediate transcriptional repression via histone deacetylation.

the corepressor complex and recruitment coregulators that induce histone acetylation and transcriptional activation. Some recent studies, based on chromatin affinity precipitation (ChAP)-seq and DNase hypersensitivity analysis showed extensive chromatin remodeling in the presence of T_3 stimulation in accord with the classical model of THs action (Ayers et al., 2014; Chatonnet et al., 2012; Grontved et al., 2015; Ramadoss et al., 2014). These studies have also established that RXR overlaps TR DNA binding, endorsing the fact that TRs act like heterodimers with retinoid X receptor (RXR) (Ramadoss et al., 2014). New studies demonstrated that the molecular mechanisms of T_3 actions are more complicated and the classical model does not explain the exact mechanism of action; in particular, the T_3 negative regulation remains a paradox, in the liver TRs down-regulate more genes in the presence of T_3 than they activate. Several hypotheses have been suggested to explain the molecular mechanism of T_3 negative gene regulation. One of these asserts the possibility that coregulators act as a repressor on promoters of negative regulated genes (Costa-e-Sousa & Hollenberg, 2012), but it is possible that numerous models will be required to explain the exact molecular mechanism of TRs action. To support the direct effects of THs on sexual function it is important to demonstrate that TRs are expressed in male and female genitalia.

3. Thyroid hormone receptors expression in sexual organs

3.1. Male

3.1.1. Thyroid hormone receptors in testis

T_3 regulates the maturation and growth of the testis, controlling Sertoli and, with much lower impact, Leydig cell proliferation and differentiation during testicular development in human and other mammalian species (Francavilla et al., 1991; Holsberger and Cooke, 2005; Jannini et al., 1999; Mendis-Handagama and Siril Ariyaratne, 2005). The first study describing the selective presence of specific TH nuclear binding sites in Sertoli cells of developing rat testis opened a new research avenue (Jannini et al., 1990), since these findings changed the classical view of the testis as a thyroid unresponsive organ (Barker and Klitgaard, 1952). Ontogenetic expression of TRs in human and in rat testis established that the fetal and prepubertal ages of Sertoli cells represent the period of maximal expression, but the binding capacity is not completely absent in the adult (Buzzard et al., 2000; Canale et al., 2001; Jannini et al., 1993, 1995, 1999, 2000; Tagami et al., 1990; Ulisse et al., 1992). In rats, TRs were present in different testicular cell types and their expression changes during testis maturation (Table 1). $TR_{\alpha 1}$ expression was very high in Sertoli cells during their proliferative

phase, until day 15 post-natal, then decreases and remains very low in adulthood (Jannini et al., 1994, 1999). $TR_{\alpha 1}$ was also found in spermatogenic germ cells from intermediate spermatogonia to mid-cycle pachytene spermatocytes (Buzzard et al., 2000), as well in the interstitial compartment in immature testis, probably in the Leydig cells, but at very low levels (Buzzard et al., 2000). The same laboratory found that the $TR_{\beta 1}$ expression in immature Sertoli cells and spermatogenic germ cells was also very low in relation to $TR_{\alpha 1}$ expression. This confirmed our findings, which previously showed that TR_{β} mRNA is below the threshold of detectability of both Northern and ribonuclease (RNase) protection analyses (Jannini et al., 1995). In fact, Palmero et al. (1995) demonstrated that $TR_{\beta 1}$ is detectable only after 30 cycles of PCR amplification. According to our evidences using biochemical and molecular techniques of detection of TRs, studies on transgenic mice lacking either TR_{α} or TR_{β} have shown that the effect of T_3 on Sertoli cell proliferation is mediated through TR_{α} isoform (Holsberger et al., 2005). Furthermore, Fumel et al. (2015), using a TR_{α} dominant negative isoform selectively expressed in Sertoli and Leydig cells, confirmed the role of $TR_{\alpha 1}$ in Sertoli cells during post-natal life and pointed out that T_3 does not regulate steroidogenic activity in a direct manner on Leydig cells, also fitting the spatio-temporal distribution of TRs previously demonstrated. Thyroid hormones modulates the activity of glucose transporters and aromatase in Sertoli cells to regulate the function of seminiferous epithelium (Carosa et al., 2005; Ulisse et al., 1994, 1998).

3.1.2. Thyroid hormone receptors in epididymis

The presence of $TR_{\alpha 1}$ and $\beta 1$ isoforms was more recently demonstrated in epithelial cells derived from adult rat epididymis (De Paul et al., 2008) (Table 1). However, further studies are necessary to understand the role of TRs, if any, in epididymis function in light of the fact that TR proteins are localized in the cytosol of the cells, suggesting a possible active role for THs in the epithelial cells of epididymis acting through a non-genomic fashion (Bassett et al., 2003).

3.1.3. Thyroid hormone receptors in corpora cavernosa

All isoforms of TRs, i.e. $\alpha 1$, $\alpha 2$ and $\beta 1$ are expressed in rat and human penis (Table 1). In particular, $TR_{\alpha 1}$ and $TR_{\alpha 2}$ are present from perinatal to adult age, while the $\beta 1$ form is almost entirely absent in fetal cells while present in the adult. Immunohistochemistry experiments using an anti- $TR_{\alpha 1}$ antibody demonstrated a widespread staining for this TR in both endothelial and muscular cells of the corpora cavernosa and corpus spongiosum of the human adult penis (Carosa et al., 2010).

3.2. Female

3.2.1. Thyroid hormone receptors in ovary

Thyroid function has been associated with female reproduction also in ancient medicine (Niazi et al., {Krassas, #9215}), but only very recently to the female sexual function. Thyroid diseases have been associated with menstrual disturbance, miscarriages and infertility (Dittrich et al., 2011; Romano et al., 1991) (Table 2). Several studies have demonstrated the presence of TRs in all ovary cell types. $TR_{\alpha 1}$ and $\beta 1$ are expressed in primordial, primary and secondary follicles (Aghajanova et al., 2009), and they are also present in follicular fluid and in granulosa cells (Aghajanova et al., 2009; Wakim et al., 1993, 1994b). Mature oocytes from women undergoing IVF express $TR_{\alpha 1}$, $TR_{\alpha 2}$, $TR_{\beta 1}$, and $TR_{\beta 2}$ mRNA, thus supporting the hypothesis that T_3 has a direct effect on the human oocyte (Stavreus Evers, 2012; Zhang et al., 1997). Furthermore, the corpus luteum of rats, cows and rabbits shows immunostaining for all isoforms of TRs (Aghajanova et al., 2011; Fedail et al., 2014;

Table 1
Thyroid hormone nuclear receptors (TRs) Expression in male genital tissues.

Tissue/Cell Type	Receptor isoform	Expression		References
		Fetal/Pre- pubertal	Adult	
Whole testis	TR α ₁	++	+	(Jannini et al., 1999) (Buzzard et al., 2000; Jannini et al., 2000)
	TR α ₂	++	++	(Buzzard et al., 2000; Jannini et al., 1999; Jannini et al., 2000)
	TR β ₁	±	–	(Buzzard et al., 2000; Jannini et al., 2000)
Sertoli cells	TR α ₁	++	++	(Buzzard et al., 2000; Jannini et al., 2000)
	TR α ₂	?	?	
	TR β ₁	?	±	(Palmero et al., 1995)
Leydig cells	TR α ₁	?	±	(Buzzard et al., 2000)
	TR α ₂	?	?	
	TR β ₁	?	?	
Germ cells	TR α ₁	?	+	(Buzzard et al., 2000)
	TR α ₂	?	?	
	TR β ₁	?	±	(Buzzard et al., 2000)
Epididymis	TR α ₁	?	+	(De Paul et al., 2008)
	TR α ₂	?	?	
	TR β ₁	?	+	(De Paul et al., 2008)
Penis	TR α ₁	++	++	(Carosa et al., 2010)
	TR α ₂	++	++	(Carosa et al., 2010)
	TR β ₁	–	++	(Carosa et al., 2010)

Table 2
Thyroid hormone nuclear receptors (TRs) expression in female genital tissues.

Tissue/Cell type	Receptor isoform	Expression	References	
Ovary	Stroma TR α ₁	++	(Rodriguez-Castelan et al., 2017) (Aghajanova et al., 2009; Wakim et al., 1994b)	
	Epithelium TR α ₂	++	(Rodriguez-Castelan et al., 2017)	
	Primary follicle TR β ₁	++	(Aghajanova et al., 2009; Rodriguez-Castelan et al., 2017) (Wakim et al., 1994b)	
	Corpus luteus TR α ₁	++	(Rodriguez-Castelan et al., 2017) (Stavreus Evers, 2012) (Navas et al., 2014) (Stavreus Evers, 2012)	
Oviduct	TR α ₂	++		
	TR β ₁	++		
	TR α ₁	++	(Rodriguez-Castelan et al., 2017)	
	TR α ₂	++		
Uterus	Endometrium	TR α ₁	++	(Rodriguez-Castelan et al., 2017) (Aghajanova et al., 2011) (Stavreus Evers, 2012)
		TR α ₂	+	(Aghajanova et al., 2011)
		TR β ₁	++	(Rodriguez-Castelan et al., 2017) (Aghajanova et al., 2011) (Aghajanova et al., 2011) (Stavreus Evers, 2012)
	Myometrium	TR α ₁	+	(Rodriguez-Castelan et al., 2017) (Hulchiy et al., 2012)
TR α ₂		+	(Rodriguez-Castelan et al., 2017)	
Vagina	Epithelium	TR α ₁	++	(Rodriguez-Castelan et al., 2017)
		TR α ₂	++	
		TR β ₁	+	
Clitoris	Smooth muscle cells	TR α ₁	++	(Rodriguez-Castelan et al., 2017)
		TR α ₂	++	
		TR β ₁	++	
Corpus cavernosum	TR β ₁	++		
		++		

Rodriguez-Castelan et al., 2017). Positive staining of ovarian stromal cells and in ovarian surface epithelium was also observed for both TR α ₁ and β ₁ (Aghajanova et al., 2009; Wakim et al., 1994a).

3.2.2. Thyroid hormone receptors in oviduct

Papers that have studied the presence of TRs in oviduct are scarce. The presence of TR α ₁ was described in the epithelium and smooth musculature of ampulla and isthmus in rat (Navas et al., 2014) (Table 2). Recently, the expression of all TR isoforms was shown in the luminal and glandular epithelium and in smooth muscle cells of whole oviductal regions in rabbit (Rodriguez-Castelan et al., 2017).

3.2.3. Thyroid hormone receptors in uterus

Extensive studies were conducted in uterus during the normal menstrual cycle (Aghajanova et al., 2011) and during pregnancy (Stavreus Evers, 2012). The localization of TR α ₁ and TR β ₁ has been found in the luminal epithelium, glandular epithelium and stroma

of human endometrium throughout menstrual cycle (Aghajanova et al., 2011), however no staining was present for the α ₂ isoform (Table 2). Despite the fact that the presence of binding sites for T₃ in the myometrium was shown in 1983 (Kirkland et al., 1983), only in recent years has TR α ₁ been described in the myometrium from macaques (Hulchiy et al., 2012), rats (Navas et al., 2014), and rabbits (Rodriguez-Castelan et al., 2017).

3.2.4. Thyroid hormone receptors in vagina

The presence of TRs in the vagina has been studied very recently (Rodriguez-Castelan et al., 2017) (Table 2). Both the vaginal epithelium and the smooth and striated musculatures of all portions of the vagina were positive for TR α ₁₋₂ and β ₁. Notably, the positive immunostaining for all TR isoforms are found in smooth muscle cells of the corpus cavernosum of the clitoris and the epithelial cells of the Skens's gland, both involved in the female pleasure (Jannini et al., 2014). This finding is interesting because it reveals an additional connection between male and female

genitalia. Indeed, the presence of TRs in the erectile tissues of men and women could emphasize a similar involvement of T_3 action in the sexual function of both genders.

3.3. Brain

The brain may also be considered a sexual organ for regulating sexual desire and arousal (Ciocca et al., 2016). TH actions in the brain are extremely complex because they are fundamentals for brain development (Moog et al.). Indeed, hypothyroidism during fetal life is associated with cretinism, a severe form of mental retardation (Bernal, 2007). The expression of TRs in the brain has been extensively studied in mammals: in mouse brain, TRs α and β isoforms are expressed in almost all parts of the brain with an overlapping distribution. Some differences between α and β isoform expression are observed in the hippocampus, amygdala and hypothalamus (Fig. 3) (Bernal, 2000).

In the human brain, TR mRNAs are detected at around 8 weeks of gestation (Iskaros et al., 2000). Concerning the receptor protein, there is only one study in which receptor concentrations were quantitated by ligand-binding assays (Bernal and Pekonen, 1984). The receptor protein concentration was found to be very low at 10 weeks of gestation, and increased by a factor of 10 up to the 16th week, coinciding with the period of neuroblast multiplication. During this time, the brain gains in weight and DNA content by about five-fold, so that the total brain T_3 receptor content increases 500 times. This coincides with the period of active neuroblast proliferation (Dobbing and Sands, 1970). In neural cell cultures, T_3 receptors have been detected in neurons, astrocytes and oligodendrocytes (Luo et al., 1986; Yusta et al., 1988).

4. Thyroid and sexual function in male

The association between thyroid diseases and sexual dysfunction in men has not been systematically studied until recently. Indeed, in the last 10 years, an increasing number of studies have correlated different aspects of sexual performance and thyroid functions. The first report that demonstrated a close correlation between specific sexual disorders in males with thyroid hypo- and hyperfunction was published in 2005 (Carani et al., 2005). In this study four major sexual alterations – HSDD (hypoactive sexual desire disorder), PE (premature ejaculation), DE (delayed

ejaculation), and ED (erectile dysfunctions) – were associated with hyper- or hypothyroid status (Fig. 4). The main sexological complaint found in that study in hyperthyroid patients was PE, whereas in hypothyroid subjects, it was DE. Remarkably, both ejaculatory dysfunctions very frequently reverted upon





Fig. 4. Prevalence of sexual dysfunction in men with thyroid diseases. Modified from Carani et al. (2005).

TR expression in mouse brain

TR α 1



TR β 1




Fig. 3. TRs expression in mouse brain (Bernal, 2000). This image has been made freely available under a Creative Commons (CC-BY-NC-ND) license. A copy of the license can be viewed at <http://creativecommons.org/licenses/by-nc-nd/2.0/>.

achievement of euthyroidism in the absence of any other treatment for the sexual symptom. Although these can be considered secondary effects of treatment on mood, such a response on PE and DE was beyond all expectations, suggesting a direct involvement of thyroid hormones on the physiology of ejaculation. Now, it is known that TRs are located in the male genitals (Carosa et al., 2010), so it can be hypothesized a direct T₃ effect on erectile tissues and smooth muscle cells of the genital tract. Nevertheless, the molecular mechanisms involved in the T₃ action at this level remain unclear. Below we report all the evidences published to date about the influence of T₃ on the various aspects of male sexual function.

4.1. Thyroid and ejaculation

Several studies have confirmed the strong association between DE and PE with hypothyroidism and hyperthyroidism (Carani et al., 2005; Cihan et al., 2009a; Corona et al., 2011, 2012a, 2006b), respectively. This connection has been widely documented both in animal models and humans (Cahangirov et al., 2011; Cihan et al., 2009b). Cahangirov et al. (Cahangirov et al., 2011) demonstrated that hyperthyroidism in rats increases seminal vesicle contraction frequency and bulbospongiosus muscle contractile activity, which indicates that hyperthyroidism can affect ejaculatory emission and expulsion phases. In that study, after a 28-day washout period (to determine spontaneous recovery from hyperthyroidism) the aforementioned alterations were reversed, confirming the direct role of THs in the control of ejaculation. These results have been replicated in several clinical studies in human patients (Carani et al., 2005; Cihan et al., 2009a). In a series of 755 consecutive men seeking medical care for sexual dysfunction, including PE, the prevalence of suppressed thyroid-stimulating hormone (TSH), which is a marker of possible hyperthyroidism, was twofold higher in patients with PE than those reporting normal ejaculatory timing (Corona et al., 2004). A recent study in a Turkish population essentially confirms an association between the hyperthyroid state and PE, which was substantially ameliorated by therapy for the thyroid disease (Cihan et al., 2009a). However, and not surprisingly, Waldinger did not find any association between TSH levels and intravaginal ejaculation latency time (IELT) in a cohort of Dutch subjects (Waldinger et al., 2005) with lifelong PE, which logically suggests that the association with hyperthyroidism is restricted to the acquired type of PE. In light of all these reports, hyperthyroidism should be considered a novel and reversible etiological risk factor for PE, and TH disorders should be suspected, by simple physical examination, in the case of ejaculatory disorders (Jannini et al., 2011, 2015; McMahon et al., 2013; Sansone et al., 2015).

We found high prevalence of DE in hypothyroid patients, 64.3% (Carani et al., 2005) (Fig. 4). Furthermore, TSH levels were positively related to reported ejaculatory latencies (Corona et al., 2011). In patients with hypothyroidism, a resolution of DE was obtained in half of the subjects after thyroid hormone normalization (Carani et al., 2005) (Fig. 4). The view that thyroid hormones regulate the ejaculatory reflex is emerging and hypothyroidism should be ruled out in each patient with DE as recommended by International Society of Sexual Medicine (ISSM) guidelines (Corona et al., 2016). Hence, the view that thyroid hormones regulate the ejaculatory reflex is consistently emerging (Di Sante et al., 2016).

4.2. Thyroid and erectile dysfunction

Anecdotal reports suggest that ED is frequent (up to 70%) in men with hyperthyroidism (Meikle, 2004). Small studies in Italian and Greek patients with thyroid dysfunction found that both hypothyroidism and hyperthyroidism can be associated with ED and that the correction of the underlying thyroid conditions frequently

restores erectile function (Fig. 4) (Carani et al., 2005) (Krassas et al., 2008). In Krassas et al.'s replication study (Krassas et al., 2008), 63% and 70% of the hypo- and hyperthyroid subjects, respectively, have some form of ED compared with 34% of the control group. While the prevalence of ED in hypothyroidism was similar in previous Carani et al.'s study (64% Carani et al., 2005), the reported number of hyperthyroid subjects with a pathological score of International Index of Erectile Function (IIEF) was lower (15%). In contrast, Veronelli et al. (2006) showed that 59% of men with thyroid problems had ED; interestingly, this result did not change when adjusted for age, suggesting that TH were more important than age in ED. A more recent study (Krysiak et al., 2017) documented that men with overt hypothyroidism obtained lower scores in all five domains of IIEF, while men with subclinical hypothyroidism only in erectile function domain. L-thyroxine treatment improved erectile function and normalized intercourse satisfaction, orgasmic function, sexual desire and overall satisfaction in the formed group, as well as normalized erectile function in the latter. To date, no information on the prevalence of thyroid dysfunction in ED men is available either for the general population or for men seeking medical help for sexual problems. Further research is required to understand the nature of the link between thyroid disorders and ED; for now we can only emphasize that both α and β TRs have been described in rat and human corpora cavernosa endothelial and smooth muscle cells (Carosa et al., 2010; Owen et al., 2007). Studies in animal models of hyperthyroidism indicate an impairment of nitric oxide (NO)-dependent relaxation of corpora cavernosa (Hu et al., 2009; Ozdemirci et al., 2001). In rabbit corpora cavernosa strips, both acetylcholine- and electrical field stimulation-induced relaxation were impaired, whereas sensitivity to a NO donor, sodium nitroprusside, was unchanged (Ozdemirci et al., 2001). These data suggest an effect of thyroid hormones in penile NO formation, which has been demonstrated in a rat model (Hu et al., 2009). It is therefore possible that hyperthyroidism-associated ED could be due to a direct effect of TH on their cognate receptors.

4.3. Thyroid and hypoactive sexual desire

We found that low sexual desire may be related to hypothyroidism and can be normalized by TH therapy (Carani et al., 2005) (Fig. 4). The underlying pathogenic mechanisms have not been completely clarified. It is possible that a hypothyroidism-induced prolactin rise could mediate these negative effects on sexual desire. However, a direct role of THs on the central nervous system can also be hypothesized. THs are known to interfere with the functions of the reproductive axis in men (Bagamasbad and Denver, 2011; Krassas et al., 2010), which in turn might contribute to sexual dysfunction. Furthermore, hyperthyroidism-induced increase of sex hormone binding globulin (SHBG), which binds androgens with higher affinity than estrogens, might lead to a relative hyperestrogenism, which, per se, can alter sexual responses. In addition, the increase of prolactin that is often present in association with hyperthyroidism can be associated with a decrease in libido. All these findings demonstrate a non-controversial role of THs in human sexual dysfunction. (Maggi et al., 2013).

5. Thyroid and sexual function in female

While in male are relatively rare, thyroid hypo- and hyperfunctions are frequent in female. Thus, it is very surprisingly that the incidence of sexual dysfunction in women with hypo- or hyperthyroidism is still unknown. Thyroid disorders in women are, in fact, commonly associated with abnormalities of reproductive physiology such as menstrual irregularities and infertility (Krassas, 2000), but sexual function has been rarely studied. For instance, the

prevalence of menstrual disturbances in women with hypothyroidism has been estimated at 25–70%; ovulatory function and fertility are also affected (Bhasin et al., 2007). Both hyper- and hypothyroidism are associated with fatigue, myalgias, and mood disturbances such as irritability and depression, which can contribute to sexual dysfunction (Bhasin et al., 2007). Since the incidence of hypothyroidism also peaks at the age of menopause and perimenopausal symptoms could overlap with symptoms of hypothyroidism, screening for hypothyroidism in women at this age is generally recommended (Shifren and Gass, 2014). Indeed, persistent primary hypothyroidism is occasionally associated with hyperprolactinaemia, due to the stimulatory effect of the thyrotropin-releasing hormone (TRH) on the production of prolactin (Oppo et al., 2011). Thyroxine replacement therapy with a return to the euthyroid state restores menstrual and ovulatory function in most hypothyroid and hyperthyroid women, and also normalizes prolactin levels in those with coincident hyperprolactinaemia (Krassas, 2000). Few studies have investigated sexual function in women during thyroid diseases. In 2010, Atis et al. reported that women with hyperthyroidism and hypothyroidism have significantly lower female sexual function index (FSFI) domain scores compared with age-matched healthy women (Atis et al., 2010, 2011) (Table 3 and Fig. 5). According to the proposed FSFI full-scale cut-off level of 26.55 (Wiegel et al., 2005), 60% of women with hyperthyroidism in this study may have sexual dysfunction. One study conducted by Veronelli et al. revealed that diabetic, obese, and hypothyroidic women had a reduced score in the FSFI questionnaire when compared with healthy women (Veronelli et al., 2009). The same result was found by Oppo et al. (2011), who reported that abnormal thyroid function significantly impairs female sexual function, as assessed by the FSFI questionnaire, and that restoration of the euthyroid state is associated with a rapid improvement of most FSFI domain scores (Oppo et al., 2011). In addition, Oppo showed that restoration of biochemical euthyroidism was associated with normalization of the desire, satisfaction and pain domains, while arousal/lubrication and orgasm remained significantly different from healthy euthyroid controls, in spite of some improvement in orgasm.

Hyperthyroidism has been anecdotally connected in the mind of several endocrinologists with an increase in women's sexual desire. This is not the case in all studies reported, which mirrors findings of our report in males (Carani et al., 2005) and provides objective evidences that untreated thyrotoxicosis is associated with important female sexual dysfunction.

The finding that opposite endocrine conditions are associated with similar sexual dysfunctions may appear surprising, but the mechanisms involved in determination of sexual dysfunction in women are complex (Ghizzani et al., 2003; Graziottin, 2003; Jannini and Lenzi, 2003), and the role of altered thyroid status in female sexual functions is still unclarified. However, all these studies provide clear evidence that hyper- and hypothyroidism




Fig. 5. Female Sexual Functions Index (FSFI) and Beck's Depression Inventory (BDI) questionnaire scores in women with thyroid diseases. Modified from (Atis et al., 2010); (Atis et al., 2011).

markedly impair sexual function in both women as in men (Oppo et al., 2011; Pasquali et al., 2013). Nevertheless, in women, in whom thyroid diseases are more frequent than in men, sexual function screening should be required (Balercia et al., 2007).

Emerging data suggests that, in women, sexual dysfunctions may be associated with thyroid autoimmunity, both in the presence of hypothyroidism and subclinical hypothyroidism (Krysiak et al., 2016; Pasquali et al., 2013).

Autoimmunity thyroiditis diseases are common in women. Actually, Hashimoto's thyroiditis is one of the most common human disorders, as well as the most frequent cause of hypothyroidism in developed countries (Caturegli et al., 2007). Several studies showed that women with autoimmune thyroiditis,

Table 3

FSFI (Female Sexual Functions Index) and BDI (Beck's Depression Inventory) questionnaire scores, modified from Atis et al., (2010), 2011.

	Control group	Hyper-thyroid	P	Control group	Hypo-thyroid	P
Total FSFI	29 ± 10.4	24.2 ± 9.9	>0.02	32.31 ± 3.5	23.92 ± 5.18	>0.02
Desire	4.3 ± 2.3	3.8 ± 2.1	>0.05	5.16 ± 0.74	4.01 ± 0.93	>0.02
Arousal	4.7 ± 2.2	3.4 ± 2.3	>0.02	5.37 ± 0.80	3.28 ± 1.30	>0.02
Lubrication	5.1 ± 1.6	4.3 ± 1.9	>0.02	5.40 ± 0.52	4.24 ± 0.82	>0.02
Orgasm	5.1 ± 1.5	4.0 ± 2.2	>0.02	5.46 ± 0.51	3.95 ± 1.23	>0.02
Satisfaction	4.9 ± 2.0	4.2 ± 1.9	>0.02	5.40 ± 0.54	4.06 ± 0.96	>0.02
Pain	5.0 ± 2.6	4.4 ± 2.3	>0.02	5.50 ± 0.56	4.37 ± 1.14	>0.02
BDI	11.9 ± 13.2	18.9 ± 15.4	>0.02	7.82 ± 5.20	12.8 ± 7.40	>0.05

compared with age-matched healthy controls, reported a significant increase in the prevalence of sexual dysfunction and displayed a significant decrease in sexual desire (Krysiak et al., 2016; Pasquali et al., 2013). In this respect, the availability of a shortened form of the FSFI questionnaire (Isidori et al., 2010) in all endocrine outpatient facilities would be useful to screen women with thyroid diseases. However, *mutatis mutandis*, it should be useful to screen all women complaining of sexual dysfunction for the presence of thyroid hypofunction and thyroid autoimmunity.

6. Thyroid and sexual behavior

Thyroid hormone regulates various developmental and pivotal physiological processes such as brain development. In effect, thyroid diseases are associated with altered mental processes in humans. Deficiency in TH activity during the perinatal period causes cretinism. Hypothyroidism in adults, although less dramatic, also induces cognitive dysfunction (concentration, memory psychomotor speed) and alterations in mood, with increased rates of depressive and anxiety symptoms (Bauer et al., 2008; Heinrich and Graham, 2003; Sait Gonen et al., 2004). Certainly, mood alteration correlated to thyroid dysfunction may partly justify the alterations of sexual functions found in hyper- and hypothyroid patients, mainly with respect to sexual desire. Several studies conducted in animals have revealed that THs supported estrogen in the control of sexual behavior (Bagamasbad and Denver, 2011). Estrogen plays a critical role in reproductive development, physiology and sexual behavior, acting in the brain, in particular in the hypothalamus. The interplay between TH and estrogen actions on this brain region may provide a mechanism for assessing metabolic state, thus affecting reproductive physiology and behavior. For example, in birds and mammals, TH promotes the transition to anestrus. In some mammals, exposure to cold temperatures affects reproductive behavior, and increases plasma TH levels, suggesting the presence of cross-regulation between TH and estrogen (Vasudevan et al., 2002). Molecular data have implicated that TR, specifically $TR_{\alpha 1}$, inhibit estrogen-dependent gene expression in the hypothalamus (Vasudevan et al., 2002). Thyroid hormone administration inhibited lordosis behavior in estrogen-primed female mice (Morgan et al., 2000) and rats (Dellovade et al., 1996) (Fig. 6). Surprisingly, it has been shown that female mice knockout for $TR_{\alpha 1}$ ($\alpha 1$ KO) showed poorer lordosis than wildtype ($\alpha 1$ WT), while female knockout for TR_{β} (β KO) showed greater lordosis than wildtype females (β WT) (Dellovade et al., 2000) (Fig. 6). In agreement with those data, estrogen replacement plus TH treatment decreased lordosis behavior in ovariectomized (OVX) rats and mice, compared to estrogen replacement alone (Dellovade et al., 1996; Morgan et al., 2000). Similarly, thyroid-intact, OVX female rats that received estrogen replacement showed delayed onset of lordosis behavior compared to animals that were thyroidectomized, OVX + estrogen (Dellovade et al., 1996), suggesting that the actions of TH on sexual behavior are complex. Furthermore, the $TR_{\alpha 1}$ isoform subtly decreases the sexual behavior of gonadal-intact male mice (Vasudevan et al., 2013) (Fig. 6). Several studies on estrogens regulated gene promoters demonstrated that $TR_{\alpha 1}$ can decrease the transcriptional activation mediated by the estrogen receptor isoforms; this interaction may modulate sexual behavior in mice (Dellovade et al., 1996; Morgan et al., 2000; Dellovade et al., 2000).

The role of sexual hormones in arousal has been defined in humans. In the brain, estradiol synthesis increased in areas related to sexual arousal. Furthermore, it has been described that estrogen can sustain libido acting in the preoptic area and in the anterior hypothalamus (Schulster et al., 2016). Actually, in men the levels of aromatase are highest in those brain areas, where the estrogen receptors are expressed. However, the exact role of estrogens in




Fig. 6. Sexual behavior in female and male KO mice. Data are collected from: (Dellovade et al., 2000; Ogawa et al, 1997, 1998, 1999, 2000; Vasudevan et al., 2013).

male sexual function, including libido, is difficult to determine (Carani et al., 1997; Schulster et al., 2016). In male mice, a lack of estrogen receptors is associated with the reduction or abolition of sexual behavior (Ogawa et al, 1997, 2000).

The above-mentioned data showed that the alteration of thyroid function in men and women is associated with reduced sexual desire. Animal studies show us an interaction between THs and estrogen for sexual function regulation in the brain (Fig. 6). It is possible to hypothesize that the genesis of HSDD in men and women with thyroid diseases can be mediated by the action of THs on estrogen regulated genes in the hypothalamus (Schulster et al., 2016).

7. Thyroid and sexual orientation

Several experimental findings suggest that in human sexual orientation genes and hormones play a pivotal role (Jannini et al., 2010). A recent population-based study on > 5000 Danish homosexuals found a significantly higher frequency of Graves' disease when compared to general population, also when excluding men with HIV or AIDS (Frisch et al, 2014). This has been related to possible genetic/prenatal mechanisms linking the risk of Graves' disease with possible genes or genetic markers related to homosexuality carried on the X chromosome (Hu et al., 1995). Interestingly, the mothers of homosexual men are characterized by high frequency of extreme skewing of X-inactivation (Bocklandt et al., 2006), a similar high frequency that has been reported in females with the autoimmune thyroid disease (Brix et al., 2005). The possible relationship between thyroid and sexual orientation is

further suggested by the evidence that homosexuals display lower body mass index than heterosexual men, independently of diet or exercise (Blanchard and Bogaert, 1996; Deputy and Boehmer, 2010; Frisch and Zdravkovic, 2010). Moreover, homosexual adolescents more likely have mothers with thyroid dysfunction during pregnancy than heterosexuals (Sabuncuoglu, 2015). More recently, a genome-wide association study of male sexual orientation on a primarily European ancestry sample of >1000 homosexual men identified several single nucleotide polymorphisms on chromosome 13, in the region on genes expressed in the diencephalon previously reported as differing in size in men by sexual orientation (LeVay, 1991) and on chromosome 14 in the thyroid stimulating receptor locus (Sanders et al., 2017) further indicating a possible connection between thyroid, thyroid derangements and a subset of men having sex with men. All together, these findings suggest that the relationship between genes, thyroid function and male homosexuality would provide, if confirmed by more robust data, new interesting insights in the biology of male sexual orientation.

8. Conclusion

Many evidences show that alterations in thyroid function can affect both male and female sexual function. The presence of TRs in genitals provide the basis on which to hypothesize a direct effect of THs on sexual functions. However, the effect of THs is not only confined to genitals. In fact several data showed, as in the brain, THs may affect estrogens in the triggering of desire and libido. More studies are required to better delineate the real impact of thyroid in sexual function. Currently, it can be concluded that, despite the apparent, relative disinterestedness of the large majority of clinical endocrinologists dealing with thyroid diseases, demonstrated by a lack of use of psychometric tools designed to discover the presence of a sexual dysfunction (Corona et al., 2006a). THs strongly affect the human sexual function at various levels and that the thyroid gland must be considered, along with the genitals and the brain, a sexual organ.

References

- Aghajanova, L., Lindeberg, M., Carlsson, I.B., Stavreus-Evers, A., Zhang, P., Scott, J.E., Hovatta, O., Skjoldbrand-Sparre, L., 2009. Receptors for thyroid-stimulating hormone and thyroid hormones in human ovarian tissue. *Reprod. Biomed. Online* 18 (3), 337–347.
- Aghajanova, L., Stavreus-Evers, A., Lindeberg, M., Landgren, B.M., Sparre, L.S., Hovatta, O., 2011. Thyroid-stimulating hormone receptor and thyroid hormone receptors are involved in human endometrial physiology. *Fertil. Steril.* 95 (1), 230–237, 237 e231–232.
- Atis, G., Dalkilinc, A., Altuntas, Y., Atis, A., Caskurlu, T., Ergenekon, E., 2010. Sexual dysfunction in women with clinical hypothyroidism and subclinical hypothyroidism. *J. Sex. Med.* 7 (7), 2583–2590.
- Atis, G., Dalkilinc, A., Altuntas, Y., Atis, A., Gurbuz, C., Ofluoglu, Y., Cil, E., Caskurlu, T., 2011. Hyperthyroidism: a risk factor for female sexual dysfunction. *J. Sex. Med.* 8 (8), 2327–2333.
- Ayers, S., Switnicki, M.P., Angajala, A., Lammel, J., Arumanayagam, A.S., Webb, P., 2014. Genome-wide binding patterns of thyroid hormone receptor beta. *PLoS One* 9 (2), e81186.
- Bagamasbad, P., Denver, R.J., 2011. Mechanisms and significance of nuclear receptor auto- and cross-regulation. *Gen. Comp. Endocrinol.* 170 (1), 3–17.
- Balercia, G., Boscaro, M., Lombardo, F., Carosa, E., Lenzi, A., Jannini, E.A., 2007. Sexual symptoms in endocrine diseases: psychosomatic perspectives. *Psychother. Psychosom.* 76 (3), 134–140.
- Barker, S.B., Klitgaard, H.M., 1952. Metabolism of tissues excised from thyroxine-injected rats. *Am. J. Physiol.* 170 (1), 81–86.
- Bassett, J.H., Harvey, C.B., Williams, G.R., 2003. Mechanisms of thyroid hormone receptor-specific nuclear and extra nuclear actions. *Mol. Cell Endocrinol.* 213 (1), 1–11.
- Bauer, M., Goetz, T., Glenn, T., Whybrow, P.C., 2008. The thyroid-brain interaction in thyroid disorders and mood disorders. *J. Neuroendocrinol.* 20 (10), 1101–1114.
- Bernal, J., 2000. Thyroid hormones in brain development and function. In: De Groot, L.J., Chrousos, G., Dungan, K., Feingold, K.R., Grossman, A., Hershman, J.M., Koch, C., Korbonits, M., McLachlan, R., New, M., Purnell, J., Rebar, R., Singer, F., Vinik, A. (Eds.), *Endotext*. South Dartmouth (MA).
- Bernal, J., 2007. Thyroid hormone receptors in brain development and function. *Nat. Clin. Pract. Endocrinol. Metab.* 3 (3), 249–259.
- Bernal, J., Pekonen, F., 1984. Ontogenesis of the nuclear 3,5,3'-triiodothyronine receptor in the human fetal brain. *Endocrinology* 114 (2), 677–679.
- Bhasin, S., Enzlin, P., Coviello, A., Basson, R., 2007. Sexual dysfunction in men and women with endocrine disorders. *Lancet* 369 (9561), 597–611.
- Blanchard, R., Bogaert, A.F., 1996. Biodemographic comparisons of homosexual and heterosexual men in the Kinsey interview data. *Arch. Sex. Behav.* 25 (6), 551–579.
- Bocklandt, S., Horvath, S., Vilain, E., Hamer, D.H., 2006. Extreme skewing of X chromosome inactivation in mothers of homosexual men. *Hum. Genet.* 118 (6), 691–694.
- Brix, T.H., Knudsen, G.P., Kristiansen, M., Kyvik, K.O., Orstavik, K.H., Hegedüs, L., 2005. High frequency of skewed X-chromosome inactivation in females with autoimmune thyroid disease: a possible explanation for the female predisposition to thyroid autoimmunity. *J. Clin. Endocrinol. Metab.* 90 (11), 5949–5953.
- Buzzard, J.J., Morrison, J.R., O'Bryan, M.K., Song, Q., Wreford, N.G., 2000. Developmental expression of thyroid hormone receptors in the rat testis. *Biol. Reprod.* 62 (3), 664–669.
- Cahangirov, A., Cihan, A., Murat, N., Demir, O., Aslan, G., Gidener, S., Esen, A.A., 2011. Investigation of the neural target level of hyperthyroidism in premature ejaculation in a rat model of pharmacologically induced ejaculation. *J. Sex. Med.* 8 (1), 90–96.
- Canale, D., Agostini, M., Giorgilli, G., Caglieresi, C., Scartabelli, G., Nardini, V., Jannini, E.A., Martino, E., Pinchera, A., Macchia, E., 2001. Thyroid hormone receptors in neonatal, prepubertal, and adult rat testis. *J. Androl.* 22 (2), 284–288.
- Carani, C., Isidori, A.M., Granata, A., Carosa, E., Maggi, M., Lenzi, A., Jannini, E.A., 2005. Multicenter study on the prevalence of sexual symptoms in male hypothyroid patients. *J. Clin. Endocrinol. Metab.* 90 (12), 6472–6479.
- Carani, C., Qin, K., Simoni, M., Faustini-Fustini, M., Serpente, S., Boyd, J., Korach, K.S., Simpson, E.R., 1997. Effect of testosterone and estradiol in a man with aromatase deficiency. *N. Engl. J. Med.* 337 (2), 91–95.
- Carosa, E., Di Sante, S., Rossi, S., Castri, A., D'Adamo, F., Gravina, G.L., Ronchi, P., Kostrouch, Z., Dolci, S., Lenzi, A., Jannini, E.A., 2010. Ontogenetic profile of the expression of thyroid hormone receptors in rat and human corpora cavernosa of the penis. *J. Sex. Med.* 7 (4 Pt 1), 1381–1390.
- Carosa, E., Fanelli, A., Ulisse, S., Di Lauro, R., Rall, J.E., Jannini, E.A., 1998. Ciona intestinalis nuclear receptor 1: a member of steroid/thyroid hormone receptor family. *Proc. Natl. Acad. Sci. U. S. A.* 95 (19), 11152–11157.
- Carosa, E., Radico, C., Giansante, N., Rossi, S., D'Adamo, F., Di Stasi, S.M., Lenzi, A., Jannini, E.A., 2005. Ontogenetic profile and thyroid hormone regulation of type-1 and type-8 glucose transporters in rat Sertoli cells. *Int. J. Androl.* 28 (2), 99–106.
- Caturegli, P., Kimura, H., Rocchi, R., Rose, N.R., 2007. Autoimmune thyroid diseases. *Curr. Opin. Rheumatol.* 19 (1), 44–48.
- Chatonnet, F., Guyot, R., Picou, F., Bondesson, M., Flamant, F., 2012. Genome-wide search reveals the existence of a limited number of thyroid hormone receptor alpha target genes in cerebellar neurons. *PLoS One* 7 (5), e37073.
- Cihan, A., Demir, O., Demir, T., Aslan, G., Comlekci, A., Esen, A., 2009a. The relationship between premature ejaculation and hyperthyroidism. *J. Urol.* 181 (3), 1273–1280.
- Cihan, A., Murat, N., Demir, O., Aslan, G., Demir, T., Gidener, S., Esen, A.A., 2009b. An experimental approach to the interrelationship between hyperthyroidism and ejaculation latency time in male rats. *J. Urol.* 181 (2), 907–912.
- Ciocca, G., Limoncin, E., Carosa, E., Di Sante, S., Gravina, G.L., Mollaioli, D., Gianfrilli, D., Lenzi, A., Jannini, E.A., 2016. Is testosterone a food for the brain? *Sex. Med. Rev.* 4 (1), 15–25.
- Corona, G., Isidori, A.M., Aversa, A., Burnett, A.L., Maggi, M., 2016. Endocrinologic control of Men's sexual desire and arousal/erection. *J. Sex. Med.* 13 (3), 317–337.
- Corona, G., Jannini, E.A., Lotti, F., Boddi, V., De Vita, G., Forti, G., Lenzi, A., Mannucci, E., Maggi, M., 2011. Premature and delayed ejaculation: two ends of a single continuum influenced by hormonal milieu. *Int. J. Androl.* 34 (1), 41–48.
- Corona, G., Jannini, E.A., Maggi, M., 2006a. Inventories for male and female sexual dysfunctions. *Int. J. Impot. Res.* 18 (3), 236–250.
- Corona, G., Jannini, E.A., Vignozzi, L., Rastrelli, G., Maggi, M., 2012a. The hormonal control of ejaculation. *Nat. Rev. Urol.* 9 (9), 508–519.
- Corona, G., Mannucci, E., Petrone, L., Fisher, A.D., Balercia, G., De Scisciolo, G., Pizzocaro, A., Giommi, R., Chiarini, V., Forti, G., Maggi, M., 2006b. Psychobiological correlates of delayed ejaculation in male patients with sexual dysfunctions. *J. Androl.* 27 (3), 453–458.
- Corona, G., Petrone, L., Mannucci, E., Jannini, E.A., Mansani, R., Magini, A., Giommi, R., Forti, G., Maggi, M., 2004. Psycho-biological correlates of rapid ejaculation in patients attending an andrologic unit for sexual dysfunctions. *Eur. Urol.* 46 (5), 615–622.
- Corona, G., Wu, F.C., Forti, G., Lee, D.M., O'Connor, D.B., O'Neill, T.W., Pendleton, N., Bartfai, G., Boonen, S., Casanueva, F.F., Finn, J.D., Giwercman, A., Han, T.S., Huhtaniemi, I.T., Kula, K., Lean, M.E., Punab, M., Vanderschueren, D., Jannini, E.A., Mannucci, E., Maggi, M., Group, E.S., 2012b. Thyroid hormones and male sexual function. *Int. J. Androl.* 35 (5), 668–679.
- Costa-e-Sousa, R.H., Hollenberg, A.N., 2012. Minireview: the neural regulation of the hypothalamic-pituitary-thyroid axis. *Endocrinology* 153 (9), 4128–4135.
- De Paul, A.L., Mukdsi, J.H., Pellizas, C.G., Montesinos, M., Gutierrez, S., Susperreguy, S., Del Rio, A., Maldonado, C.A., Torres, A.I., 2008. Thyroid hormone receptor alpha 1-beta 1 expression in epididymal epithelium from euthyroid and hypothyroid rats. *Histochem Cell Biol.* 129 (5), 631–642.
- Dellovade, T.L., Chan, J., Vennstrom, B., Forrest, D., Pfaff, D.W., 2000. The two thyroid

- hormone receptor genes have opposite effects on estrogen-stimulated sex behaviors. *Nat. Neurosci.* 3 (5), 472–475.
- Dellovade, T.L., Zhu, Y.S., Krey, L., Pfaff, D.W., 1996. Thyroid hormone and estrogen interact to regulate behavior. *Proc. Natl. Acad. Sci. U. S. A.* 93 (22), 12581–12586.
- Deputy, N.P., Boehmer, U., 2010. Determinants of body weight among men of different sexual orientation. *Prev. Med.* 51 (2), 129–131.
- Di Sante, S., Mollaioli, D., Gravina, G.L., Ciocca, G., Limoncin, E., Carosa, E., Lenzi, A., Jannini, E.A., 2016. Epidemiology of delayed ejaculation. *Transl. Androl. Urol.* 5 (4), 541–548.
- Dittrich, R., Beckmann, M.W., Oppelt, P.G., Hoffmann, I., Lotz, L., Kuwert, T., Mueller, A., 2011. Thyroid hormone receptors and reproduction. *J. Reprod. Immunol.* 90 (1), 58–66.
- Dobbing, J., Sands, J., 1970. Timing of neuroblast multiplication in developing human brain. *Nature* 226 (5246), 639–640.
- Fedail, J.S., Zheng, K., Wei, Q., Kong, L., Shi, F., 2014. Roles of thyroid hormones in follicular development in the ovary of neonatal and immature rats. *Endocrine* 46 (3), 594–604.
- Francavilla, S., Cordeschi, G., Properzi, G., Di Cicco, L., Jannini, E.A., Palmero, S., Fugassa, E., Loras, B., D'Armiento, M., 1991. Effect of thyroid hormone on the pre- and post-natal development of the rat testis. *J. Endocrinol.* 129 (1), 35–42.
- Friesema, E.C., Ganguly, S., Abdalla, A., Manning Fox, J.E., Halestrap, A.P., Visser, T.J., 2003. Identification of monocarboxylate transporter 8 as a specific thyroid hormone transporter. *J. Biol. Chem.* 278 (41), 40128–40135.
- Frisch, M., Zdravkovic, S., 2010. Body size at birth and same-sex marriage in young adulthood. *Arch. Sex. Behav.* 39 (1), 117–123.
- Frisch, M., Nielsen, N.M., Pedersen, B.V., 2014. Same-sex marriage, autoimmune thyroid gland dysfunction and other autoimmune diseases in Denmark 1989–2008. *Eur. J. Epidemiol.* 29 (1), 63–71.
- Fumel, B., Froment, P., Holzenberger, M., Livera, G., Monget, P., Fouchecourt, S., 2015. Expression of dominant-negative thyroid hormone receptor alpha1 in Leydig and Sertoli cells demonstrates no additional defect compared with expression in Sertoli cells only. *PLoS One* 10 (3), e0119392.
- Ghizzani, A., Razzi, S., Fava, A., Sartini, A., Picucci, K., Petraglia, F., 2003. Management of sexual dysfunctions in women. *J. Endocrinol. Invest.* 26 (3 Suppl. 1), 137–138.
- Graziottin, A., 2003. The challenge of sexual medicine for women: overcoming cultural and educational limits and gender biases. *J. Endocrinol. Invest.* 26 (3 Suppl. 1), 139–142.
- Grontved, L., Waterfall, J.J., Kim, D.W., Baek, S., Sung, M.H., Zhao, L., Park, J.W., Nielsen, R., Walker, R.L., Zhu, Y.J., Meltzer, P.S., Hager, G.L., Cheng, S.Y., 2015. Transcriptional activation by the thyroid hormone receptor through ligand-dependent receptor recruitment and chromatin remodelling. *Nat. Commun.* 6, 7048.
- Hapon, M.B., Gamarra-Luques, C., Jahn, G.A., 2010. Short term hypothyroidism affects ovarian function in the cycling rat. *Reprod. Biol. Endocrinol.* 8, 14.
- Heinrich, T.W., Grahm, G., 2003. Hypothyroidism presenting as psychosis: myxedema madness revisited. *Prim. Care Companion J. Clin. Psychiatry* 5 (6), 260–266.
- Holsberger, D.R., Cooke, P.S., 2005. Understanding the role of thyroid hormone in Sertoli cell development: a mechanistic hypothesis. *Cell Tissue Res.* 322 (1), 133–140.
- Holsberger, D.R., Kiesewetter, S.E., Cooke, P.S., 2005. Regulation of neonatal Sertoli cell development by thyroid hormone receptor alpha1. *Biol. Reprod.* 73 (3), 396–403.
- Hu, S., Pattatucci, A.M., Patterson, C., Li, L., Fulker, D.W., Cherny, S.S., Kruglyak, L., Hamer, D.H., 1995. Linkage between sexual orientation and chromosome Xq28 in males but not in females. *Nat. Genet.* 11 (3), 248–256.
- Hu, C.L., Wu, Y.D., Liu, H.T., Qin, W.B., Wang, G.Z., 2009. Effect of thyroid hormone on the contents of NOS and CO in the penile corpus cavernosum of rats. *Zhonghua Nan Ke Xue* 15 (1), 37–40.
- Hulchiy, M., Zhang, H., Cline, J.M., Hirschberg, A.L., Sahlin, L., 2012. Receptors for thyrotropin-releasing hormone, thyroid-stimulating hormone, and thyroid hormones in the macaque uterus: effects of long-term sex hormone treatment. *Menopause* 19 (11), 1253–1259.
- Isidori, A.M., Pozza, C., Esposito, K., Giugliano, D., Morano, S., Vignozzi, L., Corona, G., Lenzi, A., Jannini, E.A., 2010. Development and validation of a 6-item version of the female sexual function index (FSFI) as a diagnostic tool for female sexual dysfunction. *J. Sex. Med.* 7 (3), 1139–1146.
- Iskaros, J., Pickard, M., Evans, I., Sinha, A., Hardiman, P., Ekins, R., 2000. Thyroid hormone receptor gene expression in first trimester human fetal brain. *J. Clin. Endocrinol. Metab.* 85 (7), 2620–2623.
- Jannini, E.A., Blanchard, R., Camperio-Ciani, A., Bancroft, J., 2010. Male homosexuality: nature or culture? *J. Sex. Med.* 7 (10), 3245–3253.
- Jannini, E.A., Buisson, O., Rubio-Casillas, A., 2014. Beyond the G-spot: clitoral-ovarian complex anatomy in female orgasm. *Nat. Rev. Urol.* 11 (9), 531–538.
- Jannini, E.A., Carosa, E., Rucci, N., Screponi, E., D'Armiento, M., 1999. Ontogeny and regulation of variant thyroid hormone receptor isoforms in developing rat testis. *J. Endocrinol. Invest.* 22 (11), 843–848.
- Jannini, E.A., Ciocca, G., Limoncin, E., Mollaioli, D., Di Sante, S., Gianfrilli, D., Lombardo, F., Lenzi, A., 2015. Premature ejaculation: old story, new insights. *Fertil. Steril.* 104 (5), 1061–1073.
- Jannini, E.A., Crescenzi, A., Rucci, N., Screponi, E., Carosa, E., de Matteis, A., Macchia, E., d'Amati, G., D'Armiento, M., 2000. Ontogenetic pattern of thyroid hormone receptor expression in the human testis. *J. Clin. Endocrinol. Metab.* 85 (9), 3453–3457.
- Jannini, E.A., Dolci, S., Ulisse, S., Nikodem, V.M., 1994. Developmental regulation of the thyroid hormone receptor alpha 1 mRNA expression in the rat testis. *Mol. Endocrinol.* 8 (1), 89–96.
- Jannini, E.A., Lenzi, A., 2003. Introduction to the integrated model: medical, surgical and psychological therapies for the couple. *J. Endocrinol. Invest.* 26 (3 Suppl. 1), 128–131.
- Jannini, E.A., Maggi, M., Lenzi, A., 2011. Evaluation of premature ejaculation. *J. Sex. Med.* 8 (Suppl. 4), 328–334.
- Jannini, E.A., Mitsuhashi, T., Nikodem, V.M., 1992. Developmental expression of mRNAs from a rat C-erbA genomic locus. *Biochem. Biophys. Res. Commun.* 184 (2), 739–745.
- Jannini, E.A., Olivieri, M., Francavilla, S., Gulino, A., Ziparo, E., D'Armiento, M., 1990. Ontogenesis of the nuclear 3,5,3'-triiodothyronine receptor in the rat testis. *Endocrinology* 126 (5), 2521–2526.
- Jannini, E.A., Ulisse, S., D'Armiento, M., 1995. Thyroid hormone and male gonadal function. *Endocr. Rev.* 16 (4), 443–459.
- Jannini, E.A., Ulisse, S., Piersanti, D., Carosa, E., Muzi, P., Lazar, J., D'Armiento, M., 1993. Early thyroid hormone treatment in rats increases testis size and germ cell number. *Endocrinology* 132 (6), 2726–2728.
- Katz, D., Reginato, M.J., Lazar, M.A., 1995. Functional regulation of thyroid hormone receptor variant TR alpha 2 by phosphorylation. *Mol. Cell Biol.* 15 (5), 2341–2348.
- Kirkland, J.L., Mukku, V., Hardy, M., Young, R., 1983. Evidence for triiodothyronine receptors in human endometrium and myometrium. *Am. J. Obstet. Gynecol.* 146 (4), 380–383.
- Koenig, R.J., Lazar, M.A., Hodin, R.A., Brent, G.A., Larsen, P.R., Chin, W.W., Moore, D.D., 1989. Inhibition of thyroid hormone action by a non-hormone binding c-erbA protein generated by alternative mRNA splicing. *Nature* 337 (6208), 659–661.
- Kostrouch, Z., Kostrouchova, M., Love, W., Jannini, E., Piatigorsky, J., Rall, J.E., 1998. Retinoic acid X receptor in the diploblast, *Tripedalia cystophora*. *Proc. Natl. Acad. Sci. U. S. A.* 95 (23), 13442–13447.
- Krassas, G.E., 2000. Thyroid disease and female reproduction. *Fertil. Steril.* 74 (6), 1063–1070.
- Krassas, G.E., Poppe, K., Glinoe, D., 2010. Thyroid function and human reproductive health. *Endocr. Rev.* 31 (5), 702–755.
- Krassas, G.E., Tziomalos, K., Papadopoulos, F., Pontikides, N., Perros, P., 2008. Erectile dysfunction in patients with hyper- and hypothyroidism: how common and should we treat? *J. Clin. Endocrinol. Metab.* 93 (5), 1815–1819.
- Krysiak, R., Drosdzol-Cop, A., Skrzypulec-Plinta, V., Okopien, B., 2016. Sexual function and depressive symptoms in young women with thyroid autoimmunity and subclinical hypothyroidism. *Clin. Endocrinol. (Oxf)* 84 (6), 925–931.
- Krysiak, R., Szkrobka, W., Okopien, B., 2017. The effect of l-thyroxine treatment on sexual function and depressive symptoms in men with autoimmune hypothyroidism. *Pharmacol. Rep.* 69 (3), 432–437.
- Lazar, M.A., 2003. Thyroid hormone action: a binding contract. *J. Clin. Invest.* 112 (4), 497–499.
- Lazar, M.A., Hodin, R.A., Darling, D.S., Chin, W.W., 1988. Identification of a rat c-erbA alpha-related protein which binds deoxyribonucleic acid but does not bind thyroid hormone. *Mol. Endocrinol.* 2 (10), 893–901.
- LeVay, S., 1991. A difference in hypothalamic structure between heterosexual and homosexual men. *Science* 253 (5023), 1034–1037.
- Luo, M., Faure, R., Dussault, J.H., 1986. Ontogenesis of nuclear T3 receptors in primary cultured astrocytes and neurons. *Brain Res.* 381 (2), 275–280.
- Maggi, M., Buvat, J., Corona, G., Guay, A., Torres, L.O., 2013. Hormonal causes of male sexual dysfunctions and their management (hyperprolactinemia, thyroid disorders, GH disorders, and DHEA). *J. Sex. Med.* 10 (3), 661–677.
- McMahon, C.G., Jannini, E., Waldinger, M., Rowland, D., 2013. Standard operating procedures in the disorders of orgasm and ejaculation. *J. Sex. Med.* 10 (1), 204–229.
- Meikle, A.W., 2004. The interrelationships between thyroid dysfunction and hypogonadism in men and boys. *Thyroid* 14 (Suppl. 1), S17–S25.
- Mendis-Handagama, S.M., Siril Ariyaratne, H.B., 2005. Leydig cells, thyroid hormones and steroidogenesis. *Indian J. Exp. Biol.* 43 (11), 939–962.
- Mendoza, A., Hollenberg, A.N., 2017. New insights into thyroid hormone action. *Pharmacol. Ther.* 173, 135–145.
- Mitsuhashi, T., Tennyson, G., Nikodem, V., 1988. Nucleotide sequence of novel cDNAs generated by alternative splicing of a rat thyroid hormone receptor gene transcript. *Nucleic Acids Res.* 16 (12), 5697.
- Moog NK, Entringer S, Heim C, Wadhwa PD, Kathmann N, Buss C Influence of maternal thyroid hormones during gestation on fetal brain development. *Neuroscience* 342: 68–100
- Morgan, M.A., Dellovade, T.L., Pfaff, D.W., 2000. Effect of thyroid hormone administration on estrogen-induced sex behavior in female mice. *Horm. Behav.* 37 (1), 15–22.
- Navas, P.B., Redondo, A.L., Cuello-Carrion, F.D., Roig, L.M., Valdez, S.R., Jahn, G.A., Hapon, M.B., 2014. Luteal expression of thyroid hormone receptors during gestation and postpartum in the rat. *Thyroid* 24 (6), 1040–1050.
- Nelson, E.R., Allan, E.R., Pang, F.Y., Habibi, H.R., 2011. Auto-regulation of thyroid hormone receptors in the goldfish ovary and testis. *Gen. Comp. Endocrinol.* 172 (1), 50–55.
- Niazi AK, Kalra S, Irfan A, Islam a thyroidology over the ages. *Indian J. Endocrinol. Metab.* 15(Suppl. 2): S121–S126
- Ogawa, S., Chan, J., Chester, A.E., Gustafsson, J.A., Korach, K.S., Pfaff, D.W., 1999. Survival of reproductive behaviors in estrogen receptor beta gene-deficient

- (betaERKO) male and female mice. *Proc. Natl. Acad. Sci. U. S. A.* 96 (22), 12887–12892.
- Ogawa, S., Chester, A.E., Hewitt, S.C., Walker, V.R., Gustafsson, J.A., Smithies, O., Korach, K.S., Pfaff, D.W., 2000. Abolition of male sexual behaviors in mice lacking estrogen receptors alpha and beta (alpha beta ERKO). *Proc. Natl. Acad. Sci. U. S. A.* 97 (26), 14737–14741.
- Ogawa, S., Eng, V., Taylor, J., Lubahn, D.B., Korach, K.S., Pfaff, D.W., 1998. Roles of estrogen receptor-alpha gene expression in reproduction-related behaviors in female mice. *Endocrinology* 139 (12), 5070–5081.
- Ogawa, S., Lubahn, D.B., Korach, K.S., Pfaff, D.W., 1997. Behavioral effects of estrogen receptor gene disruption in male mice. *Proc. Natl. Acad. Sci. U. S. A.* 94 (4), 1476–1481.
- Oppo, A., Franceschi, E., Atzeni, F., Taberlet, A., Mariotti, S., 2011. Effects of hyperthyroidism, hypothyroidism, and thyroid autoimmunity on female sexual function. *J. Endocrinol. Invest.* 34 (6), 449–453.
- Owen, P.J., Sabit, R., Lazarus, J.H., 2007. Thyroid disease and vascular function. *Thyroid* 17 (6), 519–524.
- Ozdemirci, S., Yildiz, F., Utkan, T., Ulak, G., Cetinaslan, B., Erden, F., Gacar, N., 2001. Impaired neurogenic and endothelium-dependent relaxant responses of corpus cavernosum smooth muscle from hyperthyroid rabbits. *Eur. J. Pharmacol.* 428 (1), 105–111.
- Palmero, S., De Marco, P., Fugassa, E., 1995. Thyroid hormone receptor beta mRNA expression in Sertoli cells isolated from prepubertal testis. *J. Mol. Endocrinol.* 14 (1), 131–134.
- Palmero, S., Maggiani, S., Fugassa, E., 1988. Nuclear triiodothyronine receptors in rat Sertoli cells. *Mol. Cell Endocrinol.* 58 (2–3), 253–256.
- Pasquali, D., Maiorino, M.I., Renzullo, A., Bellastella, G., Accardo, G., Esposito, D., Barbatto, F., Esposito, K., 2013. Female sexual dysfunction in women with thyroid disorders. *J. Endocrinol. Invest.* 36 (9), 729–733.
- Poppe, K., Velkeniers, B., 2004. Female infertility and the thyroid. *Best. Pract. Res. Clin. Endocrinol. Metab.* 18 (2), 153–165.
- Ramadoss, P., Abraham, B.J., Tsai, L., Zhou, Y., Costa-e-Sousa, R.H., Ye, F., Bilban, M., Zhao, K., Hollenberg, A.N., 2014. Novel mechanism of positive versus negative regulation by thyroid hormone receptor beta1 (TRbeta1) identified by genome-wide profiling of binding sites in mouse liver. *J. Biol. Chem.* 289 (3), 1313–1328.
- Rodriguez-Castelan, J., Anaya-Hernandez, A., Mendez-Tepepa, M., Martinez-Gomez, M., Castelan, F., Cuevas-Romero, E., 2017. Distribution of thyroid hormone and thyrotropin receptors in reproductive tissues of adult female rabbits. *Endocr. Res.* 42 (1), 59–70.
- Romano, R., Jannini, E.A., Pepe, M., Grimaldi, A., Olivieri, M., Spennati, P., Cappa, F., D'Armiento, M., 1991. The effects of iodoprophyllaxis on thyroid size during pregnancy. *Am. J. Obstet. Gynecol.* 164 (2), 482–485.
- Romano, R.M., Gomes, S.N., Cardoso, N.C., Schiessl, L., Romano, M.A., Oliveira, C.A., 2017. New insights for male infertility revealed by alterations in spermatid function and differential testicular expression of thyroid-related genes. *Endocrine* 55 (2), 607–617.
- Sabuncuoglu, O., 2015. High rates of same-sex attraction/gender nonconformity in the offspring of mothers with thyroid dysfunction during pregnancy: proposal of prenatal thyroid model. *Ment. Illn.* 7 (2), 5810.
- Sait Gonen, M., Kisakol, G., Savas Cilli, A., Dikbas, O., Gungor, K., Inal, A., Kaya, A., 2004. Assessment of anxiety in subclinical thyroid disorders. *Endocr. J.* 51 (3), 311–315.
- Sanders, A.R., Beecham, G.W., Guo, S., Dawood, K., Rieger, G., Badner, J.A., Gershon, E.S., Krishnappa, R.S., Kolundzija, A.B., Duan, J., MGS Collaboration, Gejman, P.V., Bailey, J.M., Martin, E.R., 2017. Genome-Wide Association Study of Male Sexual Orientation. *Sci. Rep.* 7 (1), 16950.
- Sansone, A., Romanelli, F., Jannini, E.A., Lenzi, A., 2015. Hormonal correlations of premature ejaculation. *Endocrine* 49 (2), 333–338.
- Sap, J., Munoz, A., Damm, K., Goldberg, Y., Ghysdael, J., Leutz, A., Beug, H., Vennstrom, B., 1986. The c-erb-A protein is a high-affinity receptor for thyroid hormone. *Nature* 324 (6098), 635–640.
- Schulster, M., Bernie, A.M., Ramasamy, R., 2016. The role of estradiol in male reproductive function. *Asian J. Androl.* 18 (3), 435–440.
- Schwartz, C.E., May, M.M., Carpenter, N.J., Rogers, R.C., Martin, J., Bialer, M.G., Ward, J., Sanabria, J., Marsa, S., Lewis, J.A., Echeverri, R., Lubs, H.A., Voeller, K., Simensen, R.J., Stevenson, R.E., 2005. Allan-Herndon-Dudley syndrome and the monocarboxylate transporter 8 (MCT8) gene. *Am. J. Hum. Genet.* 77 (1), 41–53.
- Schweizer, U., Johannes, J., Bayer, D., Braun, D., 2014. Structure and function of thyroid hormone plasma membrane transporters. *Eur. Thyroid. J.* 3 (3), 143–153.
- Shifren, J.L., Gass, M.L., 2014. The North American Menopause Society recommendations for clinical care of midlife women. *Menopause* 21 (10), 1038–1062.
- Stavreus Evers, A., 2012. Paracrine interactions of thyroid hormones and thyroid stimulation hormone in the female reproductive tract have an impact on female fertility. *Front. Endocrinol. (Lausanne)* 3, 50.
- Sun, Z., Feng, D., Fang, B., Mullican, S.E., You, S.H., Lim, H.W., Everett, L.J., Nabel, C.S., Li, Y., Selvakumar, V., Won, K.J., Lazar, M.A., 2013. Deacetylase-independent function of HDAC3 in transcription and metabolism requires nuclear receptor corepressor. *Mol. Cell* 52 (6), 769–782.
- Tagami, T., Nakamura, H., Sasaki, S., Mori, T., Yoshioka, H., Yoshida, H., Imura, H., 1990. Immunohistochemical localization of nuclear 3,5,3'-triiodothyronine receptor proteins in rat tissues studied with antiserum against C-ERB A/T3 receptor. *Endocrinology* 127 (4), 1727–1734.
- Ulisse, S., Jannini, E.A., Carosa, E., Piersanti, D., Graziano, F.M., D'Armiento, M., 1994. Inhibition of aromatase activity in rat Sertoli cells by thyroid hormone. *J. Endocrinol.* 140 (3), 431–436.
- Ulisse, S., Jannini, E.A., Pepe, M., De Matteis, S., D'Armiento, M., 1992. Thyroid hormone stimulates glucose transport and GLUT1 mRNA in rat Sertoli cells. *Mol. Cell Endocrinol.* 87 (1–3), 131–137.
- Ulisse, S., Rucci, N., Piersanti, D., Carosa, E., Graziano, F.M., Pavan, A., Ceddia, P., Arizzi, M., Muzi, P., Cironi, L., Gnassi, L., D'Armiento, M., Jannini, E.A., 1998. Regulation by thyroid hormone of the expression of basement membrane components in rat prepubertal Sertoli cells. *Endocrinology* 139 (2), 741–747.
- Vasudevan, N., Morgan, M., Pfaff, D., Ogawa, S., 2013. Distinct behavioral phenotypes in male mice lacking the thyroid hormone receptor alpha1 or beta isoforms. *Horm. Behav.* 63 (5), 742–751.
- Vasudevan, N., Ogawa, S., Pfaff, D., 2002. Estrogen and thyroid hormone receptor interactions: physiological flexibility by molecular specificity. *Physiol. Rev.* 82 (4), 923–944.
- Veronelli, A., Masu, A., Ranieri, R., Rognoni, C., Laneri, M., Pontiroli, A.E., 2006. Prevalence of erectile dysfunction in thyroid disorders: comparison with control subjects and with obese and diabetic patients. *Int. J. Impot. Res.* 18 (1), 111–114.
- Veronelli, A., Mauri, C., Zecchini, B., Peca, M.G., Turri, O., Valitutti, M.T., dall'Asta, C., Pontiroli, A.E., 2009. Sexual dysfunction is frequent in premenopausal women with diabetes, obesity, and hypothyroidism, and correlates with markers of increased cardiovascular risk. A preliminary report. *J. Sex. Med.* 6 (6), 1561–1568.
- Wagner, M.S., Wajner, S.M., Maia, A.L., 2009. Is there a role for thyroid hormone on spermatogenesis? *Microsc. Res. Tech.* 72 (11), 796–808.
- Wajner, S.M., Wagner, M.S., Maia, A.L., 2009. Clinical implications of altered thyroid status in male testicular function. *Arq. Bras. Endocrinol. Metabol.* 53 (8), 976–982.
- Wakim, A.N., Paljug, W.R., Jasnosc, K.M., Alhakim, N., Brown, A.B., Burholt, D.R., 1994a. Thyroid hormone receptor messenger ribonucleic acid in human granulosa and ovarian stromal cells. *Fertil. Steril.* 62 (3), 531–534.
- Wakim, A.N., Polizotto, S.L., Buffo, M.J., Marrero, M.A., Burholt, D.R., 1993. Thyroid hormones in human follicular fluid and thyroid hormone receptors in human granulosa cells. *Fertil. Steril.* 59 (6), 1187–1190.
- Wakim, A.N., Polizotto, S.L., Burholt, D.R., 1994b. Alpha-1 and beta-1 thyroid hormone receptors on human granulosa cells. *Recent Prog. Horm. Res.* 49, 377–381.
- Waldinger, M.D., Zwinderman, A.H., Olivier, B., Schweitzer, D.H., 2005. Thyroid-stimulating hormone assessments in a Dutch cohort of 620 men with lifelong premature ejaculation without erectile dysfunction. *J. Sex. Med.* 2 (6), 865–870.
- Weinberger, C., Thompson, C.C., Ong, E.S., Lebo, R., Gruol, D.J., Evans, R.M., 1986. The c-erb-A gene encodes a thyroid hormone receptor. *Nature* 324 (6098), 641–646.
- Wiegel, M., Meston, C., Rosen, R., 2005. The female sexual function index (FSFI): cross-validation and development of clinical cutoff scores. *J. Sex. Marital Ther.* 31 (1), 1–20.
- You, S.H., Lim, H.W., Sun, Z., Broache, M., Won, K.J., Lazar, M.A., 2013. Nuclear receptor co-repressors are required for the histone-deacetylase activity of HDAC3 in vivo. *Nat. Struct. Mol. Biol.* 20 (2), 182–187.
- Yusta, B., Besnard, F., Ortiz-Caro, J., Pascual, A., Aranda, A., Sarlieve, L., 1988. Evidence for the presence of nuclear 3,5,3'-triiodothyronine receptors in secondary cultures of pure rat oligodendrocytes. *Endocrinology* 122 (5), 2278–2284.
- Zhang, S.S., Carrillo, A.J., Darling, D.S., 1997. Expression of multiple thyroid hormone receptor mRNAs in human oocytes, cumulus cells, and granulosa cells. *Mol. Hum. Reprod.* 3 (7), 555–562.