



Title	Molecular identification and characterization of piroplasm species in Hokkaido sika deer (<i>Cervus nippon yesoensis</i>), Japan
Author(s)	Elbaz, Elzahara; Moustafa, Mohamed Abdallah Mohamed; Lee, Kyunglee; Mohamed, Wessam Mohamed Ahmed; Nakao, Ryo; Shimozuru, Michito; Sashika, Mariko; Younis, Emad Elsayed Ahmed; El-Khodery, Sabry Ahmed; Tsubota, Toshio
Citation	Ticks and Tick-Borne Diseases, 8(5), 802-807 https://doi.org/10.1016/j.ttbdis.2017.06.007
Issue Date	2017-08
Doc URL	http://hdl.handle.net/2115/71113
Rights	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Manuscript.pdf



[Instructions for use](#)

Title: Molecular identification and characterization of piroplasm species in Hokkaido	1
sika deer (<i>Cervus nippon yesoensis</i>), Japan	2
Elzahara Elbaz ^{a1*} , Mohamed Abdallah Mohamed Moustafa ^{a2*} , Kyunglee Lee ^b ,	3
Wessam Mohamed Ahmed Mohamed ^c , Ryo Nakao ^d , Michito Shimozuru ^a , Mariko	4
Sashika ^a , Emad Elsayed Ahmed Younis ^e , Sabry Ahmed El-khodery ^e , Kyoko	5
Hayashida ^f , Chihiro Sugimoto ^f , Toshio Tsubota ^a	6
^a Laboratory of Wildlife Biology and Medicine, Department of Environmental	7
Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University,	8
Sapporo, Japan.	9
^b Cetacean Research Institute, National Institute of Fisheries Science, Ulsan, Republic	10
of Korea.	11
^c Division of bioinformatics, Hokkaido University Research Center for Zoonosis	12
Control, Sapporo, Hokkaido, Japan.	13
^d Laboratory of Parasitology, Department of Disease Control, Graduate School of	14
Veterinary Medicine, Hokkaido University, Sapporo, Japan.	15
^e Department of Internal Medicine and Infectious Diseases, Faculty of Veterinary	16
Medicine, Mansoura University, Mansoura, Egypt.	17
^f Division of Collaboration and Education, Hokkaido University Research Center for	18
Zoonosis Control, Sapporo, Japan.	19

¹ Permanent address: Department of Internal Medicine and Infectious Diseases, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

² Permanent address: Department of Animal Medicine, Faculty of Veterinary Medicine, South Valley University, Qena, Egypt.

* These first authors contributed equally to this article.

Corresponding author:	20
Toshio Tsubota	21
Laboratory of wildlife biology and medicine, department of environmental veterinary	22
sciences, graduate school of veterinary medicine, Hokkaido University, Sapporo,	23
Japan.	24
Address: 060-0818, North 18 West 9 Sapporo, Hokkaido, Japan.	25
Email: tsubota@vetmed.hokudai.ac.jp	26
	27
	28
	29
	30
	31
	32
	33
	34
	35
	36
	37
	38
	39
	40
	41
	42
	43
	44

Abstract 45

Babesia and *Theileria* species are tick-borne protozoan parasites that have a 46
veterinary and zoonotic importance. In order to investigate the prevalence and genetic 47
diversity of these parasites, a total of 269 sika deer blood DNA samples collected 48
from Hokkaido, Japan, were examined for *Babesia* and *Theileria* species by touch- 49
down PCR targeting the 18S rRNA gene. Reverse line blot (RLB) hybridization was 50
then used to detect 12 piroplasm species. The results revealed that 95.5 % (257/269), 51
94.1 % (253/269), 14.1 % (38/269), 87.7 % (236/269) and 11.5 % (31/269) of the 52
examined PCR products hybridized with the probes which were designed to detect all 53
Babesia and *Theileria* spp., all *Theileria* spp., all *Babesia* spp., *Theileria* sp. Thrivae 54
and *Babesia divergens*-like, respectively. The 18S rRNA gene partial sequences were 55
divided into *Theileria* sp. Thrivae, *T. capreoli*, *B. divergens*-like and an undescribed 56
Babesia species. This study showed the first detection of the undescribed *Babesia* sp. 57
from Japan. Therefore, more studies are required to understand the ecology of the 58
newly detected tick-borne pathogens in Hokkaido. 59

Keywords 60

Babesia; *Theileria*; Sika deer; Reverse line blot. 61

62

63

64

65

66

Introduction	67
<i>Babesia</i> and <i>Theileria</i> species are tick-borne protozoan parasites that have a	68
veterinary and zoonotic importance (Mans et al., 2015). <i>Babesia</i> and <i>Theileria</i> spp.	69
are obligatory intracellular parasites and genetically close to each other (Homer et al.,	70
2000), however, both have different life cycles. In the mammalian host, <i>Babesia</i> spp.	71
sensu stricto infect erythrocytes (Schreeg et al., 2016), while <i>Theileria</i> spp. infect both	72
leukocytes and erythrocytes (Li et al., 2014).	73
In cattle, East Coast fever and tropical theileriosis are caused by <i>T. parva</i> and <i>T.</i>	74
<i>annulata</i> , respectively, and exhibit fever, enlargement of lymph nodes and weight loss	75
(Mans et al., 2015; Olds et al., 2016). In addition, <i>B. bovis</i> and <i>B. bigemina</i> are highly	76
pathogenic in cattle and cause bovine babesiosis, a diseases which shows fever,	77
hemolytic anemia and hemoglobinuria (Decaro et al., 2013). <i>Babesia ovis</i> , <i>B. motasi</i>	78
and <i>B. crassa</i> cause sever disease in sheep which is characterized by anemia, fever,	79
icterus and hemoglobinuria (Hashemi-Fesharki, 1997).	80
To the best of our knowledge, there is no published description of clinical	81
manifestations in sika deer (<i>Cervus nippon yesoensis</i>) with piroplasmosis. However,	82
clinical piroplasmosis may occur in the case of exposure to high stressful conditions	83
such as malnutrition (Yabsley et al., 2005). The clinical symptoms of theileriosis in a	84
white-tailed deer (<i>Odocoileus virginianus</i>) fawn were previously reported from the	85
USA and included icterus and anemia (Yabsley et al., 2005).	86
One of the zoonotic <i>Babesia</i> spp. is <i>B. divergens</i> , which has been reported from	87
several deer species including reindeer, roe deer and red deer in Europe (Michel et al.,	88
2014). Recently, another zoonotic <i>B. divergens</i> -like has been detected in sika deer in	89
Japan (Zamoto-Niikura et al., 2014). Whereas, many <i>Theileria</i> spp. have been	90

detected in cervids, for example *T. cervi*, which is a non-pathogenic *Theileria* sp. that 91
 has been reported from white-tailed deer and elk (*Cervus canadensis*) in USA and 92
 Canada (Chae et al., 1999). However, another *T. cervi*, which is genetically separated 93
 from *T. cervi* in the USA and Canada, has been found in sika deer from China (Liu et 94
 al., 2016), but has been reported as *Theileria* sp. Thirivae in Japan (Watanabe et al., 95
 2016). Nevertheless, all *Theileria* spp. have not been recognized as zoonotic 96
 pathogens (Zanet et al., 2014). 97

The sika deer (*Cervus nippon*) is widely distributed in Asia and acts as a reproductive 98
 host for several tick species such as *Haemaphysalis longicornis*, *H. flava*, *H. Yeni*, *H.* 99
megaspinosa, *H. Formosensis*, *H. kitaokai* and *Ixodes ovatus* were detected on sika 100
 deer in Japan (Inokuma et al., 2002). In Japan, sika deer populations have an 101
 important role in maintaining several tick-borne pathogens, such as *Anaplasma* sp. 102
Ap-sd (Moustafa et al., 2015), *Theileria* sp. Thirivae (Watanabe et al., 2016), *B.* 103
divergens-like (Zamoto-Niikura et al., 2014) and *Borrelia lonestari*-like (Lee et al., 104
 2014) . In Hokkaido, sika deer population density has been expanding (Ijima et al., 105
 2015) and they live in close proximity to human settlements and livestock. 106
 Consequently, sika deer may facilitate the transmission of several tick-borne 107
 pathogens to both humans and livestock. 108

In spite of the enormous research on the role of sika deer in the transmission of tick- 109
 borne pathogens, a continuous survey is needed to detect the new emerging 110
 piroplasmids in sika deer in Japan. Previously, conventional polymerase chain 111
 reactions (PCR) have been used to detect *Babesia* and *Theileria* spp. in Japan 112
 (Watanabe et al., 2016). Here, the reverse line blot hybridization (RLB) was applied 113
 in this study for the specific detection of 12 different piroplasm species. This 114
 technique is a useful tool in identifying many tick-borne pathogens in a large number 115

of samples (Kong and Gilbert, 2006) with high specificity and less cost than 116

sequencing a large number of samples. The objective of this study was to estimate the 117

infection rates and molecular characterization of several *Babesia* and *Theileria* spp. 118

among sika deer in Hokkaido. 119

Materials and methods 120

Blood samples and DNA extraction 121

Whole blood samples were obtained from the jugular vein of 269 sika deer (*C. nippon* 122

yesoensis) that were culled for nuisance control during 2010 to 2013 from Hokkaido 123

prefecture, Japan. These blood samples were collected in EDTA-Na containing tubes 124

and stored at -20 °C until used. DNA was extracted from the blood samples by using 125

the Wizard ®Genomic DNA purification Kit (Promega, Madison, WI, USA) as per 126

manufacturer's instructions and kept at -20 °C until analysis. 127

Touch-down Polymerase Chain Reaction (PCR) 128

A touch-down PCR targeting the 18S rRNA gene of *Babesia* and *Theileria* species 129

was performed by using PCR System 9700 (Applied Biosystems, Foster City, CA), 130

QIAGEN Multiplex PCR kits (Qiagen, Hilden, Germany) and the primer pair RLB-F2 131

and RLB-R2 (Gubbels et al., 1999; Matjila et al., 2004). To amplify a 460-540 bp 132

fragment, PCR mixtures composed of 9 µl of molecular grade water, 1.25 µl of each 133

primer (10 pmol), 12.5 µl of 2 x QIAGEN Multiplex PCR Master Mix and 1 µl DNA. 134

The PCR consisted of an activation step at 94 °C for 15 min, a touch down step of 12 135

cycles at 94 °C for 30 s, 69 °C for 90 s and 72 °C for 90 s with annealing temperature 136

decreasing every cycle by 1°C. This followed by 40 cycles at 94°C for 30 s, 57°C for 137

90 s and at 72°C for 90 s and a final extension step at 72°C for 10 min. PCR products 138

were stained with 10× Midori Green Direct (Nippon Genetics, Düren, Germany) and 139

examined by electrophoresis through 1% agarose gel and visualized by UV	140
illuminator.	141
<i>Oligonucleotide probes and reverse line blot (RLB) hybridization</i>	142
A total of 8 newly designed and 9 previously published oligonucleotide probes (Table	143
1) were received from Sigma Aldrich Co., LLC, Japan. The probe designing process	144
was performed according to (Kong and Gilbert, 2006). Briefly, the 18S rRNA gene	145
sequences for the target pathogens (Table 1) were downloaded from the GenBank and	146
aligned by using MEGA software version 6 (Tamura et al., 2013). The probe	147
sequences were selected for each target and evaluated by OligoEvaluator software	148
(Sigma Aldrich, MO, USA) to ensure optimum physical characteristics. Finally, the	149
selected probe sequences were searched in the Nucleotide Basic Local Alignment	150
Search Tool (BLASTn) (Altschul et al., 1990) and tested against the positive control	151
samples to ensure its specificity.	152
RLB was carried out with the PCR products as described before (Kong and Gilbert,	153
2006; Moustafa et al., 2016) with modifications. Briefly, a 15 x 15 mm Biotodyne C	154
membrane (Pall Life Science, Ann Arbor, MI, USA) was activated for 10 min at room	155
temperature by 20 ml 16% 1-ethyl-3-(3-dimethyl-amino-propyl) carbodiimide	156
(EDAC) (Sigma Aldrich, MO, USA). The membrane was washed gently with milliQ	157
water for 2 min and placed in Miniblotter MN45 (Immunelectrics, Boston, Massa-	158
chusetts). A total of 17 oligonucleotide probes (Table 1) linked with C6 amino linker	159
were received from Sigma Aldrich Co., LLC, Japan, and reconstituted into 100 pmol	160
solutions. Ten microliters of each 100 pmol/μl probes were diluted in 0.5 M NaHCO ₃	161
to a final volume of 170 μl. The openings of the miniblotter were filled with a 150 μl	162
of each diluted oligonucleotide and the membrane was incubated at room temperature	163
for 5 min. The membrane was inactivated in 250 ml 0.1 M NaOH with gentle shaking	164

for exactly 8 min at room temperature, then washed at 60°C with prewarmed 250 ml 165
of 2× SSPE/0.1% SDS. Finally, the membrane was kept with 15µl of 20 mM EDTA 166
in a sealed plastic bag for the future use. 167

A total of 10 µl of each PCR product was diluted in 2× SSPE/0.1% SDS to a final 168
volume of 170 µl and denatured by heating at 99.9°C for 10 min and immediately 169
cooled on ice. The denatured mixtures were introduced to the membrane, and 170
hybridized at 60°C for 1 hr. Afterwards, this membrane was washed in 250 ml of 2× 171
SSPE/0.5% SDS twice at 52°C for 10 min and then incubated with diluted peroxidase 172
labeled NeutrAvidine (Thermo Fisher Scientific, Walyham, MA, USA) for 45-60 min 173
at 42°C. Subsequently, NeutrAvidine was washed away by 250 ml 2× SSPE/0.5% 174
SDS twice at 42°C for 10 min and 2× SSPE twice at room temperature for 5 min. To 175
visualize the results, the membrane was exposed for 5 min to 15 ml Immobilon™ 176
Western Chemiluminescent HRP Substrate (Millipore, Japan), and photographed by 177
Ez-Capture MG/ST (ATTO Corp., Japan). To remove the PCR products, the 178
membrane was washed twice in pre-warmed 1% SDS at 90°C for 30 min and washed 179
with 250 ml of 20 mM EDTA. For the future reuse, the membrane was kept with 15 180
ml of 20 mM EDTA in a sealed plastic bag at 4°C. 181

Cloning and sequencing 182

To confirm RLB results, randomly selected twenty-one RLB-positive PCR products 183
were subjected to direct sequencing. The selected PCR products included eight, one 184
and twelve samples that hybridized with the probes which were designed to detect 185
Theileria sp. Thrivae, *Babesia divergens*-like and all *Babesia* and *Theileria* spp., 186
respectively. To amplify a fragment of 460-540 bp from 18S rRNA gene of piroplasm 187
spp., KOD-Plus-Neo high fidelity DNA polymerase kit (Toyobo Co. Ltd., Japan) and 188
the above-mentioned primers were used. PCR consisted of 2.5 µl of 10× KOD-Plus- 189

Neo buffer, 2.5 μ l of dDNTPs (2mM), 1.5 μ l of 25 mM MgSO₄, 0.75 μ l of each 190
 primer (10 pmol), 0.5 μ l of KOD-Plus-Neo DNA polymerase and 1 μ l of DNA. The 191
 PCR conditions were 2 min at 94°C and 40 cycles of 10 s at 98°C, 30 s at 57°C and 30 192
 s at 68°C, followed by a final extension 7 min at 68°C. The PCR products were 193
 examined by electrophoresis and the positive bands at 460-540 bp were excised for 194
 purification by NucleoSpin® Gel and PCR clean up (Macherey-Nagel, GM & CO KG, 195
 Germany). The purified PCR products were sequenced by using the Big- Dye 196
 Terminator version 3.1 Cycle Sequencing Kit and purified by BigDye® 197
 XTerminator™ Purification Kit (Life Technologies Co., NY, USA). Subsequently, 198
 the purified products were sequenced by ABI PRISM™ 310 genetic analyzer (Life 199
 Technologies Co., NY, USA). BLASTn (Altschul et al., 1990) searches were 200
 performed to compare the obtained sequences to reference sequences recorded in 201
 GenBank for identification. 202

A total of 3 sequencing results with mixed signals were resolved by TA cloning. The 203
 purified PCR products were A-attached by mixing 0.9 μ l of 10× KOD-Plus-Neo 204
 buffer, 0.9 μ l of 2 mM dDNTPs, 0.54 μ l of 25 mM MgSO₄, 6.66 μ l of each purified 205
 PCR product and 1 μ l of 10× A-attachment mix (Toyobo Co. Ltd.) at 60°C for 10 min. 206
 One microliter of each A-attached product was cloned with pMD20 T-vector (Takara 207
 Bio, Otsu, Japan). Five colonies per sample were selected and screened by 208
 EmeraldAmp MAX PCR (Takara Bio, Otsu, Japan) as described in the manufacturer's 209
 manual and sequenced as mentioned above. When a sequence was detected from one 210
 clone, more 5 clones from the same sample were sequenced. The sequences were 211
 considered genuine when they were recovered from more than 3 clones. 212

Phylogenetic analysis 213

A total of twenty-two 18S rRNA partial sequences were submitted to DDBJ. The 214
accession numbers were: (LC271196 - LC271213) from the direct sequencing of 215
eighteen PCR products and (LC215385 - LC215388) from cloning and sequencing of 216
three other PCR products. The obtained sequences were identified by using BLASTn 217
search against the sequences of *Babesia* and *Theileria* spp. in the GenBank. The 218
divergence distances between sequences were calculated by the Maximum Composite 219
Likelihood model in MEGA software version 6 (Tamura et al., 2013). The 220
evolutionary distance matrix was visualized as a heat map (Fig. 3 & 4) that was 221
developed by using the R Stats package in the R studio software ([http://www.R-](http://www.R-project.org/) 222
[project.org/](http://www.R-project.org/)). 223

Results 224

A total of 269 sika deer blood DNA samples were molecularly screened for the 225
presence of *Babesia* and *Theileria* species by touch-down PCR and RLB 226
hybridization using 17 oligonucleotide probes (Table 1). The specificity of the newly 227
designed 8 oligonucleotide probes was ensured through examining the genomic DNA 228
samples of *B. microti*-US type, *B. microti*-Hobetsu type, *B. rodhaini*, *B. divergens*- 229
like, *B. bovis*, *B. bigemina*, *Theileria* sp. Thrivae and *T. orientalis* positive controls 230
(Fig. 1). 231

The results showed that 95.5% (257/269), 94.1% (253/269), 14.1% (38/269), 87.7% 232
(236/269) and 11.5% (31/269) of the sika deer blood DNA samples hybridized with 233
the *Babesia/Theileria* spp. catch-all, *Theileria* spp. catch-all, *Babesia* spp. catch-all, 234
Theileria sp. Thrivae and *B. divergens*-like oligonucleotide probes (Table 2), 235
respectively. In addition, 10.0% (27/269) of the PCR products hybridized with both 236
Theileria sp. Thrivae and *B. divergens*-like oligonucleotide probes, while 6.3% 237
(17/269) were positive for *Babesia / Theileria* spp. catch-all and *Theileria* spp. catch- 238

all probes, but negative for all species-specific probes (Fig. 2). Interestingly, 2.6% (7/269) hybridized with *Babesia* / *Theileria* spp. catch-all, *Theileria* spp. catch-all, *Theileria* sp. Thrivae and *Babesia* spp. 1 catch-all probes, but were negative for all *Babesia* spp. species-specific probes (Fig. 2).

The obtained 18S rRNA partial gene sequences were mainly divided into two *Theileria* and two *Babesia* spp. (Fig. 3 & 4). The similarity between the accession numbers (LC271208 - LC271213 and LC215385) was from 99 to 100%. Whereas sequence accession numbers (LC271196 - LC271207 and LC215386) were 99 - 100% similar to each other. The obtained sequence (LC215385) was identical to *T. cervi* (KT863532) from sika deer in China and 99% similar to *Theileria* sp. (AB012199 & AB012194) from sika deer in Japan, whereas sequence (LC215386) was identical to *Theileria* sp. (AB012189) from sika deer in Hokkaido, Japan and 99% similar to *T. capreoli* (KJ188219) from red deer in China. In addition, the obtained sequence (LC215387) was identical to *B. divergens*-like (KC465973 & KC465975) from sika deer in Japan, whereas sequence (LC215388) was 98% similar to *Babesia* sp. Kh-Hj42 (GU057380) from *Haemaphysalis japonica* in Russia.

Discussion

Sika deer act as reproductive hosts for several tick species, which may transmit several tick-borne pathogens to wildlife, livestock and human beings (Watanabe et al., 2016). During the last two decades, numbers of sika deer (*C. nippon*) populations in Hokkaido have been increasing (Ijima et al., 2015). In Japan, previous studies stated that sika deer act as reservoir hosts for *Theileria* sp. Thrivae and *B. divergens*-like (Ikawa et al., 2011; Inokuma et al., 2004; Zamoto-Niikura et al., 2014). In this study, we examined sika deer from Hokkaido for infection with 12 different piroplasm species by touch-down PCR and RLB hybridization. This study identified four

piroplasm species; *B. divergens*-like, undescribed *Babesia* sp., *Theileria* sp. Thrivae 264
 and *T. capreoli*. Unfortunately, no species-specific oligonucleotide probes were 265
 designed for the detected *Babesia* sp. and *T. capreoli*; both organisms were detected 266
 in this study by cloning and sequencing. The respective infection rates of the *B.* 267
divergens-like (11.5%) and *Theileria* sp. Thrivae (87.7%) detected in this study were 268
 similar to and higher than previously published reports from Japan (Ikawa et al., 2011; 269
 Watanabe et al., 2016; Zamoto-Niikura et al., 2014). This is possibly due to the higher 270
 sensitivity of RLB technique than conventional PCR assays. The hybridization results 271
 were confirmed by sequencing, which showed that all obtained sequences have the 272
 same hybridization sites as their species-specific probes. However, a total of 7 and 17 273
 PCR products, which were hybridized with the respective catch-all probes for *Babesia* 274
 and *Theileria* species, were negative for all other *Babesia* and *Theileria* species- 275
 specific probes. 276

The obtained partial 18S rRNA gene sequences were divided into two *Babesia* and 277
 two *Theileria* species. One *Babesia* sp. clustered with the previously published *B.* 278
divergens-like (Fig. 3) from sika deer in Japan (Zamoto-Niikura et al., 2014). The 279
 second *Babesia* sp. sequence was completely separated from the other *Babesia* spp. 280
 and clustered with a *Babesia* sp. which was detected in *Haemaphysalis japonica* from 281
 Russia (Fig. 3). Hence, additional research on this *Babesia* sp. is required to fully 282
 understand its molecular taxonomy and to identify its vector in Japan. 283

The phylogenetic position of the detected *Theileria* spp. in the sika deer of this study 284
 was among the benign theileriosis group, which includes *T. orientalis* and *T. buffeli* 285
 (Inokuma et al., 2008; Li et al., 2014). One *Theileria* spp. cluster was very similar to 286
Theileria sp. Thrivae (Fig. 4), which was reported from sika deer in Japan (Watanabe 287
 et al., 2016) and *T. cervi* from sika deer in China (Liu et al., 2016). However, this 288

cluster was completely divergent from *T. cervi* (U97054) that was detected in white-tailed deer and elk (*Cervus canadensis*) from the USA and Canada (Fig. 4), respectively (Chae et al., 1999). The second *Theileria* spp. cluster was closely related to *T. capreoli* (Fig. 4). Previously, *T. capreoli* 18S rRNA gene sequence (AB012189) was deposited into the GenBank from Japan, however, this study was not published. In addition, *Theileria* sp. Sola, which is 97.8% similar to *T. capreoli*, and was previously reported from Japan (Watanabe et al., 2016), was not detected in this study.

Conclusions

This study identified the role of sika deer in maintaining 4 different tick-borne protozoan parasites in Hokkaido, Japan with *Theileria* sp. Thrivae having the highest infection rates and *T. capreoli* needing more examinations to understand its ecology. Furthermore, a potentially novel *Babesia* sp. was detected in the sika deer samples of this study. However, this study lacks information about the prevalence of the undescribed *Babesia* sp. and *T. capreoli* in sika deer in Hokkaido. Future investigations will also be needed to identify the vector hosts for these organisms to fully understand the ecology of piroplasm spp. in Japan.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

This study was supported in part by the Program for Leading Graduate Schools “Fostering Global Leaders in Veterinary Science for Contributing to One Health” grant no. (F01), MEXT, Japan and the Mitsui & Co., Ltd. Environment Fund. The authors gratefully acknowledge Professor Xuenan Xuan and Dr. Paul Franck Adjou Moumouni (National Research Center for Protozoan Diseases, Obihiro University of

Agriculture and Veterinary Medicine, Japan) for providing us with the genomic DNA	313
samples of the positive controls.	314
Reference	315
Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local	316
alignment search tool. J. Mol. Biol. 215, 403-410.	317
Chae, J.S., Waghela, S.D., Craig, T.M., Kocan, A.A., Wagner, G.G., Holman, P.J.,	318
1999. Two <i>Theileria cervi</i> SSU RRNA gene sequence types found in isolates from	319
white-tailed deer and elk in North America. J. Wildl. Dis. 35, 458-465.	320
Decaro, N., Larocca, V., Parisi, A., Losurdo, M., Lia, R.P., Greco, M.F., Miccolis, A.,	321
Ventrella, G., Otranto, D., Buonavoglia, C., 2013. Clinical bovine piroplasmosis	322
caused by <i>Babesia occultans</i> in Italy. J. Clin. Microbiol. 51, 2432-2434.	323
Gubbels, J.M., de Vos, A.P., van der Weide, M., Viseras, J., Schouls, L.M., de Vries,	324
E., Jongejan, F., 1999. Simultaneous detection of bovine <i>Theileria</i> and <i>Babesia</i>	325
species by reverse line blot hybridization. J. Clin. Microbiol. 37, 1782-1789.	326
Hashemi-Fesharki, R., 1997. Tick-borne diseases of sheep and goats and their related	327
vectors in Iran. Parassitologia 39, 115-117.	328
Homer, M.J., Aguilar-Delfin, I., Telford, S.R., Krause, P.J., Persing, D.H., 2000.	329
Babesiosis. Clinical Microbiology Reviews 13, 451-469.	330
Ijima, H., Fujimaki, A., Ohta, U., Yamamura, K., Yokomizo, H., Uno, H., Matsuda,	331
H., 2015. Efficient management for the Hokkaido population of sika deer <i>Cervus</i>	332
<i>nippon</i> in Japan: accounting for migration and management cost. Population Ecology	333
57, 397-408.	334

Ikawa, K., Aoki, M., Ichikawa, M., Itagaki, T., 2011. Occurrence of two distinct	335
<i>Theileria</i> lineages in sika deer (<i>Cervus nippon</i>) of Iwate Prefecture, Japan .J. Vet.	336
Med. Sci. 73, 1371-1373.	337
Inokuma, H., Fujimoto, T., Hosoi, E., Tanaka, S., Fujisaki, K., Okuda, M., Onishi, T.,	338
2002. Tick infestation of sika deer (<i>Cervus nippon</i>) in the western part of Yamaguchi	339
prefecture, Japan. J. Vet. Med. Sci. 64, 6.617-15	340
Inokuma, H., Nagata, M., Hosoi, E., Itamoto, K., Okuda, M., 2008. Divergence of	341
p33/34 gene of <i>Theileria</i> found in <i>Cervus nippon</i> in Japan. J Vet Med Sci 70, 401-	342
405.	343
Inokuma, H., Tsuji, M., Kim, S.J., Fujimoto, T., Nagata, M., Hosoi, E., Arai, S .,	344
Ishihara, C., Okuda, M., 2004. Phylogenetic analysis of <i>Theileria</i> sp. from sika deer,	345
<i>Cervus nippon</i> , in Japan. Vet. Parasitol. 120, 339-345.	346
Kong, F., Gilbert, G.L., 2006. Multiplex PCR-based reverse line blot hybridization	347
assay (mPCR/RLB)--a practical epidemiological and diagnostic tool. Nat. Protoc. 1,	348
2668-2680.	349
Lee, K., Takano, A., Taylor, K., Sashika, M., Shimozuru, M., Konnai, S., Kawabata,	350
H., Tsubota, T., 2014. A relapsing fever group <i>Borrelia</i> sp. similar to <i>Borrelia</i>	351
<i>lonestari</i> found among wild sika deer (<i>Cervus nippon yezoensis</i>) and <i>Haemaphysalis</i>	352
spp. ticks in Hokkaido, Japan. Ticks Tick Borne Dis. 5, 841-847.	353
Li, Y., Chen, Z., Liu, Z., Liu, J., Yang, J., Li, Q., Li, Y., Cen, S., Guan, G., Ren, Q.,	354
Luo, J., Yin, H., 2014. Molecular identification of <i>Theileria</i> parasites of northwestern	355
Chinese <i>Cervidae</i> . Parasites & vectors 7, 225.	356

Liu, J., Yang, J., Guan, G., Liu, A., Wang, B., Luo, J., Yin, H., 2016. Molecular	357
detection and identification of piroplasms in sika deer (<i>Cervus nippon</i>) from Jilin	358
Province, China. Parasit Vectors 9, 156.	359
Mans, B.J., Pienaar, R., Latif, A.A., 2015. A review of <i>Theileria</i> diagnostics and	360
epidemiology. Int J Parasitol Parasites Wildl 4, 104-118.	361
Matjila, P.T., Penzhorn, B.L., Bekker, C.P., Nijhof, A.M., Jongejan, F., 2004 .	362
Confirmation of occurrence of <i>Babesia canis vogeli</i> in domestic dogs in South Africa.	363
Vet. Parasitol. 122, 119-125.	364
Michel, A.O., Mathis, A., Ryser-Degiorgis, M.P., 2014. <i>Babesia</i> spp. in European	365
wild ruminant species: parasite diversity and risk factors for infection. Veterinary	366
research 45, 65.	367
Moustafa, M.A., Lee, K., Taylor, K., Nakao, R., Sashika, M., Shimozuru, M., Tsubota,	368
T., 2015. Molecular characterization and specific detection of <i>Anaplasma</i> species	369
(AP-sd) in sika deer and its first detection in wild brown bears and rodents in	370
Hokkaido, Japan. Infect. Genet. Evol. 36, 268-274.	371
Moustafa, M.A., Taylor, K., Nakao, R., Shimozuru, M., Sashika, M., Rosa, R., Thu,	372
M.J., Rizzoli, A., Tsubota, T., 2016. Dynamics, co-infections and characteristics of	373
zoonotic tick-borne pathogens in Hokkaido small mammals, Japan. Ticks Tick Borne	374
Dis. 7, 922-928.	375
Olds, C.L., Paul, T., Scoles, G.A., 2016. Detection of <i>Theileria parva</i> in tissues of	376
cattle undergoing severe East Coast fever disease show significant parasite DNA	377
accumulation in the spleen. Vet. Parasitol. 232, 32-35.	378

Schreeg, M.E., Marr, H.S., Tarigo, J.L., Cohn, L.A., Bird, D.M., Scholl, E.H., Levy, M.G., Wiegmann, B.M., Birkenheuer, A.J., 2016. Mitochondrial genome sequences and structures aid in the resolution of <i>Piroplasmida</i> phylogeny. PLoS One 11, e0165702.	379 380 381 382
Tamura, K., Stecher, G., Peterson, D., Filipinski, A., Kumar, S., 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol. Biol. Evol. 30, 2725-2729.	383 384
Watanabe, Y., Fukumoto, S., Harasawa, R., 2016 .Prevalence of tick-borne hemolytic microbes in Free-living sika deer (<i>Cervus nippon</i>) captured in a deer-overcrowded area. Japanese Journal of Zoo and Wildlife Medicine 21, 17-27.	385 386 387
Yabsley, M.J., Quick, T.C., Little, S.E., 2005. Theileriosis in a white-tailed deer (<i>Odocoileus virginianus</i>) fawn. J. Wildl. Dis. 41, 806-809.	388 389
Zamoto-Niikura, A., Tsuji, M., Imaoka, K., Kimura, M., Morikawa, S., Holman, P.J., Hirata, H., Ishihara, C., 2014. Sika deer carrying <i>Babesia</i> parasites closely related to <i>B. divergens</i> , Japan. Emerg Infect Dis 20, 1398-1400.	390 391 392
Zanet, S., Trisciuglio, A., Bottero, E., de Mera, I.G., Gortazar, C., Carpignano, M.G., Ferroglio, E., 2014. Piroplasmosis in wildlife: <i>Babesia</i> and <i>Theileria</i> affecting free-ranging ungulates and carnivores in the Italian Alps. Parasites & Vectors 7, 70.	393 394 395 396 397 398 399 400

Probe	Sequence (5'-3')	Reference
<i>Babesia</i> / <i>Theileria</i> all	TAATGGTTAATAGGARCRGTWG	Moustafa et al. (2016)
<i>Theileria</i> spp. all	ATTAGAGTGCTCAAAGCAGGC	Matjila et al. (2008)
<i>Babesia</i> spp. all 1	ATTAGAGTGTTC AAGCAGAC	Matjila et al. (2008)
<i>Babesia</i> spp. all 2	ACTAGAGTGTTC A A A C A G G C	Matjila et al. (2008)
<i>B. microti</i> all	GRCTTGGC A T C W T C T G G A	Matjila et al. (2008)
<i>B. microti</i> Otsu and Hobetsu	GGGTACTGTTTCCAGGGT	This study
<i>B. microti</i> US	GGGTACTATTTCCAGGAT	This study
<i>B. gibsoni</i>	TACTTGCCTTGTCTGGTTT	Matjila et al. (2008)
<i>B. divergens</i> -like	TTAATCATAACWGATGTTTTG	This study
<i>B. rodhaini</i>	TGTGGATTAGTGCGCAAG	This study
<i>B. bigemina</i>	CTCGTAGTTGTATTT C A G C C T	This study
<i>B. bovis</i>	GAGCATGGAATAACCTTGTAT	This study
<i>B. venatorum</i>	GAGTTATTGACTCTTGTCTTTAA	Gigandet et al. (2011)
<i>B. divergens</i> & <i>B. capreoli</i>	GGTGTTAATATTGACTRATGTCGAG	Moustafa et al. (2016)
<i>B. duncani</i>	AGTTGAACTTCTGCCGCTT	Moustafa et al. (2016)
<i>Theileria</i> sp. Thrivae	ACGAGTGTCTGTATTGCG	This study
<i>Theileria orientalis</i> / <i>buffeli</i> / <i>sergenti</i>	TTTGAGTTTGTTATTGTGG	This study

TABLE 1 List of *Babesia* and *Theileria* spp. and their oligonucleotide probes used

for its detection

401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417

Total	No. of positive deer samples (%)
-------	----------------------------------

		<i>Babesia</i> / <i>Theileria</i> all	<i>Theileria</i> spp. all	<i>Babesia</i> spp. all 1	<i>Theileria</i> sp. <i>Thrivae</i>	<i>B. divergens</i> -like
Fawn	108	102 (94.4)	98 (90.7)	26 (24.1)	86 (79.6)	25 (23.1)
Adult	161	155 (96.3)	155 (96.3)	12 (7.5)	150 (93.2)	6 (3.7)
Total	269	257 (95.5)	253 (94.1)	38 (14.1)	236 (87.7)	31 (11.5)

418

TABLE 2 Summary of *Babesia* and *Theileria* spp. detection in sika deer blood

419

samples.

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

FIG 1 Validation of the newly designed oligonucleotide probes in this study. RLB 437
 result of *Babesia* and *Theileria* spp. genomic DNA derived 18S rRNA gene partial 438
 sequence (1-8). Oligonucleotide probes are listed on the left of the figure. 439






FIG 2 A representative RLB result. 1- 41: Sika deer blood derived 18S rRNA gene partial sequence. Oligonucleotide probes are listed on the left of the figure.



459
460
461
462
463
464
465
466
467
468
469
470
471
472

FIG 3 Phylogenetic relationship between 18S rRNA gene sequences of *Babesia* spp. 473
 from GenBank. Sequences from the current study are labeled in bold. The DNA 474
 sequences from this study was deposited to DDBJ as accession number LC215387 475
 and LC215388. The evolutionary distance matrix is visualized as a heat map. In the 476
 heat map, numbers in the table are presented as colored shades scaled from light to 477
 dark purple according to the degree of homology between sequences. The darker color 478
 the higher homology (less evolutionarily distant). White colors represent the same 479
 sequence 480



481

482

483


484

485

486

487

FIG 4 Phylogenetic relationship between 18S rRNA gene sequences of *Theileria* spp. 488
 from GenBank. Sequences from the current study are labeled in bold. The DNA 489
 sequences from this study was deposited to DDBJ as accession number LC215385 490
 and LC215386. The evolutionary distance matrix is visualized as a heat map. In the 491
 heat map, numbers in the table are presented as colored shades scaled from light to 492
 dark blue according to the degree of homology between sequences. The darker color 493
 the higher homology (less evolutionarily distant). White colors represent the same 494
 sequence 495



496

497

498

499