

# Antonie van Leeuwenhoek Journal of Microbiology

## Kroppenstedtia pulmonis sp. nov. and Kroppenstedtia sanguinis sp. nov., isolated from human patients --Manuscript Draft--

<b>Manuscript Number:</b>	ANTO-D-15-00548R1
<b>Full Title:</b>	Kroppenstedtia pulmonis sp. nov. and Kroppenstedtia sanguinis sp. nov., isolated from human patients
<b>Article Type:</b>	Original Article
<b>Keywords:</b>	Kroppenstedtia species, Kroppenstedtia pulmonis, Kroppenstedtia sanguinis, polyphasic taxonomy, 16S rRNA gene, thermoactinomycetes
<b>Corresponding Author:</b>	Melissa E Bell, MS Centers for Disease Control and Prevention Atlanta, Georgia UNITED STATES
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Centers for Disease Control and Prevention
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Melissa E Bell, MS
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Melissa E Bell, MS Brent A. Lasker, PhD Hans-Peter Klenk, PhD Lesley Hoyles, PhD Catherine Spröer Peter Schumann June Brown
<b>Order of Authors Secondary Information:</b>	
<b>Funding Information:</b>	
<b>Abstract:</b>	<p>Three human clinical strains (W9323T, X0209T and X0394) isolated from lung biopsy, blood and cerebral spinal fluid, respectively, were characterized using a polyphasic taxonomic approach. Comparative analysis of the 16S rRNA gene sequences showed the three strains belonged to two novel branches within the genus Kroppenstedtia: 16S rRNA gene sequence analysis of W9323T showed closest sequence similarity to Kroppenstedtia eburnea JFMB-ATE T (95.3 %), Kroppenstedtia guangzhouensis GD02T (94.7 %) and strain X0209T (94.6 %); sequence analysis of strain X0209T showed closest sequence similarity to K. eburnea JFMB-ATE T (96.4 %) and K. guangzhouensis GD02T (96.0 %). Strains X0209T and X0394 were 99.9 % similar to each other by 16S rRNA gene sequence analysis. The DNA-DNA relatedness was 94.6 %, confirming that X0209T and X0394 belong to the same species. Chemotaxonomic data for strains W9323T and X0209T were consistent with those described for the genus Kroppenstedtia: whole-cell peptidoglycan contained LL-diaminopimelic acid; the major cellular fatty acids were iso-C15 and anteiso-C15; and the major menaquinone was MK-7. Different endospore morphology and carbon utilization profiles of strains W9323T and X0209T supported by phylogenetic analysis enabled us to conclude that the strains represent two new species within the genus Kroppenstedtia, for which the names Kroppenstedtia pulmonis sp. nov. (type strain W9323T =DSM 45752 T) and Kroppenstedtia sanguinis sp. nov. (type strain X0209T =DSM 45749T=CCUG 38657 T) are proposed.</p>

Responses to reviewer's comments:

Editors comment: it appears from title page that Catherine Spröer and Peter Schumann should have been included as authors but their names are missing from the author list. Please correct.

Response: The author list has been corrected.

Reviewer #1:

Due to strains X0209T and X0394 belong to the same species, I would suggest the strain X0394 will be deleted from the manuscript.

[Editors comment: please retain consideration of strain X0394 - descriptions of taxa are better when based on more than one strain]

Response: We are agreement with the Editor and will leave the X0394 in the manuscript.

The abstract is too rough; please give the key characteristic differences between w9323T and X0209T.

Response: A key characteristic for differentiation between the two novel species is 16S rRNA gene sequence similarity which is stated in the abstract. In the manuscript, the details of phenotypic differences between the strains is discussed in depth. Due to word limitation in the abstract, we have elected not add additional differential phenotypic characteristics to the abstract.

Reviewer #3

Author's need to ensure all culture collection certificates are available for the two new species.

[Editors comment: please supply these on resubmission]

Response: The certificates are provided here with the revised manuscript.

This is correct in line 60 "plastic surface in a contract manufacturing"

Response: This line is correct.

The main phylogenetic tree should also have the maximum parsimony algorithm included. In addition in the figure legend the references for the algorithms used should be included. Finally in instead of using black dots the initials of the algorithms which the branches were recovered would be more appropriate.

Response: We have edited the tree accordingly and added the reference to the figure legend.

It is not evident if the phenotypic analysis was carried out with type strains from the genus *Kroppenstedtia* or the work was carried out on just the novel taxa. If the later happened then the work needs to be repeated alongside a type strain from the genus.

Response: As noted in the footnote of Table 2, all data is from the present study except for *K. gangzhouensis*. This is due to the inability of our lab to acquire the type strain from the culture collections of China and Korea. Unfortunately, the type strain is unavailable from any other resource.

Several references in the reference list are not present in the manuscript please remove:

Addou et al., 2012; Felsenstein, 1981; Kampfer et al., 2004; Klude et al., 1969; Mesbah et al., 1989; Rhuland et al., 1955; Saitou & Nei, 1989; Sneath, 1989; Tamura et al., 2011

Response: The references have been edited accordingly.

September 30, 2015

Antonie van Leeuwenhoek

To Whom It May Concern,

Please find enclosed our manuscript entitled “*Kroppenstedtia pulmonis* sp. nov. and *Kroppenstedtia sanguinis* sp. nov., isolated from human patients” for consideration for publication as an original article in the journal *Antonie van Leeuwenhoek*. We believe that the enclosed description of these two novel, clinically relevant, species expands our knowledge of the epidemiology of the genus.

All authors have seen and approved the manuscript and I assure that it has not previously been published nor is it under consideration for publication elsewhere. There are no conflicts of interest among the authors. Thank you for your consideration of our manuscript.

Please direct correspondence to: Melissa E. Bell, Bacterial Special Pathogens Branch, Centers for Disease Control and Prevention, 1600 Clifton Road, Atlanta, GA 30333 or at [jqv7@cdc.gov](mailto:jqv7@cdc.gov), phone: (404) 639-1348, facsimile (404) 639-3022.

Sincerely,

Melissa E. Bell M.Sc.  
Bacterial Special Pathogens Branch  
Division of High Consequence Pathogens and Pathology  
Centers for Disease Control and Prevention

1 *Kroppenstedtia pulmonis* sp. nov. and *Kroppenstedtia sanguinis* sp. nov., isolated from human patients

2 Melissa E. Bell, Brent A. Lasker, Hans-Peter Klenk, Lesley Hoyles, Catherine Spröer, Peter Schumann, June M. Brown

3

4 **Keywords:** *Kroppenstedtia* species, *Kroppenstedtia pulmonis*, *Kroppenstedtia sanguinis*, polyphasic taxonomy, 16S rRNA  
5 gene, thermoactinomycetes

6

7 **Corresponding Author**

8 M. E. Bell

9 Centers for Disease Control and Prevention, Building 17, Room 2209, Mailstop D-11, 1600 Clifton Road, Atlanta, GA

10 30333, USA

11 Phone: 404-639-1348

12 FAX: 404-639-3022

13 e-mail: jqv7@cdc.gov

14

15 B. A. Lasker, J. M. Brown

16 Bacterial Special Pathogens Branch, Division of High-Consequence Pathogens and Pathology, National Center for Emerging  
17 and Zoonotic Infectious Disease, Centers for Disease Control and Prevention, Atlanta, GA 30333, USA

18

19 H.-P. Klenk

20 Head of School, School of Biology, Ridley Building 2, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

21

22 C. Spröer, P. Schumann

23 Leibniz-Institute DSMZ-German Collection of Microorganisms and Cell Cultures, D-38124 Braunschweig, Germany

24

25 L. Hoyles

26 Department of Biomedical Sciences, Faculty of Science & Technology, University of Westminster, 115 New Cavendish  
27 Street, London, W1W 6UW, UK

28 **Abstract**

29 Three human clinical strains (W9323<sup>T</sup>, X0209<sup>T</sup> and X0394) isolated from lung biopsy, blood and cerebral spinal fluid,  
30 respectively, were characterized using a polyphasic taxonomic approach. Comparative analysis of the 16S rRNA gene  
31 sequences showed the three strains belonged to two novel branches within the genus *Kroppenstedtia*: 16S rRNA gene  
32 sequence analysis of W9323<sup>T</sup> showed closest sequence similarity to *Kroppenstedtia eburnea* JFMB-ATE<sup>T</sup> (95.3 %),  
33 *Kroppenstedtia guangzhouensis* GD02<sup>T</sup> (94.7 %) and strain X0209<sup>T</sup> (94.6 %); sequence analysis of strain X0209<sup>T</sup> showed  
34 closest sequence similarity to *K. eburnea* JFMB-ATE<sup>T</sup> (96.4 %) and *K. guangzhouensis* GD02<sup>T</sup> (96.0 %). Strains X0209<sup>T</sup> and  
35 X0394 were 99.9 % similar to each other by 16S rRNA gene sequence analysis. The DNA-DNA relatedness was 94.6 %,  
36 confirming that X0209<sup>T</sup> and X0394 belong to the same species. Chemotaxonomic data for strains W9323<sup>T</sup> and X0209<sup>T</sup> were  
37 consistent with those described for the genus *Kroppenstedtia*: whole-cell peptidoglycan contained LL-diaminopimelic acid;  
38 the major cellular fatty acids were *iso*-C<sub>15</sub> and *anteiso*-C<sub>15</sub>; and the major menaquinone was MK-7. Different endospore  
39 morphology, carbon utilization profiles, and whole cell wall sugar patterns of strains W9323<sup>T</sup> and X0209<sup>T</sup> supported by  
40 phylogenetic analysis enabled us to conclude that the strains represent two new species within the genus *Kroppenstedtia*, for  
41 which the names *Kroppenstedtia pulmonis* sp. nov. (type strain W9323<sup>T</sup> = DSM 45752<sup>T</sup> = CCUG 68107<sup>T</sup>) and  
42 *Kroppenstedtia sanguinis* sp. nov. (type strain X0209<sup>T</sup> = DSM 45749<sup>T</sup> = CCUG 38657<sup>T</sup>) are proposed.

43

---

**44 Introduction**

45

46 The family *Thermoactinomycetaceae* was established to accommodate new taxa (*Mechercharimyces* spp.) and the previously  
47 described genera *Thermoactinomyces*, *Laceyella*, *Thermoflavimicrobium*, *Seinonella* and *Planifilum* (Matsuo et al. 2006).

48 The family now encompasses 19 genera, with representatives isolated from clinical specimens (e.g. *Desmospora*,  
49 *Marinithermofilum*, *Hazenella*) or environmental sources (e.g. *Mechercharimyces*, *Lihuaxuella*, *Geothermomicrobium* and  
50 *Risungbinella*) (Matsuo et al. 2006; Yassin et al. 2009; Yu et al. 2012; Buss et al. 2013; Zhou et al. 2014; Kim et al. 2015;  
51 Zhang et al. 2015). Salient properties of the family *Thermoactinomycetaceae* include thermotolerant growth up to 60 °C,  
52 Gram positivity, non-acid fastness and formation of single spores that may be sessile or on simple sporophores with the  
53 structure and properties of endospores. The description of the family was emended by establishing the 16S rRNA gene  
54 sequence signature nucleotides as required for suprageneric assignment (Yassin et al. 2009). Von Jan et al. (2011) further  
55 emended the description of the family *Thermoactinomycetaceae* (to contain either LL-diaminopimelic acid or *meso*-  
56 diaminopimelic acid) when they described a new genus and species, *Kroppenstedtia eburnea*, that contained the isomer LL-  
57 diaminopimelic acid in its whole-cell peptidoglycan.

58

59 *Kroppenstedtia* species have been isolated from environmental and clinical sources. Although the type strain of *K. eburnea*  
60 was isolated from an environmental source (plastic surface in a contract manufacturing company in Germany), Barker et al.  
61 (2012) identified 14 strains of this species in clinical (blood, skin, peritoneal fluid, cerebral spinal fluid) samples in the  
62 United States. Another species, *Kroppenstedtia guangzhouensis*, was isolated from soil in south China (Yang et al. 2013).  
63 Three human clinical strains of *Kroppenstedtia* species were identified in a retrospective evaluation of 16S rRNA gene  
64 sequences at the Special Bacteriology Reference Laboratory (SBRL). Genotypic and phenotypic data suggest that strain  
65 W9323<sup>T</sup> represents a new species, for which we propose the name *Kroppenstedtia pulmonis* sp. nov., and that strains X0209<sup>T</sup>  
66 and X0394 both represent another new species, for which we propose the name *Kroppenstedtia sanguinis* sp. nov.

67

68

---

## 69 **Materials and methods**

70

71 Bacterial strains

72

73 Strain X0209<sup>T</sup> was obtained from the Culture Collection University of Göteborg (CCUG), Göteborg, Sweden as74 '*Thermoactinomyces sanguinis*' (CCUG 38657<sup>T</sup>). Two strains, W9323<sup>T</sup> and X0394, were sent to the SBRL, Centers for75 Disease Control and Prevention for identification. *K. eburnea* DSM 45196<sup>T</sup> was used as a phenotypic, chemotaxonomic and76 genetic control and *K. guangzhouensis* GD02<sup>T</sup> and *Melghirimyces algeriensis* NariEX<sup>T</sup> were used as chemotaxonomic and

77 genetic controls throughout this study. The GenBank accession numbers for the 16S rRNA gene sequences and the patients'

78 data associated with these strains are given in Table 1.

79

80 Phenotypic analyses

81

82 Morphological, cultural, physiological and biochemical analyses

83

84 To examine morphologic features, strains were grown aerobically using brain heart infusion (BHI) broth, heart infusion agar

85 (HIA; Becton, Dickinson and Co, Sparks, MD) supplemented with 5 % rabbit blood, HIA slants, trypticase soy agar (TSA)

86 supplemented with 5 % sheep blood (Becton, Dickinson and Co) at 35 and 45 °C for 3 to 7 days and then examined for

87 microscopic and macroscopic features. Gram and modified Kinyoun acid-fast staining were conducted as described

88 previously (Berd 1973). Cultures were examined for optimal growth at 35, 45, 50 and 60 °C for 7 days on HIA slants with 5

89 % rabbit blood. Optimal growth was determined by comparative observation of the amount of cell mass production at each

90 temperature. All phenotypic studies were performed under optimal growth conditions (at 45 °C in air).

91

92 Decomposition tests for adenine, casein, esculin, hypoxanthine, tyrosine, urea and xanthine, utilization of 22 carbohydrates as

93 sole source of carbon, utilization of acetamide and citrate, arylsulfatase production and nitrate reduction and growth in the

94 presence of lysozyme were performed as described previously (Conville and Witebsky 2007; Conville et al. 2008; Weyant et

95 al. 1996; Yassin et al. 1995). The decomposition of casein was compared with casein plus 0.5 % NaCl as described by von

96 Jan et al. (2011).

97

98

## 99 Antimicrobial susceptibility testing

100

101 Since no guidelines were available for the genus *Kroppenstedtia*, the MICs to 10 antimicrobial agents were determined  
102 following interpretative breakpoints as recommended for aerobic actinomycetes (Clinical and Laboratory Standards Institute  
103 2011) for amikacin, amoxicillin-clavulanate, ceftriaxone, ciprofloxacin, clarithromycin, imipenem, linezolid, minocycline,  
104 moxifloxacin and trimethoprim-sulfamethoxazole; the interpretive breakpoint for ampicillin used was that recommended for  
105 *Enterobacteriaceae* (Clinical and Laboratory Standards Institute 2015). The preparation of the inoculum suspension  
106 followed the guidelines as described previously (Clinical and Laboratory Standards Institute 2003).

107

## 108 Chemotaxonomic analyses

109

110 Assays of diaminopimelic acid stereoisomers and whole-cell sugars were performed by thin-layer chromatography using  
111 methods described previously (Lechevalier and Lechevalier 1970; Rhuland et al. 1955; Staneck and Roberts 1974).  
112 Isoprenoid quinones and polar lipids were extracted and purified and analyzed by the method of Minnikin et al. (1984).  
113 Analysis of isoprenoid quinones by HPLC was performed as described by Kroppenstedt (1982, 1985). Cellular fatty acids  
114 were prepared by the method of Klatte et al. (1994) and the fatty acid methyl esters were then separated as described by  
115 Sasser (1990) using the Microbial Identification System (MIDI, Inc., Sherlock version 6.1). Standardization of the  
116 physiological age of the study and reference strains was obtained by choosing the sector from a quadrant streak of culture  
117 plates.

118

## 119 Genetic analyses

120

## 121 16S rRNA gene sequence analysis

122

123 Purification of whole-cell DNA, amplification of near full-length 16S rRNA gene fragments, primers and nucleotides for  
124 PCR, purification of amplicons and DNA cycle sequencing were described previously (Lasker et al. 2011). Consensus 16S  
125 rRNA gene sequences were assembled and edited using Sequencher version 4.10.1 software. To identify related gene  
126 sequences in the GenBank database, consensus sequences were submitted to GenBank using BLASTn software



127 (<https://www.ncbi.nlm.nih.gov/blast/>). A multiple sequence alignment was created using Clustal W (within Geneious 8.1.4),  
128 from which gaps and 5' and 3' ends were trimmed. The distance matrix was calculated using DNADIST (Kimura 2-correction  
129 parameter) (Felsenstein 1989). Phylogenetic trees were constructed using the neighbor-joining (NEIGHBOR), maximum  
130 likelihood (DNAML with global rearrangements) and maximum parsimony (DNAPARS with global rearrangements)  
131 methods available in the PHYLIP package (Felsenstein, 1989). Stability of groupings within the neighbor-joining tree was  
132 estimated by bootstrap analysis (1000 replications) using the programs SEQBOOT, DNADIST, NEIGHBOR and  
133 CONSENSE (Felsenstein 1989). The phenograms were visualized by using the program DRAWGRAM (Felsenstein 1989).  
134 The sequence of the type strain of *Bacillus subtilis* was used as the outgroup.

135

136 DNA-DNA hybridization and DNA G + C content

137

138 Cells for hybridization and G+C mol% determination were disrupted using a Constant Systems TS 0.75 KW (IUL  
139 Instruments, Germany). DNA in the crude lysates was purified on hydroxyapatite by chromatography as described by  
140 Cashion et al. (1977). DNA-DNA hybridization was carried out in duplicate as described by De Ley et al. (1970) as modified  
141 by Huss et al. (1983) using a model Cary 100 Bio UV/VIS-spectrophotometer equipped with a Peltier-thermostat 6 x 6  
142 multicell changer and a temperature controller with *in-situ* temperature control. DNA-DNA hybridization studies were  
143 between the clinical strains X0209<sup>T</sup> and X0394 to confirm they were the same genomic species. DNA-DNA hybridization  
144 studies between X0209<sup>T</sup> and W9323<sup>T</sup> and their respective closest phylogenetically related neighbors were not conducted  
145 because of the low probability inferred from 16S rRNA gene similarities observed (Meier-Kolthoff et al. 2013).

146

147 DNA G+C content

148

149 The method of Mesbah et al. (1989) was performed to determine the G+C content of the novel type strains.

150

## 151 **Results and discussion**

152 Strains W9323<sup>T</sup>, X0209<sup>T</sup> and X0394 were aerobic, mesophilic to thermophilic, Gram-positive bacteria but were not acid fast.  
153 Substrate hyphae were filamentous and branched, and could be seen as fringes around the colony margins; no aerial hyphae  
154 were observed. Rare elongated (paddle shaped) endospores on long, unbranched sporophores were seen on Gram-stained  
155 smears of W9323<sup>T</sup>; single globose endospores on unbranched sporophores were seen on Gram-stained smears of X0209<sup>T</sup> and

156 X0394. Colonies of all three strains showed beta hemolysis on TSA with 5 % sheep blood at 45 °C. Growth occurred at 35  
157 and 45 °C but not at 50 °C with optimum growth at 45 °C; the optimal growth at 45 °C was consistent with the type strain of  
158 *K. eburnea* but differed from the optimal growth at 50 °C of the *K. guangzhouensis* type strain. Pale yellow colonies on 7-day  
159 TSA with 5 % sheep blood had irregular margins with random surface ridges at 35 and 45 °C. Except for the production of  
160 paddle-shaped endospores on long, unbranched sporophores of strain W9323<sup>T</sup>, the microscopic morphology was consistent  
161 with the type strains of *K. eburnea* and *K. guangzhouensis*. The macroscopic morphology of W9323<sup>T</sup>, X0209<sup>T</sup> and X0394  
162 was consistent with that described for the type strains of *K. eburnea* and *K. guangzhouensis*, however, none of the three  
163 strains produced aerial hyphae as reported by von Jan et al. (2011) and Yang et al (2013).

164

165 Table 2 gives the differential phenotypic, chemotaxonomic and genetic characteristics of the study strains and their closest  
166 phylogenetic relatives. Strains X0209<sup>T</sup> and X0394 both were able to utilize D-mannitol; W9323<sup>T</sup> was the only strain able to  
167 utilize D-glucose and sucrose. Strain X0209<sup>T</sup> utilized cellobiose and salicin but the related strain X0394 did not.

168

169 Results of antimicrobial susceptibility testing showed strains W9323<sup>T</sup>, X0209<sup>T</sup> and X0394 were resistant (MIC,  $\geq 8$   $\mu\text{g/ml}$ ) to  
170 clarithromycin but were susceptible to amikacin (MIC,  $\leq 8$ ), amoxicillin-clavulanate (MIC,  $\leq 8/4$   $\mu\text{g/ml}$ ), ampicillin (MIC,  $\leq$   
171 4  $\mu\text{g/ml}$ ), ceftriaxone (MIC,  $\leq 8$   $\mu\text{g/ml}$ ), ciprofloxacin (MIC,  $\leq 1$   $\mu\text{g/ml}$ ), imipenem (MIC,  $\leq 4$   $\mu\text{g/ml}$ ), linezolid (MIC,  $\leq 8$   
172  $\mu\text{g/ml}$ ), minocycline (MIC,  $\leq 1$   $\mu\text{g/ml}$ ), moxifloxacin (MIC,  $\leq 1$   $\mu\text{g/ml}$ ) and trimethoprim-sulfamethoxazole (MIC,  $\leq 2/38$   
173  $\mu\text{g/ml}$ ). Except for resistance to ampicillin of *K. eburnea* JFMB-ATE<sup>T</sup>, susceptibility results of the three study strains were  
174 consistent with *K. eburnea* JFMB-ATE<sup>T</sup>. The antimicrobial susceptibility test results in our study were consistent with the  
175 MIC results of 14 clinical strains of *K. eburnea* reported by Barker et al. (2012); for example, all their strains were  
176 susceptible to all antimicrobial agents tested except for clarithromycin.

177

178 Whole-cell wall hydrolysates contained LL-diaminopimelic acid, ribose and traces of galactose (X0209<sup>T</sup> and X0394); *K.*  
179 *eburnea* JFMB-ATE<sup>T</sup> contained ribose and glucose. The predominant menaquinones of W9323<sup>T</sup> were MK-7 (95 %) and MK-  
180 8 (5 %); the predominant menaquinones of X0209<sup>T</sup> and X0394 were identified as MK-7 (97 %) and MK8 (3 %). Polar lipids  
181 were phosphatidylglycerol (PG), diphosphatidylglycerol (DPG), phosphatidylethanolamine (PE),  
182 phosphatidylmethylethanolamine (PME), two unknown phospholipids (PL) (4PL, W9323<sup>T</sup> and X0394; 5PL, X0209<sup>T</sup>) and  
183 one unknown glycolipid (GL) (1GL, W9323<sup>T</sup>, X0209<sup>T</sup> and X0394). The fatty acid profile of the three strains consisted  
184 predominantly of *iso*-C<sub>15:0</sub> (74-75 %) and *anteiso*-C<sub>15:0</sub> (11-13 %).

185

186 DNA-DNA hybridization studies between the clinical strains X0209<sup>T</sup> and X0394 confirmed they were the same genomic  
187 species with relatedness value of 93.6% ± 6.5%. The genomic DNA G+C content of strains W9323<sup>T</sup> and X0209<sup>T</sup> was 45.9  
188 and 50.6 mol%, respectively. These values fall within the range of genomic DNA G+C content reported for the genus  
189 *Kroppenstedtia* as it exists presently (46 to 56 mol%) (von Jan et al. 2011, Yang et al 2013).

190

191 Based on 16S rRNA gene sequence analysis, the clinical strains were assigned within the subclade for the genus  
192 *Kroppenstedtia*, within the family *Thermoactinomycetaceae* (Fig. 1). The highest sequence similarity with strain W9323<sup>T</sup>  
193 was to *K. eburnea* JFMB-ATE<sup>T</sup> (95.3 % sequence similarity), *K. guangzhouensis* GD02<sup>T</sup> (94.7 % sequence similarity),  
194 *Melghirimyces thermohalophilus* Nari11A<sup>T</sup> (94.5 % sequence similarity), *M. algeriensis* DSM 45474<sup>T</sup> (94.3 % similarity),  
195 and *Desmospora activa* DSM 45169<sup>T</sup> (94.2 % similarity). The 16S rRNA gene sequences for strains X0209<sup>T</sup> and X0394 were  
196 99.9 % identical to each other. Strain X0209<sup>T</sup> showed the highest sequence similarity to *K. eburnea* JFMB-ATE<sup>T</sup> (96.4 %  
197 similarity), *K. guangzhouensis* GD02<sup>T</sup> (96.0 % similarity), *M. algeriensis* DSM 45474<sup>T</sup> (94.3 % similarity), *M.*  
198 *thermohalophilus* Nari11A<sup>T</sup> and *D. activa* DSM 45169<sup>T</sup> (93.7 % similarity). The 16S rRNA gene sequences of strain X0209<sup>T</sup>  
199 and *K. eburnea* JFMB-ATE<sup>T</sup> differed by 59 bp; the sequences of strain W9323<sup>T</sup> and *K. eburnea* JFMB-ATE<sup>T</sup> differed by 66  
200 bp.

201

202 From the results of our phenotypic, chemotaxonomic and genetic studies, it is proposed that strains W9323<sup>T</sup> and X0209<sup>T</sup> be  
203 classified in the genus *Kroppenstedtia* as *Kroppenstedtia pulmonis* sp. nov. and *Kroppenstedtia sanguinis* sp. nov.,  
204 respectively.

205

#### 206 **Description of *Kroppenstedtia pulmonis* sp. nov.**

207 *K. pulmonis* (N. L. fem. adj. *pul.mo'nis*. L. n. pulmo-onis, lung; L. gen. n. pulmonis of a lung, isolated from a lung).

208 Cells are Gram-positive, non-acid-fast and filamentous. Elongated spores (paddle shaped) are formed singly on sessile  
209 sporophores on substrate hyphae. Pale yellowish-gray colonies are wrinkled with random ridges in 3 to 7 days at 35 and 45  
210 °C on trypticase soy agar with 5 % sheep blood. Colonies are beta hemolytic but do not form aerial hyphae. Growth occurs at  
211 35 and 45 °C but not at 50 °C. D-Glucose and sucrose are utilized but i-adonitol, L-arabinose, cellobiose, citrate, dulcitol, i-  
212 erythritol, D-fructose, D-galactose, glycerol, i-*myo*-inositol, lactose, maltose, D-mannitol, mannose, melibiose, raffinose, L-  
213 rhamnose, salicin, D-sorbitol, trehalose and xylose are not utilized. Grows in the presence of lysozyme but has no

214 arylsulfatase activity and does not reduce nitrate. Hydrolyses casein with 0.5 % NaCl but does not hydrolyse acetamide,  
215 adenine, casein, hypoxanthine, tyrosine, urea or xanthine. Esculin hydrolysis is positive by browning but negative by UV  
216 light absorption. Major fatty acids (>10 %) are *iso*-C<sub>15:0</sub> (75 %) and *anteiso*-C<sub>15:0</sub> (11 %). Whole-cell hydrolysates contain  
217 LL-diaminopimelic acid and the sugar ribose. The predominant menaquinones of strain W9323<sup>T</sup> are MK-7 (95 %) and MK-8  
218 (5 %). Polar lipids are phosphatidylglycerol (PG), diphosphatidylglycerol (DPG), phosphatidylethanolamine (PE),  
219 phosphatidylmethylethanolamine (PME), one unknown phospholipid (4PL) and one unknown glycolipid (1GL). The type  
220 strain (W9323<sup>T</sup> = DSM 45752<sup>T</sup> = CCUG 68107<sup>T</sup>) was isolated from lung biopsy of a 78-year-old male patient from New  
221 York, USA. The G+C content of its genomic DNA is 45.9 mol%.

222

### 223 **Description of *Kroppenstedtia sanguinis* sp. nov.**

224 *K. sanguinis* (san'gui.nis L. n. sanguis-inis, blood; L. gen. n. sanguinis, of blood).

225 Cells are Gram-positive, non-acid-fast and filamentous. Globose spores are formed singly on sessile sporophores on substrate  
226 hyphae. Pale yellowish gray colonies are wrinkled with random ridges in 3 to 5 days at 35 and 45 °C on trypticase soy agar  
227 with 5 % sheep blood. Colonies are beta hemolytic but do not form aerial hyphae. Growth occurs at 35 and 45 °C but not at  
228 50 °C. Cellobiose, D-mannitol, mannose and salicin are utilized but i-adonitol, L-arabinose, citrate, dulcitol, i-erythritol, D-  
229 fructose, D-galactose, D-glucose, glycerol, *i-myo*-inositol, lactose, maltose, melibiose, raffinose, L-rhamnose, D-sorbitol,  
230 sucrose, trehalose and D-xylose are not utilized. Grows in the presence of lysozyme but has no arylsulfatase activity and does  
231 not reduce nitrate. Hydrolyses casein with 0.5 % NaCl but does not hydrolyse acetamide, adenine, casein, hypoxanthine,  
232 tyrosine, urea or xanthine. Esculin hydrolysis is positive by browning but negative by UV light absorption. The predominant  
233 menaquinone is MK-7 (95 %). Whole-cell hydrolysates contain LL-diaminopimelic acid and traces of the sugar galactose.  
234 Major fatty acids (>10 %) are *iso*-C<sub>15:0</sub> (74 %) and *anteiso*-C<sub>15:0</sub> (12 %). Whole-cell hydrolysates contain LL-diaminopimelic  
235 acid and traces of galactose and ribose. Polar lipids are phosphatidylglycerol (PG), diphosphatidylglycerol (DPG),  
236 phosphatidylethanolamine (PE), phosphatidylmethylethanolamine (PME), one unknown phospholipid (5PL), and one  
237 unknown glycolipid (1GL). The type strain (X0209<sup>T</sup> = DSM 45749<sup>T</sup> = CCUG 38657<sup>T</sup>) was isolated from the blood of a 59-  
238 year-old male from Gävle, Sweden. The G+C content of its genomic DNA is 50.6 mol%.

239

240 Acknowledgements We thank Jean Euzéby for nomenclatural advice.

241

242

243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269

---

## References

- Barker AP, Simmons KE, Cohen S, Slechta ES, Fisher MA, Schlager R (2012) Isolation and identification of *Kroppenstedtia eburnea* isolates from multiple patient samples. *J Clin Microbiol* 50:3391-3394
- Berd D (1973) Laboratory identification of clinically important aerobic actinomycetes. *Appl Microbiol* 25:665-681
- Buss SN, Cole JA, Hannett GE, Nazarian EJ, Nazarian L, Coorevits A, Van Landschoot A, De Vos P, Schumann P, Musser KA, Wolfgang WJ (2013) *Hazenella coriacea* gen. nov., sp. nov., isolated from clinical specimens. *Int J Syst Evol Microbiol* 63:4087-4093
- Cashion P, Hodler-Franklin MA, McCully J, Franklin M (1977) A rapid method for base ratio determination of bacterial DNA. *Anal Biochem* 81:461-466
- Clinical and Laboratory Standards Institute (2003) Susceptibility testing of mycobacteria, nocardiae, and other aerobic actinomycetes; approved standard M24-A. National Committee for Clinical Laboratory Standards, Wayne, PA
- Clinical and Laboratory Standards Institute (2011) Susceptibility testing of mycobacteria, nocardiae, and other aerobic actinomycetes; approved standards M24-A2. Clinical Laboratory Standards Institute, Wayne, PA
- Clinical and Laboratory Standards Institute (2015) Performance standards for antimicrobial susceptibility testing; twenty-fifth information supplement M100-S25. National Committee for Clinical Laboratory Standards, Wayne, PA
- Conville PS, Witebsky FG (2007) *Nocardia*, *Rhodococcus*, *Gordonia*, *Actinomadura*, *Streptomyces*, and other aerobic actinomycetes. In: *Manual of Clinical Microbiology*, 9<sup>th</sup> edn, pp. 515-542. PR Murray, EJ Baron, JH Jorgensen, ML Landry and MA Tenover, (eds) Washington, DC: American Society for Microbiology

- 270 Conville PS, Brown JM, Steigerwalt AG, Brown-Elliott BA, Witebsky FG (2008) *Nocardia wallacei* sp. nov., and *Nocardia*  
271 *blaklockiae* sp. nov., human pathogens and members of the “*Nocardia transvalensis* complex”. J Clin Microbiol 46:1178-  
272 1184  
273
- 274 De Ley J, Cattoir H, Reynaerts A (1970) The quantitative measurement of DNA hybridization from renaturation rates. Eur J  
275 Biochem 12:133-142  
276
- 277 Felsenstein J (1989) PHYLIP – Phylogeny inference package (version 3.2). Cladistics 5:164-166.  
278
- 279 Huss VAR, Festl H, Schleifer KH (1983) Studies on the spectrophotometric determination of DNA hybridization from  
280 renaturation rates. Syst Appl Microbiol 4:184-192  
281
- 282 Kim M, Kim T, Ri S, Jiang F, Chang X, Liu Y, Ren L, Qiu X, Deng S, Fang C, Zheng C, Peng F (2015) *Risunghinella*  
283 *pyongyangensis* gen. nov., sp. nov., a mesophilic strain of the family *Thermoactinomycetaceae* isolated from agricultural soil  
284 sample. Int J Syst Evol Microbiol doi:10.1099/ijms.0.000328  
285
- 286 Klatte S, Kroppenstedt RM, Rainey FA (1994) *Rhodococcus opacus* sp. nov., an unusual nutritionally versatile *Rhodococcus*  
287 species. Syst Appl Microbiol 17:355-360  
288
- 289 Kroppenstedt RM (1982) Separation of bacterial menaquinones by HPLC using reverse phase (RP18) and a silver loaded ion  
290 exchanger as stationary phases. J Liq Chromatogr 5: 2359-2367  
291
- 292 Kroppenstedt RM (1985) Fatty acid and menaquinone analysis of actinomycetes and related organisms. In: Chemical  
293 Methods in Bacterial Systematics, no. 20 SAB Technical Series, pp. 173-199. Edited by Goodfellow M & Minnikin DE.  
294 London: Academic Press  
295
- 296 Lasker BA, Moser BD, Brown JM (2011) *Gordonia*. In: Molecular Detection of Human Bacterial Pathogens, pp. 95-110.D  
297 Liu, (ed) Boca Raton, FL: CRC Taylor & Francis Group  
298

- 299 Lechevalier MP, Lechevalier HA (1970) Chemical composition as a criterion in the classification of aerobic actinomycetes.  
300 Int J Syst Bacteriol 20:435-443  
301
- 302 Matsuo Y, Katsuta A, Matsuda S, Shizurti Y, Yokota A, Kasai H (2006) *Mechercharimyces mesophilus* gen. nov. and sp. nov.  
303 *Mechercharimyces asporophorigenens* sp. nov., antitumor substance-producing marine bacteria, and description of  
304 *Thermoactinomycetaceae* fam. nov. Int J Syst Evol Microbiol 56:2837-2842  
305
- 306 Meier-Kolthoff JP, Göker M, Spröer C, Klenk HP (2013) When should a DDH experiment be mandatory in microbial  
307 taxonomy? Arch Microbiol 195:413-418  
308
- 309 Mesbah M, Premachandran U, Whitman WB (1989) Precise measurement of the G+C content of deoxyribonucleic acid by  
310 high-performance liquid chromatography. Int J Syst Bacteriol 39:159-167  
311
- 312 Minnikin DE, O'Donnell AG, Goodfellow G, Alderson G, Athalye M, Schaal A, Parlett JH (1984) An intergrated procedure  
313 for the extraction of bacterial isoprenoid quinones and polar lipids. J Microbiol Methods 2:233-241  
314
- 315 Rhuland LE, Work, E, Denman RF, Hoare DS (1955) The behavior of the isomers of  $\alpha,\epsilon$ -diaminopimelic acid on paper  
316 chromatograms. J Am Chem Soc 77:4844-4846  
317
- 318 Sasser M (1990) Identification of bacteria by gas chromatography of cellular fatty acids. USFCC Newsl 20, 16  
319
- 320 Staneck JL, Roberts GD (1974) Simplified approach to identification of aerobic actinomycetes by thin-layer chromatography.  
321 Appl Microbiol 28:226-231  
322
- 323 von Jan M, Riegger N, Pötter G, Schumann P, Verborg S, Spröer C, Rohde M, Lauer B, Labeda DP, Klenk HP (2011)  
324 *Kroppenstedtia eburnea* gen. nov., sp. nov., a thermoactinomycete isolated by environmental screening, and emended  
325 description of the family *Thermoactinomycetaceae* Matsuo *et al.* 2006 emend. Yassin *et al.* 2009. Int J Syst Evol Microbiol  
326 61:2304-2310  
327


- 328 Weyant RS, Moss CW, Weaver RE, Hollis DG, Jordan JG, Cook EC, Daneshvar MI (1996) Identification of Unusual  
329 Pathogenic Gram-Negative and Facultatively Anaerobic Bacteria, 2<sup>nd</sup> edn. Baltimore, MD: Williams and Wilkins  
330
- 331 Yang G, Qin D, Wu C, Yuan Y, Zhou S, Cai Y (2013) *Kroppenstedtia guangzhouensis* sp. nov., a thermoactinomycete  
332 isolated from soil. Int J Syst Evol Microbiol 61:2304-2310  
333
- 334 Yassin AF, Hupfer H, Klenk HP, Siering C (2009) *Desmospora activa* gen. nov., sp. nov., a thermoactinomycete isolated  
335 from sputum of a patient with suspected pulmonary tuberculosis, and emended description of the family  
336 *Thermoactinomycetaceae* Matsuo *et al.* 2006. Int J Syst Bacteriol 59:454-459  
337
- 338 Yassin AF, Rainey FA, Brzezinka H, Burghardt J, Lee HJ, Schaal KP (1995) *Tsukamurella inchonensis* sp. nov. Int J Syst  
339 Bacteriol 45:522-527  
340
- 341 Yu TT, Zhang BH, Yao JC, Tang SK, Zhou EM, Yin YR, Wei DQ, Ming H, Li WJ (2012) *Lihuaxuella thermophila* gen.  
342 nov., sp. nov., isolated from a geothermal soil sample in Tengchong, Yunnan, south-west China. Antonie Van Leeuwenhoek  
343 102:711-718  
344
- 345 Zhang Y, Li J, Tian X, Zhang S (2015) *Marinithermofilum abyssi* gen. nov., sp. nov. and *Desmospora profundinema* sp. nov.  
346 isolated from a deep-sea sediment, and emended description of the genus *Desmospora* Yassin *et al.* 2009. Int J Syst Evol  
347 Microbiol doi: 10.1099/ijms.0.000314  
348
- 349 Zhou EM, Yu TT, Liu L, Ming H, Yin YR, Dong L, Tseng M, Nie GX, Li WJ (2014) *Geothermomicrobium terrae* gen. nov.,  
350 sp. nov., a novel member of the family *Thermoactinomycetaceae*. Int J Syst Evol Microbiol 64:2998-3004  
351  
352  
353  
354



---

355 Figure Legend

356 Figure 1. Neighbor joining tree showing the positions of *Kroppenstedtia sanguinis* sp. nov. and *Kroppenstedtia pulmonis* sp.  
357 nov. within the family *Thermoactinomycetaceae*. The tree was constructed based on an analysis of ~1480 nt. Bootstrap values  
358 shown at the nodes are expressed as a percentage of 1000 replications (neighbor joining); only values >70 % are shown. ML,  
359 nodes common to the neighbor joining and maximum likelihood analyses; MP, nodes common to the neighbor joining and  
360 maximum parsimony analyses (Felsenstein 1989). The sequence of the type strain *Bacillus subtilis* was used as the outgroup.  
361



**Table 1** Strains used in this study

Strain	Date received or reference	Source	Geographic source	GenBank accession number of 16S rRNA gene
<i>Kroppenstedtia eburnea</i> (DSM 45196 <sup>T</sup> )	von Jan et al. (2011)	Plastic surface	Germany	FN665656
<i>K. guangzhouensis</i> (GD02 <sup>T</sup> )	Yang et al. (2013)	Soil	China	KC311557
<i>Melghirimyces algeriensis</i> (NariEX <sup>T</sup> )	Addou et al. (2012)	Soil from salt lake Lung, 78-year-old male	Algeria	HQ383683
W9323 <sup>T</sup>	2008	Blood, 59-year-old male	New York, USA	KC875234
X0209 <sup>T</sup>	1997	CSF, 16-year-old female	Gävle, Sweden	AJ251778
X0394	2010		Quebec, Canada	KC875233

<b>Table 2</b> Phenotypic, chemotaxonomic and genotypic characteristics that differentiate the study strains from other <i>Kroppenstedtia</i> species					
Characteristic	<i>K. pulmonis</i> W9323 <sup>T</sup>	<i>K. sanguinis</i> X0209 <sup>T</sup>	<i>K. sanguinis</i> X0394	<i>K. eburnea</i> DSM 45196 <sup>T</sup>	<i>K. guangzhouensis</i> GD02 <sup>T a</sup>
Endospore morphology	Elongated paddle shaped	Globose	Globose	Globose	Globose
Aerial hyphae	-	-	-	- (+ <sup>b</sup> )	NT (+ <sup>c</sup> )
Optimal growth on TSA	45 °C	45 °C	45 °C	45 °C	50 °C
sheep blood at 7 days					
Beta hemolysis on TSA	+	+	+	+	NT
sheep blood					
Utilization of:					
Cellobiose	-	+	-	+	-
Dulcitol	-	-	-	+	-
D-Fructose	-	-	-	+	-
D-Glucose	+	-	-	-	-
D-Mannitol	-	+	+	-	-
Mannose	-	+	-	-	NT
Salicin	-	+	-	+	NT
Sucrose	+	-	-	-	-
Whole cell-wall sugars	Ribose Glucose	Ribose Trace of galactose	Ribose Trace of galactose	Ribose	NT
Cellular fatty acids (> 5 %)	<i>iso</i> -C <sub>15</sub> (75.0 %) <i>anteiso</i> -C <sub>15</sub> (11.0 %) <i>iso</i> -C <sub>17</sub> (8.0 %) <i>iso</i> -C <sub>16</sub> (2.5 %)	<i>iso</i> -C <sub>15</sub> (74.0 %) <i>anteiso</i> -C <sub>15</sub> (12.0 %) <i>iso</i> -C <sub>17</sub> (7.0 %) <i>iso</i> -C <sub>16</sub> (1.4 %)	<i>iso</i> -C <sub>15</sub> (75.0 %) <i>anteiso</i> -C <sub>15</sub> (13.0 %) <i>iso</i> -C <sub>17</sub> (8.0 %) <i>iso</i> -C <sub>16</sub> (-)	<i>iso</i> -C <sub>15</sub> (73.0 %) <i>anteiso</i> -C <sub>15</sub> (13.0 %) <i>iso</i> -C <sub>17</sub> (-) <i>iso</i> -C <sub>16</sub> (4.5 %)	<i>iso</i> -C <sub>15</sub> (63.5 %) <i>anteiso</i> -C <sub>15</sub> (7.0 %) <i>iso</i> -C <sub>17</sub> (8 %) <i>iso</i> -C <sub>16</sub> (12.6 %)
Menaquinones	MK-7 (95.0 %) MK-8 (5.0 %)	MK-7 (97.0 %) MK-8 (3.0 %)	MK-7 (97.0 %) MK-8 (3.0 %)	MK-7 (97.0 %) MK-8 (3.0 %)	MK-7 (98.6 %) MK-8 (1.4 %)
Polar lipids	PG, DPG, PE, PME, 4 unknown PL, 1 unknown GL	PG, DPG, PE, PME, 5 unknown PL, 1 unknown GL	PG, DPG, PE, PME, 4 unknown PL, 1 unknown GL	PG, DPG, PE, PME, 2 unknown PL	PG, DPG, PE, PME, 2 unknown L
DNA G + C mol%	45.9	50.5	50.6	54.6	56.3

All strains tested were negative for utilization of acetamide, adonitol, L-arabinose, citrate, i-erythritol, D-galactose, glycerol, i-myoinositol, lactose, maltose, melibiose, raffinose, L-rhamnose, D-sorbitol, trehalose and D-xylose; production of arylsulfatase and urease, hydrolysis of adenine, aesculin by fluorescence, hypoxanthine, tyrosine, xanthine and reduction of nitrate; all strains tested were positive for browning of aesculin, growth in lysozyme and hydrolysis of casein with 0.5 % NaCl. No growth on casein. PG, phosphatidylglycerol; DPG, diphosphatidylglycerol; PE, phosphatidylethanolamine; PME, phosphatidylmethylethanolamine; PL, phospholipid, GL, glycolipid; L, lipid; -, negative; +, positive; NT, not tested; TSA, trypticase soy agar.

<sup>a</sup> All phenotypic data for *K. guangzhouensis* from Yang et al. (2013). All other data including cellular fatty acids, menaquinones, polar lipids and DNA G + C content were generated from present study.

<sup>b</sup> Aerial hyphae reported by von Jan et al. (2011).

<sup>c</sup> Aerial hyphae reported by Yang et al. (2013).

CCUG: CULTURE COLLECTION UNIVERSITY OF GOTHENBURG  
 Department of Clinical Bacteriology  
 Sahlgrenska University Hospital  
 Sahlgrenska Academy of the University of Gothenburg  
 Box 7193, SE-402 34 Göteborg, Sweden



CCUG  
 Culture Collection  
 University of Gothenburg

Accreditation Registration Number: 1240 (SWEDAC)

To Depositor:

Dr. Melissa Bell  
 US CDC  
 Mailstop D-11  
 1600 Clifton Rd.  
 Atlanta, GA 30329  
 USA

Tel: +1-404-639-1348

Fax: +1-404-639-3022

E-mail: [jqv7@cdc.gov](mailto:jqv7@cdc.gov)

Date: January 27, 2016

### CERTIFICATE CONFIRMING DEPOSIT AND AVAILABILITY OF A TYPE STRAIN

This Certificate confirms that the listed strain has been deposited with the CCUG and has passed the viability, purity and authentication controls. The strain is archived irrevocably, except in the case that it is found to be unsuitable.

In accordance with the International Code of Nomenclature (the Bacteriological Code), the strain has been assigned a collection accession number; this number may not be subsequently altered.

The strain will be made available to the international scientific community upon the date of valid publication of the species/sub-species name(s) in the International Journal of Systematic and Evolutionary Microbiology. The depositor may request an earlier release.

Upon valid publication, the information on the strain will be transferred to the CCUG open database, on internet: <http://www.ccug.se>

*Edward Moore*

Edward Moore, Curator – CCUG

CCUG 68107 T

*Kroppenstedtia pulmonis sp. nov.*

HIS: 20151109 <- Bell M, CDC, Atlanta GA, USA  
 OCC: W9323 = DSM 45752-T  
 AUTH: 16S rRNA gene sequence (Accession Number: KC875234)  
 RESTR: # Restricted distribution until validly published or 3 years

Dr. Edward Moore, PhD; Professor of Bacteriology

Tel: +46-31-342-4696; Fax: +46-31-82 96 17; E-mail: [curator@ccug.se](mailto:curator@ccug.se); URL: <http://www.ccug.se>

CCUG: CULTURE COLLECTION UNIVERSITY OF GOTHENBURG  
Department of Clinical Bacteriology  
Sahlgrenska University Hospital  
Sahlgrenska Academy of the University of Gothenburg  
Box 7193, SE-402 34 Göteborg, Sweden



CCUG  
Culture Collection  
University of Gothenburg

Accreditation Registration Number: 1240 (SWEDAC)

To Depositor:

Dr. Melissa Bell  
US CDC  
Mailstop D-11  
1600 Clifton Rd.  
Atlanta, GA 30329  
USA

Tel: +1-404-639-1348

Fax: +1-404-639-3022

E-mail: [jqv7@cdc.gov](mailto:jqv7@cdc.gov)

Date: January 27, 2016

### CERTIFICATE CONFIRMING DEPOSIT AND AVAILABILITY OF A TYPE STRAIN

This Certificate confirms that the listed strain has been deposited with the CCUG and has passed the viability, purity and authentication controls. The strain is archived irrevocably, except in the case that it is found to be unsuitable.

In accordance with the International Code of Nomenclature (the Bacteriological Code), the strain has been assigned a collection accession number; this number may not be subsequently altered.

The strain will be made available to the international scientific community upon the date of valid publication of the species/sub-species name(s) in the International Journal of Systematic and Evolutionary Microbiology. The depositor may request an earlier release.

Upon valid publication, the information on the strain will be transferred to the CCUG open database, on internet: <http://www.ccug.se>

*Edward Moore*

Edward Moore, Curator – CCUG

CCUG 38657 T

*Kroppenstedtia sanguinis sp. nov.*

HIS: 19971028 <- PHL, Gävle SE  
OCC: X0209 = DSM 45749-T  
AUTH: 16S rRNA gene sequence (Accession Number: AJ251778)  
RESTR: # Restricted distribution until validly published or 3 years

Dr. Edward Moore, PhD; Professor of Bacteriology

Tel: +46-31-342-4696; Fax: +46-31-82 96 17; E-mail: [curator@ccug.se](mailto:curator@ccug.se); URL: <http://www.ccug.se>

Leibniz-Institut  
DSMZ-Deutsche Sammlung von  
Mikroorganismen und Zellkulturen GmbH



**Confirmation of the availability of a strain for the purpose of valid publication of a new name according to the Bacteriological Code**

The following information is confidential and serves only to allow the International Journal of Systematic and Evolutionary Microbiology to confirm that a strain has been deposited and will be available from the DSMZ in accordance with the Rules of the Bacteriological Code (1990 revision) as revised by the ICSP at the plenary sessions in Sydney and Paris.

Strain *Kroppenstedtia* sp. X0209 has been deposited in the DSMZ under the number

**DSM 45749**

This strain is available in the publicly accessible section of the DSMZ and restrictions have not been placed on access to information concerning the presence of this strain in the DSMZ. It will be included in published and online catalogues after publication of this number by the authors.

This strain has been checked for viability in the DSMZ and is stored using one of the standard methods used in the DSMZ.

**!! The DSMZ is not responsible for differences between the properties of the strain deposited in the DSMZ and properties given in the literature/databases !!**

**It is the sole responsibility of the depositor to ensure that type strains deposited in the DSMZ conform to the requirements of the appropriate Rules governing prokaryotes nomenclature and the deposition of type strains (Rules 18a, 27, & 30 of the ICNB/ICNP, including changes made at plenary sessions of the JC/ICSP).**

Dr. E. Atasayar, Curator Actinomycetales

Geschäftsführer/  
Managing Director:  
Prof. Dr. Jörg Overmann  
Aufsichtsratsvorsitzender/Head of  
Supervisory Board: RD Dr. David Schnieders

Braunschweigische Landessparkasse  
(NORD/LB) Kto.-Nr./Account: 2 039 220  
BLZ/Bank Code: 250 500 00  
IBAN DE22 2505 0000 0002 0392 20  
SWIFT (BIC) NOLADE 2 H

Handelsregister/  
Commercial Register:  
Amtsgericht Braunschweig  
HRB 2570  
Steuer-Nr. 13/200/24030



Leibniz-Institut  
DSMZ-Deutsche Sammlung von  
Mikroorganismen und Zellkulturen GmbH



**Confirmation of the availability of a strain for the purpose of valid publication of a new name according to the Bacteriological Code**

The following information is confidential and serves only to allow the International Journal of Systematic and Evolutionary Microbiology to confirm that a strain has been deposited and will be available from the DSMZ in accordance with the Rules of the Bacteriological Code (1990 revision) as revised by the ICSP at the plenary sessions in Sydney and Paris.

Strain *Kroppenstedtia* sp. **W9323** has been deposited in the DSMZ under the number  
**DSM 45752**

This strain is available in the publicly accessible section of the DSMZ and restrictions have not been placed on access to information concerning the presence of this strain in the DSMZ. It will be included in published and online catalogues after publication of this number by the authors.

This strain has been checked for viability in the DSMZ and is stored using one of the standard methods used in the DSMZ.

**!! The DSMZ is not responsible for differences between the properties of the strain deposited in the DSMZ and properties given in the literature/databases !!**

**It is the sole responsibility of the depositor to ensure that type strains deposited in the DSMZ conform to the requirements of the appropriate Rules governing prokaryotes nomenclature and the deposition of type strains (Rules 18a, 27, & 30 of the ICNB/ICNP, including changes made at plenary sessions of the JC/ICSP).**



Dr. E. Atasayar, Curator Actinomycetales

Geschäftsführer/  
Managing Director:  
Prof. Dr. Jörg Overmann  
Aufsichtsratsvorsitzender/Head of  
Supervisory Board: RD Dr. David Schnieders

Braunschweigische Landessparkasse  
(NORD/LB) Kto.-Nr./Account: 2 039 220  
BLZ/Bank Code: 250 500 00  
IBAN DE22 2505 0000 0002 0392 20  
SWIFT (BIC) NOLADE 2 H

Handelsregister/  
Commercial Register:  
Amtsgericht Braunschweig  
HRB 2570  
Steuer-Nr. 13/200/24030

