

Age-related changes in hippocampal-neocortical connectivity during successful associative retrieval

Gaby Pfeifer*^{1,2}, Natasha Sigala^{1,3}

¹ Clinical Imaging Sciences Centre, Brighton and Sussex Medical School

² School of Psychology, University of Sussex

³ Sackler Centre for Consciousness Science, University of Sussex

*E-mail: g.pfeifer@bsms.ac.uk




Acknowledgements

This work was funded by the Brighton and Sussex Medical School, an MRC Ph.D. studentship, and the Brighton and Sussex Universities NHS Hospital Trust. Ethical approval was obtained by the RGEC of the Brighton and Sussex Medical School.

Background

- **Successful associative retrieval** critically depends on the hippocampus¹.
- **Hippocampal activity in older adults** varies across studies, either showing *hyper*²- or *hypo*-activation³ during memory tasks, despite comparable task performance with young adults.
- **Functional connectivity measures** of the hippocampus with other cortical regions can provide a window into the underlying neuronal compensation mechanisms in the aging brain.
- **Compensatory mechanisms** in older adults are often reflected by increased frontal coupling⁴. However, as task demand increases, older adults reach a resource ceiling⁵, showing reduced flexibility in functional coupling to meet demands.



Predictions

- **Successful associative retrieval was expected to lead to:**
 - 1) increased hippocampal coupling with frontal regions in older *vs.* young adults, even when memory load was low.
 - 2) undifferentiated coupling in older adults from the low to the high memory load condition due to deficient resource allocation.

Methods


Participants

- 19 young adults (21-32 years; $M = 24.06$)
- 19 older adults (59-81 years; $M = 66.21$)

Tasks:

1. Self-paced learning (pre-scanning session)


- Participants learned 8 fractal pair-associates to 87% correct performance criterion on a computer outside the scanner. Memory load was manipulated by visual similarity.



2. Memory retrieval (scanning session)

- Memory was tested in the scanner using a delayed pair-associative retrieval task.


Delayed pair-associative retrieval task



fMRI protocol & Analyses

Event-related design, including 32 similar and 32 dissimilar pair trials

Trial structure:



Data acquisition parameters:

Data were collected on a 1.5 T Siemens Magnetom Avanto MRI Scanner with a 32-channel head coil. 644 volumes were obtained using a T2* weighted EPI sequence; TR=2.62s, TE=42ms, flip angle=90 degrees


Functional connectivity

- Region of interest (ROI): an anatomical seed region of the entire L hippocampus was created.
- Temporal pre-processing: spurious effects of white matter, cerebrospinal fluid, head motion and task conditions were removed; a band-pass filter was set at $0.008\text{Hz} < f < 0.09\text{Hz}$.
- Seed-to-voxel analysis: Using the CONN fMRI connectivity toolbox in SPM8 (v13o, 2011, <http://www.nitrc.org/projects/conn>), the mean BOLD time-series of the ROI was regressed with all other voxels in the brain.

Univariate analyses

- Pre-processing: All functional images were slice-time and motion corrected, unwarped, coregistered to participants' individual structural volume, and spatially normalised.
- Hits & Correct rejections of similar & dissimilar pairs were analysed using the GLM. Events were convolved with a canonical HDR function, time-locked to cue onset with a 1s duration (modelling cue-period).
- Contrast images were computed for the *Similar* and *Dissimilar* conditions. Results were subjected to a group analysis (2-level random effects analyses). A conjunction analysis was performed with trials averaged across similarity. This yielded activity in the left (L) hippocampus.


Behavioural Results



Results summary


- We found a significant interaction between young and older adults and the number of successfully retrieved trials:
- Young and older adults showed a comparable number of retrieved similar pair-associates (low memory load) but older adults retrieved significantly fewer trials of the dissimilar pair-associates than young adults.

fMRI Results



Results summary

- **In the low memory load condition**, young adults show hippocampal coupling with the basal ganglia & parahipp. gyrus, while older adults recruit a frontal-parietal network.
- **In the high memory load condition**, young adults show hippocampal coupling with medial occ.-temp. regions & the orbitofrontal cortex, while older adults continue to recruit a frontal-parietal network.



Conclusions

- Our results demonstrate that in healthy aging, associative retrieval involves hippocampal coupling with frontal-parietal brain regions, even when memory load is low. The memory network in older adults remains largely invariant to changing memory load.
- By contrast, younger adults show more automatic retrieval mechanisms during low memory load, supported by subcortical brain structures. They flexibly switch to visual and orbitofrontal brain regions with increasing memory load to aid associative retrieval.
- Our results support the account of a deficient resource allocation in aging.

References

- ¹ Squire, L.R., Zola-Morgan, M., 1991. The Cognitive Neuroscience of Human Memory Since HM. *Annu Rev Neurosci* 14, 299-324.
- ² Yassa, M.A., Lacy, J.W., Stark, S.M., Albert, M.S., Gallagher, M., Stark, C.E.L., 2011. Pattern Separation Deficits Associated With Increased Hippocampal CA3 and Dentate Gyrus Activity in Nondemented Older Adults. *Hippocampus* 21, 968-979.
- ³ Cabeza, R., Daselaar, S.M., Dolcos, F., Prince, S.E., Budde, M., Nyberg, L., 2004. Task-independent and task-specific age effects on brain activity during working memory, visual attention and episodic retrieval. *Cerebral Cortex* 14, 364-375.
- ⁴ Grady, C.L., Protzner, A.B., Kovacevic, N., Strother, S.C., Afshin-Pour, B., Wojtowicz, M., Anderson, J.A.E., Churchill, N., McIntosh, A.R., 2010. A Multivariate Analysis of Age-Related Differences in Default Mode and Task-Positive Networks across Multiple Cognitive Domains. *Cerebral Cortex* 20, 1432-1447.
- ⁵ Geerlings, L., Sallasi, E., Renken, R.J., Maurits, N.M., Lorist, M.M., 2013. Flexible Connectivity in the Aging Brain Revealed by Task Modulations. *Hum Brain Mapp*.